Biomarkers of Sepsis: Time for a Reappraisal

Charalampos Pierrakos

Brugmann University Hospital

Dimitrios Velissaris

University of Patras

Max Bisdorff

Université libre de Bruxelles

John C Marshall

St Michael's Hospital

Jean-Louis Vincent (✉ jlvincent@intensive.org)

Erasmus University Hospital <https://orcid.org/0000-0001-6011-6951>

Research

Keywords: procalcitonin, C-reactive protein, diagnosis, prognosis, infection, validation

Posted Date: March 4th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-15792/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Critical Care on June 5th, 2020. See the published version at <https://doi.org/10.1186/s13054-020-02993-5>.

Abstract

Introduction: Sepsis biomarkers can have important diagnostic, therapeutic, and prognostic functions. In a previous review, we identified 3370 references reporting on 178 different biomarkers related to sepsis. In the present review, we evaluate the progress in the research of sepsis biomarkers.

Methods: Using the same methodology as in our previous review, we searched the PubMed database from 2009 until September 2019 using the terms “Biomarker” AND “Sepsis”. There were no restrictions by age or language and all studies, clinical and experimental, were included.

Results: We retrieved a total of 5367 new references since our previous review. We identified 258 biomarkers, 80 of which were new compared to our previous list. The majority of biomarkers have been evaluated in fewer than 5 studies, with 81 (31%) being assessed in just a single study. Apart from studies of C-reactive protein (CRP) or procalcitonin (PCT), only 26 biomarkers have been assessed in clinical studies with more than 300 participants. Forty biomarkers have been compared to PCT and/or C-reactive protein CRP for their diagnostic value; 9 were shown to have a better diagnostic value for sepsis than either or both of these biomarkers. Forty-four biomarkers have been evaluated for a role in answering a specific clinical question rather than for their general diagnostic or prognostic properties in sepsis.

Conclusions: The number of biomarkers being identified is still increasing although at a slower rate than in the past. Most of the biomarkers have not been well-studied; in particular the clinical role of these biomarkers needs to be better evaluated.

Introduction

Biomarkers have been evaluated for several applications in patients with sepsis including diagnosis of infection, prognostication and therapeutic guidance. Sepsis is a common and severe condition [1, 2], responsible for high mortality and morbidity rates and also for reduced quality of life [1-4]. Sepsis biomarkers may provide information beyond what is available using other metrics, and could therefore help inform clinical decision making and potentially improve patient management. For example, more timely and appropriate antibiotic therapy could be administered and unnecessary antibiotics avoided if biomarkers are available that can accurately diagnose sepsis early. Similarly, biomarkers could help physicians monitor the effectiveness of therapeutic decisions and adjust treatment if necessary [5]. Many potential sepsis biomarkers have been proposed, procalcitonin (PCT) and C-reactive protein (CRP) being the most frequently studied. The Surviving Sepsis Campaign guidelines for the management of sepsis mention that sepsis biomarkers can complement clinical evaluation [6] but in the Sepsis-3 definition consensus, the role of biomarkers in sepsis diagnosis remains undefined [7].

In 2010, we published a literature review of biomarkers that had been studied for their potential diagnostic or prognostic role in sepsis [8]. We concluded that none of the 178 biomarkers identified had “sufficient specificity or sensitivity to be routinely employed in clinical practice” [8]. In this narrative review, we evaluate the progress that has been made in identifying new sepsis biomarkers since that report and reappraise the utility of such research in the management of patients with sepsis.

Methods

We searched the Medline database via the PubMed portal between February 2009 and September 2019 using (“biomarker” AND “sepsis”) as keywords to identify all studies that evaluated a biomarker in sepsis. There were no restrictions by age or language and all studies, clinical and experimental, were included. The reference lists from all relevant retrieved manuscripts were further reviewed in order to identify additional studies. For each identified biomarker, the PubMed database was searched again using the biomarker name and the key word 'biomarker'.

Newly found biomarkers were added to our previous database. Details related to the methodology used in each study were collected, namely: (1) type of study (mono- vs. multicenter, prospective vs. retrospective, experimental vs. clinical); (2) study population (intensive care unit [ICU], emergency room, other population); (3) number of studied subjects; (4) reference non-sepsis population; and (5) purpose of study or use of biomarker being tested (diagnostic, prognostic, other clinical role). Results of receiver operating characteristic (ROC) curve analysis were noted where this technique was used to assess biomarker specificity and sensitivity. For

each biomarker, the main pathophysiological role was recorded. We also reported separately biomarkers that had been compared with PCT and/or CRP.

Results

A total of 5367 studies met our search criteria for the period 2009 to 2019 compared with the 3370 studies retrieved in our previous study [3]. A total of 80 new biomarkers (54 assessed in clinical studies, 23 in clinical and experimental studies, and 3 in only experimental studies) were added to the list of 178 biomarkers that had previously been identified. Despite a steady increase in the number of published studies related to sepsis biomarkers over time, the number of publications reporting *new* biomarkers has decreased since our prior review (Figure 1).

The full list of biomarkers with selected references and major findings are shown in Additional File 1, Tables S1–9. Of the 258 biomarkers, 69 (27%) were assessed primarily for their diagnostic value, 100 (39%) for their prognostic value and 89 (34%) for both diagnostic and prognostic purposes. A validation population was used in just 12 studies. Most of the biomarkers ($n=216$ [84%]) have been assessed in fewer than five studies, and 81 (31%) have been studied only once. CRP and PCT are the biomarkers that have been studied most frequently, followed by interleukin (IL)-6, presepsin and CD64 in 31, 25 and 21 studies, respectively.

Apart from CRP and PCT, only 26 biomarkers have been evaluated in studies that enrolled more than 300 patients (Tables 1 and 2). In 15 of these 24 studies (63%), sepsis was defined using either the 1992 ACCP/SCCM [9] or the 2001 International Sepsis Definitions Conference [10] definitions. In one study, the Sepsis-3 definition [7] was used. Other studies used definitions based on clinical signs compatible with sepsis or positive blood cultures. Of the 10 biomarkers evaluated for their diagnostic value in more than 300 patients, 6 (60%) were evaluated using receiver operating characteristic (ROC) curve analysis; the area under the curve (AUC) was >0.8 for just three of the biomarkers (for inter-alpha inhibitor proteins [11], CD64 [12] and IL-6 [13]). Of the 18 biomarkers evaluated for their prognostic value in more than 300 patients, mortality was the primary study endpoint for 14 (78%); prediction of circulatory failure or organ dysfunction and failure of antibiotic therapy were the primary endpoints in the other studies. ROC curve analysis was used in the analysis of 9 of the 18 biomarkers (50%): the AUC for predicting mortality was >0.8 only for pro-adrenomedullin, with a high specificity (specificity: 92%, sensitivity: 75%). In two studies, combining a sepsis biomarker with a severity score improved the predictive value (urokinase plasminogen activator receptor [uPAR] + APACHE II AUC: 0.83 [14], adrenomedullin + Mortality in Emergency Department Sepsis (MEDS) score AUC: 0.81 [15]).

Forty biomarkers have been compared with CRP and/or PCT for their diagnostic value (Table 3); 9 were shown to have better diagnostic value and 11 improved the diagnostic value of CRP and/or PCT when used in combination with one of these two biomarkers. In 10 of the 23 studies in which these results were reported (43%), patients with SIRS without infection were selected as the reference group; two studies used patients after major surgery as the reference group. A validation group of healthy volunteers was used in 5 studies (22%).

Forty-four biomarkers were tested in 55 clinical studies for their use in answering specific, clinically relevant questions rather than simply for diagnosis and/or prognosis of sepsis in general (Table 4): 20 were assessed for use to diagnose infection in specific groups of critically ill patients where diagnosis may be difficult based on clinical evaluation and laboratory values; 8 were assessed for diagnosis of ARDS or associated endothelial damage in patients with sepsis; 6 were tested for their ability to identify specific infections or type of microorganism; 6 were studied for use in the diagnosis of disseminated intravascular coagulation; 4 were assessed for use in deciding which patients with hematological malignancy or neutropenia had a low risk of infection; 3 were assessed for their ability to diagnose infection before any clinical symptoms; 2 were evaluated for use in assessing the risk of delirium or encephalopathy in patients with sepsis; and 1 to differentiate between sepsis and graft rejection.

Discussion

Our literature search illustrates that, although new biomarkers have been proposed, little real progress has been made in identifying biomarkers with clinical significance. Using a similar method of searching for sepsis biomarkers to that of our previous study, we noted that the number of publications related to sepsis biomarkers has increased considerably over the years. The proportion of new biomarkers being identified has decreased, but this may reflect publication bias with journals becoming more selective in deciding what merits publication as the volume of these studies increases. Because of the complexity of the sepsis response with multiple mediators, and the improved sensitivity of many tests enabling identification of smaller concentrations of substances than in the

past, it is likely that our list of biomarkers will expand further in the future. However, the potential utility of creating an ever expanding list of potential biomarkers without a more rigorous framework to evaluate them is questionable. An improved methodological approach is needed in order to assess the utility of sepsis biomarkers in daily clinical practice.

Accurate evaluation of the possible clinical utility of a biomarker requires assessment in a large number of patients [5], but we identified only a few biomarkers that have been assessed in studies of more than 300 patients. Moreover, many of the biomarkers have been assessed in only a limited number of clinical studies and one third in just a single study. Patients with sepsis represent a very heterogeneous population and potential biomarkers need to be assessed in studies with a significant number of patients to ensure random distribution of risk factors that may affect the results of the study (e.g., age, organ dysfunction, type of infection, comorbidities). However, the number of patients enrolled in a study is not the only factor to consider when evaluating the potential role of a sepsis biomarker and, of note, none of the large multicenter studies were able to draw conclusions about the biomarker under study that could change clinical practice.

There was considerable diversity in the methods used to assess sepsis-related biomarkers. Most biomarkers were proposed as being useful for diagnosing sepsis simply because they were increased or decreased to a larger extent in septic than in non-septic patients or healthy individuals. Many studies have assessed the sensitivity and specificity of the biomarker for sepsis diagnosis, but identification of sepsis was often based on the commonly used constellation of non-specific clinical and laboratory findings; in the absence of a “gold standard” diagnostic tool, this method cannot therefore conclusively demonstrate the value of the biomarker with respect to diagnosing sepsis. Other parameters, including positive and negative predictive value or likelihood ratios, can provide greater insight into how well a biomarker performs but these were rarely provided [16]. Similarly, many biomarkers have been used to evaluate sepsis severity using all-cause mortality as the primary end-point. Importantly, the majority of the studies that evaluated sepsis biomarkers using this method showed only a limited value; it seems highly unlikely that mortality in septic patients is related to only one pathophysiologic process that could be reflected by abnormal levels of a biomarker. Furthermore, the need for another prognostic test can be questioned because clinical data and other laboratory test results, including blood lactate levels, can already reflect severity and the risk of death in septic patients [17]. Prognostic biomarkers may be useful to triage patients in special environments, such as in the emergency room, when the information provided can help clinicians to decide whether hospitalization is necessary and, if so, on the ICU or on the regular floor. However, in a multicenter trial (TRIAGE III) in which emergency room physicians were asked to incorporate the prognostic information portrayed by abnormal uPAR levels into their triage decisions, there was no effect on mortality rates compared to standard practice without uPAR levels [18].

To be of value in clinical practice, a biomarker must be shown to provide an answer to a specific, clinically relevant question, rather than just having diagnostic or prognostic value in general. We identified just 55 studies in which a sepsis biomarker was shown to have a potentially useful role by answering a specific clinical question. For example, biomarkers that could identify specific types of infection may help in guiding a more targeted antibiotic therapy, and a biomarker able to identify septic patients at risk of ARDS may influence fluid management in such patients, reducing risks of fluid overload. Further study needs to better evaluate the potentially utility and beneficial effects on outcomes of using biomarkers to answer specific clinical questions.

We attempted to categorize the various biomarkers according to their pathophysiological role, although for many it was not possible to identify a clear role, and some have multiple roles. Only a few biomarkers were found to have a role specifically related to sepsis pathophysiology rather than to a more general inflammatory reaction, including presepsin (the N-terminal fragment of the macrophage lipopolysaccharide [LPS] receptor), LPS-binding protein (LBP), bactericidal/permeability increasing protein, peptidoglycan, thrombomodulin and anti-endotoxin core antibodies. Such biomarkers may help transform our understanding of sepsis from a “physiological syndrome to a group of distinct biochemical disorders” [19] and advance our search for adjunctive sepsis therapies.

CRP and PCT are by far the most widely used and studied biomarkers. Both increase transiently during sepsis, but long enough to allow for their detection, reflecting a real-time response. Although PCT is considered superior to CRP in many studies [20, 21], it is not a definitive test for diagnosing sepsis because PCT levels can also be increased in other conditions [22]. PCT, similar to CRP, may be more useful to rule out sepsis than to diagnose it [23-25] and the combination of these two biomarkers may improve their ability to exclude sepsis [26]. Studying the time course of these biomarkers may also be helpful to evaluate an individual patient’s response to therapy. Changes in serum CRP levels during the first 48 h after antibiotic initiation can help evaluate the response to initial antimicrobial therapy [27]. Likewise, a PCT-based algorithm may help reduce antibiotic exposure in septic patients without

compromising clinical outcomes [28, 29]. However, not all studies have shown the same positive effect [30], suggesting that the effectiveness of PCT-based algorithms may depend on the physician's experience and the clinical setting. Some biomarkers have been compared to PCT and CRP, most for their diagnostic value. A few were shown to be superior to PCT and/or CRP for this purpose, for example, presepsin and CD64 [31-33].

Measuring several biomarkers concurrently may be useful to overcome the limitations of any single biomarker. Combining biomarkers that are involved in different sepsis-related pathways may be particularly attractive. A seven-biomarker panel including cellular markers and interleukins correctly identified 89% of patients with ventilator-associated pneumonia (VAP) and 100% of patients without VAP [34]. Similarly, a combination of several sepsis-related biomarkers (PCT, presepsin, galectin-3, and soluble suppression of tumorigenicity 2) was found to have better prognostic value than PCT alone [35]. However, it is not clear from the existing literature whether the biomarkers included in such panels should be selected based on pathophysiological or other criteria. The combination of a biomarker panel with clinical information may be particularly useful in the diagnosis of sepsis or in the risk stratification of patients with sepsis [36].

The study has some limitations that should be acknowledged. First, although we performed an extensive search, we cannot be sure that some studies were not missed. Nevertheless, the large number of sepsis biomarkers that we retrieved suggests that we managed to identify the majority of the biomarkers that have been studied. Second, we included studies over a long period of time, during which the definition of sepsis has changed so that it is difficult to make comparisons. Third, it is difficult to compare different biomarkers because the methods used to evaluate the biomarkers and to define sepsis and the populations studied varied across the studies.

Conclusions

Since our original search, many additional sepsis-related biomarkers have been identified. However, the precise roles of most biomarkers in the management of septic patients have not been well defined and, of the many biomarkers that have been studied, only a few have been evaluated in large or repeated studies. To help identify biomarkers that can provide valuable, clinically relevant information, a more rigorous, standardized methodology to assess sepsis biomarkers seems necessary. Such an approach could include the following factors:

1. What is the question being asked?
 - i. Greater likelihood of infection leading to administration of empiric antibiotics or performance of a diagnostic test (e.g., carcinoembryonic antigen [CEA] levels are used to detect early recurrence in patients with colon cancer, and so guide further investigations)
 - ii. Resolution of infection and therefore safety in stopping antibiotics
 - iii. Increased likelihood of benefiting from specific interventions, such as steroids or a biologic agent
 - iv. Increased risk of adverse outcome not apparent by other evidence.
 - v. Ensuring random distribution of risk factors in a randomized controlled trial
2. How is the study designed?
 - i. What is the control group
 - ii. Which patients are being studied
 - iii. How are outcomes adjudicated
 - iv. Is there a validation cohort
 - v. Uniform techniques to evaluate results – sensitivity, specificity, positive and negative predictive values, likelihood ratios, ROC analysis
3. Is the marker biologically plausible, and what do alterations tell us about the pathobiology of disease in this patient?

Consideration of these factors and their application to sepsis biomarker research may help identify new biomarkers with real clinical utility. Continuing to produce reports of novel biomarkers without developing a more rigorous framework to evaluate them and establishing a recognized purpose is futile: it is time for a re-appraisal of the possible roles of biomarkers in sepsis.

Abbreviations

AUC area under the curve

CRP C-reactive protein

ICU intensive care unit

IL interleukin

LBP lipopolysaccharide binding protein

LPS lipopolysaccharide

MEDS Mortality in Emergency Department Sepsis score

PCT procalcitonin

ROC receiver operating characteristic

uPAR urokinase plasminogen activator receptor

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

JLV is Editor-in-Chief of *Critical Care* and has no other conflicts of interest.

The other authors declare that they have no relevant financial interests.

Funding

None

Authors' contributions

CP conceived the study, performed the literature search, drafted the manuscript and approved the submitted version of the manuscript.

DV helped perform the literature search, revised the manuscript for critical content, and approved the submitted version of the manuscript.

MB helped perform the literature search, revised the manuscript for critical content, and approved the submitted version of the manuscript.

JCM revised the manuscript for critical content and approved the submitted version of the manuscript.

JLV conceived the study, revised the manuscript for critical content, and approved the submitted version of the manuscript.

References

1. Vincent JL, Marshall JC, Namendys-Silva SA, Francois B, Martin-Loeches I, Lipman J et al. Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. *Lancet Respir Med*. 2014;2:380-6.
2. Machado FR, Cavalcanti AB, Bozza FA, Ferreira EM, Angotti Carrara FS, Sousa JL et al. The epidemiology of sepsis in Brazilian intensive care units (the Sepsis PREvalence Assessment Database, SPREAD): an observational study. *Lancet Infect Dis*. 2017;17:1180-9.
3. Widmann CN, Heneka MT. Long-term cerebral consequences of sepsis. *Lancet Neurol*. 2014;13:630-6.
4. Azoulay E, Vincent JL, Angus DC, Arabi YM, Brochard L, Brett SJ et al. Recovery after critical illness: putting the puzzle together-a consensus of 29. *Crit Care*. 2017;21:296.
5. Biomarkers Definitions Working Group. Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clin Pharmacol Ther*. 2001;69:89-95.
6. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Crit Care Med*. 2017;45:486-552.
7. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315:801-10.
8. Pierrakos C, Vincent JL. Sepsis biomarkers: a review. *Crit Care*. 2010;14:R15.
9. Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest*. 1992;101:1644-55.
10. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med*. 2003;31:1250-6.
11. Chaaban H, Singh K, Huang J, Siryaporn E, Lim YP, Padbury JF. The role of inter-alpha inhibitor proteins in the diagnosis of neonatal sepsis. *J Pediatr*. 2009;154:620-2.
12. Dimoula A, Pradier O, Kassenger Z, Dalcomune D, Turkan H, Vincent JL. Serial determinations of neutrophil CD64 expression for the diagnosis and monitoring of sepsis in critically ill patients. *Clin Infect Dis*. 2014;58:820-9.
13. Henning DJ, Hall MK, Watsjold BK, Bhatraju PK, Kosamo S, Shapiro NI et al. Interleukin-6 improves infection identification when added to physician judgment during evaluation of potentially septic patients. *Am J Emerg Med*. 2019;158361.
14. Giamarellos-Bourboulis EJ, Norrby-Teglund A, Mylona V, Savva A, Tsangaris I, Dimopoulou I et al. Risk assessment in sepsis: a new prognostication rule by APACHE II score and serum soluble urokinase plasminogen activator receptor. *Crit Care*. 2012;16:R149.
15. Chen YX, Li CS. Prognostic value of adrenomedullin in septic patients in the ED. *Am J Emerg Med*. 2013;31:1017-21.
16. Vetter TR, Schober P, Mascha EJ. Diagnostic testing and decision-making: Beauty is not just in the eye of the beholder. *Anesth Analg*. 2018;127:1085-91.
17. Pletcher MJ, Pignone M. Evaluating the clinical utility of a biomarker: a review of methods for estimating health impact. *Circulation*. 2011;123:1116-24.
18. Schultz M, Rasmussen LJH, Andersen MH, Stefansson JS, Falkentoft AC, Alstrup M et al. Use of the prognostic biomarker suPAR in the emergency department improves risk stratification but has no effect on mortality: a cluster-randomized clinical trial (TRIAGE III). *Scand J Trauma Resusc Emerg Med*. 2018;26:69.
19. Marshall JC, Reinhart K. Biomarkers of sepsis. *Crit Care Med*. 2009;37:2290-8.
20. Simon L, Saint-Louis P, Amre DK, Lacroix J, Gauvin F. Procalcitonin and C-reactive protein as markers of bacterial infection in critically ill children at onset of systemic inflammatory response syndrome. *Pediatr Crit Care Med*. 2008;9:407-13.
21. Luzzani A, Polati E, Dorizzi R, Rungatscher A, Pavan R, Merlini A. Comparison of procalcitonin and C-reactive protein as markers of sepsis. *Crit Care Med*. 2003;31:1737-41.
22. Park JH, Kim DH, Jang HR, Kim MJ, Jung SH, Lee JE et al. Clinical relevance of procalcitonin and C-reactive protein as infection markers in renal impairment: a cross-sectional study. *Crit Care*. 2014;18:640.

23. Hattori T, Nishiyama H, Kato H, Ikegami S, Nagayama M, Asami S et al. Clinical value of procalcitonin for patients with suspected bloodstream infection. *Am J Clin Pathol.* 2014;141:43-51.
24. Kono T, Otsuka M, Ito M, Misawa M, Hoshioka A, Suzuki M et al. Negative C-reactive protein in children with bacterial infection. *Pediatr Int.* 1999;41:496-9.
25. Chan YL, Liao HC, Tsay PK, Chang SS, Chen JC, Liaw SJ. C-reactive protein as an indicator of bacterial infection of adult patients in the emergency department. *Chang Gung Med J.* 2002;25:437-45.
26. Han JH, Nachamkin I, Coffin SE, Gerber JS, Fuchs B, Garrigan C et al. Use of a combination biomarker algorithm to identify medical intensive care unit patients with suspected sepsis at very low likelihood of bacterial infection. *Antimicrob Agents Chemother.* 2015;59:6494-500.
27. Pova P, Teixeira-Pinto AM, Carneiro AH. C-reactive protein, an early marker of community-acquired sepsis resolution: a multi-center prospective observational study. *Crit Care.* 2011;15:R169.
28. van der Does Y, Limper M, Jie KE, Schuit SCE, Jansen H, Pernot N et al. Procalcitonin-guided antibiotic therapy in patients with fever in a general emergency department population: a multicentre non-inferiority randomized clinical trial (HiTEMP study). *Clin Microbiol Infect.* 2018;24:1282-9.
29. Wirz Y, Meier MA, Bouadma L, Luyt CE, Wolff M, Chastre J et al. Effect of procalcitonin-guided antibiotic treatment on clinical outcomes in intensive care unit patients with infection and sepsis patients: a patient-level meta-analysis of randomized trials. *Crit Care.* 2018;22:191.
30. Huang DT, Yealy DM, Filbin MR, Brown AM, Chang CH, Doi Y et al. Procalcitonin-guided use of antibiotics for lower respiratory tract infection. *N Engl J Med.* 2018;379:236-49.
31. Liu D, Su L, Han G, Yan P, Xie L. Prognostic value of procalcitonin in adult patients with sepsis: A systematic review and meta-analysis. *PLoS One.* 2015;10:e0129450.
32. Jamsa J, Ala-Kokko T, Huotari V, Ohtonen P, Savolainen ER, Syrjala H. Neutrophil CD64, C-reactive protein, and procalcitonin in the identification of sepsis in the. *J Crit Care.* 2018;43:139-42.
33. Muzlovic I, Ihan A, Stubljär D. CD64 index on neutrophils can diagnose sepsis and predict 30-day survival in subjects after ventilator-associated pneumonia. *J Infect Dev Ctries.* 2016;10:260-8.
34. Grover V, Pantelidis P, Soni N, Takata M, Shah PL, Wells AU et al. A biomarker panel (Bioscore) incorporating monocytic surface and soluble TREM-1 has high discriminative value for ventilator-associated pneumonia: a prospective observational study. *PLoS One.* 2014;9:e109686.
35. Kim H, Hur M, Moon HW, Yun YM, Di Somma S. Multi-marker approach using procalcitonin, presepsin, galectin-3, and soluble suppression of tumorigenicity 2 for the prediction of mortality in sepsis. *Ann Intensive Care.* 2017;7:27.
36. Mearelli F, Fiotti N, Giansante C, Casarsa C, Orso D, De Helsen M et al. Derivation and validation of a biomarker-based clinical algorithm to rule out sepsis from noninfectious systemic inflammatory response syndrome at emergency department admission: A multicenter prospective study. *Crit Care Med.* 2018;46:1421-9.
37. Hanna WJ, Berrens Z, Langner T, Lahni P, Wong HR. Interleukin-27: a novel biomarker in predicting bacterial infection among the critically ill. *Crit Care.* 2015;19:378.
38. Uusitalo-Seppälä R, Peuravuori H, Koskinen P, Vahlberg T, Rintala EM. Role of plasma bactericidal/permeability-increasing protein, group IIA phospholipase A(2), C-reactive protein, and white blood cell count in the early detection of severe sepsis in the emergency department. *Scand J Infect Dis.* 2012;44:697-704.
39. Hollenbach B, Morgenthaler NG, Struck J, Alonso C, Bergmann A, Kohrle J et al. New assay for the measurement of selenoprotein P as a sepsis biomarker from serum. *J Trace Elem Med Biol.* 2008;22:24-32.
40. Sakr Y, Burgett U, Nacul FE, Reinhart K, Brunkhorst F. Lipopolysaccharide binding protein in a surgical intensive care unit: a marker of sepsis? *Crit Care Med.* 2008;36:2014-22.
41. Wei S, Gonzalez Rodriguez E, Chang R, Holcomb JB, Kao LS, Wade CE. Elevated syndecan-1 after trauma and risk of sepsis: A secondary analysis of patients from the Pragmatic, Randomized Optimal Platelet and Plasma Ratios (PROPPR) trial. *J Am Coll Surg.* 2018;227:587-95.
42. Nakamura Y, Hoshino K, Kiyomi F, Kawano Y, Mizunuma M, Tanaka J et al. Comparison of accuracy of presepsin and procalcitonin concentrations in diagnosing sepsis in patients with and without acute kidney injury. *Clin Chim Acta.* 2019;490:200-6.

43. Madoiwa S, Nunomiya S, Ono T, Shintani Y, Ohmori T, Mimuro J et al. Plasminogen activator inhibitor 1 promotes a poor prognosis in sepsis-induced disseminated intravascular coagulation. *Int J Hematol.* 2006;84:398-405.
44. Novotny AR, Emmanuel K, Ulm K, Bartels H, Siewert JR, Weighardt H et al. Blood interleukin 12 as preoperative predictor of fatal postoperative sepsis after neoadjuvant radiochemotherapy. *Br J Surg.* 2006;93:1283-9.
45. Johansen ME, Johansson PI, Ostrowski SR, Bestle MH, Hein L, Jensen AL et al. Profound endothelial damage predicts impending organ failure and death in sepsis. *Semin Thromb Hemost.* 2015;41:16-25.
46. Matsubara T, Yamakawa K, Umemura Y, Gando S, Ogura H, Shiraishi A et al. Significance of plasma fibrinogen level and antithrombin activity in sepsis: A multicenter cohort study using a cubic spline model. *Thromb Res.* 2019;181:17-23.
47. Chen Y, Li C. Prognostic significance of brain natriuretic peptide obtained in the ED in patients with SIRS or sepsis. *Am J Emerg Med.* 2009;27:701-6.
48. Calfee CS, Gallagher D, Abbott J, Thompson BT, Matthay MA. Plasma angiopoietin-2 in clinical acute lung injury: prognostic and pathogenetic significance. *Crit Care Med.* 2012;40:1731-7.
49. Kinasewitz GT, Yan SB, Basson B, Comp P, Russell JA, Cariou A et al. Universal changes in biomarkers of coagulation and inflammation occur in patients with severe sepsis, regardless of causative micro-organism [ISRCTN74215569]. *Crit Care.* 2004;8:R82-R90.
50. Saeed K, Wilson DC, Bloos F, Schuetz P, van der Does Y, Melander O et al. The early identification of disease progression in patients with suspected infection presenting to the emergency department: a multi-centre derivation and validation study. *Crit Care.* 2019;23:40.
51. Linder A, Christensson B, Herwald H, Bjorck L, Akesson P. Heparin-binding protein: an early marker of circulatory failure in sepsis. *Clin Infect Dis.* 2009;49:1044-50.
52. Rodelo JR, De la Rosa G, Valencia ML, Ospina S, Arango CM, Gomez CI et al. D-dimer is a significant prognostic factor in patients with suspected infection and sepsis. *Am J Emerg Med.* 2012;30:1991-9.
53. John J, Woodward DB, Wang Y, Yan SB, Fisher D, Kinasewitz GT et al. Troponin-I as a prognosticator of mortality in severe sepsis patients. *J Crit Care.* 2010;25:270-5.
54. Kornblit B, Hellemann D, Munthe-Fog L, Bonde J, Strom JJ, Madsen HO et al. Plasma YKL-40 and CHI3L1 in systemic inflammation and sepsis-experience from two prospective cohorts. *Immunobiology.* 2013;218:1227-34.
55. Rannikko J, Seiskari T, Huttunen R, Tarkiainen I, Jylhava J, Hurme M et al. Plasma cell-free DNA and qSOFA score predict 7-day mortality in 481 emergency department bacteraemia patients. *J Intern Med.* 2018;284:418-26.
56. Garnacho-Montero J, Huici-Moreno MJ, Gutierrez-Pizarra A, Lopez I, Marquez-Vacaro JA, Macher H et al. Prognostic and diagnostic value of eosinopenia, C-reactive protein, procalcitonin, and circulating cell-free DNA in critically ill patients admitted with suspicion of sepsis. *Crit Care.* 2014;18:R116.
57. Moreira VG, Prieto B, Rodriguez JS, Alvarez FV. Usefulness of cell-free plasma DNA, procalcitonin and C-reactive protein as markers of infection in febrile patients. *Ann Clin Biochem.* 2010;47:253-8.
58. Battista S, Audisio U, Galluzzo C, Maggiorotto M, Masoero M, Forno D et al. Assessment of diagnostic and prognostic role of copeptin in the clinical setting of sepsis. *Biomed Res Int.* 2016;2016:3624730.
59. Mandi Y, Farkas G, Takacs T, Boda K, Lonovics J. Diagnostic relevance of procalcitonin, IL-6, and sICAM-1 in the prediction of infected necrosis in acute pancreatitis. *Int J Pancreatol.* 2000;28:41-9.
60. Garcia de Gadiana RL, Albaladejo Oton MD, Rebello Acebes S, Esteban TP, Hernando HA, Jimenez SE et al. Diagnostic accuracy of lipopolysaccharide-binding protein for sepsis in patients with suspected infection in the emergency department. *Ann Clin Biochem.* 2018;55:143-8.
61. Ratzinger F, Schuardt M, Eichbichler K, Tsirkinidou I, Bauer M, Haslacher H et al. Utility of sepsis biomarkers and the infection probability score to discriminate sepsis and systemic inflammatory response syndrome in standard care patients. *PLoS One.* 2013;8:e82946.
62. Kitanovski L, Jazbec J, Hojker S, Derganc M. Diagnostic accuracy of lipopolysaccharide-binding protein for predicting bacteremia/clinical sepsis in children with febrile neutropenia: comparison with interleukin-6, procalcitonin, and C-reactive protein. *Support Care Cancer.* 2014;22:269-77.

63. ten Oever J, Tromp M, Bleeker-Rovers CP, Joosten LA, Netea MG, Pickkers P et al. Combination of biomarkers for the discrimination between bacterial and viral lower respiratory tract infections. *J Infect.* 2012;65:490-5.
64. Garcia de Gadiana-Romualdo L, Berger M, Jimenez-Santos E, Rebollo-Acebes S, Jimenez-Sanchez R, Esteban-Torrella P et al. Pancreatic stone protein and soluble CD25 for infection and sepsis in an emergency department. *Eur J Clin Invest.* 2017;47:297-304.
65. Jiang YN, Cai X, Zhou HM, Jin WD, Zhang M, Zhang Y et al. Diagnostic and prognostic roles of soluble CD22 in patients with Gram-negative bacterial sepsis. *Hepatobiliary Pancreat Dis Int.* 2015;14:523-9.
66. Balci C, Sungurtekin H, Gurses E, Sungurtekin U, Kaptanoglu B. Usefulness of procalcitonin for diagnosis of sepsis in the intensive care unit. *Crit Care.* 2003;7:85-90.
67. Ayazi P, Mahyar A, Daneshi MM, Jahanihashemi H, Esmailzadehha N, Mosaferirad N. Comparison of serum IL-1beta and C reactive protein levels in early diagnosis and management of neonatal sepsis. *Infez Med.* 2014;22:296-301.
68. Stojewska M, Wasek-Buko M, Jakub B, Wisniewska-Ulfig D, Goleniowska-Krol A, Szymanska A et al. Evaluation of serum chemokine RANTES concentration as a biomarker in the diagnosis of early-onset severe infections in neonates. *Postepy Hig Med Dosw (Online).* 2016;70:272-9.
69. Sapa A, Rak A, Wybieralska M, Machon J, Krzywonos-Zawadzka A, Zawadzki K et al. Diagnostic usefulness of sCD163, procalcitonin and neopterin for sepsis risk assessment in critically ill patients. *Adv Clin Exp Med.* 2017;26:101-8.
70. Ruokonen E, Ilkka L, Niskanen M, Takala J. Procalcitonin and neopterin as indicators of infection in critically ill patients. *Acta Anaesthesiol Scand.* 2002;46:398-404.
71. Kofoed K, Andersen O, Kronborg G, Tvede M, Petersen J, Eugen-Olsen J et al. Use of plasma C-reactive protein, procalcitonin, neutrophils, macrophage migration inhibitory factor, soluble urokinase-type plasminogen activator receptor, and soluble triggering receptor expressed on myeloid cells-1 in combination to diagnose infections: a prospective study. *Crit Care.* 2007;11:R38.
72. Demirkaya M, Tugcu D, Akcay A, Aydogan G, Akici F, Salcioglu Z et al. Adrenomedullin—A new marker in febrile neutropenia: Comparison with CRP and procalcitonin. *Pediatr Hematol Oncol.* 2015;32:482-9.
73. Enguix-Armada A, Escobar-Conesa R, Garcia-De La Torre A, De La Torre-Prados MV. Usefulness of several biomarkers in the management of septic patients: C-reactive protein, procalcitonin, presepsin and mid-regional pro-adrenomedullin. *Clin Chem Lab Med.* 2016;54:163-8.
74. Gaini S, Koldkjaer OG, Moller HJ, Pedersen C, Pedersen SS. A comparison of high-mobility group-box 1 protein, lipopolysaccharide-binding protein and procalcitonin in severe community-acquired infections and bacteraemia: a prospective study. *Crit Care.* 2007;11:R76.
75. El-Maghraby SM, Moneer MM, Ismail MM, Shalaby LM, El-Mahallawy HA. The diagnostic value of C-reactive protein, interleukin-8, and monocyte chemotactic protein in risk stratification of febrile neutropenic children with hematologic malignancies. *J Pediatr Hematol Oncol.* 2007;29:131-6.
76. Matera G, Puccio R, Giancotti A, Quirino A, Pulicari MC, Zicca E et al. Impact of interleukin-10, soluble CD25 and interferon-gamma on the prognosis and early diagnosis of bacteremic systemic inflammatory response syndrome: a prospective observational study. *Crit Care.* 2013;17:R64.
77. Mihajlovic D, Brkic S, Lendak D, Mikic AN, Draskovic B, Mitic G. Endothelial biomarkers in the light of new sepsis definition. *Biomark Med.* 2019;13:341-51.
78. Gille J, Schmidt J, Kremer T, Sablotzki A. Evaluation of MR-proANP and copeptin for sepsis diagnosis after burn injury. *J Crit Care.* 2019;52:149-55.
79. Lin Q, Fu F, Shen L, Zhu B. Pentraxin 3 in the assessment of ventilator-associated pneumonia: an early marker of severity. *Heart Lung.* 2013;42:139-45.
80. Vanska M, Koivula I, Hamalainen S, Pulkki K, Nousiainen T, Jantunen E et al. High pentraxin 3 level predicts septic shock and bacteremia at the onset of febrile neutropenia after intensive chemotherapy of hematologic patients. *Haematologica.* 2011;96:1385-9.
81. Adamzik M, Eggmann M, Frey UH, Gorlinger K, Brocker-Preuss M, Marggraf G et al. Comparison of thromboelastometry with procalcitonin, interleukin 6, and C-reactive protein as diagnostic tests for severe sepsis in critically ill adults. *Crit Care.* 2010;14:R178.

82. Gao L, Yang B, Zhang H, Ou Q, Lin Y, Zhang M et al. DcR3, a new biomarker for sepsis, correlates with infection severity and procalcitonin. *Oncotarget*. 2018;9:10934-44.
83. Tan TL, Goh YY. The role of group IIA secretory phospholipase A2 (sPLA2-IIA) as a biomarker for the diagnosis of sepsis and bacterial infection in adults-A systematic review. *PLoS One*. 2017;12:e0180554.
84. Wu TW, Tabangin M, Kusano R, Ma Y, Ridsdale R, Akinbi H. The utility of serum hepcidin as a biomarker for late-onset neonatal sepsis. *J Pediatr*. 2013;162:67-71.
85. Feng L, Zhou X, Su LX, Feng D, Jia YH, Xie LX. Clinical significance of soluble hemoglobin scavenger receptor CD163 (sCD163) in sepsis, a prospective study. *PLoS One*. 2012;7:e38400.
86. Arnon S, Litmanovitz I, Regev RH, Bauer S, Shaikin-Kestenbaum R, Dolfen T. Serum amyloid A: an early and accurate marker of neonatal early-onset sepsis. *J Perinatol*. 2007;27:297-302.
87. Zhou Y, Liu Z, Huang J, Li G, Li F, Cheng Y et al. Usefulness of the heparin-binding protein level to diagnose sepsis and septic shock according to Sepsis-3 compared with procalcitonin and C reactive protein: a prospective cohort study in China. *BMJ Open*. 2019;9:e026527.
88. Hildebrand D, Decker SO, Koch C, Schmitt FCF, Ruhrmann S, Schneck E et al. Host-derived delta-like canonical Notch ligand 1 as a novel diagnostic biomarker for bacterial sepsis - results from a combinational secondary analysis. *Front Cell Infect Microbiol*. 2019;9:267.
89. Ma L, Zhang H, Yin YL, Guo WZ, Ma YQ, Wang YB et al. Role of interleukin-6 to differentiate sepsis from non-infectious systemic inflammatory response syndrome. *Cytokine*. 2016;88:126-35.
90. Lin S, Huang Z, Wang M, Weng Z, Zeng D, Zhang Y et al. Interleukin-6 as an early diagnostic marker for bacterial sepsis in patients with liver cirrhosis. *J Crit Care*. 2015;30:732-8.
91. Llewelyn MJ, Berger M, Gregory M, Ramaiah R, Taylor AL, Curdt I et al. Sepsis biomarkers in unselected patients on admission to intensive or high-dependency care. *Crit Care*. 2013;17:R60.
92. Jonsson N, Nilsson T, Gille-Johnson P, Bell M, Martling CR, Larsson A et al. Calprotectin as an early biomarker of bacterial infections in critically ill patients: an exploratory cohort assessment. *Crit Care Resusc*. 2017;19:205-13.
93. Huang L, Li J, Han Y, Zhao S, Zheng Y, Sui F et al. Serum calprotectin expression as a diagnostic marker for sepsis in postoperative intensive care unit patients. *J Interferon Cytokine Res*. 2016;36:607-16.
94. Wong HR, Cvijanovich NZ, Hall M, Allen GL, Thomas NJ, Freishtat RJ et al. Interleukin-27 is a novel candidate diagnostic biomarker for bacterial infection in critically ill children. *Crit Care*. 2012;16:R213.
95. Wong HR, Liu KD, Kangelaris KN, Lahni P, Calfee CS. Performance of interleukin-27 as a sepsis diagnostic biomarker in critically ill adults. *J Crit Care*. 2014;29:718-22.
96. Li Z, Wang H, Liu J, Chen B, Li G. Serum soluble triggering receptor expressed on myeloid cells-1 and procalcitonin can reflect sepsis severity and predict prognosis: a prospective cohort study. *Mediators Inflamm*. 2014;2014:641039.
97. Jedynak M, Siemiatkowski A, Milewski R, Mroczko B, Szmitkowski M. Diagnostic effectiveness of soluble triggering receptor expressed on myeloid cells-1 in sepsis, severe sepsis and septic shock. *Arch Med Sci*. 2019;15:713-21.
98. Liu B, Chen YX, Yin Q, Zhao YZ, Li CS. Diagnostic value and prognostic evaluation of Presepsin for sepsis in an emergency department. *Crit Care*. 2013;17:R244.
99. Iskandar A, Arthamin MZ, Indriana K, Anshory M, Hur M, Di Somma S. Comparison between presepsin and procalcitonin in early diagnosis of neonatal sepsis. *J Matern Fetal Neonatal Med*. 2019;32:3903-8.
100. Dollner H, Vatten L, Austgulen R. Early diagnostic markers for neonatal sepsis: comparing C-reactive protein, interleukin-6, soluble tumour necrosis factor receptors and soluble adhesion molecules. *J Clin Epidemiol*. 2001;54:1251-7.
101. Yang AP, Liu J, Yue LH, Wang HQ, Yang WJ, Yang GH. Neutrophil CD64 combined with PCT, CRP and WBC improves the sensitivity for the early diagnosis of neonatal sepsis. *Clin Chem Lab Med*. 2016;54:345-51.
102. Grigoras I, Branisteanu DD, Ungureanu D, Rusu D, Ristescu I. Early dynamics of leptin plasma level in surgical critically ill patients. a prospective comparative study. *Chirurgia (Bucur)*. 2014;109:66-72.
103. Angeletti S, Battistoni F, Fioravanti M, Bernardini S, Dicuonzo G. Procalcitonin and mid-regional pro-adrenomedullin test combination in sepsis diagnosis. *Clin Chem Lab Med*. 2013;51:1059-67.

104. Zeng M, Chang M, Zheng H, Li B, Chen Y, He W et al. Clinical value of soluble urokinase-type plasminogen activator receptor in the diagnosis, prognosis, and therapeutic guidance of sepsis. *Am J Emerg Med*. 2016;34:375-80.
105. Nuutila J, Jalava-Karvinen P, Hohenthal U, Laitinen I, Kotilainen P, Rajamaki A et al. CRP/CD11b ratio: a novel parameter for detecting gram-positive sepsis. *Hum Immunol*. 2009;70:237-43.
106. Juutilainen A, Hamalainen S, Pulkki K, Kuittinen T, Nousiainen T, Jantunen E et al. Biomarkers for bacteremia and severe sepsis in hematological patients with neutropenic fever: multivariate logistic regression analysis and factor analysis. *Leuk Lymphoma*. 2011;52:2349-55.
107. Kelly BJ, Lautenbach E, Nachamkin I, Coffin SE, Gerber JS, Fuchs BD et al. Combined biomarkers discriminate a low likelihood of bacterial infection among surgical intensive care unit patients with suspected sepsis. *Diagn Microbiol Infect Dis*. 2016;85:109-15.
108. Suri M, Thirupuram S, Sharma VK. Diagnostic and prognostic utility of C-reactive protein, alpha-1-antitrypsin and alpha-2-macroglobulin in neonatal sepsis: a comparative account. *Indian Pediatr*. 1991;28:1159-64.
109. Kim S, Fotiadu A, Kotoula V. Increased expression of soluble decoy receptor 3 in acutely inflamed intestinal epithelia. *Clin Immunol*. 2005;115:286-94.
110. Perrotti A, Chenevier-Gobeaux C, Ecartot F, Barrucand B, Lassalle P, Dorigo E et al. Relevance of endothelial cell-specific molecule 1 (endocan) plasma levels for predicting pulmonary infection after cardiac surgery in chronic kidney disease patients: The Endolung Pilot Study. *Cardiorenal Med*. 2017;8:1-8.
111. Djebara S, Biston P, Fosse E, Daper A, Joris M, Boudjeltia KZ et al. Time course of CD64, a leukocyte activation marker, during cardiopulmonary bypass surgery. *Shock*. 2017;47:158-64.
112. Klein HJ, Csordas A, Falk V, Slankamenac K, Rudiger A, Schonrath F et al. Pancreatic stone protein predicts postoperative infection in cardiac surgery patients irrespective of cardiopulmonary bypass or surgical technique. *PLoS One*. 2015;10:e0120276.
113. Shimizu T, Endo Y, Tabata T, Mori T, Hanasawa K, Tsuchiya M et al. Diagnostic and predictive value of the silkworm larvae plasma test for postoperative infection following gastrointestinal surgery. *Crit Care Med*. 2005;33:1288-95.
114. Duswald KH, Jochum M, Schramm W, Fritz H. Released granulocytic elastase: an indicator of pathobiochemical alterations in septicemia after abdominal surgery. *Surgery*. 1985;98:892-9.
115. Lendak DF, Mihajlovic DM, Novakov-Mikic AS, Boban JM, Ubavic M, Brkic SV. APRIL and sTACI could be predictors of multiorgan dysfunction syndrome in sepsis. *Virulence*. 2018;9:946-53.
116. Thulborn SJ, Dilpazir M, Haldar K, Mistry V, Brightling CE, Barer MR et al. Investigating the role of pentraxin 3 as a biomarker for bacterial infection in subjects with COPD. *Int J Chron Obstruct Pulmon Dis*. 2017;12:1199-205.
117. Lausevic Z, Vukovic G, Stojimirovic B, Trbojevic-Stankovic J, Resanovic V, Lausevic M. Kinetics of C-reactive protein, interleukin-6 and -10, and phospholipase A2-II in severely traumatized septic patients. *Vojnosanit Pregl*. 2010;67:893-7.
118. Bahrami S, Pelinka L, Khadem A, Maitzen S, Hawa G, van Griensven M et al. Circulating NT-proCNP predicts sepsis in multiple-traumatized patients without traumatic brain injury. *Crit Care Med*. 2010;38:161-6.
119. Simons RK, Hoyt DB, Winchell RJ, Rose RM, Holbrook T. Elevated selectin levels after severe trauma: a marker for sepsis and organ failure and a potential target for immunomodulatory therapy. *J Trauma*. 1996;41:653-62.
120. Hull MA, Jones BA, Zurakowski D, Raphael B, Lo C, Jaksic T et al. Low serum citrulline concentration correlates with catheter-related bloodstream infections in children with intestinal failure. *JPEN J Parenter Enteral Nutr*. 2011;35:181-7.
121. Ng PC, Li K, Chui KM, Leung TF, Wong RP, Chu WC et al. IP-10 is an early diagnostic marker for identification of late-onset bacterial infection in preterm infants. *Pediatr Res*. 2007;61:93-8.
122. Oude Nijhuis CSM, Vellenga E, Daenen SMGJ, van der Graaf WTA, Gietema JA, Groen HJM et al. Lipopolysaccharide-binding protein: a possible diagnostic marker for Gram-negative bacteremia in neutropenic cancer patients. *Intensive Care Med*. 2003;29:2157-61.
123. Al Shuaibi M, Bahu RR, Chaftari AM, Al Wohoush I, Shomali W, Jiang Y et al. Pro-adrenomedullin as a novel biomarker for predicting infections and response to antimicrobials in febrile patients with hematologic malignancies. *Clin Infect Dis*. 2013;56:943-50.
124. Kraft R, Herndon DN, Finnerty CC, Cox RA, Song J, Jeschke MG. Predictive value of IL-8 for sepsis and severe infections after burn injury: A clinical study. *Shock*. 2015;43:222-7.

125. Grieb G, Simons D, Piatkowski A, Bernhagen J, Steffens G, Pallua N. Macrophage migration inhibitory factor-A potential diagnostic tool in severe burn injuries? *Burns*. 2010;36:335-42.
126. Ajmani S, Singh H, Chaturvedi S, Mishra R, Rai MK, Jain A et al. Utility of neutrophil CD64 and serum TREM-1 in distinguishing bacterial infection from disease flare in SLE and ANCA-associated vasculitis. *Clin Rheumatol*. 2019;38:997-1005.
127. Deitcher SR, Eisenberg PR. Elevated concentrations of cross-linked fibrin degradation products in plasma. An early marker of gram-negative bacteremia. *Chest*. 1993;103:1107-12.
128. Rendon-Ramirez EJ, Ortiz-Stern A, Martinez-Mejia C, Salinas-Carmona MC, Rendon A, Mata-Tijerina VL et al. TGF-beta blood levels distinguish between Influenza A (H1N1)pdm09 virus sepsis and sepsis due to other forms of community-acquired pneumonia. *Viral Immunol*. 2015;28:248-54.
129. Hall MW, Geyer SM, Guo CY, Panoskaltsis-Mortari A, Jouvett P, Ferdinands J et al. Innate immune function and mortality in critically ill children with influenza: a multicenter study. *Crit Care Med*. 2013;41:224-36.
130. Sunnetcioglu A, Sunnetcioglu M, Adiyaman F, Binici I, Soyoral L. Could soluble urokinase plasminogen receptor (suPAR) be used as a diagnostic biomarker for ventilator-associated pneumonia? *Clin Respir J*. 2017;11:925-30.
131. Jones JW, Jr., Su S, Jones MB, Heniford BT, McIntyre K, Granger DK. Serum lysozyme activity can differentiate infection from rejection in organ transplant recipients. *J Surg Res*. 1999;84:134-7.
132. Ware LB, Koyama T, Zhao Z, Janz DR, Wickersham N, Bernard GR et al. Biomarkers of lung epithelial injury and inflammation distinguish severe sepsis patients with acute respiratory distress syndrome. *Crit Care*. 2013;17:R253.
133. Rubin DB, Wiener-Kronish JP, Murray JF, Green DR, Turner J, Luce JM et al. Elevated von Willebrand factor antigen is an early plasma predictor of acute lung injury in nonpulmonary sepsis syndrome. *J Clin Invest*. 1990;86:474-80.
134. Liu XW, Ma T, Cai Q, Wang L, Song HW, Liu Z. Elevation of serum PARK7 and IL-8 levels is associated with acute lung injury in patients with severe sepsis/septic shock. *J Intensive Care Med*. 2019;34:662-8.
135. Smart L, Bosio E, Macdonald SPJ, Dull R, Fatovich DM, Neil C et al. Glycocalyx biomarker syndecan-1 is a stronger predictor of respiratory failure in patients with sepsis due to pneumonia, compared to endocan. *J Crit Care*. 2018;47:93-8.
136. Zheng YJ, Xu WP, Ding G, Gao YH, Wang HR, Pan SM. Expression of HMGB1 in septic serum induces vascular endothelial hyperpermeability. *Mol Med Rep*. 2016;13:513-21.
137. Jabaudon M, Futier E, Roszyk L, Chalus E, Guerin R, Petit A et al. Soluble form of the receptor for advanced glycation end products is a marker of acute lung injury but not of severe sepsis in critically ill patients. *Crit Care Med*. 2011;39:480-8.
138. de Bont ES, Vellenga E, Swaanenburg JC, Fidler V, Visser-van Brummen PJ, Kamps WA. Plasma IL-8 and IL-6 levels can be used to define a group with low risk of septicemia among cancer patients with fever and neutropenia. *Br J Haematol*. 1999;107:375-80.
139. Diepold M, Noellke P, Duffner U, Kontny U, Berner R. Performance of Interleukin-6 and Interleukin-8 serum levels in pediatric oncology patients with neutropenia and fever for the assessment of low-risk. *BMC Infect Dis*. 2008;8:28.
140. Spasova MI, Terzieva DD, Tzvetkova TZ, Stoyanova AA, Mumdzhev IN, Yanev IB et al. Interleukin-6, interleukin-8, interleukin-10, and C-reactive protein in febrile neutropenia in children with malignant diseases. *Folia Med (Plovdiv)*. 2005;47:46-52.
141. Aquino VM, Cost C, Gomez A, Bowers DC, Ramilo O, Ahmad N et al. Predictive value of interleukin-5 and monocyte chemoattractant protein-1 for bacteremia in children with febrile neutropenia. *J Pediatr Hematol Oncol*. 2012;34:e241-e245.
142. Magudumana MO, Ballot DE, Cooper PA, Trusler J, Cory BJ, Viljoen E et al. Serial interleukin 6 measurements in the early diagnosis of neonatal sepsis. *J Trop Pediatr*. 2000;46:267-71.
143. Kuster H, Weiss M, Willeitner AE, Detlefsen S, Jeremias I, Zbojan J et al. Interleukin-1 receptor antagonist and interleukin-6 for early diagnosis of neonatal sepsis 2 days before clinical manifestation. *Lancet*. 1998;352:1271-7.
144. Vassiliou AG, Kotanidou A, Mastora Z, Maniatis NA, Albani P, Jahaj E et al. Elevated soluble endothelial protein C receptor levels at ICU admission are associated with sepsis development. *Minerva Anestesiol*. 2015;81:125-34.
145. Su CM, Cheng HH, Tsai TC, Hsiao SY, Tsai NW, Chang WN et al. Elevated serum vascular cell adhesion molecule-1 is associated with septic encephalopathy in adult community-onset severe sepsis patients. *Biomed Res Int*. 2014;2014:598762.
146. Anderson BJ, Reilly JP, Shashaty MGS, Palakshappa JA, Wysoczanski A, Dunn TG et al. Admission plasma levels of the neuronal injury marker neuron-specific enolase are associated with mortality and delirium in sepsis. *J Crit Care*. 2016;36:18-23.
147. Mosad E, Elsayh KI, Eltayeb AA. Tissue factor pathway inhibitor and P-selectin as markers of sepsis-induced non-overt disseminated intravascular coagulopathy. *Clin Appl Thromb Hemost*. 2011;17:80-7.

148. Koyama K, Madoiwa S, Nunomiya S, Koinuma T, Wada M, Sakata A et al. Combination of thrombin-antithrombin complex, plasminogen activator inhibitor-1, and protein C activity for early identification of severe coagulopathy in initial phase of sepsis: a prospective observational study. *Crit Care*. 2014;18:R13.
149. Delabranche X, Boisrame-Helms J, Asfar P, Berger A, Mootien Y, Lavigne T et al. Microparticles are new biomarkers of septic shock-induced disseminated intravascular coagulopathy. *Intensive Care Med*. 2013;39:1695-703.
150. Sivula M, Hastbacka J, Kuitunen A, Lassila R, Tervahartiala T, Sorsa T et al. Systemic matrix metalloproteinase-8 and tissue inhibitor of metalloproteinases-1 levels in severe sepsis-associated coagulopathy. *Acta Anaesthesiol Scand*. 2015;59:176-84.
151. Psuja P, Zozulinska M, Turowiecka Z, Cieslikowski W, Vinazzer H, Zawilska K. Plasma markers of hypercoagulability in patients with serious infections and risk of septic shock. *Clin Appl Thromb Hemost*. 2002;8:225-30.
152. Koyama K, Madoiwa S, Tanaka S, Koinuma T, Wada M, Sakata A et al. Evaluation of hemostatic biomarker abnormalities that precede platelet count decline in critically ill patients with sepsis. *J Crit Care*. 2013;28:556-63.

Supplementary Information

Table S1 Cytokine/chemokine biomarkers identified in the literature search

Table S2 Receptor biomarkers identified in the literature search

Table S3 Cell marker biomarkers identified in the literature search

Table S4 Coagulation-related biomarkers identified in the literature search

Table S5. Microcirculation related biomarkers identified in the literature search

Table S6. Vasodilation-related biomarkers identified in the literature search

Table S7. Biomarkers of organ dysfunction in sepsis identified in the literature search

Table S8. Acute phase proteins used as biomarkers in sepsis identified in the literature search

Table S9 Diverse sepsis biomarkers identified in the literature search

Tables

Table 1. Sepsis biomarkers, except for C-reactive protein (CRP) and procalcitonin (PCT), that have been evaluated for their diagnostic value in clinical studies with more than 300 subjects.

Biomarker [ref]	No of patients	Sepsis definition	Study population	Reference Group	Sensitivity/Specificity (%)	AUC
Interleukin (IL)-27 [37]	702	Positive blood cultures	Pediatric ICU patients with infection	Non-infected critical care patients	84/63	0.75
Inter-alpha inhibitor proteins [11]	573	Positive blood cultures	Neonates with sepsis	Neonates with risk factors for sepsis	89/99	0.9
Group II phospholipase A2 [38]	525	ACCP 1992	ED patients with sepsis	ED patients with suspected infection (with and without SIRS)	NR (logistic regression analysis)	NR
Bactericidal/permeability increasing protein [38]	525	ACCP 1992	ED with sepsis	ED patients with suspected infection (with and without SIRS)	NR (logistic regression analysis)	NR
CD64 [12]	468	International Sepsis Definitions Conference 2001	Non selected population ICU population with sepsis	ICU patients admitted without sepsis	89/87	0.94
Selenoprotein P [39]	378	ACCP 1992	Non selected population with sepsis or septic shock	Healthy individuals	NR (no test)	NR
Lipopolysaccharide-binding protein [40]	327	ACCP 1992	Surgical patients without sepsis at admission	Surgical patients with SIRS without sepsis	60/62	0.66
Syndecan-1 [41]	512	International Sepsis Definitions Conference 2001	Trauma patients (4h after admission) without sepsis	Trauma patients without sepsis	NR (logistic regression analysis)	NR
Presepsin [42]	440	Sepsis-3	ICU patients with sepsis	ICU patients without sepsis	89/59	0.76
IL-6 [13]	306	SIRS and organ dysfunction, systolic blood pressure <90 mmHg or lactate $\geq$ 4 mmol/L plus infection	ED patients with suspected sepsis	ED patients with SIRS and organ dysfunction, systolic blood pressure <90 mmHg or lactate $\geq$ 4 mmol/L without infection.	NR	0.86

ED: emergency department; ICU: intensive care unit; COPD: chronic obstructive pulmonary disease; SIRS: systemic inflammatory response syndrome; NR: not reported; AUC: area under the receiver operating characteristic curve

Table 2. Sepsis biomarkers, except for C-reactive protein (CRP) and procalcitonin (PCT), that have been evaluated for their prognostic value in clinical studies with more than 300 subjects.

Name [ref]	No of patients	Sepsis Definition	Study Population	Main finding	Sensitivity/Specificity (%)	AUC
Urokinase plasminogen activator receptor (uPAR) [14]	1914	International Sepsis Definitions Conference 2001	Critically ill patients and patients hospitalized in internal medicine ward	Levels ≥ 12 ng/mL predicted fatal outcome within 30 days	NR/ >70%,	0.708 APACHE II 0.83
Plasminogen activator inhibitor (PAI) 1 [43]	1790	ACCP 1992	Septic patients with disseminated intravascular coagulation (DIC)	Levels > 90 ng/ml predict fatal outcome within 30 days	NR (Kaplan-Meier survival functions)	NR
Interleukin (IL)-12 [44]	1444	Proven peritonitis or mediastinitis and systemic inflammation signs	Surgical patients	Pre-surgery IL-12-synthesizing capability was low in patients who had fatal sepsis after operation	NR	0.72
Thrombomodulin [45]	1103	ACCP 1992	Critically ill patients with sepsis	Levels > 14 ng/ml can predict circulatory failure or death - gray zone between 7-14 ng/ml	NR (logistic regression analysis)	NR
Syndecan-1 [45]	1103	ACCP 1992	Critically ill patients with sepsis	Levels > 240 ng/ml can predict circulatory failure or death gray zone between 70-240 ng/ml	NR (logistic regression analysis)	NR
Fibrinogen [46]	1103	ACCP 1992	Critically ill patients with sepsis	Levels <200 mg/dl related to increased risk of fatal outcome	NR (Logistic regression analysis)	NR
Antithrombin activity [46]	1103	ACCP 1992	Critically ill patients with sepsis	Decrease in activity >50% related to increased risk of fatal outcome.	NR (Logistic regression analysis)	NR
Brain natriuretic peptide (BNP) [47]	1000	International Sepsis Definition Conference 2001	ED patients	Levels > 113 pg/ml can predict fatal outcome within 28 days	86/55	0.73
Angiopietin-2 [48]	931	NR	Critically ill	Persistently	NR (Logistic	NR

			patients with ARDS	increased levels related to fatal outcome within 90 days	regression analysis)	
Prothrombin time (PT) [49]	840	Suspected infection plus ≥ 3 signs of systematic inflammatory response	Critically ill patients with sepsis	Increase in PT time within first 7 days of sepsis was higher in patients who died within 28 days	NR (no test)	NR
Adrenomedullin [15]	837	International Sepsis Definitions Conference 2001	ED patients sepsis	Levels < 34.4 ng/L predicted fatal outcome within 30 days	86/61	0.77 Mortality in Emergency Department Sepsis (MEDS) score AUC: 0.81
Pro-adrenomedullin [50]	896	Clinical suspicion of infection	ED patients with sepsis	Levels ≥ 1.6 nmol/L predicted fatal outcome within 28 days	75/92	0.89
Heparin-binding protein [51]	759	Suspected infection and at least one clinical sign of systematic inflammatory response	ED patients with sepsis	Levels > 30 ng/mL predicted any organ dysfunction development within 72 h	78/76 (cross-tabulation analysis)	NR
D-dimer [52]	684	International Sepsis Definitions Conference 2001	Emergency Department patients with sepsis	Higher in non-survivors than survivors within 28 days	NR	0.68
Troponin [53]	598	ACCP 1992	Critically ill patients	Levels > 0.06 ng/mL independent prognostic marker for 28-day mortality	NR (Logistic regression analysis)	NR
YKL-40 [54]	502	ACCP 1992	Critically ill patients	Levels ≤ 505 ng/ml predicted survival in 90 days	53/76	0.64
CD64 [12]	468	International Sepsis Definition Conference 2001	Critically ill patients	Sustained elevated levels were related to non-appropriate antibiotic therapy.	93/48	0.74

Cell-free DNA [55]	481	International Sepsis Definitions Conference 2001	ED patients	Levels >1.6 µg/ml predicted short term fatal outcome.	70/76	0.77
--------------------	-----	--	-------------	---	-------	------

ARDS: acute respiratory distress syndrome; NR: not reported; IL: interleukin; SOFA: sequential organ failure assessment

AUC: area under the receiver operating characteristic curve

Table 3. Sepsis biomarkers that were compared with procalcitonin (PCT) and/or C-reactive protein (CRP) for sepsis diagnosis

Diagnostic performance similar to or worse than that of PCT and/or CRP			
Biomarker	Study Group	Reference Group	Comment [refs]
Cell-free DNA (cfDNA)	ICU patients with sepsis	ICU patients with SIRS	No better than PCT [56, 57]
Copeptin	ED patients with sepsis	ED patients with SIRS	No better than PCT [58]
ICAM-1	Patients with necrotic pancreatitis	Patients with sterile necrosis	No better than PCT [59]
Lipopolysaccharide-binding protein	ED patients with sepsis	ED patients with infection	No better than PCT [60]
	Non-critically ill patients with sepsis	Non-critically ill patients with infection	No better than PCT [61]
	Children with neutropenia and clinical sepsis and/or bacteremia,	Children with febrile neutropenia without infection	No better than PCT [62]
	patients with proved bacterial lower respiratory infection	Patients with proven viral lower respiratory infection	No better than CRP [63]
	Patients treated in internal medicine ward,	Healthy control	No better than PCT [16]
Pancreatic stone protein	ED patients with sepsis	ED patients without infection	No better than PCT [64]
sCD22	Surgical patients with infection after major operation	Surgical patients without infection but SIRS	Equal value to PCT [65]
Interleukin (IL)-2	ICU patients with sepsis	ICU patients with SIRS without infection	No better than CRP [66]
IL-1b	Neonates with infection and sepsis	Neonates with infection without sepsis	No better than CRP [67]
RANTES	Neonates with infection	Healthy neonates controls	No better than CRP [68]
Neopterin	ICU patients with sepsis	ICU patients without sepsis	Less accurate than PCT [69, 70]
Macrophage migration inhibitory factor (MIF)	Patients with infection in medical ward or ED	No bacterial infection	No better than PCT [71]
Adrenomedullin	Neutropenic patients with sepsis	Neutropenic patients with fever and clinically documented infection	No better than PCT [72]
Pro-adrenomedullin	Sepsis with organ dysfunction and or shock	Patients admitted to coronary unit without infection	No better than PCT [73]
High-mobility group-box 1 protein (HMGB1)	Infected patients admitted in the ward	Healthy individuals	No better than CRP or PCT [74]
IL-8	Neutropenic children with blood culture positive, and/or fever periods with a documented clinical sepsis and/or local infection	Neutropenic children with fever and no infection	No better than CRP in diagnosis of sepsis [75]
IL-10	Patients with bacteremia and SIRS ,	Patients with SIRS without bacteremia	Comparable with PCT [76]
Endocan	Critically ill patients with sepsis and organ dysfunction	Critically ill patients with infection and SIRS	Comparable with PCT [77]
Pro-atrial natriuretic peptide (ANP)	Burned patients that received antibiotics and had either microbiological confirmation of infection or antibiotics led to an improvement in clinical situation.	Burned patients without infection	Comparable with PCT [78]
Pentraxin 3	Mechanically ventilated patients with ventilator associated pneumonia	Mechanically ventilated patient >48h without VAP	No better than CRP [79]

	Hematological patients with bacteremia and/or septic shock	Hematological patients with fever without infection	No better than CRP [80]
Better diagnostic value than PCT and/or CRP			
Thromboelastometry lysis index	Patients with severe sepsis	Patients after operation without sepsis	Better than PCT [81]
Decoy receptor 3	ICU patients with sepsis	ICU patients with SIRS	Positive when PCT was negative [82]
group II Phospholipase A2 (PLA2-II)	ED patients with sepsis and organ dysfunction	ED patients with SIRS without infection	Better than CRP [83]
Hepcidin	Infants with sepsis and or bacteremia	Infants with SIRS and not sepsis	Better than CRP [84]
sCD163	Patients with sepsis admitted to ICU	Patients with SIRS without sepsis	Better than PCT [85]
CD64	ICU patients with sepsis	ICU patients without sepsis	Better than PCT and CRP [32]
	Patients with ventilator associated pneumonia and sepsis	Patients with ventilator associated pneumonia without sepsis	Better than PCT and CRP [33]
Serum amyloid A	Full term infants with sepsis	Full term infants with risk for sepsis but without sepsis	Earlier increase in neonates with early onset sepsis than CRP [86]
Heparin binding protein	Patients with Sepsis for less than 48h	Patients with infection without sepsis	Better than CRP and PCT [87]
Delta-like canonical Notch ligand 1 (DLL1)	Patients with abdominal infection or surgical site associated infection	Surgical patients, trauma patients without infection and healthy volunteers	Better than CRP and PCT [88]
Conflicting findings			
IL-6	Critically ill patients with Sepsis	Patients with SIRS without infection	IL-6 was not found to have lower diagnostic utility compared to PCT (meta-analysis) [89]
	Cirrhotic patients with infection at admission to ICU	Cirrhotic patients without sepsis	IL-6 was found to increase earlier than PCT in cirrhotic patients [90]
sCD25	ED patients with infection	ED patients with suspected infection but finally infection excluded	Equal diagnostic value to PCT for diagnosis of infection in ED [64]
	Patients admitted in ICU with infection and SIRS	Patients with SIRS without sepsis	Better performance than PCT to identify Sepsis I at ICU admission [91]
Calprotectin	ICU patients with infection	ICU patients without sepsis	Better than CRP and PCT [92]
	Patients after major operation who developed sepsis	Patients after major operation that did not developed sepsis	Similar value to PCT [93]
IL-27	Critically ill children with sepsis	Children with SIRS without infection	Better than PCT [94]
	ICU patients with sepsis	ICU patients without sepsis	No better than PCT [95]
sTREM	ICU patients with sepsis	ICU patients with SIRS	Better than PCT [96]
	ICU patients with sepsis	ICU patients with SIRS	No better than PCT and CRP [97]
Presepsin (CD14)	ED patients with sepsis	ED patients with at least two	Better than PCT in diagnosis

		criteria of SIRS without sepsis	of sepsis in ED [98]
	Critically ill patients with sepsis and organ dysfunction	Critically ill patients without infection	Not better than PCT regardless of the presence or not of AKI [42]
	Neonates with SIRS and positive blood cultures	Neonates with SIRS with negative blood cultures	Better than PCT [99]
Better performance when combined with PCT and/or CRP			
IL-6	Neonates with infection within the first week of life	Neonates with suspicion of infection but finally excluded within the first week of sepsis	Combination with CRP in neonates with suspected sepsis [100]
CD64	Neonates with sepsis	healthy controls	Combination with PCT and CRP for diagnosis of neonatal sepsis [101]
Leptin	Patients with Community acquired pneumonia with sepsis or Complicated intraabdominal infection	SIRS without infections healthy controls	Combination with CRP [102]
Pro-adrenomedullin	Septic patients	Patients with SIRS without sepsis	Combination to PCT [73, 103]
suPAR	Septic patients admitted to ICU	Critically ill patients with SIRS without infection and healthy controls	Combination with PCT to diagnosis of sepsis on day 1 of sepsis [104]
CD11b	Patients with Gram (+) infection	Patients with Gram (-) infection	Combination with CRP for differentiation from Gram (-) infection [105]
Fibrinogen	Neutropenic patients with sepsis	Neutropenic patients with fever without infection	Combination with CRP for diagnosis of sepsis [106]
BNP and antithrombin	Neutropenic patients with fever and bacteremia	Neutropenic patients with fever without infection	Combination with PCT for diagnosis of Gram (-) bacteremia [106]
IL-27	Pediatric patients with sepsis,	Pediatric patient with SIRS without infection	Improvement of diagnostic accuracy of PCT for diagnosis of sepsis [95, 107]
α -2 macroglobulin	Surgical patients with sepsis	Surgical patients with SIRS without sepsis	Combination with PCT for exclude sepsis in surgical patients [108]
Decoy receptor 3 and uPAR	Patients with sepsis	Patients with SIRS without infection , healthy volunteers	Combination with PCT for diagnosis of sepsis [109]

sTREM: soluble triggering receptor expressed on myeloid cells; RANTES: regulated on activation, normal T-cell expressed, and secreted

Table 4. Some examples of biomarkers that have been assessed for use in specific clinical situations

Situation	Biomarker
<i>To diagnose infection in patients with a particular pathology/condition</i>	
After cardiac surgery	Endocan [110], CD64 [111], pancreatic stone protein [112]
After major surgery	Peptidoglycan [113], elastase [114], leptin [102], calprotectin [93], a proliferation-inducing ligand [115], α -2 macroglobulin [107], lipopolysaccharide-binding protein [40]
COPD	Pentraxin 3 [116]
Cirrhosis	Interleukin (IL)-6 [90]
Trauma	IL-10 [117], NT-proCNP [118], P-selectin [119]
Catheter-related infections	Citrulline [120]
Infants with necrotic enterocolitis	IP-10 [121]
Neutropenic patients	Lipopolysaccharide-binding protein [122], pro-adrenomedullin [123]
Burns	IL-8 [124], MIF [125]
Autoimmune diseases	CD64 [126]
<i>To diagnose specific types of infection</i>	
Gram (-) vs Gram (+)	Fibrin degradation products [127], lipopolysaccharide-binding protein [122], CD11b [105]
Virus vs bacterial infection or co-infection	Transforming growth factor (TGF- β) [128], tumor necrosis factor (TNF)- α [129]
VAP	suPAR [130]
<i>Diagnosis of specific conditions</i>	
Sepsis vs graft rejection	Lysozyme [131]
Diagnosis of ARDS	Club cell secretory protein (CC)-16 [132], surfactant protein [132]
Vascular leakage risk in ARDS	von Willebrand factor [133], angiopoietins (1&2) [48], IL-8 [134], syndecan-1 [135], HMGB-1 [136]
Recovery from ARDS - endothelial repair	sRAGE [137]
Identification of low-risk of infection in hematological/oncological patients	IL-6 [138, 139], IL-8 [138-140], MCP-1 [75], IL-5 [141]
Identification of infection before clinical symptoms	IL-6 [142], IL-1ra [143], soluble protein C receptors [144]
Risk of encephalopathy/delirium	VCAM [145], neuron-specific enolase [146]
Disseminated intravascular coagulation	P-selectin [147], protein C [148], microparticles [149], matrix-metalloproteinases [150], thrombin-antithrombin complex [151], a2PI [152]

COPD: chronic obstructive pulmonary disease; ARDS: acute respiratory distress syndrome; TNF: tumor necrosis factor; VAP: ventilator-associated pneumonia; NT-ProCNP: N-terminal pro-C-type natriuretic peptide; MIF: macrophage migration inhibitory factor; VCAM: vascular cell adhesion molecule; IP: interferon-gamma-inducible protein; sUPAR: soluble urokinase plasminogen receptor; IL-1ra: IL-1 receptor antagonist; MCP: monocyte chemoattractant protein; sRAGE: soluble receptor for advanced glycation end products; HMGB: high-mobility group-box 1 protein

Figures

Figure 1

Changes over time in a) number of references meeting our search criteria; and b) number of new biomarkers referred to in identified references

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Sepsisbiomarkerssupplfinal.pdf](#)