

A common variant of the *NOTCH4* gene modulates functional connectivity of the occipital cortex and its relationship with schizotypal traits

xiaohui xie

First Affiliated Hospital of Anhui Medical University <https://orcid.org/0000-0002-6324-2731>

Meidan Zu

First Affiliated Hospital of Anhui Medical University

Long Zhang

First Affiliated Hospital of Anhui Medical University

Tongjian Bai

First Affiliated Hospital of Anhui Medical University

Ling Wei

First Affiliated Hospital of Anhui Medical University

Wanling Huang

First Affiliated Hospital of Anhui Medical University

Gong-Jun Ji

Anhui Medical College

Bensheng Qiu

China University of Science and Technology

Panpan Hu

First Affiliated Hospital of Anhui Medical University

Yanghua Tian

First Affiliated Hospital of Anhui Medical University

Kai Wang (✉ wangkai1964@126.com)

Research article

Keywords: NOTCH4, rs204993, functional connectivity strength, occipital cortex, schizotypal traits

Posted Date: April 27th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-15827/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Psychiatry on July 9th, 2020. See the published version at <https://doi.org/10.1186/s12888-020-02773-z>.

Abstract

Background: Schizotypal traits are considered as inheritable traits and the endophenotype for schizophrenia. A common variant in the *NOTCH4* gene, rs204993, has been linked with schizophrenia, but the neural underpinnings are largely unknown.

Methods: In present study, we compared the differences of brain functions between different genotypes of rs204993 and its relationship with schizotypal traits among 402 Chinese Han healthy volunteers. The brain function was evaluated with functional connectivity strength (FCS) using the resting-state functional magnetic resonance image (rs-fMRI). The schizotypal traits were measured by the schizotypal personality questionnaire (SPQ).

Results: Our results showed that carriers with the AA genotype showed reduced FCS in the left occipital cortex when compared with carriers with the AG and GG genotypes, and the carriers with the AG genotype showed reduced FCS in the left occipital cortex when compared with carriers with the GG genotype. The FCS values in the left occipital lobe were negatively associated with the SPQ scores and its subscale scores within the carriers with the GG genotype, but not within the carriers with AA or AG genotype.

Conclusion: Our results suggested that the common variant in the *NOTCH4* gene, rs204993, modulates the function of the occipital cortex, which may contribute to schizotypal traits. These findings provide insight for genetic effects on schizotypal traits and its potential neural substrate.

Background

Schizophrenia is a chronically severe, mental disorder, affecting approximately 1% of the worldwide population [1]. In addition, some individuals present preclinical symptoms of schizophrenia called schizotypal traits [2]. Schizotypal traits refer to a set of traits, continually distributed in the general population, that resemble some of the symptoms of schizophrenia. Schizotypal traits provide important insights into understanding the origins of schizophrenia [3]. Raine et al. concluded that structural and functional alterations in the brain induced by genetic and early environmental influences contribute to the development of schizotypy [4]. Indeed, schizotypal traits are more frequent in first-degree relatives of patients with schizophrenia, suggesting a genetic link between schizotypal traits and schizophrenia [5]. The schizophrenia-risk gene also contributes to the presence of schizotypal traits [6, 7]. Hence, the schizophrenia-risk gene may have potential modulatory effects on schizotypal traits.

Although previous studies have found that some genes indicate susceptibility for schizophrenia [8, 9], only a small percentage of the genes have been consistently studied among different racial populations. The *NOTCH4* (neurogenic locus notch homolog protein4) gene, located in the major histocompatibility complex (MHC) region of 6p21.3 in humans, is highly associated with schizophrenia among different racial populations [10-12]. Wei and Hemmings found that the *NOTCH4* locus was involved in susceptibility to schizophrenia in a British population [12]. The relationship between schizophrenia and the common variants of *NOTCH4* gene was also found among different racial populations (the

rs2071287 and rs3131296 in European and Japanese population) [13, 14]. However, these SNPs were not confirmed among Chinese populations. Recently, a study among 218 Taiwanese families reported another SNP, rs204993 with a significant relationship with schizophrenia, found by testing the association of the entire genomic region of *NOTCH4* [15]. Specially, the AA genotype of rs204993 was associated with a higher risk for schizophrenia in the Chinese Han population [16]. These findings suggested the common variant of the NOTCH4 gene, SNP rs204993 may be a potential genetic basis for schizophrenia in the Chinese population.

Recent evidences suggested the intermediary role of brain function for the genetic effect on schizophrenia or schizotypal traits. However, the effect of the common variant of the NOTCH4 gene, SNP rs204993, on brain function and its relationship with schizophrenia trait has not been investigated. Imaging genetics may provide a neural basis to explain how the gene affects the occurrence and development of diseases via connecting the link between gene and brain diseases [17]. Resting-state functional magnetic resonance imaging (rsfMRI), as a task-free and noninvasive measurement tool, has been used in several mental disorders [18]. Functional connectivity strength (FCS), which identifies functional hubs of human brain networks based on the graph theory method [19], is a developed method to test the connectivity of each voxel with all other voxels in the brain [20]. The FCS analysis has been used to characterize the neural mechanism in various psychiatric disorders [21], including schizophrenia and schizotypal traits [22, 23]. For example, decreased FCS in the bilateral occipital cortex and the right sensorimotor cortex as well as increased FCS in the bilateral temporal cortex, hippocampus, and the left prefrontal cortex were observed in schizophrenia [20]. Moreover, schizotypal traits presented with abnormal connectivity in the default mode network compared with healthy controls[23].

In present study, we aimed to investigate the genetic effect of the schizophrenia-related gene (*NOTCH4* rs204993 genotype) on brain function and its relationship with schizotypal traits among healthy volunteers. Based on the above findings, we propose that the schizophrenia-risk rs204993 genotype (AA genotype) may be linked with attenuated function in schizophrenia-related brain regions, which contributes to schizotypal traits.

Methods

Participants

We recruited 641 eligible healthy Chinese Han participants from the Anhui Medical University in Hefei, Anhui province. The inclusion criteria included: 1) undergraduate students; 2) the participant and whose parents were the Chinese Han populations; 3) age 18-25 years old; 4) Voluntary to be taken blood sample and MRI scanning; 5) no contraindication for MRI scanning. All subjects were accepted using the Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition [24], to exclude any psychiatric disorders. Other exclusion criteria included: neurological diseases, alcohol or drug abuse, brain injury, visible brain structure abnormalities, or excessive motion during MRI scanning (> 2.0 mm, 2.0°). Participants with a first-degree relative with a neurological or psychiatric disorder were also excluded. Of

these participants, a total of 239 participants were excluded because of *NOTCH4* genotyping failure (12 participants) or poor MRI data (198 participants with excessive head motion and 29 participants with artifact in their structural images). Finally, a total of 402 participants were included in the data analysis of the current study. The study was approved by the Anhui Medical University Ethics Committee, and all participants signed informed consent forms after being given a thorough description of the study.

SNP genotyping

Blood samples were collected to obtain DNA using standard procedures. The genotyping of all participants was determined in comparison with control DNA confirmed by sequencing in the SNP pattern. The *NOTCH4* rs204993 was then genotyped by the SNP scan technique (Genesky Biotechnologies Inc., Shanghai, China) according to a previous study [25]. Genotype screening for rs204993 was conducted through the small amplicon genotyping method. The PCR primers were designed to flank the one base pair just adjacent to the rs204993: 5'-TTCGGGACTTCTGTTCAGCC-3'(forward),5'-TCCTGGAAGCACTCGTTGAC-3'(reverse). The primer was designed with the LightScanner Primer Design software program (Idaho Technology, UT, USA). DNA amplification was performed with a 96-well plate at a 10- μ L final volume containing 4 μ L of 2.5 \times high-sensitivity genotyping master mix (Idaho Technology), 1 μ M of each primer, and 20 ng of genomic DNA. Through DNA electrophoresis, we repeated the experiment three times and found positive bands. Finally, the genotype of *NOTCH4* rs204993 was divided into the AA, GA, and GG genotypes.

Schizotypal trait evaluations

The schizotypal traits were evaluated using the schizotypal personality questionnaire (SPQ) [26]. The SPQ includes a 74-item with a "yes/no" response to assess schizotypal traits in the general population. The questionnaire subscale scores reflect different schizotypal dimensional features, including interpersonal, cognitive perceptual, and disorganized factors [27]. In the present study, we used the Chinese version SPQ, which has good internal consistency (0.95) and reliability value (0.86) for the Chinese population [28].

Neuroimaging data acquisition

We acquired rsfMRI images of participants at the University of Science and Technology of China, Hefei, Anhui Province. During scanning, all participants were instructed to keep their eyes closed without moving their body, and not to think of anything in particular. Functional images were obtained with a 3.0 T MRI scanner (Signa HDxt 3.0 T; GE Healthcare, Buckinghamshire, UK) composed of 217 echo-planar imaging volumes with the following parameters: TR = 2,400 ms; TE = 30 ms; flip angle = 90°; matrix size = 64 \times 64; field of view = 192 \times 192 mm²; slice thickness = 3 mm; and 46 continuous slices (voxel size = 3 \times 3 \times 3 mm³). T1-weighted anatomic images with 188 slices were also acquired in sagittal orientation (TR = 8.16 ms; TE = 3.18 ms; flip angle = 12°; field of view = 256 \times 256 mm²; slice thickness = 1 mm; and voxel size = 1 \times 1 \times 1 mm³).

Functional image preprocessing

The Data Processing Assistant for Resting-State Functional MRI imaging toolkit (DPARSF) was used for the functional image preprocessing, which was based on the following software: the Resting State Functional MR Imaging Toolkit and statistical parametric mapping software package (SPM8) [29, 30].

We discarded the first five volumes of data to ensure stable longitudinal magnetization, and the remaining volumes were corrected for slice timing and head motion. Next, we normalized the structural T1 image to the Montreal Neurological Institute space based on the T1 image unified segmentation with a 12 parameter nonlinear transformation. Finally, the data were nuisance regressed with 24 Friston motion parameters, white matter high signals, cerebrospinal fluid signals, and global signals as regressors, and filtered with a temporal band-pass of 0.01–0.1 Hz.

Whole brain FCS analysis

A voxel-wise, whole-brain functional connectivity analysis was performed on the preprocessed rsfMRI data as in our previous study [31]. First, Pearson's correlations between the residual time series of all pairs of brain voxels were computed and a whole brain connectivity matrix for each participant was constructed. The individual correlation matrices were then transformed to a z-score matrix using a Fisher r-to-z transformation. For a given voxel, FCS was computed as the sum of the z-values between the given voxel and all other voxels. We restricted our analyses to correlations above a threshold of $r = 0.1$ to eliminate weak correlations possibly arising from noises. These FCS maps were smoothed with a 4 mm full-width at a half-maximum (FWHM) Gaussian kernel [19, 32].

Statistical analysis

The χ^2 test was used to compare the sex differences, and one-way analysis of variance (ANOVA) was used to check for differences in age, educational years and SPQ among the three genotype groups (AA, AG and GG). Although there is study has suggested a recessive effect (AA) of rs204993, there is no consensus about analytically grouping the genotypes. Hence, we performed with an unbiased approach that of three possible effects were all considered: additive (GG versus GA versus AA genotypes), dominant (A allele carriers versus GG); recessive (AA versus G allele carriers). For additive effect, one-way ANOVA was performed to determine the changed regions of FCS for healthy participants among the three genotype groups. For dominant or recessive effect, two sample t-tests were performed (A allele carriers versus GG, AA versus G allele carriers). These tests were constrained in a gray matter mask, and multiple comparison corrections were based on a Gaussian random field theory (voxel-level $p < 0.001$; cluster-level $p < 0.05$). The FCS value of the changed regions was extracted and compared with one-way ANOVA among the three genotype groups, and post hoc analysis was conducted using the LSD method. Spearman's correlation analyses were performed to assess the correlation between the values of FCS and the behavior scores (total SPQ scores and its sub-domains) for each *NOTCH4* genotype group (the significance level: $p < 0.05/5$, two-tail).

Results

Demographic information and the genetic effect on behavior performance

Table 1 shows the demographic, genetic, and behavioral data for participants included in the final analysis. The genotype distribution of *NOTCH4* rs204993 in the current study (AA = 142, AG = 188, and GG = 72) was consistent with a previous report of variations of this gene in the healthy Chinese Han population [16]. Genotype groups did not significantly deviate from Hardy-Weinberg equilibrium ($\chi^2 = 0.505$, $p = 0.477$). There was no significant difference in sex, educational years, and age between the genotype groups. There was also no significant difference of SPQ between the genotype groups.

The genetic effect of rs204993 on FCS

After controlling for demographic factors, only the additive effect (GG versus GA versus AA genotypes) was survived after corrected using Gaussian random field theory (voxel-level $p < 0.001$; cluster-level $p < 0.05$). The results showed that FCS was significantly abnormal in the left occipital cortex among the *NOTCH4* genotypes (Figure 1). The FCS values of the above cluster were then extracted in the three genotype groups. We found a main effect of the genotype on FCS values ($F = 15.354$, $p < 0.001$). Post hoc analysis showed AA individuals having reduced values when compared with AG and GG participants ($p < 0.001$ for both) and AG individuals having reduced values when compared with GG participants ($p = 0.05$) (Figure 1 and Table 1). No cluster was survived after GRF multiple comparison corrections based dominant or recessive statistical model.

Correlation analyses between SPQ and FCS

The FCS values in the left occipital cortex were negatively associated with the total SPQ scores ($r = -0.391$, $p = 0.001$), cognitive-perceptual scores ($r = -0.411$, $p < 0.001$), interpersonal scores ($r = -0.342$, $p = 0.003$) and disorganized scores ($r = -0.233$, $p = 0.049$) among GG participants (Figure 2). No significant relationship was found between the values of FCS and the behavior scores (total SPQ scores and its sub-domains) for the AA or AG group (Figure 3).

Discussion

In the current study, we aimed to investigate the genetic effect of the schizophrenia-related gene (*NOTCH4* rs204993 genotype) on brain function and its relationship with schizotypal traits among healthy volunteers. Using a large sample size, we found that functional connectivity in the occipital cortex was the highest in GG individuals and decreased from GG and AG individuals to AA individuals. Notably, the values of FCS in the left occipital cortex were negatively correlated with total SPQ scores and its subset scores (cognitive-perceptual and interpersonal scores) in GG individuals.

Previous studies have reported that *NOTCH4* polymorphism of rs204993 is closely related to schizophrenia [11, 12]. However, as a candidate gene for schizophrenia, we did not find a direct effect of

NOTCH4 rs204993 on the schizotypal traits among healthy volunteers. Even so, it was notable that we found the modulatory effect of *NOTCH4* rs204993 on brain function (the functional connectivity of occipital cortex), which was associated with schizotypal traits. The impact of the *NOTCH4* gene on the central nervous system has been consistently reported [33]. The *NOTCH4* gene is one member of the *NOTCH* family, which may code for the notch protein, which was identified as a *Drosophila* neurogenic protein during embryogenesis [34]. Functionally, the *NOTCH* family mainly controls the cell fate in the process of neurodevelopment, which promotes proliferative signaling [35, 36]. Therefore, the *NOTCH4* gene variants may potentially affect the function or expression level of the Notch4 protein, which in turn influences the neurodevelopment of certain psychiatric disorders, such as schizophrenia and schizotypal traits. In the present study, we found an aberrant functional connectivity in the left occipital cortex among carriers with the *NOTCH4* rs204993 schizophrenia-risk genotypes.

The occipital cortex plays an important role in the neural circuits of perceptual processing, such as visual and auditory processing [37, 38]. Considerable studies have confirmed that schizophrenia patients and their unaffected siblings present with impaired perceptual performance, which is associated with impaired structural and functional connectivity in the occipital cortex [39-41]. Furthermore, the occipital lobe has also been gradually recognized as the neural basis for perceiving personal communication and social interactions [42-44]. Of course, some studies have shown that negative symptoms in patients with schizophrenia are associated with abnormal changes in the occipital cortex [45-47]. For example, abnormal connectivity between the occipital cortex and ventral tegmental area is related to the negative symptoms in schizophrenia [45]. In general, the occipital cortex may be a common neural pathway for the processing of positive and negative symptoms in schizophrenia. Consistent with this possibility, our study showed that the occipital-cortex functional connectivity was related with cognitive-perceptual and interpersonal subscores, which represented the positive and negative symptoms in schizotypal traits. We therefore speculate that the *NOTCH4* gene may induce the occurrence and development of diseases by affecting the neurodevelopment of occipital cortex.

It is worth noting that we found that the FCS of the occipital cortex was degressive in sequences of the GG, AG, and AA genotype groups that suggest an additive effect of rs204993. This is not in complete agreement with a recessive effect revealed by the previous results that the AA genotype of rs204993 is associated with a higher risk for schizophrenia in the Chinese Han population [16]. The difference between schizophrenia and schizotypal traits may explain this discrepancy. As our results suggested, individuals with the AG genotype also presented lower FCS in the occipital cortex, when compared with the GG genotype carriers, which may indicate the risk of schizotypal traits in the AG genotype carriers, but not the risk of schizophrenia.

It is interesting that the correlation between occipital-cortex function and SPQ is restricted to specific genotype. Detailedly, individual occipital-cortex FCS was inversely correlated with individual SPQ and its sub-domains (cognitive-perceptual and interpersonal scores) only for GG carriers who may present lower risk for schizophrenia. These results implicate a modulation of occipital-cortex function with cognitive-perceptual and interpersonal ability that the individuals with higher occipital-cortex FCS present more

normal cognitive-perceptual and interpersonal ability, which are featured impairments in schizophrenia. The disappearance of such correlation may suggest an impaired modulation of occipital-cortex function with cognitive-perceptual and interpersonal ability in among relative high-risk individuals (AA and AG carriers). Similarly, this genotype-dependent correlation has been reported by previous genetic imaging study [48]. Another interesting finding of the present study was that the significant difference among groups was only located at the left occipital cortex and not the right. We speculate that the left-hemisphere dominant is the potential factor underlying this lateralization. Although we did not collect the handedness data, based on a survey reported about 87.7% - 94.8% right hand domination in Chinese population [49], we believe that most participants in present study are left-hemisphere dominant. The left occipital cortex may prefer to involve in mental and emotional processing, hence, left occipital-cortex connective decline may more contribute to schizophrenia symptoms.

Several potential limitations of our study need to be considered. First, we did not find a direct modulatory effect of the *NOTCH4* gene on schizotypal traits (no significant difference between *NOTCH4* genotype groups). The gross feature of the schizotypal-trait assessment (SPQ) may partly explain this defect. Second, the absence of long scanning durations reduced the reliability of measuring functional connectivity. Third, our study did not reveal the cognitive mental mechanism of *NOTCH4* gene modulating the occipital cortex and its relationship with schizotypal traits. Fourth, previous studies confirmed the relationship between schizophrenia and *NOTCH4* polymorphisms, including single nucleotide polymorphisms (SNPs) (e.g., rs520692, rs3131296, rs204993, and rs2071287). But we only explored the relationship between FCS of left occipital cortex and the *NOTCH4* rs204993 genotype in populations with schizotypal traits. Further studies are needed to analyze the associations with other SNPs of the *NOTCH4* gene. Finally, we did not characterize the modulatory effect of the *NOTCH4* gene on local and long-range functional connectivities, respectively, which may represent distinct biological differences.

Conclusion

Our results suggested that the common variant in the *NOTCH4* gene, rs204993, modulates the function of the occipital cortex, which may contribute to schizotypal traits. These findings provide insight for genetic effects on schizotypal traits and its potential neural substrate.

Declarations

Acknowledgements

We thank all the participants from Anhui Medical University and operators of the Center for Biomedical Engineering, University of Science and Technology of China for their support of this study.

Funding

This study was supported by the National Natural Science Foundation of China (91432301, 31571149, 81171273, 31970979, 91232717 to K.W., 81671354 and 91732303 to Y.T., and 31800909 to L.Z.). The funding agency had no role in the study design, data collection and analysis, or preparation of the manuscript.

Availability of data and materials

Data in the manuscript will not be shared. This was inconsistent with the written informed consent of the participants.

Author's Contributors

XHX, MDZ, LZ, and KW designed the study. LZ, TJB, WLH, and PPH acquired the behavioral and imaging data. LW and GJJ collected and analyzed the genetic data. TJB, GJJ, BSQ, and YHT analyzed the imaging data. XHX, MDZ, LZ, and KW wrote this article, which all authors reviewed. All authors approved the final version to be published, and we certify that no other individuals not listed as authors have made substantial contributions to the paper.

Competing interests

All authors declare no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

All participants signed the informed consent. The study adhered to the tenets of the Declaration of Helsinki and was approved by the Biomedical Ethics Committee of Anhui Medical University.

Author's information

¹Department of Neurology, The First Affiliated Hospital of Anhui Medical University, Hefei, 230022, China; ²Collaborative Innovation Center for Neuropsychiatric Disorders and Mental Health, Anhui Medical University, Hefei, 230022, China; ³Anhui Province Key Laboratory of Cognition and Neuropsychiatric Disorders, Hefei, 230022, China; ⁴Department of Medical Psychology, Chaohu Clinical Medical College, Anhui Medical University, Hefei, China; ⁵Center for Biomedical Engineering, University of Science and Technology of China, Hefei, Anhui, China.

Abbreviations

FCS, functional connectivity strength; SPQ, Schizotypal Personality Questionnaire; rs-fMRI, resting-state functional magnetic resonance image; *NOTCH4*, neurogenic locus notch homolog protein4; MHC, major

histocompatibility complex; SNPs, single nucleotide polymorphisms; FCS, functional connectivity strength; DPARSF, Data Processing Assistant for Resting-State Functional; SPM8, statistical parametric mapping software package; REST, Resting State Functional MR Imaging Toolkit; FWHM, full-width at half-maximum.

References

1. Andreasen NC. Symptoms, signs, and diagnosis of schizophrenia. *Lancet*. 1995; 346(8973):477-481.
2. Rado S. Dynamics and classification of disordered behavior. *Am J Psychiatry*. 1953;110(6):406-416.
3. Gottesman, II, Gould TD. The endophenotype concept in psychiatry: etymology and strategic intentions. *Am J Psychiatry*. 2003;160(4):636-645.
4. Raine A. Schizotypal personality: neurodevelopmental and psychosocial trajectories. *Annu Rev Clin Psychol*. 2006;2:291-326.
5. Vollema MG, Postma B. Neurocognitive correlates of schizotypy in first degree relatives of schizophrenia patients. *Schizophr Bull*. 2002;28(3):367-377.
6. Baker KD, Skuse DH. Adolescents and young adults with 22q11 deletion syndrome: psychopathology in an at-risk group. *Br J Psychiatry*. 2005;186:115-120.
7. Stefanis NC, Van Os J, Avramopoulos D, Smyrnis N, Evdokimidis I, Hantoumi I, Stefanis CN. Variation in catechol-o-methyltransferase val158 met genotype associated with schizotypy but not cognition: a population study in 543 young men. *Biol Psychiatry*. 2004;56(7):510-515.
8. Ikeda M, Aleksic B, Kinoshita Y, Okochi T, Kawashima K, Kushima I, Ito Y, Nakamura Y, Kishi T, Okumura T et al. Genome-wide association study of schizophrenia in a Japanese population. *Biol Psychiatry*. 2011;69(5):472-478.
9. Ripke S, Sanders AR, Kendler KS, Levinson DF, Sklar P, Holmans PA, Lin DY, Duan J, Ophoff RA, Andreassen OA et al. Genome-wide association study identifies five new schizophrenia loci. *Nat Genet*. 2011;43(10):969-976.
10. Aberg KA, Liu Y, Bukszar J, McClay JL, Khachane AN, Andreassen OA, Blackwood D, Corvin A, Djurovic S, Gurling H et al. A comprehensive family-based replication study of schizophrenia genes. *JAMA psychiatry*. 2013;70(6):573-581.
11. Luo XG, Klempan TA, Lappalainen J, Rosenheck RA, Charney DS, Erdos J, van Kammen DP, Kranzler HR, Kennedy JL, Gelernter J. NOTCH4 gene haplotype is associated with schizophrenia in African Americans. *Biol Psychiatry*. 2004; 55(2):112-117.
12. Wei J, Hemmings GP. The NOTCH4 locus is associated with susceptibility to schizophrenia. *Nat Genet*. 2000;25(4):376-377.
13. Ikeda M, Aleksic B, Yamada K, Iwayama-Shigeno Y, Matsuo K, Numata S, Watanabe Y, Ohnuma T, Kaneko T, Fukuo Y et al. Genetic evidence for association between NOTCH4 and schizophrenia supported by a GWAS follow-up study in a Japanese population. *Molecular psychiatry*. 2013;18(6):636-638.

14. Tochigi M, Zhang X, Ohashi J, Hibino H, Otowa T, Rogers M, Kato T, Okazaki Y, Kato N, Tokunaga K. Association study between the TNXB locus and schizophrenia in a Japanese population. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*. 2007;144(3):305-309.
15. Liu CM, Liu YL, Fann CSJ, Chenk WJ, Yang WC, Ouyang WC, Chen CY, Jou YS, Hsieh MH, Liu SK et al. Association evidence of schizophrenia with distal genomic region of NOTCH4 in Taiwanese families. *Genes Brain Behav*. 2007;6(6):497-502.
16. Zhang B, Fan QR, Li WH, Lu N, Fu DK, Kang YJ, Wang N, Li T, Wen XP, Li DX. Association of the NOTCH4 Gene Polymorphism rs204993 with Schizophrenia in the Chinese Han Population. *Biomed Res Int*. 2015;2015:408096.
17. Chen J, Xu Y, Zhang J, Liu ZF, Xu C, Zhang KR, Shen Y, Xu Q. A combined study of genetic association and brain imaging on the DAOA gene in schizophrenia. *Am J Med Genet B*. 2013;162B(2):191-200.
18. Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proc Natl Acad Sci U S A*. 2001;98(2):676-682.
19. Buckner RL, Sepulcre J, Talukdar T, Krienen FM, Liu HS, Hedden T, Andrews-Hanna JR, Sperling RA, Johnson KA. Cortical Hubs Revealed by Intrinsic Functional Connectivity: Mapping, Assessment of Stability, and Relation to Alzheimer's Disease. *Journal Of Neuroscience*. 2009;29(6):1860-1873.
20. Zhu JJ, Zhuo CJ, Xu LX, Liu F, Qin W, Yu CS. Altered Coupling Between Resting-State Cerebral Blood Flow and Functional Connectivity in Schizophrenia. *Schizophr Bull*. 2017;43(6):1363-1374.
21. Wang L, Xia MR, Li K, Zeng YW, Su YN, Dai WJ, Zhang QG, Jin Z, Mitchell PB, Yu X et al. The Effects of Antidepressant Treatment on Resting-State Functional Brain Networks in Patients With Major Depressive Disorder. *Hum Brain Mapp*. 2015; 36(2):768-778.
22. Chen X, Ji GJ, Zhu C, Bai X, Wang L, He K, Gao Y, Tao L, Yu F, Tian Y et al. Neural Correlates of Auditory Verbal Hallucinations in Schizophrenia and the Therapeutic Response to Theta-Burst Transcranial Magnetic Stimulation. *Schizophr Bull*. 2019;45(2):474-483.
23. Zhang Q, Shen J, Wu JL, Yu X, Lou WT, Fan HY, Shi L, Wang DF. Altered default mode network functional connectivity in schizotypal personality disorder. *Schizophrenia research*. 2014;160(1-3):51-56.
24. Angell M. *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition. *Alzheimer Disease & Associated Disorders*. 1996;10(2):20-22.
25. Zhang B, Li DX, Lu N, Fan QR, Li WH, Feng ZF. Lack of Association between the TSPAN18 Gene and Schizophrenia Based on New Data from Han Chinese and a Meta-Analysis. *Int J Mol Sci*. 2015;16(6):11864-11872.
26. Raine A. The SPQ: a scale for the assessment of schizotypal personality based on DSM-III-R criteria. *Schizophr Bull*. 1991;17(4):555-564.
27. Raine A, Reynolds C, Lencz T, Scerbo A, Triphon N, Kim D. Cognitive-perceptual, interpersonal, and disorganized features of schizotypal personality. *Schizophr Bull*. 1994;20(1):191-201.

28. Chen WJ, Hsiao CK, Lin CC. Schizotypy in community samples: the three-factor structure and correlation with sustained attention. *J Abnorm Psychol.* 1997;106(4):649-654.
29. Chao-Gan Y, Yu-Feng Z. DPARSF: A MATLAB Toolbox for "Pipeline" Data Analysis of Resting-State fMRI. *Front Syst Neurosci.* 2010;4:13.
30. Song XW, Dong ZY, Long XY, Li SF, Zuo XN, Zhu CZ, He Y, Yan CG, Zang YF. REST: a toolkit for resting-state functional magnetic resonance imaging data processing. *PloS one.* 2011;6(9):e25031.
31. Wei Q, Bai T, Chen Y, Ji G, Hu X, Xie W, Xiong Z, Zhu D, Wei L, Hu P et al. The Changes of Functional Connectivity Strength in Electroconvulsive Therapy for Depression: A Longitudinal Study. *Front Neurosci.* 2018;12:661.
32. Liang X, Zou Q, He Y, Yang Y. Coupling of functional connectivity and regional cerebral blood flow reveals a physiological basis for network hubs of the human brain. *Proc Natl Acad Sci U S A.* 2013;110(5):1929-1934.
33. Justice NJ, Jan YN. Variations on the Notch pathway in neural development. *Curr Opin Neurobiol.* 2002;12(1):64-70.
34. Fortini ME, Artavanis-Tsakonas S. Notch: neurogenesis is only part of the picture. *Cell.* 1993;75(7):1245-1247.
35. Fiuza UM, Arias AM. Cell and molecular biology of Notch. *J Endocrinol.* 2007;194(3):459-474.
36. Sestan N, Artavanis-Tsakonas S, Rakic P. Contact-dependent inhibition of cortical neurite growth mediated by notch signaling. *Science (New York, NY).* 1999; 286(5440):741-746.
37. Benetti S, Novello L, Maffei C, Rabini G, Jovicich J, Collignon O. White matter connectivity between occipital and temporal regions involved in face and voice processing in hearing and early deaf individuals. *NeuroImage.* 2018;179:263-274.
38. Cappadocia DC, Monaco S, Chen Y, Blohm G, Crawford JD. Temporal Evolution of Target Representation, Movement Direction Planning, and Reach Execution in Occipital-Parietal-Frontal Cortex: An fMRI Study. *Cereb Cortex.* 2017;27(11):5242-5260.
39. Oestreich LKL, McCarthy-Jones S, Whitford TJ, Bank ASR. Decreased integrity of the fronto-temporal fibers of the left inferior occipito-frontal fasciculus associated with auditory verbal hallucinations in schizophrenia. *Brain imaging and behavior.* 2016;10(2):445-454.
40. van de Ven V, Jagiela AR, Oertel-Knochel V, Linden DEJ. Reduced intrinsic visual cortical connectivity is associated with impaired perceptual closure in schizophrenia. *Neuroimage-Clin.* 2017;15:45-52.
41. Zalesky A, Pantelis C, Cropley V, Fornito A, Cocchi L, McAdams H, Clasen L, Greenstein D, Rapoport JL, Gogtay N. Delayed Development of Brain Connectivity in Adolescents With Schizophrenia and Their Unaffected Siblings. *JAMA psychiatry.* 2015;72(9):900-908.
42. Che XW, Wei DT, Li WF, Li HJ, Qiao L, Qiu J, Zhang QL, Liu YJ. The correlation between gray matter volume and perceived social support: A voxel-based morphometry study. *Soc Neurosci-Uk.* 2014;9(2):152-159.

43. Lingnau A, Downing PE. The lateral occipitotemporal cortex in action. *Trends Cogn Sci.* 2015;19(5):268-277.
44. Quadflieg S, Gentile F, Rössion B. The neural basis of perceiving person interactions. *Cortex.* 2015;70:5-20.
45. Giordano GM, Stanziano M, Papa M, Mucci A, Prinster A, Soricelli A, Galderisi S. Functional connectivity of the ventral tegmental area and avolition in subjects with schizophrenia: a resting state functional MRI study. *Eur Neuropsychopharm.* 2018;28(5):589-602.
46. Rigucci S, Rossi-Espagnet C, Ferracuti S, De Carolis A, Corigliano V, Carducci F, Mancinelli I, Cicone F, Tatarelli R, Bozzao A et al. Anatomical substrates of cognitive and clinical dimensions in first episode schizophrenia. *Acta Psychiatr Scand.* 2013;128(4):261-270.
47. Taylor SF, Chen AC, Tso IF, Liberzon I, Welsh RC. Social appraisal in chronic psychosis: Role of medial frontal and occipital networks. *Journal of psychiatric research.* 2011;45(4):526-538.
48. Liu B, Zhang XL, Hou B, Li J, Qiu CX, Qin W, Yu CS, Jiang TZ. The Impact of MIR137 on Dorsolateral Prefrontal-Hippocampal Functional Connectivity in Healthy Subjects. *Neuropsychopharmacol.* 2014;39(9):2153-2160.
49. Xin-tian L. The distribution of left and right handedness in Chinese people. *Acta Psychologica Sinica.* 1983;03:268-276.

Table

Table 1 Demographic and behavioral-neuroimaging information for participants grouped by rs204993 genotype (mean \pm SD).

	AA	AG	GG	Statistics	<i>P</i> value
Number of participants	142	188	72	/	/
Gender (male/female)	68/74	88/100	41/31	2.250	0.325
Age (years)	20.761 ± 1.104	20.942 ± 1.061	21.097 ± 1.115	2.490	0.084
SPQ	26.930 ± 13.853	26.553 ± 14.038	31.167 ± 14.470	2.955	0.051
FCS in occipital cortex	0.379 ± 0.461	0.612 ± 0.531	0.755 ± 0.542	15.354	<0.001

Note: SPQ, schizotypal personality questionnaire; FCS, functional connectivity strength; SD, standard deviation.

Figures




Figure 1

Modulatory effect of rs204993 on the functional strength (FCS). The X-axis shows the different genotypes, and the Y-axis shows the mean \pm SD of the FCS in the occipital cortex. Group comparisons revealed that the FCS in the occipital cortex was the highest in GG individuals and decreased from GG and AG individuals to AA individuals. All thresholds for comparisons were set as a whole brain Gaussian random field (GRF) correction (voxel $P < 0.001$, cluster $P < 0.05$).




Figure 2

Relationship between the functional strength (FCS) in the occipital cortex and schizotypal traits among the GG genotype carriers. The FCS in the occipital cortex was negatively correlated with the score of schizotypal personality questionnaire (SPQ) and its subscale scores (cognitive perceptual and interpersonal) among the GG genotype carriers). Note: *** $p < 0.001$, * $p < 0.05$.




Figure 3

Relationship between the functional strength (FCS) in the occipital cortex and schizotypal traits among the AA and AG genotype carriers. No significant correlation was found between the FCS in the occipital cortex and the schizotypal personality questionnaire scores or its subscale scores.