

Clinical impact of severe chronic kidney disease in patients with non-HIV pulmonary cryptococcosis

Hiroki Tashiro (✉ si3222@cc.saga-u.ac.jp)

Saga Daigaku Igakubu <https://orcid.org/0000-0001-9699-414X>

Tetsuro Haraguchi

Saga Daigaku Igakubu

Koichiro Takahashi

Saga Daigaku Igakubu

Hironori Sadamatsu

Saga Daigaku Igakubu

Ryo Tajiri

Saga Daigaku Igakubu

Ayako Takamori

Saga Daigaku Igakubu

Shinya Kimura

Saga Daigaku Igakubu

Naoko Aragane

Saga Daigaku Igakubu

Research article

Keywords: Pulmonary cryptococcosis, Chronic kidney disease, Clinical manifestations, Dissemination

Posted Date: March 3rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-15876/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on April 29th, 2020. See the published version at <https://doi.org/10.1186/s12890-020-1149-3>.

Abstract

Background

Pulmonary cryptococcosis is an uncommon infectious disease that can develop in both immunocompromised and immunocompetent patients. Severity of chronic kidney disease (CKD) was reported to be one of the risk factors for pulmonary cryptococcosis, but its clinical characteristics have not been fully assessed. The purpose of this study is to clarify clinical characteristics of severe CKD in patients with pulmonary cryptococcosis.

Methods

The present study retrospectively investigated 56 patients who had pulmonary cryptococcosis with non-human immunodeficiency virus (HIV) infection and were treated at Saga University Hospital between 2005 and 2018. The clinical characteristics were evaluated and compared between patients with estimated glomerular filtration rate (eGFR) >45 mL/min/1.73m² (n = 42) (preserved eGFR) and those with eGFR <45 mL/min/1.73m² (n = 14) (declined eGFR).

Results

Compared with patients with preserved eGFR, those with declined eGFR had significantly higher rate of disseminated cryptococcosis (p = 0.03); higher percentage of patients who did not recover after treatment (p = 0.02); and more frequent clinical features of fever (p <0.01), pleural effusion (p = 0.03), high white blood cell count (p <0.01) and C-reactive protein (p <0.01), and low level of albumin (p <0.01). Multivariate analysis adjusted by immunosuppressive drug use indicated significant differences of fever (p <0.01), high white blood cell count (p = 0.03), C-reactive protein (p = 0.01) and low level of albumin (p <0.01) in patients with eGFR <45 mL/min/1.73m².

Conclusion

Severe CKD is associated with poor clinical characteristics and outcomes in patients with non-HIV pulmonary cryptococcosis.

Background

Cryptococcosis is an uncommon infectious disease caused by *Cryptococcus neoformans* and *Cryptococcus gattii* (1, 2). The fungus mainly infects the lungs and central nervous system, but it can less frequently affect other organs, including the eyes, prostate, skin, and bone, in both immunocompetent and immunocompromised patients (3–5). Occasionally, cryptococcus can disseminate to several organs and lead to mortality, depending on the host immunity (6–8).

Pulmonary cryptococcosis is important, because the respiratory tract is the most common portal of entry (9, 10). Several studies have reported that the clinical manifestations, including symptoms, laboratory

data, radiologic findings, dissemination, and outcome, of pulmonary cryptococcosis were different in patients with immunocompromised comorbidities, such as human immunodeficiency virus (HIV) infection, diabetes mellitus, malignancy, organ transplantation, immunosuppressive treatment use, and chronic kidney disease (CKD) (5, 11–14). CKD is one of the essential health conditions and is defined by an estimated glomerular filtration rate (eGFR) of $< 60 \text{ mL/min/1.73 m}^2$ with positive findings of urinary protein that persists for 3 months (15, 16). A large cohort study identified that an eGFR $< 45 \text{ mL/min/1.73 m}^2$ was related with all-cause and cardiovascular mortality (17). There is increasing evidence that CKD is closely linked with the host immune systems (18) and the risk for infections (19). Some reports identified that severity of CKD was involved with the clinical characteristics of pulmonary cryptococcosis (20, 21); however the clinical features have not been fully assessed.

The purpose of this study is to clarify clinical characteristics of severe CKD in patients with pulmonary cryptococcosis. We investigated 56 patients with non-HIV pulmonary cryptococcosis and compared the clinical manifestations between 42 patients with $\text{eGFR} > 45 \text{ mL/min/1.73 m}^2$ and 14 patients with $\text{eGFR} < 45 \text{ mL/min/1.73 m}^2$ with positive findings of urinary protein. We identified that the symptoms, laboratory findings, radiologic patterns, and outcome, including dissemination, were different between the 2 groups and were worse in patients with pulmonary cryptococcosis and severe CKD. These results may practically contribute to the clinical risk evaluation for pulmonary cryptococcosis.

Methods

Patients and setting

We retrospectively evaluated 56 patients diagnosed as pulmonary cryptococcosis without HIV infection at Saga University Hospital between 2005 and 2018. This study was approved by the ethics committee of Saga University Hospital (approval number: 2019-09-06, approval date: Nov 25, 2019) and was performed in accordance with the 1964 Declaration of Helsinki.

The assessment of pulmonary cryptococcosis was made by 1) clinical diagnosis, based on serum anticryptococcal antigen and chest radiologic abnormalities; 2) histologic diagnosis, based on the presence of cryptococcal pathogen on Grocott's staining of lung histologic samples and chest radiologic abnormalities; and 3) pathogenic diagnosis, based on the detection of cryptococcus species on airway, blood, or cerebrospinal fluid culture and chest radiologic abnormalities. All of the species detected from the cultured samples were *C. neoformans*. Disseminated cryptococcosis was defined as isolation of *C. neoformans* from blood, sterile body fluid, or any extrapulmonary site (8). Upon the diagnosis of pulmonary cryptococcosis, the clinical manifestations, including comorbidities, symptoms, laboratory data, and radiologic findings, were collected from the medical records and eGFR was calculated by formula as followed; $194 \times \text{Serum creatinine}^{-1.094} \times \text{Age}^{-0.287} \times 0.739$ (if female) (22). All patients with $\text{eGFR} < 45 \text{ mL/min/1.73 m}^2$ were verified for the persistence of the same renal function for more than 3 months with positive findings of urinary protein, according to the CKD guidelines (15, 17).

To address the clinical impact of severe CKD on pulmonary cryptococcosis, we divided 56 patients with pulmonary cryptococcosis into those with eGFR > 45 mL/min/1.73 m² (preserved eGFR) and those with eGFR < 45 mL/min/1.73 m² and positive findings of urinary protein (declined eGFR). The cutoff of 45 mL/min/1.73 m² for eGFR was set, based on a previous report that identified a relationship between mortality and eGFR < 45 mL/min/1.73 m² (17). Treatment, including the intake of antifungal drugs or surgery, was based on the physician's judgment. The outcome of non-recovery from pulmonary cryptococcosis after antifungal drug treatment for 3 months was defined as death or worsening radiologic and clinical findings.

Comorbidities

The predisposing factors, including pulmonary diseases, diabetes mellitus, connective tissue diseases, malignancy, immunosuppressive drug use, and hemodialysis, were collected from the medical records. There were no patients who received organ transplantation.

Radiologic Findings

In the present study, high resolution computed tomography (HRCT) of the chest upon the diagnosis of pulmonary cryptococcosis was evaluated by 2 pulmonologists, who referred to the official reports from a radiologist. The most dominant chest radiologic finding was classified based on the following patterns: (1) single nodule (< 30 mm), (2) multiple nodules, (3) mass (> 30 mm), (4) cavitation, (5) consolidation, (6) ground glass attenuation, or (7) pleural effusion.

Statistical analysis

The clinical data were analyzed by Student's t-test for continuous variables or chi-square test for categorical variables. The present study found that immunosuppressive drug was used more often by patients with declined eGFR than by those with preserved eGFR. Because this might affect the current results, we performed multivariate analysis for the variables, including fever, white blood cell count, albumin, and C-reactive protein (CRP), which were significantly different between patients with preserved eGFR and those with declined eGFR in the univariate analysis (data not shown in tables). Disseminated cryptococcosis, pleural effusion, and recovery after treatment were not assessed because of the small sample size. The comparative results were adjusted by the predisposing variable of immunosuppressive drug use. For multivariate analyses, logistic regression analysis for categorical variables and multiple linear regression analysis for continuous variables were performed. Quantitative data were presented as mean ± standard deviation (SD), and odds ratio or coefficient β value were calculated. The two-tailed significance level was set at $p < 0.05$. Statistical analysis was performed with JMP Pro version 14.2.0 software (SAS Institute Inc., Cary, NC, USA).

Results

Clinical characteristics and comorbidities in patients with pulmonary cryptococcosis

There were 42 patients with preserved eGFR and 14 patients with declined eGFR. Compared with the group with preserved eGFR, the group with declined eGFR had similar age, sex, and body mass index; tended to have higher number of patients with pathogenic diagnosis; had significantly higher rate of disseminated pulmonary cryptococcosis; similar comorbidities of pulmonary diseases, diabetes mellitus, connective tissue diseases, and malignancy; and more frequently used immunosuppressive drugs. In the present study, only 1 patient in the declined eGFR group was on hemodialysis (Table 1).

Table 1
Epidemiologic and clinical characteristics of patients pulmonary nocardiosis, according to eGFR (n = 56)

	eGFR > 45	eGFR ≤ 45	p value
n	42	14	
Age (years)	67.4 ± 2.5	67.7 ± 2.9	0.95
Gender (Male/Female)	21/21	6/8	0.64
BMI (kg/m ²)	22.2 ± 0.7	22.2 ± 1.7	0.96
Diagnosis			
Clinical	24 (57.1%)	7 (50.0%)	0.64
Histological	10 (23.8%)	2 (14.3%)	0.44
Pathogenic	10 (23.8%)	7 (50.0%)	0.07
Disseminated cryptococcosis			
	1 (7.1%)	3 (21.4%)	0.03
Comorbidity			
Pulmonary diseases	4 (9.5%)	1 (7.1%)	0.78
Diabetes mellitus	12 (28.6%)	6 (42.9%)	0.33
Connective tissue diseases	4 (9.5%)	2 (14.3%)	0.63
Malignancy	12 (28.6%)	2 (14.3%)	0.26
Immunosuppressive drug use	14 (33.3%)	9 (64.3%)	0.04
On hemodialysis	0 (0.0%)	1 (7.1%)	0.09
eGFR; estimated glomerular filtration rate, BMI; body mass index			

Comparison of symptoms and laboratory findings in patients with pulmonary cryptococcosis

Compared with the group with preserved eGFR, the group with declined eGFR had similar percentage of asymptomatic patients; similar pulmonary cryptococcosis-related symptoms of cough, sputum, chest pain, and dyspnea; significantly higher percentage of patients who have fever (> 37.5 °C) (57.1% vs. 19.0%, p < 0.01); and on laboratory data, significantly higher white blood cell count and CRP level but

significantly lower albumin ($p < 0.01$ for all) and similar lymphocyte count, immunoglobulin G, and anticryptococcal antigen (Table 2).

Table 2
Comparison of the symptoms and laboratory data in patients with pulmonary cryptococcosis, according to eGFR

	eGFR > 45	eGFR ≤ 45	p value
Asymptomatic	15 (35.7%)	2 (14.3%)	0.11
Cough	12 (28.6%)	6 (42.9%)	0.32
Sputum	6 (14.3%)	4 (28.6%)	0.24
Chest pain	4 (9.5%)	1 (7.1%)	0.78
Fever (> 37.5 °C)	8 (19.0%)	8 (57.1%)	< 0.01
Dyspnea	2 (2.4%)	1 (7.1%)	0.74
Laboratory data			
White blood cell (/ml)	6928.6 ± 445.1	9885.7 ± 1470.9	< 0.01
Lymphocyte cell (/ml)	1292.0 ± 110.3	1375.2 ± 267.3	0.73
Albumin (g/dl)	3.7 ± 0.1	3.0 ± 0.2	< 0.01
Immunoglobulin G (mg/dl)	1018.5 ± 110.2	1185.6 ± 206.3	0.44
CRP (mg/dl)	1.0 ± 0.3	2.8 ± 0.7	< 0.01
Anti-cryptococcus antigen	90.7 ± 31.8	137.7 ± 67.0	0.51
CRP; C reactive protein, eGFR; estimated glomerular filtration rate			

Comparison of the radiologic findings on HRCT in patients with pulmonary cryptococcosis

According to previous reports on the different distribution and patterns of radiologic findings in pulmonary cryptococcosis, depending on immune status (12, 23), we evaluated and compared the area and features of pulmonary abnormalities between patients with preserved eGFR and those with declined eGFR. Compared with the group with preserved eGFR, the group with declined eGFR had lower number of patients whose pulmonary abnormalities were limited to 1 lobe ($p = 0.06$) and were distributed in only a unilateral lung field ($p = 0.07$); similar patterns of single nodule, multiple nodules, masses, cavitation,

consolidation, and ground glass attenuation; and significantly higher number of patients with pleural effusion ($p = 0.03$) (Table 3).

Table 3
Comparison of the radiologic features on high resolution computed tomography in patients with pulmonary cryptococcosis, according to eGFR

	eGFR > 45	eGFR ≤ 45	p value
Limited in one lobe	24 (57.1%)	4 (28.6%)	0.06
Unilateral lung field	32 (76.2%)	7 (50.0%)	0.07
Single nodule	6 (14.3%)	1 (7.1%)	0.46
Multiple nodules	27 (64.3%)	10 (33.3%)	0.62
Mass	3 (7.1%)	1 (7.1%)	1
Cavitation	6 (14.3%)	1 (7.1%)	0.46
Consolidation	8 (19.0%)	3 (21.4%)	0.85
Ground glass attenuation	0 (0.0%)	1 (7.1%)	0.09
Pleural effusion	1 (7.1%)	3 (21.4%)	0.03
eGFR; estimated glomerular filtration rate			

Treatment and outcome of patients with pulmonary cryptococcosis

The number of patients who took antifungal drugs and the duration of antifungal drug treatment were not different between the 2 groups. Azole was the antifungal drug used by 94.1% of patients with preserved eGFR and 75.0% of patients with declined eGFR. Surgery tended to be performed more frequently for patients with preserved eGFR than for those with declined eGFR ($p = 0.08$). Evaluation of the clinical outcomes of 40 patients with preserved eGFR and 11 patients with declined eGFR showed that the rate of recovery after treatment was significantly higher in patients with preserved eGFR than in those with declined eGFR (92% vs. 63.6%, $p = 0.02$) (Table 4).

Table 4
Comparison of the treatment and outcome of patients with
pulmonary cryptococcosis patients, according to eGFR

Treatment	eGFR > 45	eGFR ≤ 45	p value
No treatment	3 (7.1%)	2(14.3%)	0.44
Antifungal drugs	34 (81.0%)	12 (85.7%)	0.68
Treatment duration	8.9 ± 1.7	11.0 ± 2.8	0.53
Surgery	5 (11.9%)	0 (0.0%)	0.08
Outcome			
Recovered	37/40	7/11	0.02
eGFR; estimated glomerular filtration rate			

Multivariate analysis of the clinical impact of CKD on pulmonary cryptococcosis Previous studies reported that immunocompromising comorbidities, such as diabetes mellitus, malignancy, and immunosuppressive drug use, were associated with clinical characteristics (5, 11, 12). The current study found that immunosuppressive drug was used more often by patients with declined eGFR than by those with preserved eGFR (Table 1). Because this might affect the current results, we performed multivariate analysis for the variables which were significantly different between patients with preserved eGFR and those with declined eGFR. Disseminated cryptococcosis, pleural effusion, and recovery after treatment were not assessed because of the small sample size. On multivariate analysis, fever, white blood cell count, albumin, and CRP remained significantly different between patients with preserved eGFR and those with declined eGFR, even after adjustment (Table 5).

Table 5

Multivariate analysis adjusted according to eGFR \leq 45 mL/min/1.73 m² and use of immune suppressive drug

Multivariate analysis				
	eGFR \leq 45		Immunosuppressive drug use	
	OR or β	p value	OR or β	p value
Fever (> 37.5 °C)	6.4	< 0.01	0.7	0.59
Laboratory data				
White blood cell (/ml)	1293.2	0.03	598.9	0.25
Albumin (g/dl)	-0.34	< 0.01	-0.06	0.52
C reactive protein (mg/dl)	0.89	0.01	0.12	0.7
Fever, white blood cell count, albumin and CRP which were significantly different between patients with higher eGFR and those with lower eGFR, and the comparative results were adjusted by predisposing variable of immunosuppressive drug use.				
eGFR; estimated glomerular filtration rate, CRP; C reactive protein, OR; odds ratio, β ; standardized β value,				

Discussion

In the present study, we analyzed 56 patients with non-HIV pulmonary cryptococcosis and evaluated the clinical characteristics depending on the CKD severity. We identified that the rates of disseminated cryptococcosis and non-recovery after treatment were significantly higher in patients with declined eGFR than in those with preserved eGFR. For the clinical features, fever, pleural effusion, high white blood cell count and CRP, and low level of albumin were more frequently seen in patients with declined eGFR than in those with preserved eGFR. To our best knowledge, this was the first report to clarify the clinical manifestations of severe CKD in patients with non-HIV pulmonary cryptococcosis.

CKD is one of the risk factors for pulmonary cryptococcosis; this statement is supported by several individual cases of severe clinical manifestations, such as dissemination and mortality, in chronic renal failure, especially when under dialysis (14, 20). Pyrgos V et al reported that patients hospitalized for cryptococcus meningitis had significantly more frequent comorbidities of acute (28.5%) and chronic renal failure (14.3%), compared with those in all hospitalized patients (24). Moreover, Hung MS et al identified that fever and pleural effusion were significantly more frequent in disseminated pulmonary cryptococcosis than in localized pulmonary cryptococcosis (8). Similarly, our findings showed higher rates of disseminated cryptococcosis and non-recovery after treatment in severe CKD patients with pulmonary cryptococcosis complicated by fever and pleural effusion.

The mechanisms of the association between CKD severity and the risk for pulmonary cryptococcosis remain unclear. However, some possible mechanisms, particularly the effect of the T cell response on the risk for pulmonary cryptococcosis and the pathophysiology of CKD, had been reported. HIV, which is characterized by a decline in CD4⁺ T cells, is one of the major causes of cryptococcosis (25). Moreover, the clinical characteristics and outcome of pulmonary cryptococcosis in patients with immunocompromising conditions, such as underlying malignancy, immunosuppressive drug use, and diabetes mellitus, were reported to be different from those in immunocompetent patients (12, 13, 23). Likewise, CKD has been associated with inactivation of the T cell response and induction of T cell apoptosis (26, 27), which causes immune dysfunction and increases the risk for infection (18, 19). Notably, host response to cryptococcal infection involves helper T cell response with production of cytokines, including tumor necrosis factor, interferon- γ , and interleukin-2 (28, 29). According to these data, we considered that the immune dysfunction in severe CKD might have affected the clinical characteristics and outcomes of patients with pulmonary cryptococcosis.

The clinical characteristics, including symptoms, laboratory abnormalities, and radiologic features, in pulmonary cryptococcosis are variable (30). Consistent with our results, the absence of symptoms was reported to be relatively frequent in immunocompetent patients, whereas the pulmonary cryptococcosis-associated symptoms, especially fever and chest pain, were markedly seen in immunocompromised host (12, 31). The radiologic patterns and distribution in pulmonary cryptococcosis have been closely related with host immunity. Nodules have been more frequently observed in immunocompetent patients, whereas pleural effusion, cavitation, and consolidation in a large lung area have been markedly observed in immunocompromised patients (31–34). Our results of more frequent pulmonary CT abnormalities that were limited to 1 lobe and in a unilateral lung field in patients with preserved eGFR than in those with declined eGFR supported these previous data and the involvement of immunodeficiency from CKD. In addition, we clarified that high white blood cell count and CRP and low albumin might be biomarkers for risk evaluation in patients with pulmonary cryptococcosis and severe CKD.

There were 3 limitations in the present study. First, we used 45 mL/min/1.73 m² as the cutoff between preserved and declined eGFR, based on the results of a previous study (17). However, CKD is defined as eGFR < 60 mL/min/1.73 m² (15), and patients with mild CKD severity were included in the group with preserved eGFR; this might have influenced the results of this study. Second, treatment including the intake of antifungal drugs or surgery was elected depending on the physician's judgment, which might affect to the clinical outcomes. Third, the present study was performed on a small number of patients at a single hospital with limited ethnic diversity. To confirm the validity of our results, multicenter prospective studies with larger number of patients should be performed.

Conclusions

Compared with preserved eGFR, declined eGFR was associated with significantly higher rates of disseminated cryptococcosis and non-recovery after treatment and more frequent clinical features of

fever, pleural effusion, high white blood cell count and CRP, and low level of albumin. These results may practically contribute to the clinical risk evaluation of patients with pulmonary cryptococcosis.

Declarations

Ethics approval and consent to participate

This study was approved by the ethics committee of Saga University Hospital (approval number: 2019-09-06, approval date: Nov 25, 2019) and was performed in accordance with the 1964 Declaration of Helsinki.

Consent for publications

Not applicable

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author Contributions

HT, TH, KT, and NA conceived the project. HT, KT, HS and NA designed the clinical research and interpreted the data. RT and AT advised for statistical analysis. HT and KT prepared the manuscript with input from all other authors. KT, SK and NA performed final check of manuscript.

Acknowledgements

Not applicable

References

1. Knoke M, Schwesinger G. One hundred years ago: the history of cryptococcosis in Greifswald. Medical mycology in the nineteenth century. *Mycoses*. 1994;37(7-8):229-33.
2. Perfect JR, Casadevall A. Cryptococcosis. *Infect Dis Clin North Am*. 2002;16(4):837-74, v-vi.

3. MacDougall L, Fyfe M, Romney M, Starr M, Galanis E. Risk factors for *Cryptococcus gattii* infection, British Columbia, Canada. *Emerg Infect Dis*. 2011;17(2):193-9.
4. Marr KA, Datta K, Pirofski LA, Barnes R. *Cryptococcus gattii* infection in healthy hosts: a sentinel for subclinical immunodeficiency? *Clin Infect Dis*. 2012;54(1):153-4.
5. Maziarz EK, Perfect JR. Cryptococcosis. *Infect Dis Clin North Am*. 2016;30(1):179-206.
6. Denham ST, Brown JCS. Mechanisms of Pulmonary Escape and Dissemination by *Cryptococcus neoformans*. *J Fungi (Basel)*. 2018;4(1).
7. Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, et al. Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. *Lancet Infect Dis*. 2017;17(8):873-81.
8. Hung MS, Tsai YH, Lee CH, Yang CT. Pulmonary cryptococcosis: Clinical, radiographical and serological markers of dissemination. *Respirology*. 2008;13(2):247-51.
9. Brizendine KD, Baddley JW, Pappas PG. Pulmonary cryptococcosis. *Semin Respir Crit Care Med*. 2011;32(6):727-34.
10. Setianingrum F, Rautemaa-Richardson R, Denning DW. Pulmonary cryptococcosis: A review of pathobiology and clinical aspects. *Med Mycol*. 2019;57(2):133-50.
11. Hu Z, Chen J, Wang J, Xiong Q, Zhong Y, Yang Y, et al. Radiological characteristics of pulmonary cryptococcosis in HIV-infected patients. *PLoS One*. 2017;12(3):e0173858.
12. Chang WC, Tzao C, Hsu HH, Lee SC, Huang KL, Tung HJ, et al. Pulmonary cryptococcosis: comparison of clinical and radiographic characteristics in immunocompetent and immunocompromised patients. *Chest*. 2006;129(2):333-40.
13. Zhang Y, Li N, Zhang Y, Li H, Chen X, Wang S, et al. Clinical analysis of 76 patients pathologically diagnosed with pulmonary cryptococcosis. *Eur Respir J*. 2012;40(5):1191-200.
14. Hong N, Chen M, Fang W, Al-Hatmi AMS, Boekhout T, Xu J, et al. Cryptococcosis in HIV-negative Patients with Renal Dialysis: A Retrospective Analysis of Pooled Cases. *Mycopathologia*. 2017;182(9-10):887-96.
15. Levey AS, Atkins R, Coresh J, Cohen EP, Collins AJ, Eckardt KU, et al. Chronic kidney disease as a global public health problem: approaches and initiatives - a position statement from Kidney Disease Improving Global Outcomes. *Kidney Int*. 2007;72(3):247-59.
16. Stenvinkel P. Chronic kidney disease: a public health priority and harbinger of premature cardiovascular disease. *J Intern Med*. 2010;268(5):456-67.
17. Stengel B, Metzger M, Froissart M, Rainfray M, Berr C, Tzourio C, et al. Epidemiology and prognostic significance of chronic kidney disease in the elderly—the Three-City prospective cohort study. *Nephrol Dial Transplant*. 2011;26(10):3286-95.
18. Tecklenborg J, Clayton D, Siebert S, Coley SM. The role of the immune system in kidney disease. *Clin Exp Immunol*. 2018;192(2):142-50.

19. Syed-Ahmed M, Narayanan M. Immune Dysfunction and Risk of Infection in Chronic Kidney Disease. *Adv Chronic Kidney Dis.* 2019;26(1):8-15.
20. Inaba A, Okada A, Yoshida T, Itoyama S, Nakai T, Hisada T, et al. Disseminated Cryptococcosis with Rapidly Growing Lung Nodules in an End-stage Renal Disease Patient. *Intern Med.* 2017;56(3):377-80.
21. Kao CC, Wu VC, Sun HY, Wu CH, Group N. Paradoxical cryptococcal immune reconstitution inflammatory syndrome in advanced chronic kidney disease. *Int Urol Nephrol.* 2013;45(5):1505-9.
22. Matsuo S, Imai E, Horio M, Yasuda Y, Tomita K, Nitta K, et al. Revised equations for estimated GFR from serum creatinine in Japan. *Am J Kidney Dis.* 2009;53(6):982-92.
23. Yanagawa N, Sakai F, Takemura T, Ishikawa S, Takaki Y, Hishima T, et al. Pulmonary cryptococcosis in rheumatoid arthritis (RA) patients: comparison of imaging characteristics among RA, acquired immunodeficiency syndrome, and immunocompetent patients. *Eur J Radiol.* 2013;82(11):2035-42.
24. Pyrgos V, Seitz AE, Steiner CA, Prevots DR, Williamson PR. Epidemiology of cryptococcal meningitis in the US: 1997-2009. *PLoS One.* 2013;8(2):e56269.
25. Park BJ, Wannemuehler KA, Marston BJ, Govender N, Pappas PG, Chiller TM. Estimation of the current global burden of cryptococcal meningitis among persons living with HIV/AIDS. *AIDS.* 2009;23(4):525-30.
26. Brinkkoetter PT, Marinaki S, Gottmann U, Fleckenstein S, Stump C, Van Der Woude FJ, et al. Altered CD46-mediated T cell co-stimulation in haemodialysis patients. *Clin Exp Immunol.* 2005;139(3):534-41.
27. Stachowski J, Pollok M, Burrichter H, Spithaler C, Baldamus CA. Signalling via the TCR/CD3 antigen receptor complex in uremia is limited by the receptors number. *Nephron.* 1993;64(3):369-75.
28. Neal LM, Qiu Y, Chung J, Xing E, Cho W, Malachowski AN, et al. T Cell-Restricted Notch Signaling Contributes to Pulmonary Th1 and Th2 Immunity during *Cryptococcus neoformans* Infection. *J Immunol.* 2017;199(2):643-55.
29. Fa Z, Xu J, Yi J, Sang J, Pan W, Xie Q, et al. TNF-alpha-Producing *Cryptococcus neoformans* Exerts Protective Effects on Host Defenses in Murine Pulmonary Cryptococcosis. *Front Immunol.* 2019;10:1725.
30. Kishi K, Homma S, Kurosaki A, Kohno T, Motoi N, Yoshimura K. Clinical features and high-resolution CT findings of pulmonary cryptococcosis in non-AIDS patients. *Respir Med.* 2006;100(5):807-12.
31. Liu K, Ding H, Xu B, You R, Xing Z, Chen J, et al. Clinical analysis of non-AIDS patients pathologically diagnosed with pulmonary cryptococcosis. *J Thorac Dis.* 2016;8(10):2813-21.
32. Khoury MB, Godwin JD, Ravin CE, Gallis HA, Halvorsen RA, Putman CE. Thoracic cryptococcosis: immunologic competence and radiologic appearance. *AJR Am J Roentgenol.* 1984;142(5):893-6.
33. Rozenbaum R, Goncalves AJ. Clinical epidemiological study of 171 cases of cryptococcosis. *Clin Infect Dis.* 1994;18(3):369-80.

34. Xie LX, Chen YS, Liu SY, Shi YX. Pulmonary cryptococcosis: comparison of CT findings in immunocompetent and immunocompromised patients. *Acta Radiol.* 2015;56(4):447-53.