

Activation of the SA-Associated Plant Defense Pathway Alters the Composition of Soil Bacterial Communities

Jing Zhang (✉ j.zhang@biology.leidenuniv.nl)

Universiteit Leiden Instituut Biologie Leiden

Peter G.L. Klinkhamer

Universiteit Leiden Instituut Biologie Leiden

Klaas Vrieling

Universiteit Leiden Instituut Biologie Leiden

T. Martijn Bezemer

Universiteit Leiden Instituut Biologie Leiden

Research Article

Keywords: Metatranscriptomics, Soil microbial community, Taxonomy, Plant-soil interactions, Microbial diversity, Rhizosphere soil, Salicylic acid

Posted Date: February 3rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-159617/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background and aims

Many plant species grow better in sterilized than in live soil. Foliar application of SA mitigates this negative effect of live soil on the growth of the plant *Jacobaea vulgaris*. To examine what causes the positive effect of SA application on plant growth in live soils, we analyzed the effects of SA application on the composition of active rhizosphere bacteria in the live soil.

Methods

We studied this over four consecutive plant cycles (generations), using mRNA sequencing of the microbial communities in the rhizosphere of *J. vulgaris*.

Results

Our study shows that the composition of the rhizosphere bacterial communities of *J. vulgaris* greatly differed among generations. Application of SA resulted in both increases and decreases in a number of active bacterial genera in the rhizosphere soil, but the genera that were affected by the treatment differed among generations. In the first generation, there were no genera that were significantly affected by the SA treatment, indicating that induction of the SA defense pathway in plants does not lead to immediate changes in the soil microbial community. 89 species out of the total 270 (32.4%) were present in all generations in all soils of SA-treated and control plants suggesting that these make up the “core” microbiome. On average in each generation, 72.9% of all genera were present in both soils. Application of SA to plants significantly up-regulated genera of *Caballeronia*, unclassified *Cytophagaceae*, *Crinalium* and *Candidatus Thermofonsia Clade 2*, and down-regulated genera of *Thermomicrobiales*, unclassified *Rhodobacterales*, *Paracoccus* and *Flavihumibacter*. While the functions of many of these bacteria are poorly understood, bacteria of the genus *Caballeronia* play an important role in fixing nitrogen and promoting plant growth, and hence this suggests that activation of the SA signaling pathway in *J. vulgaris* plants may select for bacterial genera that are beneficial to the plant.

Conclusions

Overall, our study shows that aboveground activation of defenses in the plant affects soil microbial communities and, as soil microbes can greatly influence plant performance, this implies that induction of plant defenses can lead to complex above-belowground feedbacks. Further studies should examine how activation of the SA signaling pathway in the plant changes the functional genes of the rhizosphere soil bacterial community.

Introduction

Plants encounter a myriad of threats from the surrounding environment, including both abiotic and biotic stresses (Suzuki et al. 2014). Biotic stresses are mostly due to herbivory and pathogen infestation both

below- and above-ground (Adair and Douglas 2017; Pieterse and Dicke 2007). Microbes in the soil can have a beneficial, pathogenic or neutral effect on the host plant. For example, soil bacteria such as *Rhizoctonia species*, often strongly negatively affect plant growth and survival (Issac et al. 1971). On the other hand, plant growth-promoting rhizobacteria (PGPR), such as *Pseudomonas* and *Burkholderia* species are beneficial for the plant, e.g. via suppressing the growth of soil-borne pathogens or increasing nutrient availability (Bhattacharyya and Jha 2012). However, the overall net effect of soil microbial communities on plant growth is often negative (Nijjer et al. 2007). Most plant species grow less well in soils that contain a natural microbial community than in sterilized soils. This might be due to e.g. competition between plants and microbes for available nutrients or due to soil-borne plant pathogens (Berendsen et al. 2012; Callaway et al. 2004; Cesarano et al. 2017; Mazzoleni et al. 2015).

Systemic acquired resistance (SAR) is one of the most common defensive strategies of plants against biotrophic pathogenic microbes. Foliar application of salicylic acid to plant tissues can activate SAR and boost the innate immune system of a plant (Reymond and Farmer 1998). Cultivars with a higher sensitivity to SA are often better defended against the pathogens. For example, in tomato, exogenous application of SA can be effective against the pathogens *Oidium neolycopersici* and *Botrytis cinerea*, which cause powdery mildew and grey mould diseases (Achuo et al. 2004; Seskar et al. 1998). In agriculture, application of SA is now used to suppress pathogenic microbial effects in e.g. tomato, pepper and pea crops (Barilli et al. 2010; Choi and Hwang 2011; Esmailzadeh et al. 2008). How SA application to the plant affects the microbial community in the soil is not fully uncovered.

Because plants alter the composition of the microbial community in the soil in which they grow, and SAR protects plants against pathogens, an important question is how activation of SAR alters the plant's effect on the soil microbial community. Several studies have demonstrated that the activation of SAR indeed altered the composition of the soil microbial community and that SA can play a key role in shaping root bacterial communities (Hein et al. 2008; Kniskern et al. 2007; Lebeis et al. 2015). However, several other studies reported that foliar application of SA does not affect the bacterial composition in the soil (Doornbos et al. 2011; Wang et al. 2015). These studies on the effects of SAR on soil bacterial composition were mostly limited to the model plant species *Arabidopsis*. As plant species differ greatly in the way and magnitude in which they influence the soil bacterial community (Hannula et al. 2019; Pineda et al. 2020; Wubs and Bezemer 2018), we may expect that the effects of SA application on the soil microbial community also differ among plant species.

Several studies have shown that the composition of the soil bacterial microbial community varies greatly over time (e.g. Hannula et al. 2019). In a study on temporal variation in three land-use types, the number of taxa present in the soil showed strong temporal variability, and these changes over time were considerably larger than the variation associated with land-use types (Lauder et al. 2013). Shade et al. (2012) demonstrated that soil microbial communities have clear successional trajectories. If generally true this would imply that application of SA to plants could also cause directed changes in the soil microbial community over time. An important question is therefore how activation of SAR will alter the soil microbial community over time.

Previous studies showed that inoculation of a sterilized soil with natural, live soil reduced plant growth in comparison with that in sterilized soil for the plant species *Jacobaea vulgaris* (Joosten et al. 2009; Kos et al. 2015). Interestingly, applying SA to the leaves mitigated these negative effects (Jing et al. unpublished data). This implies that activation of SA-induced resistance may potentially suppress microbial pathogens present in live soil. If this is the case, an important question is whether the repeated foliar application of SA during consecutive generations of plant growth will increase this effect and hence, whether there is a selection for a more beneficial bacterial community. Conceptually, the temporal dynamics of foliar application of SA can follow different trajectories (Fig. 1). First, it is possible that both foliar application of SA and the effect of different generations do not alter the soil bacterial composition (Fig. 1-i). Second, foliar application of SA may lead to different bacterial communities independent of time (Fig. 1-ii). Third, bacterial communities may differ among generations but are not influenced by the SA application (Fig. 1-iii). Fourth, foliar application of SA may influence bacterial communities but these effects may differ among generations (Fig. 1-iv).

In this study, we sequenced the mRNA from rhizosphere soil samples of both SA-treated and control plants during four consecutive generations of growth of *J. vulgaris*. In each consecutive generation soil from the previous plant growth period was used. Using mRNA instead of DNA or rRNA enabled us to focus on the active soil microbial community (Gilbert et al. 2008). In this study, we focus on the bacterial community. Twenty-four rhizosphere soils were sequenced with an Illumina sequencing platform. The goal of this study is to answer the following questions: (1) How does the foliar application of SA in *J. vulgaris* affect the bacterial composition in the rhizosphere and is there a time effect or an interactive time × SA effect on the bacterial community? (2) What is the “core” bacterial community in the soils of plants exposed to the SA treatment and of control plants? (3) How does the application of SA influence the bacterial community in each generation? Are the SA effects consistent over time?

Materials And Methods

Multi-generational plant growth experiment

The current study focuses on the effect of foliar SA application on the composition of the bacterial community in the rhizosphere in the inoculated soil. The effects of foliar application of SA on plant growth in the four generations are described elsewhere (Jing et al. submitted). Details of the experiment are described below.

J. vulgaris (common ragwort) seeds were collected at the dunes of Meijendel (a calcareous sandy area from a coastal dune area north of The Hague, The Netherlands, 52°11'N, 4°31'E). Prior to germination, all seeds were surface sterilized (shaken for 2 min in 70% ethanol, then rinsed with sterilized water, put for 12 min in 2% bleach, and then rinsed again four times with sterilized water to minimize influences of seed-borne microbes (Bakker et al. 2015). The soil was also collected at Meijendel. The topsoil was collected to a depth of 15 cm after removing the grassland vegetation and the organic layer of the surface. The soil was sieved using a 5 mm sized mesh, homogenized with a concrete mixer, and then stored into 20-liter

plastic bags (Nasco Whirl-Pak Sample Bag). Bags were either sterilized by 35-K Gray gamma-irradiation (Synergy Health Company, Ede, The Netherlands) or kept at 4°C for inoculation.

Surface sterilized seeds were germinated in sterile Petri dishes on filter paper. After one week, seedlings were randomly transferred individually to 500 ml pots consisting of a mixture of 90% sterilized soil and 10% live soil. Prior to potting but after mixing, the soil was kept in bags and left in the climate room for 14 days so that the mixed soil could settle and microbial communities could colonize the sterilized soil. After potting the seedlings, pots were randomly distributed over a climate room (relative humidity 70%, light 16h at 20°C, dark 8h at 20°C). Plants were watered regularly with Milli-Q and 5 ml Steiner nutrient solution was added per plant on day seven after planting, 10 ml Steiner nutrient solution (Steiner 1979) was added on day 13, and 20 ml Steiner nutrient solution was added on days 19, 28, 37, 42. The Steiner nutrient solution was prepared from seven different stock solutions (106.2 g $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, 29.3 g KNO_3 , 13.6 g KH_2PO_4 , 49.2 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 25.2 g K_2SO_4 and 2.24 g KOH, 3.29 g Fe-EDTA added to 1 liter demineralized water, and a stock solution with micro elements (a mixed solution of 0.181 g $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 0.286 g H_3BO_3 , 0.022 g $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.0078 g $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and 0.0126 g $\text{NaMoO}_4 \cdot 2\text{H}_2\text{O}$ added to 1 liter demineralized water). Ten ml of each stock solution was diluted in 1 liter of demineralized water before use.

Plants were allocated to either a hormonal treatment (SA) or served as control (only solvent). Both treatments were replicated ten times. During plant growth, the phytohormone SA was applied through foliar application three times a week for four consecutive weeks. The first application was given when plants were 14 days old. About 0.75 ml of 100 μM SA was sprayed on the leaves while carefully avoiding spillover to the soil. One week later the treatment was repeated with 1.50 ml of SA. In the next week, the treatment was repeated with 2.25 ml of SA. SA solvent (purchased from Sigma-Aldrich, $\geq 99.0\%$) was made by dissolving 6.91 mg in 69.10 μl of ethanol. Milli-Q water was then added until a final volume of 500 ml. Control plants were sprayed with sterile water with the same solvent (ethanol in Milli-Q water).

After six weeks, plants were gently removed from the pots. The rhizosphere soil for each treatment was harvested for each pot individually by gently shaking three times to remove the loosely adhering soil, after which rhizosphere soil was collected onto a sterile filter paper by removing the remnant soil with a fine sterilized brush. Rhizosphere soil was put in a 2 ml Eppendorf tube and stored at -80°C for further RNA extraction. The remaining rhizosphere soil and adhering soil of the ten pots were mixed and used as inoculum (live soil) for the next generational of plant growth. The inoculum soil (10%) was mixed with 90% sterilized soil.

The set-up was repeated for another three generations under the same conditions as described above so that there were four generations of plant growth. For the second, third, and fourth generation, the soil inoculum was derived from the previous generation from the same treatment and was a mixture of rhizosphere soil and the loosen adhering soil surrounding the roots. Again, after mixing, the soil was kept in bags and left in the climate room for 14 days. Hereafter, pots were filled with soil and a *J. vulgaris* seedling was planted into each pot. All replicate soils from the SA or control treatment were

mixed before the inoculation. The SA treatment was carried out as described above in each generation. Fifty-four days after planting, all plants were harvested each time.

RNA extraction and metatranscriptomic sequencing

For each treatment, the three successively labeled samples (No. 1, 2, 3, No. 4, 5, 6 and No. 7, 8, 9) were mixed and used as one composed replicate, Hence, three composed replicates were generated and used for RNA extraction for each treatment in each generation and a total of 24 soil samples were used for RNA extraction (3 replicates × 2 treatments × 4 generations). Total RNA was extracted with the RNeasy PowerSoil Total RNA kit (Qiagen). RNA concentration and quality were assessed by running 1 µl of the extracted raw RNA on the 4200 TapeStation (Agilent). Subsequently, unwanted DNA, salts and buffers were removed with the RNeasy minElute Cleanup Kit (Qiagen). The Ribo-Zero Magnetic kit for bacteria (Illumina) was used for mRNA enrichment and a RNA Clean & Concentrator kit (Zymoresearch) was used to clean additional buffers and proteins of the rRNA-depleted RNA. All the steps in extracting and cleaning RNA were according to the supplier's instructions. Double-stranded cDNA was generated from the cleaned RNA obtained in the final step. Library preparation (Illumina Nextera XT DNA library), processing and sequencing were performed by FG Technologies (Leiden, The Netherlands) with paired-end (PE) 150 bp templates. Raw sequencing reads were deposited in NCBI database (accession number: SUB8738755). Twenty-four metatranscriptomic libraries were generated, the size of each library is indicated in Table S1 and Fig. S1.

Bioinformatics processing

Trimmomatic 0.39 was used for the removal of adapters of paired-end raw reads (Bolger et al. 2014). FastQC was applied to check the qualities, the bases with a threshold lower than 30 were cut off with Trimmomatic (Andrews 2010). Ribosomal RNAs of all 24 metatranscriptomic libraries were filtered with SortMeRNA (Sorting ribosomal RNA) (Kopylova et al. 2012). Eight rRNA representative databases (silva-bac-16s-id90, silva-arc-16s-id95, silva-euk-18s-id95, silva-bac-23s-id98, silva-arc-23s-id98, silva-euk-28s-id98 rfam-5s-id98, rfam-5.8s-id98) were derived from the SILVA SSU and LSU databases (release 119) and the RFAM databases with HMMER 3.1b1 and SumaClust v1.0.00 were used for fast filtering of rRNA from eukaryote, prokaryote and archaea. Then, all reads of the 24 metatranscriptomic libraries were combined into one set, which was the input of a de novo assembly. Trinity with default parameters was used for the metatranscriptomic assembly (Haas et al. 2013). Later, the quality of assembled contigs was assessed with Trinity scripts. The CD-HIT-EST algorithm was used to remove the duplicates of each transcript and reads with shorter than 300 bps were removed with a script modified from Li and Godzik (2006), after which reads of each library were mapped back to transcriptome with Bowtie2 (Langmead and Salzberg 2012). The isoform IDs per sample were extracted with Seqkit (Shen et al. 2016). Contigs of each sample were generated and then aligned against the NCBI NR (non-redundant) database by DIAMOND with a cut off e-value at 1e-5 (Buchfink et al. 2015). The closest match with an identity higher than 80% was kept for mapping. The output file of Blastx was further analyzed with the lowest common ancestor (LCA) algorithm in MEGAN (version 6.0) with all default parameters (Camon et al. 2005; Huson

et al. 2016). MEGAN was used to compute the data at different taxonomic levels and in this process NCBI taxonomy was employed to summarize results. The detailed workflow is described in Huson et al. (2007). A count table of microbial species was obtained with read counts assigned directly to taxon for the 24 samples. The number of assigned reads per taxa was extracted at species, genus, family and phylum level respectively. The number of identified phyla, families, genera and species were counted, and the composition and the percentage of reads used for each classification level were calculated.

Statistical analysis

Differences in the number of total reads and the number of non-rRNA reads over four generations were presented as mean \pm SD. A Heinrich's triangle figure was generated to visualize microbial composition at different phylogenetic levels of all the identified microbes from the 24 rhizosphere soil samples. Log₁₀-transformed hit numbers of each genus were plotted as a function of the ranked genus abundance numbers, including all species, and a cut-off was performed with an abundance larger than 0.01% of the total reads. A Shapiro-Wilk test was used to test for differences between the distribution of abundance between the SA and control treatment.

The Shannon-diversity index was calculated for each of the 24 samples and differences between the Shannon-diversity of soils of SA treated plants and soils of control plants were tested with a student t-test. Subsequently, abundance at genus level was used for to construct NMDS (nonmetric multidimensional scaling), PCA (Principal component analysis), OPLS-DA (orthogonal projection to latent structures discriminant analysis), and Venn diagrams, and Pearson distance and the Ward clustering algorithm statistical analysis was calculated since most of the reads were identified at the genus level.

Two-factor Venn diagrams were constructed to illustrate the numbers of unique and common genera in soil samples within each generation for the SA and control treatments, and a four-factor Venn diagram including all generations was performed for the SA and the control treatment separately (Heberle et al. 2015).

PCA and OPLS-DA were performed with SIMACA 15.0 using relative abundance at genus level. The relative abundance was calculated using the absolute abundance number of one genus divided by the total abundance of all bacterial genera in the sample. Before performing OPLS-DA analysis, we checked that our data fitted the model with a cross-validated residual (CV)-ANOVA significance testing ($n = 270$, $P < 0.02$).

To visualize the compositional changes among different treatment and time categories, a NMDS using the Bray-Curtis index as a measure of dissimilarity was generated using relative abundances. To verify changes in composition due to the SA treatment and time effect, a PERMANOVA test was performed using the *Adonis* function (number of permutations = 999) in R within the "vegan" package.

Local "immigration" and "extinction" in the rhizosphere soil of SA-treated or control plants over generations at genus level was calculated and the numbers were presented in Venn diagram. A Student's

t-test was used to identify bacterial genera that were significantly enriched in soil samples of SA-treated or control plants. *P* values were adjusted for false discovery rates (FDR).

Spearman's rank correlation without multiple comparison tests were performed to identify the genera that were significantly positively or negatively correlated with generation within the SA or control treatment. Genera with *P* values smaller than 0.05 were selected to create a heatmap for all 24 samples. Hierarchical clustering analysis was done for the 24 samples together, based on the relative abundance to show the similarity. The row-centered relative abundance of each genus was used to construct the color key (Chong et al. 2018). Heatmaps for only SA and only control treatments were also generated.

Results

Metatranscriptomic sequence data

A total of 898.4 million raw sequence reads were obtained from the 24 metatranscriptomic libraries, the smallest and largest library contained 25.0 and 52.0 million raw sequence reads, respectively (supplementary data Table S1). 846.9 million reads were kept after removing adapters and quality filtering control with FastQC. In total, 775.3 million reads were removed with the SortMeRNA program as ribosomal RNA (rRNA) reads when aligning them against eight rRNA representative databases (silva-bac-16s-id90, silva-arc-16s-id95, silva-euk-18s-id95, silva-bac-23s-id98, silva-arc-23s-id98, silva-euk-28s-id98 rfam-5s-id98, rfam-5.8s-id98), and 71.6 million reads were used as non-rRNA reads for further de novo assembly with Trinity (Fig. S1). For this set, the smallest library contained 1.5 million reads and the largest one 5.9 million reads. Reads for de novo assembly were normalized with Trinity in silico normalization algorithm. The average guanine-cytosine (GC) content for the 24 libraries was 60.10%. After assembly, 0.99 million contigs were removed because their length was shorter than 300 bps. A total of 1.3 million unique contigs were identified after removing duplicates with CD-HIT-EST. In total, 392.4 million bases were assembled. After we checked the quality of the contigs in all samples by realigning all contigs back to the assemblies using Bowtie2, the average mapping rate for proper pairs was 45.41%.

Overview of the assigned reads at differential microbial classification levels

When we aligned the 1.3 million unique contigs against the NR (non-redundant) database with DIAMOND and MEGAN 6.0, 0.39 million contigs were taxonomically classified, while the others did not provide a match with the available taxonomic information. Based on the analysis in MEGAN, the identified contigs were assigned at different classification levels. Twenty-two different bacterial phyla were identified, 283 families and 382 bacterial genera and 1081 bacterial species (Fig. S2). At the phylum, family, genus and species level 23.4%, 23.4%, 20.4% and 14.9% of the total number of contigs were assigned, respectively. Bacteria were the most prevalent in the microbial community taking up 98.3% of the total number of reads (Fig. S3a). Eukaryotes, with algae taking the largest proportion, were the second dominant, but Eukaryotes only covered 1.5% of the total number of reads (Fig. S3b).

SA application and time effects on bacterial community diversity and composition

From the total of 408 bacterial genera, 270 genera were included in the analysis (contigs with more than 0.01% of the total number of reads Fig. S5). The genera in both soils showed significantly different abundance curves (Shapiro-Wilk test, $df=407$, $P<0.0001$; Fig. S5), the abundance curve in the SA soil was lower than that in the control soil. Application of SA did not significantly increase or decrease the Shannon diversity at genus level within each generation (t-test for the 1st generation: $t=-0.63$, $df=5$, $P=0.27$; 2nd generation: $t=0.07$, $df=5$, $P=0.47$; 3rd generation: $t=0.67$, $df=5$, $P=0.26$; 4th generation: $t=0.50$, $df=5$, $P=0.31$).

The NMDS plot showed that the bacterial communities of the same generation clustered together (Fig. 2a), PERMANOVA $R^2=0.30$, $P=0.001$). The SA and control separated in the NMDS plot (Fig. S6) but this was not significant (PERMANOVA $R^2=0.05$, $P=0.18$). Similar patterns were observed in a principal component analysis (PCA; Fig. S7). The OPLS-DA analysis showed clusters for replicates within each generation, and clear separation for the SA effect but only in the 2nd, 3rd and 4th generation (Fig. 2b). However, the generation effect was more evident than the SA effect.

Core bacterial community

Eight-nine species out of the total of 270 (32.4%) were present in all generations in at least two out of the three replicates of the soils of SA-treated and control plants suggesting that these make up the “core” microbiome (Fig. 3a). On average in each generation, 72.9% of all the genera were present in both soils (Fig. 3b). In the first generation, both soils shared about 74.2% of the genera while 7.7% only occurred in the SA-treatment and 18.0% only in the control (Fig. 3b-1). The percentage of shared genera by the two soils in the 2nd, 3rd and 4th generation was 67.6%, 72.9% and 76.8% (Fig. 3b-2, 3, 4). For soils of the control treatment, 49.5% of the genera were shared over all four generations; while 45.1% of genera were shared in soils of the SA treated plants over four generations (Fig. S2c; Table S2). Immigration (i.e genera not present in the soil in the previous generation) was somewhat higher in the SA treatment (on average 42 new genera) than in the control (on average 34 new genera) while the opposite was true for extinction rates (genera present in the previous but not in the current generation). On average, 31 genera went extinct in the SA treatment and 33 in the control treatment; Fig. 4. The information of Archaea, virus and eukaryote is listed in the supplementary information (Fig. S4).

SA selection of soil bacteria

When analyzed per generation, in total eight genera differed among the SA treatment and control (Fig. 5). No genus was significantly affected in more than one generation and no genera were significantly affected in the first generation. Most of the significant genera were only present in either the control or SA treatment. A Spearman's rank correlation showed that 41 (out of 240) genera in the rhizosphere soil of SA-treated plants were significantly increasing or 31 genera were decreasing over generations. For the control soils these numbers were 47 and 27, respectively out of a total of 239 genera (Table S3). The heatmap including all 24 samples showed a clear generation effect, but no clear SA effect (Fig. 6). A heatmap representing the patterns of all identified genera in the 12 rhizosphere soils of SA-treated plants

showed that replicates within a generation clustered and that the 2nd, 3rd and 4th generation showed a higher similarity than the 1st generation (Fig S8a). For the control plants, the samples from the first generation differed from the three other generations (Fig S8b).

Discussion

In this study, we examined how the activation of SA-induced resistance in the plant impacts the microbial composition in the rhizosphere, and how this effect changes over generations of plant growth. Our study shows that the composition of the active rhizosphere bacteria communities of the plant *J. vulgaris* *changed significantly over generations*, but that neither the effects of activation of SA-associated plant defense pathways nor the interaction between generation number and SA on the bacterial composition had a significant on the composition. Within generations the application of SA selected for different bacterial genera in the rhizosphere soil, but these selected genera differed from generation to generation. There were no SA-mediated changes in active bacterial genera in the first generation, suggesting that there are no immediate effects of activation of the SA defense pathway on the soil microbial composition. The majority (76.1%) of the bacterial genera that we detected was present in all soils and represents the “core” bacterial microbiome.

Our study showed that aboveground activation of SA-associated plant defense pathways influenced different bacterial genera in the second, third and fourth generations. Effects of SA-induced resistance on the soil microbial community have been reported in several other studies. For example, Hein et al. (2008) compared the effect of SA application on the composition of rhizosphere bacterial communities in several *Arabidopsis* mutants with terminal restriction fragment length polymorphism (T-RFLP) analysis. They found that SA-induced resistance changed the structure of bacterial communities in the rhizosphere. In addition, Lebeis et al. (2015) demonstrated that SA application modulates colonization of the root microbiome by specific bacterial taxa. SA in plants is associated with the expression of pathogenesis-related proteins (PRPs). These PRPs possess antimicrobial activities resulting in suppression of microbial pathogens, consequently changing the microbial composition (Van Loon and Van Strien 1999; Yalpani et al. 1991). Alternatively, hormonal-induced resistance in the plant may promote beneficial bacteria and fungi. However, the impact of SA-induced resistance on soil microbial communities is still debated. For instance, Wang et al. (2015) and Doornbos et al. (2011) both demonstrated that activation of SA-induced resistance did not significantly affect the composition and diversity of the rhizosphere bacterial community.

Even though the experimental conditions and plant genotypes remained the same throughout the experiment, the effects of SA application on the bacterial community differed among generations. In this context, it is important to note that for each generation we used an inoculum, which means that we placed a subset of the microbial community in a sterile background. This may explain why we saw so much variation temporally as in each generation a different subset of the microbial community may have been activated. It is also possible that the composition of the bacterial community is variable over time

within each generation and as a consequence also among generations (Gilbert et al. 2009; Hannula et al. 2019; Hickey et al. 2013; Lauber et al. 2013).

Of the four potential models, our data confirmed the third hypothesis (Fig. 1-iii), showing that the bacterial communities did differ among generations but were not strongly influenced by SA application. This is in line with studies showing that the composition of the soil bacterial microbial community exhibits large fluctuations over time (Hannula et al. 2019; Lauder et al. 2013). Moreover, our data also shows that the application of SA selects for different bacterial genera in the rhizosphere soil but that these selected genera differ from generation to generation. This suggests that the effects of SA application to plants on the soil microbial community are not consistent over time and that it will be difficult to predict the effects of activation of plant defenses on soil microbes, and ultimately how this will influence the interactions between plants and microbes in the rhizosphere.

Interestingly, in soils of SA-treated plants, we found an increase of *Caballeronia*, unclassified *Cytophagaceae*, *Crinalium* and *Candidatus Thermofonsia Clade 2*. The *Caballeronia* genus is often reported to play an important role in fixing nitrogen and promoting plant growth. Species in this genus are predominantly endophytic diazotrophic bacteria and N-fixing bacteria (Padda et al. 2018; Puri et al. 2018; Puri et al. 2020). This suggests that activation of SA signaling pathways in *J. vulgaris* plants benefited bacteria that were more beneficial to plant growth, but further studies are needed to confirm this. The functions of the other species of which their abundance differentially increases are poorly understood. It is noteworthy, though, that *Crinalium* is a genus that is often isolated from sandy dune soils so it is not surprising that we detected this genus as we used dune soils in our experiment (Mikhailyuk et al. 2019). Further studies should extract the information of these detected genera at the species level.

In conclusion, we provide evidence that the composition of bacterial communities in the rhizosphere significantly differed between plant cycles (generation), but we found no evidence that application of SA altered this pattern. However, application of SA influenced different bacterial genera in the rhizosphere, but the responsive genera varied between generations. No bacterial genera were detected that responded to SA application in the first generation suggesting that there are no immediate responses of bacteria in the rhizosphere to SA application to plants. This would question the so-called ‘cry for help’ hypothesis (Biere and Bennett 2013; Pineda et al. 2013; Rasmann et al. 2017), but further studies are required before we can make firm conclusions about this. Our results provide a new perspective on the effects of plant hormones on temporal changes in the soil microbial community.

Declarations

Author contributions

JZ, MB, KV and PK designed the experiment, JZ carried out the experiment, JZ, KV, MB and PK performed the data analysis. JZ, KV, MB and PK wrote the first draft of the manuscript, and all the authors contributed to revisions.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We would like to thank Connor Philippo for his practical assistance in extracting RNA and quality control of the raw reads, Bing Xie for his generous supporting in sharing his private server and help in troubleshooting, Fons Verbeek for sharing public resource of the server in the Leiden institute of advanced computer science (LIACS). We also thank Martine Huberty for discussing PCA statistical analysis. We thank Robin van Schendel for assisting in the Trinity assembly on Shark cluster, Meciael Liem and Alexander Kloosterman for troubleshooting coding errors in Linux, Niels van Tol for helping introducing people who can help us with performing Trinity assembly. Jing Zhang would also like to thank the China Scholarship Council for financial support.

References

- Achuo EA, Audenaert K, Meziane H, Höfte M (2004) The salicylic acid-dependent defence pathway is effective against different pathogens in tomato and tobacco. *Plant Pathol* 53:65-72
- Adair KL, Douglas AE (2017) Making a microbiome: the many determinants of host-associated microbial community composition. *Curr Opin Microbiol* 35:23-29
- Andrews S (2010) FastQC: a quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Bakker MG, Chaparro JM, Manter DK, Vivanco JM (2015) Impacts of bulk soil microbial community structure on rhizosphere microbiomes of *Zea mays*. *Plant Soil* 392:115-126
- Barilli E, Sillero JC, Rubiales D (2010) Induction of systemic acquired resistance in pea against rust (*Uromyces pisi*) by exogenous application of biotic and abiotic inducers. *J Phytopathol* 158:30-34
- Berendsen RL, Pieterse CM, Bakker PA (2012) The rhizosphere microbiome and plant health. *Trends Plant Sci* 17:478-486
- Bhattacharyya PN, Jha DK (2012) Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World J Microb Biot* 28:1327-1350
- Biere A, Bennett AE (2013) Three-way interactions between plants, microbes and insects. *Funct Ecol* 27:567-573
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30: 2114-2120

- Buchfink B, Xie C, Huson DH (2015) Fast and sensitive protein alignment using DIAMOND. *Nat Methods* 12:59
- Callaway RM, Thelen GC, Rodriguez A, Holben WE (2004) Soil biota and exotic plant invasion. *Nat* 427:731-733
- Camon EB, Barrell DG, Dimmer EC, Lee V, Magrane M, Maslen J, Binns D, Apweiler R (2005) An evaluation of GO annotation retrieval for BioCreAtIvE and GOA. *BMC bioinformatics* 6:S17
- Cesarano G, Zotti M, Antignani V, Marra R, Scala F, Bonanomi G (2017) Soil sickness and negative plant-soil feedback: A reappraisal of hypotheses. *J Plant Pathol* 99:545-570
- Choi HW, Hwang BK (2011) Systemic acquired resistance of pepper to microbial pathogens. *J Plant Pathol* 159:393-400
- Chong J, Soufan O, Li C, Caraus I, Li S, Bourque G, Wishart DS, Xia J (2018) MetaboAnalyst 4.0: towards more transparent and integrative metabolomics analysis. *Nucleic Acids Res* 46:W486-W494
- Doornbos RF, Geraats BP, Kuramae EE, Van Loon LC, Bakker PA (2011) Effects of jasmonic acid, ethylene, and salicylic acid signaling on the rhizosphere bacterial community of *Arabidopsis thaliana*. *Mol Plant-microbe Interact* 24:395-407
- Esmailzadeh M, Soleimani MJ, Rouhani H (2008) Exogenous applications of salicylic acid for inducing systemic acquired resistance against tomato stem canker disease. *J Biol Sci* 8:1039-1044
- Gilbert JA, Field D, Huang Y, Edwards R, Li W, Gilna P, Joint I (2008) Detection of large numbers of novel sequences in the metatranscriptomes of complex marine microbial communities. *PLoS One* 3:e3042
- Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, MacManes MD (2013) De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc* 8:1494
- Hannula SE, Kielak AM, Steinauer K, Huberty M, Jongen R, Jonathan R, Heinen R, Bezemer TM (2019) Time after time: temporal variation in the effects of grass and forb species on soil bacterial and fungal communities. *MBio* 10:e02635-19
- Heberle H, Meirelles GV, da Silva FR, Telles GP, Minghim R (2015) InteractiVenn: a web-based tool for the analysis of sets through Venn diagrams. *BMC Bioinformatics* 16: 169
- Hein JW, Wolfe GV, Blee KA (2008) Comparison of rhizosphere bacterial communities in *Arabidopsis thaliana* mutants for systemic acquired resistance. *Microb Ecol* 55:333-343
- Hickey RJ, Abdo Z, Zhou X, Nemeth K, Hansmann M, Osborn III TW, Wang F, Forney LJ (2013) Effects of tampons and menses on the composition and diversity of vaginal microbial communities over

time. *BJOG* 120:695-706

Huson DH, Auch AF, Qi J, Schuster SC (2007) MEGAN analysis of metagenomic data. *Genome Res* 17:377-386

Huson DH, Beier S, Flade I, Górska A, El-Hadidi M, Mitra S, Ruscheweyh HJ, Tappu R (2016) MEGAN community edition-interactive exploration and analysis of large-scale microbiome sequencing data. *PLoS Comput Biol* 12:e1004957

Isaac I, Fletcher P, Harrison JAC (1971) Quantitative isolation of *Verticillium spp.* from soil and moribund potato haulm. *Ann Appl Biol* 67:177-183

Jing Z, Klinkhamer PGL, Vrieling K, Bezemer TM (2020) Activation of hormone-associated plant defense pathways alters the effects of live soils on plant performance BAAE submitted

Joosten L, Mulder PP, Klinkhamer PG, van Veen JA (2009) Soil-borne microorganisms and soil-type affect pyrrolizidine alkaloids in *Jacobaea vulgaris*. *Plant Soil* 325:133

Kniskern JM, Traw MB, Bergelson J (2007) Salicylic acid and jasmonic acid signaling defense pathways reduce natural bacterial diversity on *Arabidopsis thaliana*. *Mol Plant-microbe Interact* 20:1512-1522

Kopylova E, Noé L, Touzet H (2012) SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics* 28:3211-3217

Kos M, Tuijl MA, de Roo J, Mulder PP, Bezemer TM (2015) Plant-soil feedback effects on plant quality and performance of an aboveground herbivore interact with fertilisation. *Oikos* 124:658-667

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357

Lebeis S. L, Paredes S. H, Lundberg D. S, Breakfield N, Gehring J, McDonald M, Malfatti S, Del Rio TG, Jones CD, Tringe SG, Dangl JL (2015) Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Sci* 349:860-864

Lauber CL, Ramirez KS, Aanderud Z, Lennon J, Fierer N (2013) Temporal variability in soil microbial communities across land-use types. *ISME J* 7:1641-1650

Li W, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* 22:1658-1659

Mazzoleni S, Bonanomi G, Incerti G, Chiusano ML, Termolino P, Mingo A, Senatore M, Giannino F, Carteni F, Rietkerk M, Lanzotti V (2015) Inhibitory and toxic effects of extracellular self-DNA in litter: a mechanism for negative plant-soil feedbacks? *New Phytol* 205:1195-1210

Mikhailyuk T, Vinogradova O, Holzinger A, Glaser K, Samolov E, Karsten U (2019) New record of the rare genus *Crinalium* Crow (Oscillatoriales, Cyanobacteria) from sand dunes of the Baltic Sea, Germany:

epitypification and emendation of *Crinalium magnum* Fritsch et John based on an integrative approach. *Phytotaxa* 400:165-179

Nijjer S, Rogers WE, Siemann E (2007) Negative plant-soil feedbacks may limit persistence of an invasive tree due to rapid accumulation of soil pathogens. *Proc Royal Soc* 274:2621-2627

Padda KP, Puri A, Chanway CP (2018) Isolation and identification of endophytic diazotrophs from lodgepole pine trees growing at unreclaimed gravel mining pits in central interior British Columbia, Canada. *Can J For Res* 48:1601-1606

Pieterse CM, Dicke M (2007) Plant interactions with microbes and insects: from molecular mechanisms to ecology. *Trends Plant Sci* 12:564-569

Pineda A, Dicke M, Pieterse CM, Pozo MJ (2013) Beneficial microbes in a changing environment: are they always helping plants to deal with insects? *Funct Ecol* 27:574-586

Pineda A, Kaplan I, Hannula SE, Ghanem W, Bezemer TM (2020) Conditioning the soil microbiome through plant-soil feedbacks suppresses an aboveground insect pest. *New Phytol* 226:595-608

Puri A, Padda KP, Chanway CP (2018) Evidence of endophytic diazotrophic bacteria in lodgepole pine and hybrid white spruce trees growing in soils with different nutrient statuses in the West Chilcotin region of British Columbia, Canada. *For Ecol Manag* 430:558-565

Puri A, Padda KP, Chanway CP (2020) Can naturally-occurring endophytic nitrogen-fixing bacteria of hybrid white spruce sustain boreal forest tree growth on extremely nutrient-poor soils? *Soil Biol Biochem* 140:107642

Rasmann S, Bennett A, Biere A, Karley A, Guerrieri E (2017) Root symbionts: Powerful drivers of plant above-and belowground indirect defenses. *Insect Sci* 24:947-960

Reymond P, Farmer EE (1998) Jasmonate and salicylate as global signals for defense gene expression. *Curr Opin Plant Biol* 1:404-411

Seskar M, Shulaev V, Raskin I (1998) Endogenous methyl salicylate in pathogen-inoculated tobacco plants. *Plant Physiol* 116:387-392

Shade A, Peter H, Allison SD, Baho D, Berga M, Bürgmann H, Huber DH, Langenheder S, Lennon JT, Martiny JB, Matulich KL (2012) Fundamentals of microbial community resistance and resilience. *Front Microbiol* 3:417

Shen W, Le S, Li Y, Hu, F (2016) SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file manipulation. *PLoS One* 11:e0163962

Steiner AA (1979) The selective capacity of plants for ions and its importance for the composition and treatment of the nutrient solution. In Symposium on Research on Recirculating Water Culture 98:87-98

Suzuki N, Rivero RM, Shulaev V, Blumwald E, Mittler R (2014) Abiotic and biotic stress combinations. *New Phytol* 203:32-43

Van Loon LC, Van Strien EA (1999) The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiological and Molecular Plant Pathol* 55: 85-97


Wang B, Guo XW, Li K, Han X, Xu SJ, Liu ZD, Guo YS, Xie HG (2015) Effects of Salicylic acid on grape plants and the soil microbial community. *Allelopathy* J36:49-61

Wubs EJ, Bezemer TM (2018) Plant community evenness responds to spatial plant-soil feedback heterogeneity primarily through the diversity of soil conditioning. *Funct Ecol* 32:509-521


Yalpani N, Silverman P, Wilson TM, Kleier DA, Raskin I (1991) Salicylic acid is a systemic signal and an inducer of pathogenesis-related proteins in virus-infected tobacco. *Plant Cell* 3:809-818

Figures

i) No effect of SA and time




ii) SA effect only



iii) Time effect only



iv) SA × time effect



generation

- 1
- 2
- 3
- 4

treatments

- control
- ▲ SA

Figure 1

Conceptual figure showing the potential effects of SA application on *J. vulgaris* over four generations. i) No effect of SA and time. The bacterial community does not differ between SA-treated and control plants, and does not differ over time. ii) SA effect only. The bacterial community is affected by SA application but the effect does not differ over time. iii) Time effect only. The bacterial community changes over time, but is not affected by the SA treatment. iv) SA × time effect. The bacterial community is affected by SA application but these effects differ among generations.




Figure 2

Multivariate analysis of the bacterial community in soil samples from SA-treated and control plants grown in four generations. Shown are sample scores from a Nonmetric Multi-Dimensional Scaling (NMDS) plot (a), and an Orthogonal Projections to Latent Structures Discriminant analysis (OPLS-DA) plot (b) from the 24 rhizosphere soil samples.




Figure 3

Venn diagrams showing the number of unique and shared bacterial genera in the rhizosphere soil samples of SA-treated and control plants. The diagram in (a) is based on an analysis of genera that occur in all generations of the SA treatment, in (b) each generation is analyzed separately, (c) shows the diagram for all generations combined for the SA and control treatment.




Figure 4

Local “immigration” and “extinction” of bacterial genera in the rhizosphere soil of SA-treated and control plants over time. For each of two consecutive generations, shown are the number of genera present only in the first of those generations (i.e. representing genera that go extinct), present in both generations, and present only in the second of those generations (i.e. representing genera that immigrate). Genera were considered present in a treatment when present in at least two of the tree replicates. G1, G2, G3 and G4 represent the 1st, 2nd, 3rd and 4th generation.




Figure 5

Bar chart showing the relative abundance expressed as % (mean \pm SE) in control (left) or SA treated (right) plants of the genera that were significantly up or down regulated in at least one generation. The relative abundance is shown for all four generations. The significance is based on a student t-test with a false discovery rate (FDR) adjusted P values (< 0.05). Genus names and generation code are listed on the left. RA represents relative abundance of read counts.




Figure 6

Heatmap with a hierarchical clustering analysis of all the genera of rhizosphere soil of SA-treated plants and control plants in the 24 samples. The hierarchical clustering was calculated with Pearson distance and the Ward clustering algorithm based on the relative abundance of each genus. The color code represents the row-centered relative abundance. SA1, SA2, SA3 and SA4 represent SA treatments from the 1st generation, 2nd generation, 3rd generation and 4th generation. Control 1, control2, control3 and

control4 represent control treatments from the 1st generation, 2nd generation, 3rd generation and 4th generation.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarytable123.docx](#)
- [s1.jpg](#)
- [s2.jpg](#)
- [s3.jpg](#)
- [s4.jpg](#)
- [s5.jpg](#)
- [s6.jpg](#)
- [s7.jpg](#)
- [s8.jpg](#)