Use of Shockwave intravascular lithotripsy for the treatment of symptomatic and strongly calcified superior mesenteric artery stenosis

Oscar Balboa Arregui (✉ obalboa@saludcastillayleon.es)

Hospital de León: Complejo Asistencial Universitario de León <https://orcid.org/0000-0002-5850-0898>

Carmen Seoane Pose

Hospital Clínico Universitario de Santiago de Compostela

Maria Balboa Alonso

Hospital San Juan de Alicante

Teresa Bolaño Pampín

Hospital Clínico Universitario de Santiago de Compostela

Research Article

Keywords: Intravascular lithotripsy, Superior mesenteric artery stenosis, Chronic mesenteric ischemia

Posted Date: February 11th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-159948/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: We present the use of intravascular lithotripsy for the definitive treatment of highly calcified superior mesenteric artery stenosis.

Case presentation: A 67-year-old diabetic man had chronic postprandial abdominal pain and weight loss. Computed tomography angiography revealed highly calcified stenosis of the superior mesenteric artery. Selective angiography confirmed severe stenosis. A Shockwave lithotripsy balloon catheter was successfully used via brachial access to modify calcified plaque and increase vascular lumen. After twelve months of follow-up the patient had gained weight and had no abdominal postprandial pain.

Conclusion: Intravascular lithotripsy could be considered a new treatment modality to modify calcified lesions in the visceral arteries. More controlled studies are needed to demonstrate the efficacy, safety and feasibility of this new technology.

Level of Evidence: 4, Case Series

Background

Clinical manifestations of chronic mesenteric ischemia (CMI) may present widespread symptoms such as postprandial abdominal pain and weight loss when there is a 60–75% reduction in blood flow. Superior mesenteric artery (SMA) stenosis is a common finding in elderly patients with atherosclerosis. Hansen found that the prevalence of significant stenosis in subjects over 65 was 17.5% [1]. Image tests as computed tomography angiography (CTA) or lateral projection angiography are needed to visualize the narrowing of the SMA. If CMI is diagnosed there are two therapeutic options: surgical revascularization or percutaneous transluminal angioplasty (PTA) with or without stenting [2, 3]. The rate of survival shows that both treatments are successful. PTA, less invasive procedure, has associated complications such as thrombosis, restenosis and dislocation of stent [4].

Intravascular lithotripsy (IVL) therapy was applied as a definitive treatment for vascular stenosis in a patient who had postprandial abdominal pain and weight loss due to intense calcification and severe stenosis in the proximal segment of SMA. We confirm the written informed consent of the patient before the start of treatment.

Case Presentation

The symptoms of the 67-year-old man included 9 months of postprandial abdominal pain, most severe in the last 2 months and 10 kilograms of weight loss. He comorbidities include diabetes and revascularized ischemic cardiomyopathy 7 years ago. CTA revealed an extensive atherosclerotic disease involving aortoiliac, celiac and SMA calcification. Selective angiography revealed severe stenosis in the proximal segment of the SMA and hepatic artery originating in the same SMA (Fig. 1).

The patient was with antiplatelet therapy. Percutaneous endovascular intervention of SMA was performed under sedation. Vascular access was established in the right brachial artery with 6 Fr flexor guide sheath 80 cm long (Cook Medical, Bloomington, IN, USA) advanced towards the abdominal aorta with distal end in front of the SMA origin. A 110 cm long Shockwave Lithotripsy M5 6x60 mm balloon catheter (Shockwave Medical Inc., Santa Clara, California) was then placed through the SMA ostium, and 5 of 1-pulse treatments were delivered to activate only the distal emitter and facilitate calcium fragmentation and advancement of the IVL balloon catheter (Fig. 2). Once the balloon catheter had been advanced and covered all stenosis, 5 complete cycles of 30 pulses were applied each cycle with good angiographic result (Fig. 3). The patient tolerated the procedure without complications and went home 24 hours later. A month later he reported no complications after percutaneous endovascular intervention and absence of abdominal pain. Twelve months after the procedure, the patient gained weight and has not complained again of postprandial abdominal pain.

Discussion

The Shockwave Lithoplasty® System IVL consists of three components: the balloon catheter for performing the IVL, the connecting cable for pulse therapy, and a power generator. The IVL catheter consists of an OTW balloon catheter, which has, inside, several emitters of sonic shock waves whose diffusion generates a pressure of 50 atm. Activation of the device applies pulsed mechanical energy transmitted in the form of shock waves that pass through soft tissue without altering it and without causing damage to the vessel. Upon reaching the calcified plaque, the transmitted energy generates microfractures, modifying the calcium and therefore the morphology of the artery, allowing a posterior dilation with the balloon at low pressure (6 atm).

The application of IVL technology has been reported in coronary and peripheral endovascular procedures, including the iliac, femoral, and renal arteries. Currently, several multicenter studies are being conducted to evaluate the safety (DISRUPT CAD II) and efficacy (DISRUPT CAD / PAD III) of IVL in the coronary and peripheral territories [5–8]. The first studies on its peripheral use show promising results confirming a latency of 100% after 30 days, 70–80% at 6 months and 70% at 12 months, with a 95% clinical success rate, indicating less than 50% residual stenosis after the procedure with no evidence of acute adverse cardiac events, embolizations, thrombus formations, or perforations (6,8).

Treatment of symptomatic CMI is widely accepted to prevent acute mesenteric ischemia. There is no documentation on the use of IVL in the treatment of atherosclerotic disease presenting as mesenteric ischemia. Open surgical repair has survival rates of approximately 80–90% [9]. Intravascular repair has become an alternative to open surgery. One study showed that although primary latency rates were below 45%, secondary latency rates were comparable to 94%. The presence of highly calcified ostial occlusions represents an important limitation of endovascular interventions [10].

This report with the application of IVL therapy demonstrates successful revascularization in lesions that combine calcium thickness and a large calcium arc (270° of the vessel circumference), making dilation

difficult with conventional therapy.

Conclusion

IVL is an emerging treatment modality that dilates calcified lesions endovascularly. Its use in the treatment of highly calcified SMA in patients with postprandial pain should be considered as a novel approach for patients whose traditional endovascular treatment modalities are not optimal. More controlled studies are needed to demonstrate the efficacy, safety, and feasibility of this new technology.

Abbreviations

CMI: chronic mesenteric ischemia.

SMA: superior mesenteric artery.

CTA: computed tomography angiography.

PTA: percutaneous transluminal angioplasty.

IVL: intravascular lithotripsy.

Declarations

Funding: This study was not supported by any funding.

Conflict of interest: The authors declare that they have no conflict of interest.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. Institutional review board approval was obtained. This article does not contain any studies with animals performed by any of the authors.

Informed consent: Informed consent was obtained from all individual participants included in the study.

Consent for publication: Consent for publication was obtained for every individual person's data included in the study.

Availability of data and materials: Not applicable.

Authors' contributions:

All authors contributed to the conception of the work, writing/revision of the manuscript and approved the final version of the manuscript. No funding for any author.

Corresponding author: Correspondence to Oscar Balboa Arregui

Acknowledgements: Not applicable.

References

1. -Hansen KJ, Wilson DB, Craven TE, Pearce JD, English WP, Edwards MS et al (2004 Jul;40(1):45–52) Mesenteric artery disease in the elderly. *J Vasc Surg.* doi: 10.1016/j.jvs.2004.03.022. PMID: 15218461
2. -Leke MA, Hood DB, Rowe VL, Katz SG, Kohl RD, Weaver FA (2002 Dec;68(12):1088-92) Technical consideration in the management of chronic mesenteric ischemia. *Am Surg.* PMID: 12516815
3. –Sreenarasimhaiah J Chronic mesenteric ischemia. *Curr Treat Options Gastroenterol* (2007 Feb;10(1):3–9) doi: 10.1007/s11938-007-0051-x. PMID: 17298759
4. -Sakorafas GH, Sarr MG, Peros G (2008 Feb;206(2):349 – 56) Celiac artery stenosis: an underappreciated and unpleasant surprise in patients undergoing pancreaticoduodenectomy. *J Am Coll Surg.* doi: 10.1016/j.jamcollsurg.2007.09.002. PMID: 18222391
5. -Adams G, Shammass N, Mangalmurti S, Bernardo NL, Miller WE, Soukas PA et al. Intravascular Lithotripsy for Treatment of Calcified Lower Extremity Arterial Stenosis: Initial Analysis of the Disrupt PAD III Study. *J Endovasc Ther.* 2020 Jun;27(3):473–480. doi: 10.1177/1526602820914598. PMID: 32242768
6. -Brinton TJ, Ali ZA, Hill JM, Meredith IT, Maehara A, Illindala U et al (2019) Feasibility of Shockwave Coronary Intravascular Lithotripsy for the Treatment of Calcified Coronary Stenoses. *Circulation.* Feb 5;139(6):834-6. doi: 10.1161/CIRCULATIONAHA.118.036531. PMID: 30715944
7. -Brodmann M, Werner M, Brinton TJ, Illindala U, Lansky A, Jaff MR et al. Safety and performance of lithoplasty for treatment of calcified peripheral artery lesions. *J Am Coll Cardiol* 2017;70:908–10. doi:10.1016/j.jacc.2017.06.022. PMID: 28797363
8. -Armstrong EJ, Soukas PA, Shammass N, Chamberlain J, Pop A, Adams G et al (2020 Oct;21(10):1262–1268) Intravascular Lithotripsy for Treatment of Calcified, Stenotic Iliac Arteries: A Cohort Analysis From the Disrupt PAD III Study. *Cardiovasc Revasc Med.* doi: 10.1016/j.carrev.2020.02.026. PMID: 32147133
9. - Lejay A, Georg Y, Tartaglia E, Creton O, Lucereau B, Thaveau F et al. Chronic mesenteric ischemia: 20 year experience of open surgical treatment. *Eur J Vasc Endovasc Surg* 2015;49:587–92. doi: 10.1016/j.ejvs.2015.01.004. PMID: 25728455
10. -Van Petersen AS, Kolkman JJ, Beuk RJ, Huisman AB, Doelman CJA, Geelkerken RH et al. Open or percutaneous revascularization for chronic splanchnic syndrome. *J Vasc Surg* 2010;51:1309–16. doi: 10.1016/j.jvs.2009.12.064. PMID: 20304586

Figures

Figure 1

CTA revealed highly calcified stenosis in the proximal segment of SMA. Selective brachial access angiography confirmed severe SMA stenosis.

Figure 2

Shockwave Lithotripsy M5-6 x 60 mm, balloon catheter placed first through the SMA ostium, and total subsequent coverage of stenosis. Balloon inflated to 4 atm during energy emission.

Figure 3

Control arteriography post-treatment with IVL.