

Identification of *ATP2C1* mutations in the patients of Hailey-Hailey disease

xiaoli li

Xi'an Jiaotong University

dingwei zhang

Xi'an Jiaotong University

jiahui ding

Xi'an Jiaotong University

li li

Xi'an Jiaotong University

zhenghui wang (✉ ehui4298@163.com)

Xi'an Jiaotong University <https://orcid.org/0000-0002-5630-7277>

Research article

Keywords: Familial benign chronic pemphigus, ATP2C1, gene mutation, p63

Posted Date: May 21st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-16069/v6>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Medical Genetics on June 1st, 2020.
See the published version at <https://doi.org/10.1186/s12881-020-01056-4>.

Abstract

Background: Familial benign chronic pemphigus, also known as Hailey-Hailey disease (HHD), is a clinically rare bullous Dermatitis. However the mechanism has not been clarified. The study aim to detect novel mutations in exons of *ATP2C1* gene in HHD patients; to explore the possible mechanism of HHD pathogenesis by examining the expression profile of hSPCA1, miR-203, p63, Notch1 and HK α proteins in the skin lesions of HHD patients.

Methods: Genomic DNA was extracted from peripheral blood of HHD patients. All exons of *ATP2C1* gene in HHD patients were amplified by PCR and the products were purified and sequenced. All related signaling proteins of interest were stained by using skin lesion tissues from HHD patients and miR-203 levels were also determined.

Results: One synonymous mutation c.G2598A (in exon 26), one nonsense mutation c.C635A and two missense mutations c.C1286A (p.A429D) and c. A1931G (p. D644G) were identified. The nonsense mutation changed codon UCG to stop codon UAG, causing a premature polypeptide chain of the functional region A. The two missense mutations were located in the region P (phosphorylation region) and the Mn binding site of hSPCA1. The level of hSPCA1 was significantly decreased in HHD patients compared to the normal human controls, accompanied by an increase of miR-203 level and a decrease of p63 and HK α levels. **Conclusion:** In our study, we found four mutations in HHD. Meanwhile we found increase of miR-203 level and a decrease of p63 and HK α levels. In addition, Notch1, which was negatively regulated p63, is downregulated. These factors may be involved in the signaling pathways of HHD pathogenesis. Our data showed that both p63 and miR-203 may have significant regulatory effects on Notch1 in the skin.

Background

Familial benign chronic pemphigus, also known as Hailey-Hailey disease (HHD), is a clinically rare bullous Dermatitis, and mainly manifested as erosion, erythema, accompanied with blisters and pimples. The incidence of this disease is approximately 0.002% [1], and has no significant difference between men and women[2]. HHD is currently not curable, and symptomatic treatment is the main strategy taken so far to reduce symptoms and prevent disease recurrence[3-6], using cortisol to reduce inflammation[7], and using topical vitamins[8,9] to promote the differentiation of epidermal keratinocytes and regulate intercellular Ca²⁺ concentrations [10,11].

HHD is caused by mutations in the *ATP2C1* gene. *ATP2C1* is expressed in the skin as well as the brain, skeletal muscle, placenta, heart, and lungs. In humans, *ATP2C1* mutations rarely cause skin tumors, squamous cell carcinoma and basal cell carcinoma[12-13] in HHD lesions. Liver failure with HHD was ever reported[14]. There are also some cases that have emotional disorders have no desire in almost all activities[13]. It is believed that HHD development is due to insufficient gene dose of *ATP2C1*. The protein level of hSPCA1 (the human secretory pathway Ca²⁺/Mn²⁺-ATPase protein 1, hSPCA1) encoded by the

ATP2C1 in the Golgi apparatus is strikingly decreased, resulting in an increase in the intracellular Ca^{2+} concentration. As an important messenger, Ca^{2+} has significant impact on the maturation and differentiation of keratinocytes. Increased Ca^{2+} concentration eventually destroy cell-to-cell connections[1].

In this study, we searched for novel mutations of *ATP2C1* by sequencing the gene from 2 different HHD pedigrees and 2 sporadic cases. We analyzed the impact of these mutations on the structure and function of hSPCA1 by using bioinformatics tool. Meanwhile the expression levels of hSPCA1, miR-203, p63, Notch1, and HK α in the skin lesion tissues of HHD patients were examined.

Methods

Patients

The study subjects were HHD patients enrolled in the dermatology clinic of the Second Affiliated Hospital of Xi'an Jiaotong University, and pathological biopsy was performed on the patient's typical skin lesions to further confirm the diagnosis. All procedures were approved by the Institutional Human Experiment and Ethics Committee of the Second Hospital of Xi'an Jiaotong University. The samples from 2 patients in pedigree I, II and 2 sporadic patients were collected. The patient's information such as gender, age, and skin lesion performance, time of onset, and family history was collected.

HE staining

Tissues were selected from a typical site of the patient's skin lesions following a biopsy under local anesthesia. The removed tissues were placed in 4% formaldehyde solution and fixed at 4 °C for >4 hours; the tissues were then rinsed under water for 10 minutes and then dehydrated in a biological tissue dehydrator following standard procedures. The tissues were then immediately put into 60 °C paraffin, and poured into the embedding box. After the paraffin is completely cooled down, the tissue was sectioned with thickness of 5 μm . The tissue sections were deparaffined following to standard procedures[15].

DNA extraction

A total of 2 ml of peripheral venous blood was drawn from 2 patients with HHD from two different pedigrees and 2 sporadic patients following to the patient's consent. As controls, 2 ml of peripheral blood were collected from the patient's healthy family members and 100 normal people who were not biological related. The blood was collected in a 2% Ethylene Diamine Tetraacetic Acid(EDTA) anticoagulant tube. The DNA was immediately extracted or stored in a -20 °C refrigerator. DNA extraction from peripheral blood was performed according to the following protocols: 200 μl of proteinase K solution was added to 200 μl of fresh or thawed blood and mixed well, and then 200 μl of binding solution(CB) was added and mixed thoroughly by vortexing, and incubated at 70 °C for 10 minutes; 100 μl of isopropanol was added after the samples cooled down, vortexed immediately to mix thoroughly, and then put it into the AC

column, centrifuged at 13,000 rpm for 60 seconds, and then discarded the flow-through in the collection tube; the inhibitor remover (IR) and the rinse solution (WB) were added sequentially, centrifuged at 12,000 rpm for 30 seconds and discarded the flow-through. 100 µl of pre-warmed elution buffer EB was added onto the center of the adsorption membrane, and then centrifuged at 12,000 rpm to collect DNA solution; the DNA product was stored at 4 °C, or at -20 °C for long-term use.

ATP2C1 primers and PCR reactions

Primer sequences were designed and verified using Primer 5.0. All upstream and downstream primers cover some intron sequences on both sides of the exon. Primer synthesis and purification of PCR product were completed by Beijing Aoke Dingsheng Biotechnology Co., Ltd. PCR reaction: 2 × Taq Master MIX, 25 µl; DNA template, 5 µl; forward primer, 1 µl; reverse primer 1 µl; double distilled water, 18 µl; total volume, 50 µl. PCR protocol was summarized in Table 1. Protocol I was used for exon 1, protocol III was used for exon 5, 7, 12, 27 and 28 and protocol II was used for other exons[15]. The sequencing data of each exon was read by Chromas 2.4 and aligned with the original sequence of human *ATP2C1* gene.

IHC staining

The skin tissues collected from patients with HHD were cut into thickness of 4-5 µm by using a paraffin microtome. The sections were deparaffined and rehydrated following to standard procedures. Hydrogen peroxide (0.3%) was added onto the sections to block endogenous peroxidase. The sections were then incubated with 0.1% saponin for 30 min; 10% normal goat serum was added onto the sections and incubated at 37 °C for 30 min to cover non-specific sites; the blocking serum was then discarded and the primary antibody (hSPCA1, p63, Notch1, HK α ; 1:50 dilution) was added and incubated at 4 °C for 24-72 hours; secondary antibodies (goat anti-rabbit or goat anti-mouse) labeled with horseradish peroxidase was added and incubated at 37 °C for 30~40 min and then washed with PBS for 3 times, 5~10 min each time[15].

Real time-PCR

Total RNA was extracted and cDNA was synthesized according to the kit instructions. The primers were synthesized by Beijing Aoke Dingsheng Biotechnology Co., Ltd. according to the following sequences. miR-203: Forward 5'- ttgagttagggttattttgtgt -3 ' ; Reverse 5'-ctaaccaaccaattttccaa -3'. The PCR reaction: 2 × Taq Master Mix, 10 µl; cDNA, 2 µl; forward primer; 0.2 µl; reverse primer, 0.2 µl; DEPC water, 7.4 µl; total volume, 20 µl. PCR protocol: pre-denaturation, 95 °C, 3 min; denaturation, 95 °C, 12 s; annealing, 62 °C, 35 s; extension, 74 °C, 3 s; 40 cycles; extension, 74 °C, 5 min. The expression level of each gene was determined by relative quantification, and GAPDH included in the kit was used as an endogenous control. The relative expression level was determined by using the Ct value. Relative expression level = $2^{-\Delta\Delta CT}$.

Results

Pedigree analysis and clinical data of familial benign chronic pemphigus

Pedigree analysis and clinical data of pedigree Ⅱ

Among 4 generations of the pedigree, there are 2 HHD patients (1 male and 1 female) who were present in the 2nd and 3rd generations, consistent with the inheritance pattern of autosomal dominant genetic disease (Figure 1A). The proband is a 50 year-old woman whose parents were not biological related. Physical exam showed that she had good health condition in general, and no abnormality was found in other body systems. Clinical manifestations included erythema, blisters, erosions under bilateral axilla, left groin and umbilicus, and slight hypertrophy of skin in the left groin (Figure 1B). Histopathological examination of typical skin lesions in the left groin revealed cracks in the epidermis and partial loss of the stratum corneum. The spines in the lower and middle layers of epidermis were loosened and cells of spinous layer were visible (Figure 1C).

Pedigree analysis and clinical data of family Ⅱ

Among 3 generations of the pedigree, there were 3 patients, one male and two females. The patients were present in two consecutive generations Ⅱ and Ⅲ, consistent with the inheritance pattern of autosomal dominant genetic disease (Figure 1D). The proband was a 43 year-old male whose parents were not blood related. Physical exam showed that he had good general health condition without abnormality found in other body system. Clinical manifestations included flaky erythematous papule under the left armpit and left groin, perianal area, left popliteal fossa, adjacent popliteal fossa, obvious erosion in addition to some erythema (Figure 1E). The left axillary skin lesions were taken for pathological examination. The results showed that the spines in the lower and middle layers of epidermis were loosened, and were overflowed on the top with red blood cells. A single layer of basal cells covered on the dermal papilla and formed a villi (Figure 1F).

Clinical data of sporadic case I

Among 3 generations of the pedigree, there was only 1 patient. No other patients in the pedigree were seen. No obvious sign of familial genetic disease was observed. The patient was a 68 year-old male whose parents were not blood related. Physical exam showed that this patient was generally in a good health condition without obviously abnormalities found in other body systems. Clinical manifestations included erythema on both sides of the groin and anus, scattered papules in addition to the anal erythema, obvious erosion and exudation in the skin lesions (Figure 2A). A piece of skin from lesions on the left groin was taken for histological exam. The results showed that spines in the middle and lower layers of the epidermis were loosen and formed fissures in which loosening cells were presented (Figure 2B).

Clinical data of sporadic cases Ⅱ

Among 3 generations of the pedigree, there was 1 patient only. No other patients in the pedigree were seen. No obvious inheritance pattern of familial genetic diseases was shown. The patient was a 42 years old male whose parents were not blood related. Physical exam showed that this patient

had good general health condition without other abnormalities found in other body systems. Clinical manifestations included slight erythema and desquamation on the back of right ear, erythematous papular rash on both sides of the groin, obvious erosion and exudation in addition to some erythema, and epidermal lesions on the left groin covered with crust (Figure 2C). Histopathological examination of the typical skin lesions from the left groin showed that the spinous layer in the epidermis formed fissures in which the loosening cells were scattered around (Figure 2D).

Gene sequencing of *ATP2C1*

PCR amplification for the exons of *ATP2C1*

Exon 26 of *ATP2C1* in pedigree I, exon 15 of *ATP2C1* in pedigree II, exon 8 of *ATP2C1* in sporadic patient I, and exon 22 of *ATP2C1* in sporadic patient II were amplified (Figure 3A). The sizes of PCR products were 312 bp, 383 bp, 425 bp and 298 bp respectively.

The sequencing result of exon 26 of *ATP2C1* in pedigree I

A synonymous mutation c.G2598A in exon 26 was found in the proband of pedigree I, which was further confirmed by reverse sequencing that showed a complementary nucleotide mutation. The mutation did not change amino acid code of lysine. We further verified this mutation was actually a nonsense mutation that was not previously reported (Figure 3B). Such mutation was not found in the normal members of this pedigree or other people who were not blood related (Figure 3C, D, E).

Sequencing results of *ATP2C1* exon 15 in pedigree II

A missense mutation c.C1286A was detected in the exon 15 of the proband in pedigree II, which was further confirmed by reverse sequencing that showed a complementary nucleotide mutation. This mutation changed the original alanine to aspartic acid (p.A429D). We further verified that it was actually an unreported missense mutation (Figure 4A-D). Such mutation was not found in the normal members of this pedigree or other people who were not blood related (Figure 4B, C).

Sequencing results of *ATP2C1* exon 8 in sporadic case I

A nonsense mutation c.C635A in exon 8 was detected in the proband of sporadic case I. This mutation was further confirmed by reverse sequencing that showed a complementary nucleotide mutation. This mutation resulted in a premature stop codon in the polypeptide chain and was previously reported (Figures 5A-D). Such mutation was not seen in the normal human controls (Figure 5B, C).

Sequencing results of *ATP2C1* exon 22 in sporadic case II

A missense mutation c. A1931G was detected in exon 22 of the proband in sporadic case II, which was further confirmed by reverse sequencing that showed a complementary nucleotide mutation. This mutation changed the original 644th aspartic acid (GAU) to be Glycine (GGU) (p. D644G). We noticed that

this mutation was previously reported (Figure 6A-D). Such mutation in the gene was not found in the normal members in the pedigree and other people who were not blood related (Figure 6B, C).

IHC staining for hSPCA1

hSPCA1 is mainly expressed in the cytoplasm of keratinocytes in the epidermal layer with some on the cell membrane. Positive staining appeared as yellow or tan particles or clumps. In normal human skin tissues (Figure 7A-a), hSPCA1 showed strong positive signals, that is, a dark brown stained band in the epidermis. The negative control in which the hSPCA1 antibody was replaced with PBS (Figure 7A-a, right corner inset) did not show any nonspecific staining. The expression levels of hSPCA1 in the skin tissues of all patients with HHD (Figure 7A-b/c/d, E) were significantly lower than that of the positive control, especially at the typical skin lesions (acanthosis). No obvious staining signal was seen even in some local areas (Figure. 7A-c).

IHC staining for proteins in relevant signaling pathways

IHC for p63

P63 is an intranuclear protein and can be stained in the nucleus of keratinocytes in the epidermal layer. In normal human skin tissues, p63 staining was strongly positive in the nucleus. In the area adjacent to the stratum corneum the nucleus gradually disappeared with differentiation, therefore, the staining is mainly present in the basal layer (Figure 7B-a). PBS was used as a negative control, which did not show any non-specific staining (Figure 7B-a, right corner inset). The expression level of p63 protein in the nucleus of HHD patients was significantly reduced compared to the positive control, especially in the typical lesions (Figure 7B-b/c/d, E). No obvious staining was seen in some local areas of the tissue.

IHC for Notch1

Notch1 is expressed in the cytoplasm of keratinocytes in the epidermal layer. In normal human skin tissues, Notch1 was strongly stained, appeared as brown or tan particles or clumps (Figure 7C-a). In the entire epidermal layer, Notch1 showed a even deep-stained band. In the skin tissues of all HHD patients, Notch1 signals (Figure 7C-b/c/d, E) were weaker compared to the positive control. In contrast, the negative control by using PBS instead of Notch1 antibody did not show any non-specific staining (Figure 7C-a, right corner inset).

IHC for HK α protein

HK α is expressed in the cytoplasm of the keratinocytes in the epidermal layer. In normal human skin tissues, HK α was strongly stained (Figure 7D-a) and showed a even deep-stained band in the entire epidermal layer. In contrast, the expression level of HK α in the cytoplasm of skin lesion tissues was significantly reduced compared to the normal human positive control (Figure 7D-b/c/d, E). The negative control by using PBS other than HK α antibody (Figure 7D-a, right corner inset) did not show obviously non-specific staining.

Real time PCR for miR-203

RNA was extracted from the skin lesion tissues of HHD patients and real-time PCR was performed to detect the expression level of miR-203 in these tissues. Normal human skin tissue was used as a control. The data showed that the levels of miR-203 in the skin lesions of HHD patients were significantly upregulated compared to that in the normal human control (Figure 8).

Discussion

In this study, all patients (including all pedigree cases and sporadic cases) we collected have typical clinical manifestations and histopathological characteristics. Pedigree I and pedigree II showed vertical transmission and presented in an inheritance pattern of autosomal dominant genetic disease. By using gene sequencing and exon alignment of *ATP2C1* from 4 patients, we found 4 different heterozygous mutations, including 1 synonymous mutation c.G2598A, 2 missense mutations c.C1286A and c. A1931G, and 1 nonsense mutation c.C635A.

In the patient of pedigree I, the synonymous mutation c.G2598A in exon 26 changed the codon from AAG to AAA, but did not change the amino acid (lysine). In the patient of pedigree II, the missense mutation c.C1286A (p.A429D) in exon 15 changed the alanine to aspartic acid. Protein structural analysis showed that this mutation was located in the P region (phosphorylation region), the most important functional site of hSPCA1 to which $\text{Ca}^{2+}/\text{Mn}^{2+}$ and ATP are bound. This mutation changed the primary structure of the protein and disrupted the catalytic site of ATPase, phosphorylation function and protein conformation, leading to loss of protein function and transport failure of $\text{Ca}^{2+}/\text{Mn}^{2+}$ and abnormal concentrations of intracellular $\text{Ca}^{2+}/\text{Mn}^{2+}$ in the cells.

In keratinocytes, hSPCA1 not only transports Ca^{2+} , but also competitively transports one Mn^{2+} from the cytoplasm to the Golgi apparatus, which is nonreplaceable by other calcium pumps. It is known that the cells are very sensitive to Mn^{2+} concentration and a slight change of Mn^{2+} concentration may affect cell metabolism seriously[16]. Excessive intracellular Mn^{2+} may prevent Mg^{2+} binding to proteins, compromise the fidelity of DNA polymerase, and disrupt the transport function of cell membranes, thus affecting the physiological conditions of human body[17,18]. In sporadic case I, a nonsense mutation c.C635A in exon 8 changed the serine codon UCG (position 212) to a stop codon UAG. This mutation is located in the A region of regulatory element and results in premature polypeptide chain. In addition, the mRNA with this nonsense mutation is easily degraded, reducing the expression level of truncated hSPCA1 that lacks important structures in the calcium pump. In sporadic case II, a missense mutation c. A1931G in exon 22 changed aspartate (GAU, at position 644) to glycine (GGU) (p. D644G). This mutation is located at the Mn^{2+} binding site of hSPCA1[19] and disrupts the binding of hSPCA1 to Mn^{2+} , thus interfere the metabolism of intracellular Mn^{2+} [20], affect the activity of DNA polymerase and the transport function of cell membranes, and increase the potential of mutation in other genes[21]. hSPCA1 is highly expressed in the skin and kidney tissues, localized in cytoplasmic Golgi apparatus in the cells. The clinical symptoms of *ATP2C1* mutation are mainly limited to the skin of the patients. Therefore, we chose

skin lesions to investigate dysregulated signaling pathways in HHD patients. Immunohistochemistry data showed that the expression levels of hSPCA1 in the skin tissues of all HHD patients was significantly downregulated compared to normal human controls.

MiR-203 is a family member of tumor suppressor genes. It is a type of short, non-coding single-stranded miRNA [22]. MiR-203 has the highest abundance in the skin, and is exclusively expressed in keratinocytes, indicating that it may be a key to various biological functions of keratinocytes[23]. When keratinocytes treated with high concentration of Ca^{2+} , miR-203 was upregulated > 3-fold[23,24]. In the keratinocytes of HHD patients, Ca^{2+} concentration is strikingly increased due to the dysfunction of the calcium pump. In our study, miR-203 in the skin lesions of patients showed a significant upregulation.

The transcription factor p63 is a member of the p53 tumor suppressor family[25] and plays an important role in the regulation of epidermal growth, keratinocyte differentiation, cell adhesion and cell migration[26]. During keratinization, the expression level of *ATP2C1* is strikingly decreased, so does p63. Numerous studies have shown that p63 regulates downstream target genes and also plays roles in a variety of intracellular signaling pathways[27,28]. p63 and Notch1, two important factors that regulate cell proliferation and differentiation, cell adhesion and other functions in keratinocytes, are negatively regulated to each other[23]. Mature miR-203 binds to p63 3'-UTR and downregulate p63[29]. In animal experiments, miR-203 affected the "stemness" of mouse skin stem cells by targeting p63 expression, thus enhancing the differentiation capacity of basal cells[30,31]. MiR-203 can also inhibit Notch1 expression by downregulating ligand Jagged1 in the Notch1 pathway[32,33]. In our study we found p63 was significantly downregulated compared to the normal human controls. From previous studies and ours, it is believed that increased intracellular Ca^{2+} concentration is actually accompanied by upregulation of miR-203 and downregulation of p63. We also found that the expression level of Notch1, which is negatively regulated by p63, only showed a slight decrease. Our data suggest that p63 and miR-203 are both associated with Notch1 in the skin lesions.

Hexokinase is a key enzyme in the first step of glycolytic pathway, catalyzing glucose into glucose 6-phosphate. In human cells, hexokinase has four subtypes (I-IV), of which HKII is the one present in human keratinocytes and regulates metabolism of glucose and lactate. Therefore, inhibiting HKII activity can effectively block sugar metabolism. HKII gene is a target gene of p63, and p63 downregulation can reduce 80% of HKII mRNA. In keratinocytes, the p63-HKII axis is an important pathway regulating cell metabolism and proliferation. Downregulation of HKII decreased ATP production, which in turn further compromised hSPCA1 function that is greatly dependent on energy supply[26]. Our result showed that the level of HKII in the tissues of HHD patients were obviously decreased to about 60% of the normal value and demonstrates that increased Ca^{2+} concentration caused by *ATP2C1* mutation does result in decreased HKII, which reduces ATP production and affects normal physiological function of the human body.

Conclusion

In this study, we detected four *ATP2C1* mutations in HHD patients, which has further enriched the pathogenic mutation pool of HHD. We showed that miR-203 was upregulated in HHD patients, which in turn might downregulate p63 and Notch1, and downregulation of p63 might further decrease the level of HKK. The exact mechanism underlying HHD pathogenesis remains unclear and needs further investigation.

Declarations

Ethics approval and consent to participate

The study was approved by ethical committee of the Second Hospital of Xi'an Jiaotong University(number: ECX2013054). All participants signed informed consent.

Consent for publication

Written informed consent for publication of personal details and clinical details was obtained from all of the participants.

Availability of data and materials

The datasets generated and/or analysed during the current study are available in the https://www.jianguoyun.com/p/DS_N9qkQ7Ym5CBjJ-ZkD. All data generated or analysed during this study can be easily acquired from corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interest.

Funding

The work was funded by Shaanxi Province Government, Grant/Award Number:2014KJXX-32; 2020SF-177. Natural Science Foundation of China, Grant/Award Numbers: 30901297. Fundamental Research Funds for the Central Universities of Xi'an Jiaotong University, Grant/Award Numbers: 2019-53. The funders had no role in the study design, data collection, data analysis, interpretation of results, and writing of the manuscript.

Authors' contributions

XLL, DWZ and ZHW designed the study. JHD and LL performed the molecular analysis. XLL collected patients' data. DWZ performed statistical analysis. XLL, DWZ and ZHW prepared the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We thanked the technical support of Yanfei Zhang and Xiaoge Zhao.

Author details

¹Department of Dermatology, The Second Affiliated Hospital, Xi'an Jiaotong University, Xi'an, 710004, China; ²Department of Otolaryngology-Head and Neck Surgery, The Second Affiliated Hospital, Xi'an Jiaotong University, Xi'an, China, 710004.

Abbreviations

Hailey-Hailey disease (HHD), the human secretory pathway $\text{Ca}^{2+}/\text{Mn}^{2+}$ -ATPase protein 1 (hSPCA1), Ethylene Diamine Tetracetic Acid (EDTA), inhibitor remover (IR), the rinse solution (WB), hematoxylin-eosin staining (HE), Polymerase Chain Reaction (PCR) and Immunohistochemistry (IHC).

References

1. Zhang DW, Huo J, Li RL, Zhang YF, Wang ZH, Li XL. Altered levels of focal adhesion and extracellular matrix-receptor interacting proteins were identified in Hailey-Hailey disease by quantitative iTRAQ proteome analysis. *J Cell Biochem.* 2019;120(3):3801-12.
2. Cheng Y, Cheng YM, Zhao G, Jia MC. A novel missense mutation of the ATP2C1 gene in a Chinese patient with Hailey-Hailey disease. *Biochem Biophys Res Commun.* 2011;406:420-2.
3. Voisset C, Garcia-Rodriguez N, Birkmire A, Blondel M, Wellinger RE. Using yeast to model calcium-related diseases: example of the Hailey-Hailey disease. *Biochimica et biophysica acta.* 2014;1843:2315-21.
4. Burge SM. Hailey-Hailey disease: the clinical features, response to treatment and prognosis. *Br J Dermatol.* 1992;126:275-82.
5. Le Sache-de Peufeuilhoux L, Raynaud E, Bouchardeau A, Fraitag S, Bodemer C. Familial benign chronic pemphigus and doxycycline: a review of 6 cases. *J Eur Acad Dermatol Vene.* 2014; 28:370-3.
6. Almeida L, Grossman ME. Benign familial pemphigus complicated by herpes simplex virus. *Cutis.* 1989;44:261-2.
7. Ikeda S, Suga Y, Ogawa H. Successful management of Hailey-Hailey disease with potent topical steroid ointment. *J Dermatol Sci.* 1993;5:205-11.
8. Jeon SY, Ha SM, Ko DY, Song KH, Kim KH. Hailey-hailey disease treated with topical tacalcitol. *Ann Dermatol.* 2013;25:389-90.
9. Rajpara SM, King CM. Hailey-Hailey disease responsive to topical calcitriol. *Br J Dermatol.* 2005;152:816-7.
10. Vanderbeck KA, Giroux L, Murugan NJ, Karbowski LM. Combined therapeutic use of oral alitretinoin and narrowband ultraviolet-B therapy in the treatment of hailey-hailey disease. *Dermatol Rep.* 2014; 6:5604.

11. Norman R, Greenberg RG, Jackson JM. Case reports of etanercept in inflammatory dermatoses. *J Am Acad Dermatol.* 2006;54:S139-42.
12. Micaroni M, Giacchetti G, Plebani R, Xiao GG, Federici L. ATP2C1 gene mutations in Hailey-Hailey disease and possible roles of SPCA1 isoforms in membrane trafficking. *Cell death & disease.*2016; 7:e2259.
13. Holst VA, Fair KP, Wilson BB, Patterson JW. Squamous cell carcinoma arising in Hailey-Hailey disease. *J Am Acadof Dermatol.*2000;43:368-71.
14. Amagai M, Kobayashi M, Wakabayashi K, et al. A case of generalized Hailey-Hailey disease with fatal liver injury. *Keio J Med.* 2001;50:109-16.
15. Zhang DW, Li XL, Wang ZH, Zhang YF, Guo K, Wang S, Tu C, Huo J, Xiao SX. Hailey–Hailey disease: investigation of a possible compensatory SERCA2 up-regulation and analysis of SPCA1, p63, and IRF6 expression. *Arch Dermatol Res.* 2015;307:143–9.
16. Gao S, Chen X, Jin H, Ren S, Liu Z, Fang X, et al. Overexpression of ErbB2 renders breast cancer cells susceptible to 3-BrPA through the increased dissociation of hexokinase II from mitochondrial outer membrane. *Oncology letters.*2016; 11:1567-73.
17. Towler MC, Prescott AR, James J, Lucocq JM, Ponnambalam S. The manganese cation disrupts membrane dynamics along the secretory pathway. *Exp Cell Res.*2000;259:167-79.
18. Monk S, Sakuntabhai A, Carter SA, Bryce SD, Cox R, Harrington L, et al. Refined genetic mapping of the darier locus to a <1-cM region of chromosome 12q24.1, and construction of a complete, high-resolution P1 artificial chromosome/bacterial artificial chromosome contig of the critical region. *Am J Hum Genet.*1998;62:890-903.
19. Foggia L, Aronchik I, Aberg K, Brown B, Hovnanian A, Mauro T.M. Activity of the hSPCA1 Golgi Ca²⁺ pump is essential for Ca²⁺-mediated Ca²⁺ response and cell viability in Darier disease. *J Cell Sci.*2006;119:671-9.
20. Behne MJ, Tu CL, Aronchik I, Epstein E, Bench G, Bikle DD, et al. Human keratinocyte ATP2C1 localizes to the Golgi and controls Golgi Ca²⁺ stores. *J Inves Dermatol.*2003; 121:688-94.
21. Brini M, Carafoli E. Calcium pumps in health and disease. *Physiol Rev.* 2009;89:1341-78.
22. Wang K, Zhang ZW. Expression of miR-203 is decreased and associated with the prognosis of melanoma patients. *Int J Clin Exp Pathol.*2015; 8:13249-54.
23. Hildebrand J, Rutze M, Walz N, Gallinat S, Wenck H, Deppert W, et al. A comprehensive analysis of microRNA expression during human keratinocyte differentiation in vitro and in vivo. *J Invest Dermatol.*2011;131:20-9.
24. Nguyen BC, Lefort K, Mandinova A, Antonini D, Devgan V, Della Gatta G, et al. Cross-regulation between Notch and p63 in keratinocyte commitment to differentiation. *Genes & development.* 2006;20:1028-42.
25. Wu N, Rollin J, Masse I, Lamartinea J, Gidrol X. p63 regulates human keratinocyte proliferation via MYC-regulated gene network and differentiation commitment through cell adhesion-related gene

network. J Bio Chem.2012; 287:5627-38.

26. Viticchie G, Agostini M, Lena AM, Mancini M, Zhou H, Zolla L, et al. p63 supports aerobic respiration through hexokinase II. P Nat Acad Sci USA .2015;112:11577-82.
27. Truong AB, Kretz M, Ridky TW, Kimmel R, Khavari P.A. p63 regulates proliferation and differentiation of developmentally mature keratinocytes. Genes & development.2006; 20:3185-97.
28. McDade SS, Patel D, McCance D.J. p63 maintains keratinocyte proliferative capacity through regulation of Skp2-p130 levels. J Cell Sci.2011;124:1635-43.
29. Ma J, Meng Y, Kwiatkowski DJ, Chen X, Peng H, Sun Q, et al. Overexpression of ErbB2 renders breast cancer cells susceptible to 3-BrPA through the increased dissociation of hexokinase II from mitochondrial outer membrane. Oncology letters. 2016;11:1567-73.
30. Lena AM, Shalom-Feuerstein R, Rivetti di Val Cervo P, Aberdam D, Knight RA, Melino G, et al. miR-203 represses 'stemness' by repressing DeltaNp63. Cell Death Differ.2008; 15:1187-95.
31. Yi R, Poy MN, Stoffel M, Fuchs E. A skin microRNA promotes differentiation by repressing 'stemness'. Nature.2008;452:225-9.
32. Sasaki Y, Ishida S, Morimoto I, Yamashita T, Kojima T, Kihara C, et al. The p53 family member genes are involved in the Notch signal pathway. J Bio Chem.2002;277:719-24.
33. Kopan R, Ilagan MX. The canonical Notch signaling pathway: unfolding the activation mechanism. Cell. 2009;137:216-33.

Table 1

Table 1. PCR protocol

94 °C 5 min					
□	94 °C 45s	60°C 45s	72°C 45s	2x	1
	94 °C 45s	59°C 45s	72°C 45s	2x	
	94 °C 45s	58°C 45s	72°C 45s	2x	
	94 °C 45s	57°C 45s	72°C 45s	2x	
	94 °C 45s	56°C 45s	72°C 45s	2x	
	94 °C 45s	55°C 45s	72°C 45s	20x	
□	94 °C 45s	55°C 45s	72°C 45s	32x	Other exon
□	94 °C 30s	54°C 30s	72°C 30s	30x	5, 7, 12, 27, 28

Figures




Figure 1

Pedigree analysis and clinical data of pedigree I and II. (A) Pedigree chart of pedigree I. (B) Clinical manifestations of the proband in pedigree I. (C) Histopathological examination of typical skin lesion for the proband in pedigree I (5×)(Arrow shows acanthosis). (D) Pedigree chart of pedigree II. (E) Clinical manifestations of the proband in pedigree II. (F) Histopathological examination of typical skin lesion for the proband in pedigree II (5×) (Arrow shows acanthosis).




Figure 2

Clinical data of sporadic case I and II. (A) Clinical manifestations of sporadic case I. Skin lesions were present in the anus (a) and groin (b). (B) Histopathological examination of typical skin lesion for sporadic case I (5×) (Arrow shows acanthosis). (C) Clinical manifestations of sporadic case II. Skin lesions were present in the right back of the ear (a) and bilateral groin (b; c, left, d, right). (D) Histopathological examination of typical skin lesion for sporadic case II (5×) (Arrow shows acanthosis).




Figure 4

Exon 15 sequencing for the ATP2C1 gene in pedigree II. (A) Sequencing result of exon 15 for the proband in pedigree II. (B) Sequencing result of exon 15 for normal member in the same pedigree. (C) Sequencing result of exon 15 for normal individual who was not blood related. (D) A missense mutation c.C1286A(Bases C→A) in exon 15 of ATP2C1 gene in the Ensemble database (indicated by arrows).




Figure 5

Exon 8 sequencing for the ATP2C1 gene in sporadic case I. (A) Sequencing result of exon 8 in sporadic case I. (B) Sequencing result of exon 8 for normal member in the same pedigree. (C) Sequencing result of exon 8 for normal individual who was not blood related. (D) A nonsense mutation c.C635A(Bases C→A) in exon 8 of ATP2C1 gene in the Ensemble database (indicated by arrows).




Figure 6

Exon 22 sequencing for the ATP2C1 gene in sporadic case II. (A) Sequencing result of exon 22 in sporadic case II. (B) Sequencing result of exon 22 for normal member in the same pedigree. (C) Sequencing result of exon 22 for normal individual without blood relation. (D) A missense mutation c. A1931G(Bases A→G) in exon 22 of ATP2C1 gene in the Ensembl database (indicated by arrows).




Figure 7

(A) IHC staining for ATP2C1 [Negative control with PBS(a) and expression levels of hSPCA1 in HHD(b,c,d)]. (B) IHC staining for p63 [Negative control with PBS(a) and p63 protein in the nucleus of HHD patients(b,c,d)]. (C) IHC staining for Notch1 in normal skin (a) and HHD (b,c,d). (D) IHC staining for HKII in normal skin(a) and HHD patients(b,c,d). (E). Quantification for IHC staining signals in A~D.




Figure 8

MiR-203 expression in HHD patients. Real-time PCR was performed to examine the mRNA expression levels of miR-203 in HHD patients.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [FigureS1.jpg](#)
- [FigureS2.jpg](#)