

# Differential Influences of Social Support on App Use for Diabetes Self-Management – a Mixed Methods Approach

**Nicola Brew-Sam**

Australian National University <https://orcid.org/0000-0001-7544-4499>

**Arul Chib** (✉ [ArulChib@ntu.edu.sg](mailto:ArulChib@ntu.edu.sg))

<https://orcid.org/0000-0002-3833-8889>

**Constanze Rossmann**

Universitat Erfurt

---

## Research article

**Keywords:** mHealth, apps, technology, diabetes, self-management, social support, shared decision-making

**Posted Date:** May 12th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-16125/v2>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on July 7th, 2020. See the published version at <https://doi.org/10.1186/s12911-020-01173-3>.

# Abstract

**Background** The impact of social support on diabetes management and health outcomes has been investigated comprehensively, with recent studies examining social support delivered via digital technologies. This paper argues that social support has an impact on the use of diabetes technologies. Specifically, we postulate differences between the impact of healthcare professional versus non-professional (family/friends) support on mobile app use for diabetes self-management.

**Methods** This research employed a triangulation of methods including exploratory semi-structured face-to-face interviews (N= 21, Study 1) and an online survey (N= 65, Study 2) with adult type 1 and type 2 diabetes patients. Thematic analysis (Study 1) was used to explore the relevance of social support (by professionals versus non-professionals) for diabetes app use. Binary logistic regression (Study 2) was applied to compare healthcare decision-making, healthcare-patient communication, and the support by the personal patient network as predictors of diabetes app use, complemented by other predictors from self-management and technology adoption theory.

**Results** The interviews (Study 1) demonstrated that (technology-supported) shared decision-making and supportive communication by healthcare professionals depended on the medical specialty of attending physicians. The personal patient network was perceived as either facilitating or hindering the use of mHealth for self-management. Binary logistic regression (Study 2) showed that the specialty of the physician significantly predicted the use of diabetes apps, with supervision by diabetes specialists increasing the likelihood of app use (as opposed to general practitioners). In addition, specialist care positively related to a higher chance of shared decision-making and better physician-patient communication. The support by the personal patient network predicted diabetes app use in the opposite direction, with less family/friend support increasing the likelihood of app use.

**Conclusion** The results emphasize the relevance of support by healthcare professionals and by the patient network for diabetes app use and disclose differences from the existing literature. In particular, we found that the use of diabetes apps may increase in the absence of social support by family or friends (e.g., compensation for lack of support), and that use of diabetes apps may decrease when such support is high (e.g., no perceived need to use technology). Implications for practice are discussed.

## Background

Prior research has suggested the impact of social support on diabetes management and health outcomes (1, 2). With technological advances in healthcare, recent studies increasingly examine social support for diabetes self-management delivered through mobile online technologies (mHealth) (e.g., 3). In this context, most studies considered social support as an outcome of mHealth use, or mHealth as a means for delivering social support. Applying a novel perspective to mHealth research, our paper investigates the influence of social support on the use of mHealth (apps), as implicated by technology adoption models.

In this paper, we distinguish between the impact of varied types of social support on diabetes self-management and outcomes in previous studies, further arguing that these sources of social support differentially influence technology-supported diabetes self-management. In particular, we identify and examine the varied influence of support by healthcare professionals and by the personal social patient network on the use of diabetes self-management apps. A separate study conducted on diabetes app quality and app features, as factors of interest, has been reported elsewhere (4, 5).

### *The relevance of social support for diabetes self-management*

Patient self-management, in which patients are decision-makers in control of their daily diabetes management (6), moved to the core of diabetes care when research revealed that patient non-compliance rates were high in care approaches based solely on healthcare professional responsibility ("doctor knows best principle") (6). Wilkinson, Whitehead, and Ritchie (7) reported that the ability to self-manage a diabetic condition (e.g., diet adherence, physical activity, medication intake, managing psychological aspects (8, 9)) is influenced by a whole series of factors (e.g., in 10, 11). Prominently, social support has been reported to influence self-management behaviors and outcomes, associated with improved patient lifestyle and clinical outcomes, and reduced psychosocial disease symptoms (2). The impact of support has been distinguished between that provided by healthcare professionals (12-14) and by the personal social patient network like relatives or peers (15-17). This impact on diabetes management and health outcomes has been examined extensively, and been compared to one another (1, 18; also see 4).

First, studies on professional support show the beneficial effects of professional support on diabetes outcomes (19), especially when health professionals empower patients by using shared decision-making and supportive communication. A collaborative process of joint decision-making with an active contribution from both parties and the involvement of patients in treatment decisions (providing choices, considering patient preferences, and encouraging them to ask questions (13, 19, 20)) have been shown to be most supportive in disease care (21).

Second, in terms of non-professional support, family and friends can deliver emotional and practical support daily in diabetes care (22-25). However, the personal social patient network has been reported to have a negative impact on diabetes self-management and health outcomes, when interactions like nagging behaviors, a lack in understanding, too much sympathy or pity expressed, or other negative behaviors occur (16, 26-29). The concept of "miscarried helping" represents some of these problems (30).

### *Social support and technology-supported diabetes self-management*

Research has shown that interventions and care approaches need to differentially address support by the professional system and by the patient's personal network for achieving improved outcomes (1). This distinction applies similarly to research related to technology-supported diabetes care (31), which includes the use of diabetes apps that are designed to help patients improve their self-management (e.g., logbook apps).

In previous research, the use of (e.g., cloud-based) technology for support by healthcare professionals was reported to mostly result in positive outcomes for diabetes self-management (32-34). Wehbe, Curcio, Gajjar, and Yadlapati (35) stated that the integration of technology into diabetes care affects physician-patient relationships positively when discussions are facilitated and when shared decision-making between physicians and patients is improved. However, according to these authors, the higher workload for physicians can also affect the relationship negatively. Abbasgholizadeh, Meneer, Robitaille, and Legare (36) reported that health apps have potential for improved patient participation in shared health decision-making, but bear risks like security concerns or increased patient anxiety.

Beneficial effects of technology use on self-management and health outcomes were typically reported in studies that addressed non-professional social support by the patient's personal networks (37, 38). The use of social media, online communities, or mobile health (mHealth) has enhanced or improved psychological empowerment (39, 40), self-care information and knowledge (38, 41), blood glucose levels (HbA1c), as well as glucose self-monitoring and physical activity (3, 42), and other health topics (43-45). The inclusion of family members or spouses in technological diabetes management systems was found to improve their understanding of self-management requirements and affected the communication about diabetes and the support of the patient positively (46). Notably, different studies reported that online social support by non-professionals might not always result in beneficial self-management and health outcomes. For example, this occurs when participation in online communities leads to mismatches between needed and received support, or when support hinders health improvements for patients with high self-efficacy (47).

*Research gap: The influence of (non-) professional social support on technology use*

Existing studies almost exclusively focused on social support as an outcome of technology use, or on technology as a means for delivering social support. However, in order to increase an understanding of factors that influence technology use for self-care, research is still required that examines how professional and non-professional social support affect (mobile) diabetes technology use. It can be assumed that social support influences the use of mobile diabetes technology (e.g., apps) just as much as it influences traditional diabetes self-management behaviors (blood glucose testing, healthy nutrition, exercise, etc.) (4).

The number of studies examining factors influencing mHealth use is steadily rising (48), particularly in the context of disease self-management (49, 50). While theoretical models of technology adoption (51) are one step towards understanding factors that influence (mobile) technology adoption and use, there is a gap in proving both significant and substantive usage effects for mHealth (52, 53). Previous studies investigated factors influencing mHealth use in specific target groups (e.g., 54, 55), for specific diseases or health topics (56-60), and for sustained mHealth engagement (61). This study aims to explicate the effect of social support.

Social support has been shown to play a role as an influencing factor on aspects of technology use in mHealth studies. Zhang et al. (59), for instance, found that the theoretical factors of social influence and

performance expectancy had the strongest direct influence on behavioral intention to use diabetes apps for disease management. Likewise, Quaasar et al. (55) found that social influence had a significant impact on intention to use mHealth, specifically in populations of higher age. We aim to address the research gap in these studies by comparing differential influences of healthcare professionals and personal patient network on technology for self-care.

We hypothesize that both professional (e.g., physician) and non-professional support (e.g., family and friends) differentially influence diabetes app use (as a technology-supported self-management behavior). Following previous diabetes research it can be assumed that: *(H1) shared decision-making styles and supportive communication of healthcare professionals are promoting technology-supported self-management, while (H2) the impact of support by the patient's personal social network on app use for self-management depends on the type of perceived support (positive or misguided).*

## Methods

We used a mixed methods approach that included qualitative and quantitative data collection and analysis to permit "a more complete and synergistic utilization of data" (62, p.1). A detailed description of the methodology can be found in Brew-Sam (4, 63).

### *Study 1: Semi-structured face-to-face interviews*

In the first study we conducted 21 semi-structured face-to-face interviews with type 1 (T1DM) and type 2 (T2DM) adult diabetes patients in Singapore. The interview guide (63) included questions on patients' previous diabetes app (non-) use, their diabetes self-management, their attitudes, the support received from their medical team, relatives, and from others, as well as other relevant factors from self-management (7) and technology adoption theories, such as Unified Theory of Acceptance and Use of Technology (UTAUT) (51).

In addition to an interview guide developed based on these theories, we used a standardized background information questionnaire to collect relevant information about the patient background (e.g., demographic data, disease information) (63). Open questions collected information about app and technology (non-) use as part of the diabetes self-management, as well as professional and non-professional social support, including shared decision-making in the patients' treatment, their communication with healthcare professionals, and the support they received from their personal social network. Regarding the latter, we focused on family and friends' support, being their strong-tie partners (64).

Ethics approval was sought and approved by the *Nanyang Technological University* Review Board (IRB-2016-01-012). Purposive sampling included patients with a variety of demographic and disease characteristics. Participants were informed about the study purpose and signed informed consent forms. All interviews were conducted by one researcher in English. The interviews were conducted and transcribed in 2015/2016 using the audio recording software "Audacity". The audio files were completely

and manually transcribed by three research assistants (explicit verbal content only); the cadence and cultural speech patterns have been retained for accuracy. The accuracy of transcription was checked, and names of interview participants were anonymized.

Thematic analysis, as described by Braun and Clarke (65), was used for data analysis, following established standards for qualitative research (66). A theme was defined as a topic resulting from different interview statements with similar content. First, preliminary categories were developed from the interviews regarding technology use, self-management, and social support using a matrix for data extracts. Broad themes were developed from the categories, which were reviewed against data extracts and the entire data set, before results were analyzed and interpreted (discussion among multiple researchers where results were ambiguous). A diabetes app (non-) user typology summarized the results on emerging themes and to map out differences in social support for different groups.

### *Study 2: Standardized online survey*

In the second study we conducted a standardized online survey with 65 type 1 and type 2 diabetes patients. The questionnaire included questions on their diabetes background, self-management behaviors, mHealth use, social support, general attitudes and feelings, psychological empowerment, and demography (63).

Previous diabetes app use or non-use, as well as the length and frequency of previous app use were measured based on the National Survey on Health App Use (67). Decision-making styles by health professionals and communication with the patient were measured using the Provider Participatory Decision-making Style Scale (13) (PDMstyle,  $M = 3.25$ ,  $SD = .08$ ,  $\alpha = .943$ ) and the Provider Communication Scale (13) (PCOM,  $M = 3.50$ ,  $SD = .37$ ,  $\alpha = .919$ ). Items from these scales, for example, asked to "specify how often the doctors or health care professionals who take care of your diabetes offered you choices in your medical care" (PDMstyle, *none of the time* to *all of the time*) and "tell us how the doctors or health care professionals who take care of your diabetes are at explaining treatment alternatives" (PCOM, *poor* to *excellent*). 5-point Likert scales were used throughout the survey. Social support by the personal social patient network was measured with a scale on support by family and friends from the Diabetes Care Profile (DCP, section V,  $M = 3.29$ ,  $SD = .19$ ,  $\alpha = .931$ ) (68, 69). Scale items included questions on support delivered by family and friends, for example, "My family or friends help and support me a lot to take my medicine" (DCP, *strongly agree* to *strongly disagree*). Diabetes background and self-management data were adapted from the Diabetes Care Profile (69), two psychological empowerment scales (70, 71), the National Survey on Health App Use (67), and the Summary of Diabetes Self-Care Activities (72).

The survey was pre-tested with sixteen diabetic and non-diabetic participants. The online survey was disseminated (snowball method) to former face-to-face interview participants, as well as through social media and diabetes support groups. A lottery participation (monetary incentives: three coupons of 100 SGD each) was offered at the end of the survey.

Binary logistic regression (enter, blockwise) was used to test H1 and H2 on social support by health professionals versus non-professionals (family and friends) influencing diabetes app use, and to compare the strength of both (and other relevant included self-management and diabetes background factors from self-management and technology adoption theory) on diabetes app use. Independent variables were metric or coded as dummy variables, while diabetes app use was used as a binary variable (previous diabetes app use or non-use). Independent factors that included data from both app users and non-users were used in binary logistic regression. Regression models were compared for model fit, prediction success, and the inclusion of significant predictors by starting with all available factors, and then removing factors step by step.

### *Study samples*

**Table 1** summarizes demographic and disease characteristics of the interview (Study 1,  $N = 21$ ) and the online survey samples (Study 2,  $N = 65$ ). Overall, the samples were diverse, including diabetes patients from various demographic and disease-related subgroups. The interview sample mostly included active patients managing their condition, with five respondents at high risk for health complications due to lacking diabetes knowledge, lacking motivation and insufficient self-management, unfavorable self-management attitudes, and/or critical health conditions (e.g., high blood glucose values). In both samples there were more type 2 diabetes patients than type 1 diabetes patients, following the global prevalence (~90%) of type 2 diabetes (73).

Table.1 Description of Study 1 and Study 2 Samples

Variable	Study 1 - Interviews						Study 2 - Survey					
	<i>n</i>	% of <i>N</i>	<i>M</i>	<i>SD</i>	<i>Min</i>	<i>Max</i>	<i>n</i>	% of <i>N</i>	<i>M</i>	<i>SD</i>	<i>Min</i>	<i>Max</i>
Age (in years)	-	-	48.35	17.46	19.00	68.00	-	-	49.74	14.67	20.00	70.00
Education												
MA graduate or higher	-	-					15	23.1				
BA graduate/college graduate	6	28.6					22	33.8				
Some college	7	33.3					12	18.5				
High school graduate	1	4.8					12	18.5				
Some high school	4	19.0					3	4.6				
Other education level	1	4.8					1	1.5				
Employment												
Full-time working	8	38.1					34	52.3				
Part-time working	3	14.3					12	18.5				
Homemaker	3	14.3					1	1.5				
Retired	3	14.3					7	10.8				
Student	3	14.3					5	7.7				
Unemployed	-	-					6	9.3				
Family status												
Married	9	42.9					32	49.2				
Never married	8	38.1					18	27.7				
Never married but relationship	-	-					4	6.2				
Separated/divorced	1	4.8					7	10.8				
Widowed	1	4.8					4	6.2				
Gender												
Men	11	52.4					32	49.2				
Women	10	47.6					33	50.8				
Nationality												
Singaporean	19	90.5					57	87.7				
Malaysian	1	4.8					4	6.2				

Other	1	4.8					4	6.2					
Diabetes Background													
Diabetes family history	12	57.1					50	76.9					
Diabetes type													
T2DM (incl. gestational)	11	52.4					50	76.9					
T1DM	9	42.9					13	20.0					
Pre-diabetes	1	4.8					2	3.1					
Diseases (other)	8	38.1					-	-					
Education on diabetes (received)	17	81.0					50	76.9					
Length of diabetes (in years)	-	-	19.89	12.07	4.00	38.00	-	-	13.73	9.81	.00	36.00	
Medication													
Insulin injection (syringe or pump)	14	66.7					25	38.5					
Oral diabetes medication	10	47.6					51	78.5					
(Self-) Management													
Check-up frequency (in months)	-	-	3.83	1.36	2.00	6.00	-	-	-	-	-	-	-
Diabetes app use	11	52.4					-	-					
Never used	-	-					34	52.3					
Previous use	-	-					17	26.2					
Current use	-	-					14	21.5					
Online health information seeking (Study 2: days per week)	19	90.5	-	-	-	-	-	-	2.70	2.27	.00	7.00	
Part of support group	15	71.4					-	-					
Offline	-	-					42	64.6					
Online	-	-					19	29.2					
Part of diabetes program	2	9.5					7	10.8					

*Note.* Table based on Brew-Sam (4); Study 1: N= 21, Study 2: N= 65

## Results

### *Interview results on social support and diabetes app use (Study 1)*

As a foundation for comprehensively examining the influence of (H1) professional and (H2) non-professional support on diabetes app use, and to specify the hypotheses, themes were extracted from the interview data, relating to the aspects "previous app use for diabetes management", "professional support", and "non-professional support". **Table 2** summarizes the results in form of a diabetes app (non-) user typology, which displays the differences in social support in differing app user and non-user groups.

Table 2. Diabetes App (Non-)Use and Social Support - Study 1

	Non-User of Diabetes Apps			Diabetes App User	
App user type	The diabetes app non-user without interest in diabetes apps	The interested non-user of diabetes apps	The dissatisfied adopter of diabetes apps	The experienced diabetes app switcher	The consistent long-term diabetes app user
Self-management	No risk group, experienced with good perceived diabetes knowledge, mainly good self-management	Diabetes risk group, lacking diabetes knowledge or misperceptions, avoidance strategies, insufficient self-management, dangerous health behaviors	No risk group, good diabetes knowledge (educated at young age), good self-management	No risk group, very good diabetes knowledge, very good self-management, experienced, strict regimen	No risk group, specialized diabetes knowledge, intense diabetes education, good self-management, strict carb counting, active
Medical specialty of healthcare professionals (HCPs)	Mainly general practitioners	Mainly general practitioners	Diabetes specialists	General practitioners or diabetes specialists	Diabetes specialists, partly other HCPs as part of a diabetes program
Perceived physician quality	Partly incontent, or content after choosing selected physicians	Incontent, physicians not supportive (with exceptions)	Content but also seeing downsides	Content or incontent, depending on medical specialty of physician	Content, physicians as "friends"
Decision-making	Independent patient decision-making	Dependent or independent patient decision-making with dangerous health behaviors	Independent patient decision-making but listening to HCP advice, shared decision-making	Independent patient decision-making	Shared decision-making with close relationships between HCP and patient
Physician communication	Partly short consultations, no engagement, patients need to ask questions to receive information	Short consultations, physicians not helpful, answer questions only (no further engagement)	Consultations also through email/online/calls, close relationships with intense communication, partly busy doctors	Short consultations (general pract.), longer consultations but more expensive (specialist), partly contact through email	Discussions similar to friends' relationships, honesty in consultations, partly long consultations
Support group participation	Support group leader or follower	No support group participation or support group follower	Support group follower or leader, volunteering for other patients, part of diabetes program	Support group leader or follower	Support group leader, part of a diabetes or app pilot program
Family/friend support	Managing without support or negative influences by family/friends (but	Managing without support, rarely support by family	Involvement/support by family only right after diagnose (beginning of the disease)	Support especially by friends, family support	Partly family support, sometimes negative family/friend influences

	perceived relevance of support)				
Interviewees in sample	(Ching Ching), Deng Li, Jie, Kang, Xin Qi	Ei Tek, Li Ting, Ming, (Rei Hong), Zhen Wei	Adit, Cheng, Pang, Sona	Bharat, Navin, Xiu Wen	Gu Fang, Henna, Kaiyan, Shi Hui

*Note.* Table based on Brew-Sam (4); names of participants changed to guarantee anonymity

*Diabetes app (non-) use in the sample – description of the dependent variable:* App (non-) use categories ranged from "no previous use", "(no) interest in apps", and "(no) knowledge about existing diabetes management apps", to "infrequent and short-term app use" to "long-term app use". Some participants had never used diabetes apps for self-management before and expressed no interest in them (e.g., Jie, Kang[1]), while others showed interest but lacked knowledge about appropriate app use (e.g., Li Ting, Zhen Wei). Most participants were familiar with available diabetes apps (apart from a few without any knowledge about app availability, e.g., Rei Hong and Zhen Wei). Reported diabetes-specific app use was almost exclusively limited to logbooks for blood glucose monitoring (e.g., DAFNE online App, MySugr, Glooko, DiabetesM) and to food databases displaying nutritional information (e.g., food database app developed by the Singaporean Health Promotion Board). Moreover, app users split into short-term users ("adopters") who were mostly unsatisfied with the current state of diabetes apps and had abandoned their use after a while (e.g., Adit, Cheng), users who constantly switched apps, using several in parallel (e.g., Bharat, Navin), and users who used one main app over a longer period of time (e.g., Gu Fang, Henna) (**Table 2**).

*Professional support – Physician-patient relationship:* The themes derived from the interviews regarding healthcare professional support in (technology-supported) diabetes management included "medical specialty of physician related to perceived care quality", "taking time for communication", "actual decision-making", "decision-making preferences", and "inclusion of apps in physician-patient relationship".

Physicians were mentioned as the main supervisors in diabetes care, with other healthcare professionals (e.g., dieticians, nurse educators, podiatrists, pharmacists) only partly included in care with considerable variation amongst respondents (Xin Qi, Bharat, Henna, Deng Li). Thus, we further focused on the physician-patient relationship.

Patients supervised by general practitioners (GPs) reported short consultations with brief physician-patient communication ("the doctor is 5-10 minutes only", Xin Qi, age 56, T2DM; "if you ask questions, they will answer. But they won't engage you for too long", Jie, age 64, T1DM), perceived GPs lacking diabetes knowledge, and a perceived lack in support ("they [physicians] are not helpful", Li Ting, age 49, T2DM). They mostly expressed dissatisfaction with the quality of supervision by their GPs. The group of dependent patients, defined as those mainly following doctors' instructions without taking diabetes-

related decisions or being less active in decision-making (e.g., Rei Hong, age 61, T2DM), mostly consulted GPs. Additionally, non-users of apps mostly consulted GPs (**Table 2**).

In contrast to GPs, diabetes specialists were reported as providing adequate time for support: "you got two types [of physicians]... one we call it family physician... one uh he charge you more, double [specialist]...the doctor will spend more time" (Bharat, age 66, T2DM). Moreover, specialists sometimes developed close relationships with their patients ("he's... more like a family friend ... than a doctor", Henna, age 60, T2DM) and mostly followed shared decision-making approaches. Yet, some patients considered their physicians as merely advisors, preferring to take diabetes care decisions independently (e.g., Sona, Bharat) (**Table 2**).

Overall, few patients reported that physicians or other healthcare professionals (nurses) talked about diabetes apps in the consultations, or shared app information with them (e.g., Xin Qi, Bharat, Henna). Some patients participated in diabetes programs (e.g., DAFNE, dose adjustment for normal eating) that included specialist supervision and an app for self-management (e.g., Sona, Shi Hui). Apart from these specialist programs, physicians hardly used apps to communicate with their patients (sometimes Email, e.g., Gu Fang).

*Non-professional support – Family & friends' support:* Self-management support by family and by friends fell into the categories of "managing alone", "negative support", "involvement only after diagnosis", and "strong involvement". Both app users and non-users reported receiving support by their family and friends in their diabetes management (**Table 2**), either after diagnosis at the beginning of self-care (e.g., Cheng, Pang) or throughout the whole process of self-care (e.g., Navin, Xiu Wen). Negative support was reported when their social contacts tempted the patients towards unhealthy lifestyles, e.g., "They always say 'never mind la! Eat la, just eat la. Only once. You don't eat this every day'" (Kang, age 67, pre-diabetes), or when involvement of the family resulted in nagging behaviors ("whenever they are with me... when the doctor tells me something... and then after ... when we go home they immediately start nagging me", Adit, age 22, T1DM). Positive attitudes towards diabetes care by family and friends were mentioned as important for self-management ("the key is that they're... not uhh... really that negative on this... they're also very positive", Xiu Wen, age 57, T1DM). Some non-users of diabetes apps said they managed their condition alone, without the involvement of family or friends ("It's myself, nobody else... No friends, no, nobody else, it is me", Jie, age 64, T1DM).

Overall, the interviews showed that aspects of professional support (style of decision-making, duration and quality of communication) related to perceptions of satisfaction and success in self-management. Moreover, this appeared to be influenced by the medical specialty of the physician supervising the patient (compare 4). Regarding non-professional support, the support by family and friends diversely related to (technology-supported) self-care, with both positive and negative influences on self-management reported. Based on the exploratory interview results we specified the theoretically derived hypotheses *H1* and *H2* as following:

*(H1a) Supportive behaviors by the supervising physician (shared decision-making styles and supportive communication) positively predict diabetes app use for self-management.*

*(H1b) The medical specialty of the supervising physician (specialist versus GP) is a predictor of diabetes app use for self-management, with specialist care promoting app use to a greater extent than care by GPs.*

*(H2a) (Positive) support by the patient's personal social network (family and friends) positively predicts diabetes app use for self-management.*

### *Online survey results on social support and diabetes app use (Study 2)*

In Study 2 we tested hypotheses (H1a), (H1b), and (H2a) using binary logistic regression. Checking for autocorrelation of all independent variables, physician decision-making and physician-patient communication were highly correlated with  $r = .772, p < .01$ . Therefore, decision-making and communication were recoded into a single variable "physician-patient relationship" (4).

Based on theoretical considerations and the interview results, we began with a binary logistic regression model that included a maximum of independent factors: physician-patient relationship, medical specialty of the physician, family/friend support, and other relevant factors from technology adoption theory (51) and from self-management theory (7) shown to be relevant predictors of technology use for disease self-management. We then compared different models by reducing independent factors to find the model with the best fit.

A model with good fit included the factors derived from the interviews *physician-patient relationship, family/friend support, medical specialty of the physician*, as well as the additional UTAUT factors' *perceived app potential* (performance and effort expectancy), *previous health information seeking online* (technological experience), and *age*; and the self-management factors *type of diabetes, length of diabetes, perceived health status, payment problems and insurance coverage* (socioeconomic), *blood glucose testing adherence* (self-management behaviors), *interest in innovation* (attitudes), *perceived diabetes knowledge, program or support group participation, and psychological empowerment*. The test of this model against the constant-only model was statistically significant, indicating that the predictors are a set reliably distinguished between diabetes app use and non-use ( $\chi^2(16) = 26.752, p < .05$ ). Nagelkerke's  $R^2 = .656$  indicated a high relationship between prediction and grouping (goodness-of-fit). Prediction success overall was 90% (80% for app non-use and 96% for app use). The Wald criterion demonstrated that only the family/friend support ( $Wald(1) = 5.315, p = .021$ ) and the medical specialty of the consulted physician (dummy GP or specialist,  $Wald(1) = 4.014, p = .045$ ) made a significant contribution to the prediction of diabetes app use. The  $Exp(\beta)$  value indicated that when the family/friend support was increased, the relative probability (odds ratio) that diabetes apps were used decreased with  $Exp(\beta) = .044, \beta = -3.131$ . The  $Exp(\beta)$  value indicated that when the patients were supervised by specialist

doctors the relative probability (odds ratio) that diabetes apps were used increased with  $Exp(\beta) = 9460.805$ ,  $\beta = 9.155$ . In contrast to non-professional support (family/friends), the physician-patient relationship was not found to be a significant predictor in the model.

A model resulting after the removal of the factors *interest in innovation*, *insurance coverage*, and *program or support group participation* (due to lacking significance) showed a slightly lower prediction success with 75% (70% for app non-use and 79% for app use), but overall model significance with  $\chi^2(13) = 26.936$ ,  $p < .05$ , and a moderate to high relationship between prediction and grouping (goodness of fit) with Nagelkerke's  $R^2 = .509$ . In this model, the Wald criterion demonstrated that the family/friend support ( $Wald(1) = 6.617$ ,  $p = .010$ ) and perceived health status ( $Wald(1) = 7.839$ ,  $p = .005$ ) made a significant contribution to the prediction of diabetes app use. Again, the  $Exp(\beta)$  value showed that when support by family/friends was increased, the relative probability (odds ratio) that diabetes apps were used decreased with  $Exp(\beta) = .283$ ,  $\beta = -1.261$ . The  $Exp(\beta)$  value also indicated that when the perceived health status was improved the relative probability (odds ratio) that diabetes apps were used increased with  $Exp(\beta) = 8.030$ ,  $\beta = 2.083$ .

Further reducing the independent factors, the models showed similar results to the last model, resulting in *family/friend support* and the *perceived health status* being significant predictors of diabetes app use. A further reduction of factors decreased the model fit, yet the *medical specialty of the physician* nearly reached significance again.

Overall, after testing various models, only the family/friend support, the medical specialty of the supervising physician, and the perceived health status significantly predicted diabetes app use. Less family/friend support was likely leading to a higher chance of diabetes app use, while the use of diabetes specialists or a better perceived health status increased the chance of app use for self-management.

Despite the physician-patient relationship lacking significance for predicting diabetes app use in the models (apart from the medical specialty of the physician as a significant predictor), additional *t*-test calculations showed that the medical specialty of the physician related to the style of decision-making and the physician-patient communication. There were differences between specialists and GPs, with specialist care positively related to higher shared decision-making and better physician-patient communication as compared to GPs (Decision-making PDMstyle: specialists:  $M = 3.72$ ,  $SD = 1.16$ ,  $n = 37$ , general practitioners:  $M = 2.65$ ,  $SD = 1.30$ ,  $n = 22$ ;  $t(57) = -3.275$ ,  $p < .01$ ,  $N = 59$ ; Communication PCOM: specialists:  $M = 3.82$ ,  $SD = 1.01$ ,  $n = 37$ , general practitioners:  $M = 3.13$ ,  $SD = 1.09$ ,  $n = 22$ ;  $t(57) = -2.472$ ,  $p < .05$ ,  $N = 59$ ) (compare 4).

[1] all names have been changed to ensure anonymity

## Discussion

In general, this two-study project found that social support from both professional and patient's personal social networks had an influence on mobile app usage in the context of diabetes self-care.

Regarding professional support, the interviews (Study 1) showed that shared decision-making and supportive communication, including support regarding diabetes technology use, depended on the medical specialty of the physician, with patients perceiving support by diabetes specialists as more helpful than by GPs. Binary logistic regression (Study 2) confirmed these results, showing that the specialty of the physician significantly predicted the use of diabetes apps by patients, with supervision by diabetes specialists increasing the likelihood of the apps being used (as opposed to GPs). Specialist care was positively related to higher shared decision-making and better physician-patient communication. These study results conform with existing research, which pointed towards differences in diabetes care depending on the medical specialty of the supervising physician. A study by Koizumi et al. (74), for example, reported that attitudes towards glucose control and self-care varied in physicians depending on their medical specialty. According to these researchers, patients' self-management behaviors can be influenced by physician's beliefs and behaviors. De Berardis et al. (75) compared diabetes consultations in 125 diabetes outpatient clinics and 103 general practices for process and intermediate outcomes (frequency of examinations, HbA1c, blood pressure, and cholesterol levels) over a period of two years. They found significant better results for the majority of the process measures, and for cholesterol levels, in the specialized diabetes outpatient clinics as compared to the general practices. In particular, care by the same specialist in a diabetes outpatient clinic ensured better quality of care in comparison with other care options. Yet, these studies did not relate to diabetes technology use. Thus, our results add preliminary evidence that differences do not just affect traditional but also technology-supported diabetes care (e.g., diabetes app use), with specialist care promoting technology-supported self-management to a greater extent than care by GPs.

Examining the use of technology in physician-patient relationships further, other studies generally discussed the usefulness of health apps for physician-patient interaction, for example, in shared decision-making (36). Abbasgholizadeh et al. (36) reported mixed results for technology-supported physician-patient interactions; for example, better accessibility to data, improved efficiency for the physician, real-time connectivity, or remote decision-making on the one hand, but diminished quality of care through overuse of mHealth, increased health disparities due to lacking mHealth access, or lack of mHealth regulation, on the other hand.

In terms of non-professional support, the patients' personal network was perceived as either facilitating or hindering the use of mobile technology for self-management, depending respectively on the presence of supportive or unsupportive behaviors (Study 1). Moreover, support by this non-professional network significantly predicted patients' use of diabetes apps, with lower family and friend support increasing the likelihood of using diabetes apps (Study 2). Potential explanations for the negative relationship could be that certain patient groups are substituting technology to compensate for a lack of perceived or actual social support (e.g., 76).

There is an emergent research field examining loneliness and Internet use (77, 78), with studies suggesting that technology can offer a solution for chronically ill patients who feel isolated with their disease and who search for a (technological) solution to overcome their lack of social connection (79). As mentioned earlier, support by relatives and friends is not necessarily always positive, for example, when unsupportive behaviors or miscarried helping occur (16, 22, 26). Thus, support can be sought elsewhere, in this case, via technology substitution. Conversely, high social support by family and friends might nullify the need for additional technology to support self-care, which might explain why higher social support accompanies less technology use. In general, our results support the extant literature showing that the patients' social network influences attitudes towards innovation, which influence innovation adoption behaviors in return (80).

### **Future research**

Future research needs to examine aspects of social support influencing mHealth use, also comparing the influence of professional versus non-professional support on mHealth use. Apart from the social influence of health professionals and of family and friends on mHealth use as studied here, previous studies show that "peer" patients also play an important role as sources of information and of empowerment for patients (38, 81). Thus, firstly, follow-up research should include the role of peers of patients. Secondly, the investigation can be extended from physicians to other healthcare professionals involved (e.g., nurse educators). Thirdly, mHealth studies should extend their theoretical foundations beyond technology adoption theories. There are few theories applied in mHealth research, which mainly relies on traditional behavior change theories (82, 83). To investigate factors influencing and predicting technology use for self-management, a broader theoretical foundation needs to be available, overcoming the limitations of a very narrow range of theories. Fourthly, even though the socioeconomic and demographic factors included in the regression analysis (e.g., age, insurance coverage) did not predict the use of diabetes apps, it has to be further examined if these factors have an impact on the results, like the association between the specialty of the physician and the use of diabetes technology, or the negative association between non-professional support and the technology use.

### **Implications for research, policy and practice**

While our study focused on diabetes apps, these results can equally inform mHealth research in other health domains involving self-management, including various chronic conditions such as heart diseases, obesity, asthma, stroke, cancer, arthritis, hypertension, multiple sclerosis, etc. (84, 85). Even though there are specific characteristics for each disease, general knowledge about technology use for self-management can be helpful across health conditions to design technology accordingly.

For diabetes policy and practice, the study results suggest that there are notable differences in patient groups regarding diabetes app use, with diabetes patients being a heterogeneous group with varying needs (86). Segmentation theory needs to be used to tailor diabetes app characteristics specifically for subgroups with varying preferences and needs regarding technology (also considering socioeconomic, cultural, and demographic differences). Moreover, influencing factors need to be considered when

designing technology for diabetes care, like the influence of aspects of social support and social influence (87). Both healthcare professional supervision and family/friend support can influence technology-supported self-management. In particular, the care provided by GPs was perceived as unsatisfactory by most diabetes patients in the Study 1 sample. This resulted in less mobile technology use as compared to specialist supervision, which deficiency should be addressed to achieve equal care for all patients. The inclusion of relatives in (technology-supported) self-care should be further investigated depending on patient preferences and the usefulness of support provided (to avoid misguided support).

## **Study limitations**

Study limitations included the small sample size of the online survey, which limited sophisticated multi-variate data analysis (4) and broad generalizability. Considerable effort was undertaken to achieve a larger sample size (extension of the field phase, seeking contact with official care organizations, seeking cooperation with healthcare professionals, repeated invitations for survey participation), yet without success. Singaporean citizens suffering from diabetes receive a large amount of invitations for study participation since the government declared "the war against diabetes" in 2016 (88), so these are a prized yet scarce resource for research. Due to the cross-sectional nature of the study design, final conclusions about causality could not be drawn. Additionally, due to the recruitment method a certain degree of self-selection was unavoidable (over-representation of support group members and educated participants). The sample was therefore not representative for all Singaporean individuals with diabetes; however, we managed to include participants of the main Singaporean cultural backgrounds in the sample (Chinese, Malay, Indian). These three groups comprise 97% of Singaporeans (89). A triangulation of the study findings with results on diabetes app features and quality (4, 5) was not undertaken here due to length limitations.

## **Conclusion**

Our study results indicated that aspects of both professional and non-professional support have an impact on the use of mobile self-management technology by diabetes patients. However, the results also showed that the effects of support are not always positive for technology use. This leads to the conclusion that social support in diabetes self-management needs to be considered and analyzed in a differentiated manner when looking into social support as a predictor of technology use.

## **List Of Abbreviations**

T1DM/T2DM – Type 1/Type 2 Diabetes Mellitus

GP – General Practitioner

HCP – Healthcare Professional

UTAUT – Unified Theory of Acceptance and Use of Technology

DAFNE – Dose Adjustment for Normal Eating

## Declarations

### Ethics approval and consent to participate

Ethical review and approval was provided by the *Nanyang Technological University* Review Board (IRB-2016-01-012). Consent for participation was signed by all study participants.

### Consent for publication

Consent for publication was obtained by all study participants.

### Availability of data and materials

The (anonymized, non-identifiable) datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request. Additional information on the methodology (questionnaires, survey) can be found online (63).

### Competing interests

Not applicable, none to be reported.

### Funding

Study funding was provided by the Nanyang Technological University (research grant number M4081081).

### Authors' contributions

NBS collected, analyzed and interpreted the data, and drafted the manuscript. CR and AC supported data collection and data analysis, reviewed the manuscript, and provided detailed comments for improvement. All authors read and approved the final manuscript.

### Acknowledgements

Not applicable.

## References

1. Rosland AM, Kieffer E, Israel B, Cofield M, Palmisano G, Sinco B, et al. When is social support important? The association of family support and professional support with specific diabetes self-

- management behaviors. *Journal of General Internal Medicine*. 2008 Dec;23(12):1992-9. PubMed PMID: 18855075. Pubmed Central PMCID: 2596525.
2. Strom JL, Egede LE. The impact of social support on outcomes in adult patients with type 2 diabetes: a systematic review. *Current diabetes reports*. 2012 Dec;12(6):769-81. PubMed PMID: 22949135. Pubmed Central PMCID: 3490012.
  3. Burner E, Lam CN, DeRoss R, Kagawa-Singer M, Menchine M, Arora S. Using mobile health to improve social support for low-income Latino patients with diabetes: a mixed-methods analysis of the feasibility trial of TExT-MED + FANS. *Diabetes Technology and Therapeutics*. 2018 Jan;20(1):39-48. PubMed PMID: 29227155. Pubmed Central PMCID: 5770080.
  4. Brew-Sam N. *App use and patient empowerment in diabetes self-management – advancing theory-guided mHealth research*. Wiesbaden: Springer; 2020.
  5. Brew-Sam, N. , & Chib, A. How do smart device apps for diabetes self-management correspond with theoretical indicators of empowerment? An analysis of app features. *International Journal of Technology Assessment in Health Care*. 2019; 35(2):150-159.
  6. Funnell MM, Anderson RM. Empowerment and self-management of diabetes. *Clinical Diabetes*. 2004;22(3):123-7.
  7. Wilkinson A, Whitehead L, Ritchie L. Factors influencing the ability to self-manage diabetes for adults living with type 1 or 2 diabetes. *International Journal of Nursing Studies*. 2014;51(1):111-22.
  8. Peyrot M, Rubin RR, Lauritzen T, Snoek FJ, Matthews DR, Skovlund SE. Psychosocial problems and barriers to improved diabetes management: results of the Cross-National Diabetes Attitudes, Wishes and Needs (DAWN) Study. *Diabetic Medicine*. 2005 Oct;22(10):1379-85. PubMed PMID: 16176200.
  9. Hunter CM. Understanding diabetes and the role of psychology in its prevention and treatment. *American Psychologist*. 2016 Oct;71(7):515-25. PubMed PMID: 27690481.
  10. Ahola AJ, Groop PH. Barriers to self-management of diabetes. *Diabetic Medicine*. 2013 Apr;30(4):413-20. PubMed PMID: 23278342.
  11. Rodriguez KM. Intrinsic and extrinsic factors affecting patient engagement in diabetes self-management: Perspectives of a certified diabetes educator. *Clinical therapeutics*. 2013 Feb;35(2):170-8. PubMed PMID: 23411000.
  12. van Dam HA, van der Horst F, van den Borne B, Ryckman R, Crebolder H. Provider–patient interaction in diabetes care: effects on patient self-care and outcomes: a systematic review. *Patient Education and Counseling*. 2003;51(1):17-28.
  13. Heisler M, Bouknight RR, Hayward RA, Smith DM, Kerr EA. The relative importance of physician communication, participatory decision making, and patient understanding in diabetes self-management. *Journal of General Internal Medicine*. 2002;17(4):243-52.
  14. Patel NJ, Datye KA, Jaser SS. Importance of Patient-Provider Communication to Adherence in Adolescents with Type 1 Diabetes. *Healthcare (Basel)*. 2018 Mar 28;6(2). PubMed PMID: 29597246. Pubmed Central PMCID: 6023481.

15. Kaselitz E, Shah M, Choi H, Heisler M. Peer characteristics associated with improved glycemic control in a randomized controlled trial of a reciprocal peer support program for diabetes. *Chronic Illness*. 2018 Jan 1;1742395317753884. PubMed PMID: 29343089.
16. Grant RW, Schmittiel JA. Adults with diabetes who perceive family members' behaviour as unsupportive are less adherent to their medication regimen. *Evidence Based Nursing*. 2013 Jan;16(1):15-6. PubMed PMID: 22952019.
17. Armour TA, Norris SL, Jack L, Jr., Zhang X, Fisher L. The effectiveness of family interventions in people with diabetes mellitus: a systematic review. *Diabetic Medicine*. 2005 Oct;22(10):1295-305. PubMed PMID: 16176186.
18. Heisler M, Vijan S, Makki F, Piette JD. Diabetes control with reciprocal peer support versus nurse care management: a randomized trial. *Annals of internal medicine*. 2010 Oct 19;153(8):507-15. PubMed PMID: 20956707. Pubmed Central PMCID: 4117390.
19. Gensichen J, Von Korff M, Rutter CM, Seelig MD, Ludman EJ, Lin EH, et al. Physician support for diabetes patients and clinical outcomes. *BMC Public Health*. 2009 Sep 29;9:367. PubMed PMID: 19788726. Pubmed Central PMCID: 2762989.
20. Kaplan SH, Greenfield S, Ware JE, Jr. Assessing the effects of physician-patient interactions on the outcomes of chronic disease. *Medical Care*. 1989 Mar;27(3 Suppl):S110-27. PubMed PMID: 2646486.
21. Emanuel EJ, Emanuel LL. Four models of the physician-patient relationship. *The Journal of the American Medical Association, JAMA*. 1992;267(16):2221.
22. Helgeson VS, Mascatelli K, Seltman H, Korytkowski M, Hausmann LR. Implications of supportive and unsupportive behavior for couples with newly diagnosed diabetes. *Health Psychology*. 2016 Oct;35(10):1047-58. PubMed PMID: 27280364. Pubmed Central PMCID: 5033701.
23. *The Lancet*. Family matters in diabetes care. *The Lancet Diabetes & Endocrinology*. 2018;6(12):911.
24. Whitehead L, Jacob E, Towell A, Abu-Qamar M, Cole-Heath A. The role of the family in supporting the self-management of chronic conditions: A qualitative systematic review. *Journal of clinical nursing*. 2018 Jan;27(1-2):22-30. PubMed PMID: 28231630.
25. Kowitt SD, Ayala GX, Cherrington AL, Horton LA, Safford MM, Soto S, et al. Examining the Support Peer Supporters Provide Using Structural Equation Modeling: Nondirective and Directive Support in Diabetes Management. *Annals of Behavioral Medicine*. 2017 Dec;51(6):810-21. PubMed PMID: 28417438.
26. Bennich BB, Roder ME, Overgaard D, Egerod I, Munch L, Knop FK, et al. Supportive and non-supportive interactions in families with a type 2 diabetes patient: an integrative review. *Diabetology and Metabolic Syndrome*. 2017;9:57. PubMed PMID: 28736580. Pubmed Central PMCID: 5521150.
27. Carter-Edwards L, Skelly AH, Cagle CS, Appel SJ. "They care but don't understand": family support of African American women with type 2 diabetes. *Diabetes Educator*. 2004 May-Jun;30(3):493-501. PubMed PMID: 15208847.

28. Sanjari M, Peyrovi H, Mehrdad N. Managing children with diabetes within the family: Entering into the Diabetes Orbit. *Journal of Diabetes and Metabolic Disorders*. 2015;15:7. PubMed PMID: 26998445. Pubmed Central PMCID: 4797217.
29. Mayberry LS, Osborn CY. Family support, medication adherence, and glycemic control among adults with type 2 diabetes. *Diabetes Care*. 2012 Jun;35(6):1239-45. PubMed PMID: 22538012. Pubmed Central PMCID: 3357235.
30. Harris MA. The Family's Involvement in Diabetes Care and the Problem of "Miscarried Helping". *European Endocrinology*. 2006 (2):20.
31. Schiel R, Bambauer R, Steveling A. Technology in Diabetes Treatment: Update and Future. *Artificial Organs*. 2018 Nov;42(11):1017-27. PubMed PMID: 30334582.
32. Hsu WC, Lau KH, Huang R, Ghiloni S, Le H, Gilroy S, et al. Utilization of a Cloud-Based Diabetes Management Program for Insulin Initiation and Titration Enables Collaborative Decision Making Between Healthcare Providers and Patients. *Diabetes Technology and Therapeutics*. 2016 Feb;18(2):59-67. PubMed PMID: 26645932. Pubmed Central PMCID: 4753582.
33. Srivastava P, Verma A, Geronimo C, Button TM. Behavior stages of a physician- and coach-supported cloud-based diabetes prevention program for people with prediabetes. *SAGE Open Medicine*. 2019;7:2050312119841986. PubMed PMID: 31105938. Pubmed Central PMCID: 6509979.
34. Cheng C, Xin G, Xiaoli W, Chuanwei C, Xiaoming S, Yimin Z, et al. Evaluation of the implementation and effect of the healthcare cloud information platform for diabetes self-management: A case study in Shanghai. *The International Journal of Health Planning and Management*. 2019 Jul;34(3):986-97. PubMed PMID: 31368129.
35. Wehbe R, Curcio E, Gajjar M, Yadlapati A. Technology and Its Influence on the Doctor-Patient Relationship. *International Cardiovascular Forum Journal*. 2015;3:38.
36. Abbasgholizadeh RS, Menear M, Robitaille H, Legare F. Are mobile health applications useful for supporting shared decision making in diagnostic and treatment decisions? *Global Health Action*. 2017 Jun;10(sup3):1332259. PubMed PMID: 28838306. Pubmed Central PMCID: 5645724.
37. Gabarron E, Arsand E, Wynn R. Social Media Use in Interventions for Diabetes: Rapid Evidence-Based Review. *Journal of medical Internet research*. 2018 Aug 10;20(8):e10303. PubMed PMID: 30097421. Pubmed Central PMCID: 6109225.
38. Litchman ML, Edelman LS, Donaldson GW. Effect of Diabetes Online Community Engagement on Health Indicators: Cross-Sectional Study. *JMIR Diabetes*. 2018 Apr 24;3(2):e8. PubMed PMID: 30291079. Pubmed Central PMCID: 6238850.
39. Oh HJ, Lee B. The effect of computer-mediated social support in online communities on patient empowerment and doctor-patient communication. *Health Communication*. 2012;27(1):30-41. PubMed PMID: 21797714.
40. Gomez-Galvez P, Suarez Mejias C, Fernandez-Luque L. Social media for empowering people with diabetes: Current status and future trends. *Conference Proceedings, 37th Annual International*

- Conference of the IEEE, Engineering in Medicine and Biology Society (EMBC). 2015;2015:2135-8. PubMed PMID: 26736711.
41. Maki KG, O'Mally AK. Analyzing Online Social Support Within the Type 1 Diabetes Community. In: Sekalala S, Niezgodna BC, editors. *Global Perspectives on Health Communication in the Age of Social Media*. Hershey, PA: IGI Global; 2018. p. 59-84.
  42. Burner E, Mercado J, Hernandez-Saenz A, Peters A, Baezconde-Garbanati L, Arora S, et al. Design and patient characteristics of the randomized controlled trial TExT-MED+FANS A test of mHealth augmented social support added to a patient-focused text-messaging intervention for emergency department patients with poorly controlled diabetes. *Contemporary clinical trials*. 2019 May;80:1-8. PubMed PMID: 30878623. Pubmed Central PMCID: 6488230.
  43. Omboni S, Caserini M, Coronetti C. Telemedicine and M-Health in hypertension management: technologies, applications and clinical evidence. *High blood pressure & cardiovascular prevention: the official journal of the Italian Society of Hypertension*. 2016 Sep;23(3):187-96. PubMed PMID: 27072129.
  44. Fortuna KL, DiMilia PR, Lohman MC, Bruce ML, Zubritsky CD, Halaby MR, et al. Feasibility, acceptability, and preliminary effectiveness of a peer-delivered and technology supported self-management intervention for older adults with serious mental illness. *Psychiatric Quarterly*. 2018;89(2):293-305.
  45. Wang W, Su Y, Gao G. Social Determinants of mHealth Effectiveness: Evidence from a Large-Scale Experiment. *SSRN Electronic Journal*. 2018.
  46. Mayberry LS, Berg CA, Harper KJ, Osborn CY. The design, usability, and feasibility of a family-focused diabetes self-care support mHealth intervention for diverse, low-income adults with type 2 diabetes. *Journal of Diabetes Research*. 2016 2016;2016:1-13.
  47. Yan L. Good Intentions, Bad Outcomes: The Effects of Mismatches between Social Support and Health Outcomes in an Online Weight Loss Community. *Production and Operations Management*. 2018;27(1):9-27.
  48. Danaher BG, Brendryen H, Seeley JR, Tyler MS, Woolley T. From black box to toolbox: Outlining device functionality, engagement activities, and the pervasive information architecture of mHealth interventions. *Internet Interventions*. 2015 2015/03/;2(1):91-101.
  48. Zhu Z, Liu Y, Che X, Chen X. Moderating factors influencing adoption of a mobile chronic disease management system in China. *Informatics for Health and Social Care*. 2017 2017/01/09/:1-20.
  49. Azhar FAB, Dhillon JS, editors. *A systematic review of factors influencing the effective use of mHealth apps for self-care*. 3rd International Conference on Computer and Information Sciences (ICCOINS); 2016; Kuala Lumpur: IEEE.
  50. Venkatesh V, Morris MG, Davis GB, Davis FD. User acceptance of information technology: Toward a unified view. *MIS Quarterly*. 2003;27(3):425-78.
  51. Free C, Phillips G, Galli L, Watson L, Felix L, Edwards P, et al. The effectiveness of mobile-health technology-based health behaviour change or disease management interventions for health care consumers: a systematic review. *PLoS Medicine*. 2013;10(1):e1001362.

52. Chib A, van Velthoven MH, Car J. mHealth adoption in low-resource environments: a review of the use of mobile healthcare in developing countries. *Journal of Health Communication*. 2015;20(1):4-34. PubMed PMID: 24673171.
53. Hoque R, Sorwar G. Understanding factors influencing the adoption of mHealth by the elderly: an extension of the UTAUT model. *International Journal of Medical Informatics*. 2017 2017/05/;101:75-84. en.
54. Quaasar G, Hoque MR, Bao Y. Investigating Factors Affecting Elderly's Intention to Use m-Health Services: An Empirical Study. *Telemedicine Journal and e-Health*. 2018 Apr;24(4):309-14. PubMed PMID: 28976824.
55. Apolinario-Hagen J, Menzel M, Hennemann S, Salewski C. Acceptance of Mobile Health Apps for Disease Management Among People With Multiple Sclerosis: Web-Based Survey Study. *JMIR Formative Research*. 2018 Dec 12;2(2):e11977. PubMed PMID: 30684408. Pubmed Central PMCID: 6334710.
56. Breil B, Kremer L, Hennemann S, Apolinario-Hagen J. Acceptance of mHealth Apps for Self-Management Among People with Hypertension. *Studies in Health Technology and Informatics*. 2019 Sep 3;267:282-8. PubMed PMID: 31483283.
57. Apolinario-Hagen J, Hennemann S, Fritsche L, Druge M, Breil B. Determinant Factors of Public Acceptance of Stress Management Apps: Survey Study. *JMIR Mental Health*. 2019 Nov 7;6(11):e15373. PubMed PMID: 31697243.
58. Zhang Y, Liu C, Luo S, Xie Y, Liu F, Li X, et al. Factors Influencing Patients' Intentions to Use Diabetes Management Apps Based on an Extended Unified Theory of Acceptance and Use of Technology Model: Web-Based Survey. *Journal of Medical Internet Research*. 2019 Aug 13;21(8):e15023. PubMed PMID: 31411146. Pubmed Central PMCID: 6711042.
59. Liu D, Maimaitijiang R, Gu J, Zhong S, Zhou M, Wu Z, et al. Using the Unified Theory of Acceptance and Use of Technology (UTAUT) to Investigate the Intention to Use Physical Activity Apps: Cross-Sectional Survey. *JMIR Mhealth Uhealth*. 2019 Aug 22;7(9):e13127. PubMed PMID: 31507269.
60. Reading M, Baik D, Beauchemin M, Hickey KT, Merrill JA. Factors Influencing Sustained Engagement with ECG Self-Monitoring: Perspectives from Patients and Health Care Providers. *Appl Clinical Informatics*. 2018 Oct;9(4):772-81. PubMed PMID: 30304745. Pubmed Central PMCID: 6179719.
61. Wisdom J, Creswell JW. *Mixed Methods: Integrating Quantitative and Qualitative Data Collection and Analysis While Studying Patient-Centered Medical Home Models*. Rockville, MD: Agency for Healthcare Research and Quality, U.S. Department of Health and Human Services, 2013.
62. Brew-Sam N. Appendix. 2020. In: *App Use and Patient Empowerment in Diabetes Self-Management – Advancing Theory-Guided mHealth research*. Wiesbaden: Springer. Available from: <https://link.springer.com/book/10.1007/978-3-658-29357-4#toc>.
63. Parks MR. Weak and Strong Tie Relationships. In: Berger C, Roloff M, al. e, editors. *The International Encyclopedia of Interpersonal Communication*: Wiley; 2015. p. 1-5.

64. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in Psychology*. 2006;3(2):77-101.
65. Mays N, Pope C. Assessing quality in qualitative research. *BMJ*. 2000;320(7226):50-2.
66. Krebs P, Duncan DT. Health app use among US mobile phone owners: A national survey. *JMIR mHealth uHealth*. 2015 Nov 04;3(4):e101. PubMed PMID: 26537656. Pubmed Central PMCID: 4704953.
67. Fitzgerald JT, Davis WK, Connell CM, Hess GE, Funnell MM, Hiss RG. Development and validation of the Diabetes Care Profile. *Evaluation and the Health Professions*. 1996;19(2):208-30.
68. Fitzgerald JT, Davis WK, Connell CM, Hess GE, Funnell MM, Hiss RG. Diabetes Care Profile: Medical University of South Carolina; 1996 [04/13/2018]. Available from: [http://academicdepartments.musc.edu/family\\_medicine/rcmar/dcp.htm](http://academicdepartments.musc.edu/family_medicine/rcmar/dcp.htm).
69. Anderson RM, Fitzgerald JT, Gruppen LD, Funnell MM, Oh MS. The Diabetes Empowerment Scale-Short Form (DES-SF). *Diabetes Care*. 2003;26(5):1641-2.
70. Mantwill S, Fiordelli M, Ludolph R, Schulz PJ. EMPOWER – support of patient empowerment by an intelligent self-management pathway for patients: Study protocol. *BMC Medical Informatics and Decision Making*. 2015;15:18. PubMed PMID: 25890197. Pubmed Central PMCID: 4372173.
71. Toobert DJ, Hampson SE, Glasgow RE. The Summary of Diabetes Self-Care Activities Measure. *Diabetes Care*. 2000;23(7):943-50.
72. IDF. Type 2 diabetes 2019 [09/18/2019]. Available from: <https://idf.org/52-about-diabetes.html>.
73. Koizumi J, Matsukura T, Oyama O, Maeda T, Nomura H, Asano A, et al. Influence of Physician Specialty on Treatment Goals for Diabetic Patients : Results of a Survey given to the Members of the Ishikawa Medical Association. *General Medicine*. 2008;9(2):71-9.
74. De Berardis G, Pellegrini F, Franciosi M, Belfiglio M, Di Nardo B, Greenfield S, et al. Quality of care and outcomes in type 2 diabetic patients: a comparison between general practice and diabetes clinics. *Diabetes Care*. 2004 Feb;27(2):398-406. PubMed PMID: 14747220.
75. Swickert RJ, Hittner JB, Harris JL, Herring JA. Relationships among Internet use, personality, and social support. *Computers in Human Behavior*. 2002;18(4):437-51.
76. Cacioppo JT, Cacioppo S. Social Relationships and Health: The Toxic Effects of Perceived Social Isolation. *Social and personality psychology compass*. 2014 Feb 1;8(2):58-72. PubMed PMID: 24839458. Pubmed Central PMCID: 4021390.
77. Nowland R, Necka EA, Cacioppo JT. Loneliness and Social Internet Use: Pathways to Reconnection in a Digital World? *Perspectives on Psychological Science*. 2018 Jan;13(1):70-87. PubMed PMID: 28937910.
78. Rasmussen B, Dunning P, O'Connell B. Young women with diabetes: using Internet communication to create stability during life transitions. *Journal of clinical nursing*. 2007 Mar;16(3A):17-24. PubMed PMID: 17518865.

79. Talukder M, Quazi A. The impact of social influence on individuals' adoption of innovation. *Journal of Organizational Computing and Electronic Commerce*. 2011;21(2):111-35.
80. Ramirez AG, Turner BJ. The role of peer patients in chronic disease management. *Annals of internal medicine*. 2010 Oct 19;153(8):544-5. PubMed PMID: 20956712.
81. Jones KR, Lekhak N, Kaewluang N. Using mobile phones and short message service to deliver self-management interventions for chronic conditions: a meta-review. *Worldviews on evidence-based nursing / Sigma Theta Tau International, Honor Society of Nursing*. 2014 Apr;11(2):81-8. PubMed PMID: 24597522.
82. Riley WT, Rivera DE, Atienza AA, Nilsen W, Allison SM, Mermelstein R. Health behavior models in the age of mobile interventions: are our theories up to the task? *Translational Behavioral Medicine*. 2011;1(1):53-71.
83. Centers for Disease Control and Prevention C. Chronic Disease Overview 2017 [09/06/2018]. Available from: [www.cdc.gov/chronicdisease/overview/index.htm](http://www.cdc.gov/chronicdisease/overview/index.htm).
84. Council for Medical Schemes. Chronic Diseases List 2018 [09/06/2018]. Available from: [www.medicalschemes.com/medical\\_schemes\\_pmb/chronic\\_disease\\_list.htm](http://www.medicalschemes.com/medical_schemes_pmb/chronic_disease_list.htm).
85. Scheibe M, Reichelt J, Bellmann M, Kirch W. Acceptance factors of mobile apps for diabetes by patients aged 50 or older: a qualitative study. *Medicine* 20. 2015 Mar 2;4(1):e1. PubMed PMID: 25733033. Pubmed Central PMCID: 4376102.
86. Chib, A. & Lin. S. Theoretical advancements in mHealth: A systematic review of mobile apps. *Journal of Health Communication*. 2018: 23(10-11): 909-955.
87. MOH Ministry of Health, Singapore. War on Diabetes 2019 [10/24/2019]. Available from: <https://www.moh.gov.sg/wodcj>.
88. SingStat. Population Trends 2019 [2019]. Available from: <https://www.singstat.gov.sg/-/media/files/publications/population/population2019.pdf>.