

Seasonal variation of antioxidant activity and properties in *Ulva lactuca* Linnaeus, 1753: A compression study in pre and post monsoon to the northern coasts of the Oman Sea

Ali Mahdi Abkener*¹ Corresponding E-mail address: aliabkenar@gmail.com Department of Fisheries Group, Chabahar Branch, Islamic Azad University, Chabahar, Iran.

Ali Mahdi Abkener (✉ aliabkenar@gmail.com)

Research Article

Keywords: *Ulva lactuca*, Antioxidant activity, Monsoon, Oman Sea

Posted Date: May 17th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1612579/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Ulva lactuca (Ul) is a green macroalgae that contains commercially valuable molecules components. The purpose of this study was to evaluate and compare seasonal variations of antioxidant activity (AOA) of *U. l.* sample harvested from locations of Tang, Chabahar and Gowatr in the northern coast of the Oman Sea (IRAN). Sampling was carried out during April to May (spring) and October to November (autumn) 2020, representing pre- and post-monsoon, respectively. No significant difference ($P > 0.05$) were observed between samples harvested in pre- and post-monsoon from all three locations, except for the ferric reducing antioxidant power (FRAP) content. The FRAP on post-monsoon shows a significant difference between all studied locations ($P < 0.05$). In this season, the FRAP content was measured 69 ± 0.05 ; 0.60 ± 0.06 and 0.50 ± 0.06 mg GAE g^{-1} for Tang, Chabahar and Gowatr, respectively ($P < 0.05$). However, a high linear correlation (> 0.95) was calculated for parameters showing a significant contribution by β -carotene (β C) with FRAP content ($R = -0.004$) in pre-monsoon, as well as total phenolic content (TPC) with anthocyanin content (ACN) ($R = -0.013$); Half-maximal inhibitory concentration (IC_{50}) with total flavonoid content (TFC) ($R = -0.010$) and radical scavenging activity assay (DPPH) with IC_{50} content ($R = 0.022$) in post-monsoon. AOA analysis along with PCA (principal component analysis) in pre-monsoon confirm relationships and FRAP, TPC, ACN and β C determined sample distribution along PC1 (main components 1), while DPPH, IC_{50} and TFC along the PC2. However, in post monsoons all parameters determined sample distribution along PC1.

Introduction

Due to the special geographical features and ecological valuables, the Oman Sea and coasts are known as one of the rarest ecosystems in the worldwide (Al-Majed et al. 2008). Algae is one of the most important organism's lives in these Sea (Egan et al. 2013).

The findings of recent decades have been reported macroalgae are containing many potent antioxidants, antimicrobials, antiviral, anti-inflammatory, antitumor and anticancer traditional properties, which make them for suitable ingredients to possible applications such as nutraceuticals, pharmaceuticals, cosmetic industries and functional foods (dominguez and loret 2019; Arguel and Sapin 2020; Cagalj et al. 2021).

Green algae as the most diverse algae group in the world has become the focus of many studies, because they are easily accessible and extracts from them are a rich source of antioxidants, carotenoids, polyphenols and flavonoid compounds, as well as natural colorant in various natural product classes (Pérez et al. 2016; Handajani and Prabowo 2020; Prasedya et al. 2020; Fournière et al. 2021). Among the various green algae species from the genus *Ulva lactuca* L. to the Chlorophyta phylum (Ulvaceae family, order Ulvales) is one of the most common species that abound in the north coast of the Oman Sea, Iran (Sohrabipour and Rabii 1999). This species contains commercially valuable biologically active molecules components which are important in different industries.

The study of the bioactive components of algae has increased in recent years and most of the focus has been on the identification of new components (Mekinic et al. 2019; Medina-Torres et al. 2017; Sanjeewa et al. 2017). But, the high potential impact of seasonal variations on metabolite and bioactivity profiles of algae, often was ignored in analytical studies and drug discovery.

Although, the effects of various seasonal acclimation on algae bioactive compounds have been reported in various studies (Prasedya et al. 2019; Karkhaneh Yousefi et al. 2020). But, the monsoon acclimation was not well studied (Billones et al. 2016; Mahdi Abkenar et al. 2021). The monsoon has a significant impact on the environmental and ecological features of the Oman Sea and coasts (Mahdi Abkenar et al. 2021). Algal bloom is one of the most important phenomenon of the monsoon effects (Hunter 1993).

Despite the fact that antioxidant activity (AOA) and the content of phenolic compounds of algae have been widely investigated, comprehensive studies on qualitative and quantitative characteristics in various seasonal conditions using various AOA measurement methods are still lacking. In addition, information about AOA of the Ul from the northern coast of the Oman Sea is rather limited. Due to effect of environmental indices on mitigate oxidative damage of algae in this study we evaluate and compare seasonal variations of antioxidant components (total phenol, flavonoids, β -carotene and anthocyanin contents) and AOA of Ul sample harvested in three different locations of Tang, Chabahar and Gowatr in the Oman Sea coast (IRAN) at different seasonal variation of pre (spring) - and post (autumn) monsoon.

Materials And Methods

Chemicals

Sulfuric acid, Folin–Ciocalteu reagent, and aluminum chloride were from Merck (Germany); ammonium molybdate, potassium acetate, DPPH (2, 2-diphenyl-1-picrylhydrazyl), and sodium phosphate, gallic acid, rutin, ascorbic acid, were from Sigma-Aldrich (Germany).

Seaweeds collection

Seaweed *Ulva lactuca* (Ul) specimens were obtained during the low tide times (according to the tide time table obtained from www.tideforecast.com) in 2020 for two seasons during April to May (pre-monsoon) and October to November (post-Monsoon) representing the autumn and spring randomly from the northern coasts of the Gulf of Oman, Iran (Tang port: Longitude, 25° 35' 79" N; Latitude: 59° 89' 02" E, Chabahar port: Longitude, 25° 27' 71" N; Latitude: 60° 66' 77" E; and Gowatr port: Longitude, 25° 17' 59" N; Latitude: 61° 50' 22" E). Seaweed taxonomic was determined by the off-shore fisheries research center. Algae was identified by Aleem (1993), they belonged to the family: *U. lactuca* Linnaeus from Chlorophyta.

Preparation of algae methanolic extract

After the collection, algae is taken to the laboratory in a plastic bag that contains sea water to prevent evaporation. Epiphytic and foreign materials are removed by first being washed in sea water and then

distilled water to separate the potential contaminants. Dry samples were prepared by drying fresh seaweed in the air at room temperature for 7 days to dry and stored in a plastic bag for further analysis. (González del Val et al. 2001).

For the *Ul* extract preparation, first dry samples are based and powdered. Then 20 g of the *Ul* powder which was dried in air and added to 10 mL of 80% methanol and stored at 37 °C for 6 hours in an orbital shaker incubator (AXYOS, Australia) and centrifugation (Kubota, KN-70, HD8292, Japan) in 1500 × RPM for 15 minutes. After 72 hours of obtaining agent extracts centrifuged. Finally, the extract obtained was filtered by WHATMAN no.1 filter paper and was concentrated to dry to produce a rough extract residue. Extract was diluted with solvents (50 mg mL⁻¹) and stored at 4C used for further analysis (Chidambarajan et al. 2019).

Antioxidant activity (AOA)

The AOA of the *Ul* extract was determined with two different mechanisms of action, i.e., hydrogen atom transfer (HAD) (DDPH) and electron transfer (ET) (FRAB) (Granato, et al. 2018).

2, 2-diphenyl-1-picrylhydrazyl (DPPH %) radical scavenging ability

The DPPH scavenging activity assay of the *Ul* extract was investigated following Duan et al (2006). Briefly, 1 mL of 0.15 mM DPPH were added to a different extract dilution (amounted to 1 mL). Then, the reaction mixture was incubated in the dark at room temperature for 30 minutes. Its absorbance was measured at 517 nm using a spectrophotometer (ELISA Reader Epock, USA). DPPH (%) free radical referral activity is calculated using the following equation (Chidambarajan et al. 2019).

$$\text{Scavenging activity (\%)} = \frac{A_{\text{Sample}} - A_{\text{Control}}}{A_{\text{Control}}} \times 100$$

The half-maximal inhibitory concentration (IC₅₀) were calculated by the linear regression analysis and expressed as mean of three determinations. IC₅₀ was used as an index to compare the AOA of individuals. Ascorbic acid was used as positive control and reducing power was expressed as mg ascorbic acid equivalents (AAE) per mL dry weight of sample (DW).

Ferric Reducing Antioxidant Power (FRAP) Assay

The FRAP assay were determined using the Oyazu (1986) modification method. Briefly, 1 mL the *Ul* extract was added to 1 mL 0.2 M phosphate buffer (PSB, pH=6.6) and 1 mL of 1% potassium ferricyanide. The mixture was incubated at 50 °C for 20 minutes and 1 mL of 10% of TCA was added to this reaction mixture. Aliquot 1 mL from the incubation mixture were mixed with 1 mL of distilled water and 0.2 mL of 0.1% ferric chloride in the test tube. Absorbance was measured at 700 nm. Gallic acid was used as a standard power and reduction expressed as equivalent to mg gallic acid (GAE) per g dry weight sample (DW).

Total Phenolic and Flavonoid Content

Total phenolic content (TPC) of the UI extract were determined by the Folin-Ciocalteu method (Cagalj et al., 2021) using a UV spectrophotometer (Shimatzu, UV-160A, Japan) with minor modification. Briefly, 20 μL of the UI extract reacted with 1.50 mL of distilled water and 125 μL of 10% of folin-Ciocalteu reagent. The solution was mixed and after a minute, 375 μL of 20% sodium carbonate solution and 475 μL distilled water were added. The mixture was left in the dark for 2h at room temperature. Absorbance was read at 720 nm using a spectrophotometer. The Standard curve (0-500 mg L^{-1}) was prepared using gallic acid for calculation of phenolic content ($y = 0.004 \cdot x$, $r^2 = 0.9999$). The TPC were expressed as mg Gallic acid equivalents (GAE) per g dry weight sample (DW).

Total content flavonoids (TFC) of extract the UI assessed using the aluminum chloride colorimetric method described by Cagalj et al. (2021). Briefly, 250 μL the UI extract was mixed with 750 μL 96% ethanol, 50 μL 10% (w/v) aluminum chloride, 50 μL 1M sodium acetate, and 1400 μL distilled water. The mixture was stored for 30 minutes at room temperature. Absorbance was read at 415 nm using a spectrophotometer. Standard calibration curve (0-100 $\mu\text{g mL}^{-1}$) is plotted using a Quercetin solution ($Y = 0.0076 - 0.0042 \cdot x$, $r^2 = 0.9412$). TFC is expressed as mg Quercetin equivalent (QE) per g dry weight sample (DW).

Total anthocyanin content

Determination of total anthocyanin content was measured using a differential protocol spectrophotometric pH, as described by Muanda (2001) with minor modifications. Briefly, 0.5 mL of the UI extract was mixed with 3.5 mL of 0.025 M potassium chloride buffer; PH=1. The mixture was stirred and left at room temperature for 15 minutes. Then, the absorbance was measured at 515 and 700 nm against distilled water as blank. Then, the extract combined with 3.5 mL 0.025 M sodium acetate buffer; pH=4.5. Again, the mixture was stirred and left at room temperature for 15 minutes. Absorbance of mixture was measured at the same wavelength (512 and 700 nm). Total anthocyanin content was calculated using the following equation:

$$TAC (mgC - 3 - GE/100g DW) = (A \times MW \times 1000 / (\epsilon \times C))$$

Where A is absorbance= $(A_{\lambda 515} - A_{\lambda 700})$ pH 1.0 - $(A_{515} - A_{700})$ pH 4.5; MW is molecular weight for cyanidin-3-glucoside = 484.2; DF is the factor of the extract; ϵ is the molar absorbtivity of cyanidin-3-glucoside = 24825; C is the concentration of the buffer in mg mL^{-1} = 0.025. The results were expressed as milligram of cyanidin-3-glucoside equivalents (mg C-3-GE) per 100 gram of dry weight of sample (DW).

The β -Carotene content

Quantification of β -carotene were carried out following Eijckelhoff and Dekker (1995) with minor modification. Briefly, 20 g the UI crude extract was solubilized in 10 mL 85% acetone-hexane and

centrifuged at 3000 rpm for 15 minutes at 4 °C. Then, the absorbance measured at 453, 505, 645 and 663 nm using a UV spectrophotometer (Shimatzu, UV -160A, Japan). The contents of the β -carotene are calculated according to the following equations and expressed as mg per g of dry weight (DW).

$$\beta - \text{Carotene (mg } 100\text{g}^{-1} \text{ DW)} = 0.216A_{663} - 1.22A_{645} - 0.304A_{505} + 0.452A_{453}$$

Statistical analysis

All statistical analysis was conducted using SPSS Software (New York, USA) version 16.0 in a triplicate. Shapiro-Wilk and Levenes tests were used to check the normality and homogeneity of variance, respectively. To determine whether there are differences between means, ANOVA analysis and Duncan's new multiple distance test is applied to correlate the potential of antioxidants in the samples. In addition, comparison is made between monitored factors and locations in pre and post monsoon with independent sample tests. A principal component analysis (PCA) was carried out to obtain an overview of the equations and differences in antioxidant activities and properties of the UI extract to detect patterns in the data and to reduce the dimensions of information for discrimination analysis. Data is declared a means of \pm standard deviation (SD) of three replication determinations. P value <0.05.

Results

DPPH (%) scavenging activity

Table 1, indicating no significant variation of scavenging effects from *Ulva lactuca* (UI) extract tested at 1, 10, 100 and 1000 $\mu\text{g mL}^{-1}$ on the DPPH (%) of methanolic extract of the UI harvested in pre- and post-monsoon among three locations of Tang chabahar and Gowatr ($P > 0.05$).

Table 1 DPPH radical scavenging activity (%DPPH) of 1, 10, 100 and 1000 $\mu\text{g mL}^{-1}$ of the UI extracts in different seasons.

Seasons	Concentration ($\mu\text{g mL}^{-1}$)	Stations		
		Tang	Chabahar	Gowatr
Pre- (Spring)	1	39.46 \pm 1.37	35.72 \pm 1.74	37.9 \pm 1.91
	10	42.19 \pm 83	39.80 \pm 2.01	40.15 \pm 0.45
	100	47.28 \pm 3.34	45.20 \pm 1.15	46.79 \pm 0.74
	1000	50.43 \pm 1.19	48.96 \pm 1.23	50.90 \pm 3.01
Post- (autumn)	1	47.19 \pm 0.85	44.76 \pm 0.99	41.38 \pm 1.09
	10	54.27 \pm 3.45	52.96 \pm 1.36	49.76 \pm 2.59
	100	57.98 \pm 2.36	55.80 \pm 2.78	52.26 \pm 2.34
	1000	60.50 \pm 2.54	59.12 \pm 2.46	55.81 \pm 2.19
	Ascorbic acid	55.1 \pm 1.00	55.1 \pm 1.00	55.1 \pm 1.00

*Data were presented as the average of four replicates standard deviation (\pm SD).

IC₅₀

As can be seen in the Fig 3, The IC₅₀ content of ascorbic acid as a standard antioxidant has not shown a significant difference ($P < 0.05$) between both pre- (104.48 \pm 6.96 mg AAE mL⁻¹) and post- (108.41 \pm 7.89 mg AAE mL⁻¹) monsoon (sig= 0.28; t= -111; DF = 15.75). However, no significant different were observed between three locations of Tang, Chabahar and Gowatr during pre and post monsoon ($P > 0.05$). IC₅₀ content in the UI extract in pre-monsoons ranged from 94.41 to 115.40 mg AAE mL⁻¹, whereas in post monsoons were found in 98.51 to 118.71 mg AAE mL⁻¹.

Ferric Reducing Power Assay (FRAP)

The Fig 4, presents the compression of the FRAP content in the UI extract among three different locations and seasonal variations on the Northern coast of the Oman Sea. According to the results there was no significant difference found between the UI extract which was harvested in pre- (0.58 \pm 0.08 mg GAE g⁻¹ DW) and post- (0.59 \pm 0.09 mg GAE g⁻¹ DW) monsoon (sig= 0.85; t= -0.18; DF= 15.59). However, there was no significant difference found in three locations of samples harvested at pre-monsoon ($p > 0.05$); Meanwhile, a statistically significant difference was observed in the samples harvested in post-monsoon between all locations ($P < 0.05$). The highest (0.69 \pm 0.05 mg GAE g⁻¹ DW) and the lowest (0.50 \pm 0.06 mg GAE g⁻¹ DW) of the FRAP content were observed at the Tang and Gowatr site, respectively. The FRAP content measurement in the UI sample harvested from Chabahar was 0.60 \pm 0.06 mg GAE g⁻¹ DW. The content of this index in the UI extract harvested in the pre-monsoons ranges from 0.42 to 0.71 mg GEA g⁻¹ DW, and the sample harvested in the post-monsoon was 0.44 to 0.75 mg GEA g⁻¹ DW.

Total Phenolic content (TPC)

Compression of The TPC content in the Ul. extract presented in the Fig 5. Based on the results for both pre- (3.32 ± 0.77 mg GAE g^{-1} DW) and post- (3.38 ± 1.00 mg GAE g^{-1} DW) monsoon, no significant variations were observed (sig= 0.27; t= -1.13; DF= 15.97). However, no significant differences were observed between three different locations of Tang, Chabahr and Gowatr during pre and post monsoon ($P > 0.05$). The TPC compounds in extract harvested in the pre-monsoons range from 2.14 to 4.13 mg GAE g^{-1} DW, whereas for samples harvested in the post monsoon were found at 2.15 to 4.95 mg GAE g^{-1} DW.

Total Flavonoid Content (TFC)

As shown in the Fig 6, there were no significant differences between the pre- (0.38 ± 0.11 mg QE g^{-1} DW) and the post- (0.35 ± 0.07 mg QE g^{-1} DW) monsoon for methanolic the Ul extract (Sig = 0.54; T=0.62; DF=14.04), as well as between three different locations of Tang, Chabahr and Gowatr during pre and post monsoon ($p > 0.05$). The content of total flavonoid compounds in the pre-monsoons ranges from 0.22 to 0.55 QE g^{-1} DW, while the post monsoon was found at 0.25 to 0.47 mg QE g^{-1} DW.

Anthocyanin Content (ACN)

Fig 7, showing no significant ($P > 0.05$) seasonal variation for the anthocyanin content from the Ul extracts harvested in pre (3.41 ± 0.44 mg C-3-GE $100g^{-1}$ DW) and post (3.70 ± 0.37 mg C-3- GE $100g^{-1}$ DW) monsoon (Sig= 0.15; t= -1.49; DF = 15.57). However, no significant were observed for the samples harvested in three different locations of Tang, Chabahr and the Gowatr during pre and post monsoon ($P > 0.05$). The content of anthocyanin compounds in the pre-monsoons ranges from 2.87 to 3.71 mg C-3- GE $100g^{-1}$ DW, while in the post monsoon was found at 2.98 to 4.03 mg C-3-GE $100g^{-1}$ DW.

The β -Carotene content (β C)

Fig 8, showing no significant seasonal variations ($P > 0.05$) in the contents of β C the Ul extract in the pre (0.76 ± 0.10 mg g^{-1} DW) and the post (0.80 ± 0.14 mg g^{-1} DW) monsoon (Sig= 0.53; t= -0.63; DF= 14.62), and among the three differential Tang locations, chabahr and Gowatr during pre and post monsoon ($p > 0.05$). The β -carotene content in methanolic extract of the Ul harvested in pre-monsoon ranges from 0.57 to 0.88 mg g^{-1} DW, while on the post monsoon found in 0.49 to 0.97 mg g^{-1} DW.

Correlation analysis

Pearson correlation analysis was carried out to determine the relationship between each antioxidant activity index and properties in different seasonal sampling. As shown in Table 2, there is a significant negative correlation between FAR with β -carotene content ($r = -0.004$; $p < 0.05$) in pre-monsoon.

Table 2 Pearson's correlation coefficients between the variables to pre-monsoon

Indices	DDPH	IC ₅₀	FRAP	TPC	TFC	ACN	βC
DPPH	1	-0.219	0.467	-0.378	-0.105	-0.028	-0.203
IC ₅₀	-0.219	1	-0.112	-0.159	0.373	-0.115	-0.258
FRAB	0.467	-0.112	1	0.544	0.058	0.297	-0.004
TPC	-0.378	-0.159	0.544	1	-0.106	0.230	0.441
TFC	-0.105	0.373	0.058	-0.106	1	0.269	-0.246
Anthocyanin	-0.028	-0.115	0.297	0.230	0.269	1	0.181
β-carotene	-0.203	-0.258	-0.004*	0.441	-0.246	0.181	1

*=P<0.05

However, on post monsoon there were a significant negative correlation between TPC with Anthocyanin content ($r = -0.013$; $P < 0.05$), as well as between IC₅₀ with TFC content ($r = -0.010$; $P < 0.05$). There were also a significantly positive correlation between DPPH with IC₅₀ content ($r = 0.022$; $P < 0.05$).

Table 3 Pearson's correlation coefficients between the variables to post-monsoon

Variables	DPPH	IC ₅₀	FRAP	TPC	TFC	ACN	βC
DPPH	1	0.022*	0.427	-0.088	0.327	0.053	0.061
IC ₅₀	0.022	1	0.544	0.340	-0.010*	0.643	0.578
FRAP	0.427	0.544	1	0.247	0.486	0.373	0.160
TPC	-0.088	0.340	0.247	1	-0.203	-0.013*	-0.057
TFC	0.327	-0.010*	0.486	-0.203	1	0.419	-0.155
Anthocyanin.	0.053	0.643	0.373	-0.013*	0.419	1	0.687
β-carotene	0.061	0.578	0.160	-0.057	-0.155	0.687	1

*=P<0.05

Based on the PCA results in pre monsoon, the first three PCs (principal components) were extracted with the value of eigenvalue greater than 1 and together they explain 72.58% of the total variance (Table 4). PC1 explained 28.76% of data set variations and PC2 explained 22.27% (Fig 9). According to the value loaded, which expresses the correlation between PCs and variables, it can be stated that FRAP, TPC, anthocyanin and β-carotene were determined by the sample distribution along PC1, while DPPH, IC₅₀ and TFC along PC2 (Fig 9a). This was proven evidenced by high r Pearson's correlation coefficients between these variables and suggests that the AOA from the UI extract was mainly associated with phenolic

compounds. PCA analysis confirmed that the UI harvested in Tang, Chabaha and Gowatr possess high AOA and phenolic compounds, including DPPH, TPC and FRAP contents (Fig 9b).

Table 4 eigenvectors and eigenvalues of the first three principal components (PCs) for 7 qualitative pod characters in pre-monsoon

Variable	Principal components		
	PC1	PC2	PC3
Eigenvalue	2.013	1.559	1.508
Variability (%)	28.763	22.271	21.547
Cumulative (%)	28.763	51.034	72.581
Eigenvectors			
DPPH	-0.026	0.743	-0.176
IC ₅₀	-0.355	-0.172	0.446
FRAP	0.426	0.501	0.262
TPC	0.586	-0.208	0.177
TFC	-0.187	0.027	0.671
Anthocyanin	0.330	0.080	0.435
β-carotene	0.453	-0.343	-0.175

On post monsoon, the first three PCA of the UI were extracted with eigenvalue value greater than 1 and together they explained 79.49% of the total variance (Table 4). PC1 explained 38.48% of variations of data sets and PC2 explained 23.36% (Fig 10). According to the value loaded, which expresses the correlation between PCs and variables, can be stated that all studied parameters (DPPH, IC₅₀, FRAP, TPC, TFC, Anthocyanin and β-carotene) determine the distribution of the samples along PC1 (Fig. 10a). This is proven by the high R Pearson correlation coefficient between these variables and suggests that the AOA from the UI extract were mainly associated with phenolic compounds. PCA analysis confirmed that samples harvested for Tang showed high AOA and phenolic compounds, including total anthocyanin, IC₅₀, FRAP, β-carotene and TFC contents (Fig. 10b).

Table 5 eigenvectors and eigenvalues of the first three principal components (PCs) for 7 qualitative pod characters in post-monsoon

Variable	Principal components		
	PC1	PC2	PC3
Eigenvalue	2.693	1.635	1.236
Variability (%)	38.466	23.364	17.662
Cumulative (%)	38.466	61.830	79.492
Eigenvectors			
DPPH	0.192	0.494	0.088
IC ₅₀	0.504	-0.301	0.155
FRAP	0.449	0.291	0.364
TPC	0.110	-0.276	0.762
TFC	0.241	0.605	-0.094
Anthocyanin	0.521	-0.080	-0.313
β-carotene	0.407	-0.362	-0.385

Discussion

Generally, in this study antioxidant activity of the UI extract that harvested during pre- (spring) and post- (autumn) monsoon do not shows significant differences between the variables tested, except for FRAP content. Our results indicate that the FRAP content has a significant difference between seaweed harvested on post-monsoon between all three locations. The highest and lowest activity of the FRAP content were in Tang ($0.69 \pm 0.05 \text{ mg GAE g}^{-1}$) and Gowatr ($0.50 \pm 0.06 \text{ mg GAE g}^{-1}$), respectively. However, the FRAP content measured in Chabahr was $0.60 \pm 0.06 \text{ mg GAE g}^{-1}$. These results indicate that AOA does not change with an increase or decrease of temperature during the pre and post monsoon, except for the FRAP content. This observation was supported by previously studies in other seaweed which also reported a different AOA test (Fellah et al. 2017; Jelandar et al. 2017; Farasat et al. 2013, 2014; Liu et al. 2011; Horincar et al. 2011). The FRAP concentration reported in this manuscript is similar to the reported for the UI ($0.60 \pm 0.12 \text{ mg GAE g}^{-1}$) harvested in the autumn on the intertidal coast of Chabahr (Iran), using methanolic extract (Sarani Yaztapeh 2020). Moulazadeh et al (2021) reported that the Persian Gulf patched the UI extract collected in the autumn had $64.89 \pm 11.33 \text{ } \mu\text{mol Fe}^{2+}/\text{g}$. Sivaramakrishnan et al (2017) measure AOA and phenolic compounds of four green macroalgae (*Halimeda Tuna*, *Halimeda Macroloba*, *enteromorpha* sp. and *Acetabularia Acetabulum*) were collected from the coast of South Andaman (India) in three different locations. According to the author reporting the FRAP content was around from 152.89 ± 2.53 to $164.3 \pm 1.04 \text{ } \mu\text{g g}^{-1}$. Likewise, seasonal variations of the FRAP content followed the patterns reported for *Halopetrical Scoparia* in the same study as significant differences found in the spring and fall (Fellah et al. 2017). Karkhaneh Yousefi et al (2020) by measured the FRAP content in four different brown algae (*Dictyota indica*, *Padina tenuis*, *Colpomenia*

sinuosa and *Iyengaria stellata*) harvested from Qeshm Island (Persian Gulf) reported that Antioxidant activities of all summer collected seaweeds were lower than the winter ones which were in accordance with their fucoxanthin contents. Contrary to our results, Trigui et al (2013) observed the highest free radical scavenging activity in *U. rigida* extract collected at the end of winter (February) and the beginning of spring (March) compared to the end of summer (August) and the beginning of fall (September-October). Based on the results, the authors suggested AOA increase with a decrease of temperature (Trigui et al. 2013).

However, variations of the FRAP concentration obtained from the UI and other seaweed species can be associated with environmental factors such as various species compositions, seasonal variations, salinity, irradiation and inorganic and phosphorus nitrogen and the type of extraction solvent used (Oucif et al. 2017; Aruelles et al. 2018; Aruelles et al. 2021). Increased solar radiation and UV exposure during summer have also been associated with an increase in the production of antioxidant compounds in intertidal macroalgae at the monsoon (Garcia-Vaquero et al. 2021). In addition, inorganic nutrition released from shrimp farms in Gowatr is considered to be ignored compared to nitrogen background concentrations that occur naturally (Marinho et al. 2015c).

To investigate whether AOA from the UI extract studied, a significant correlation was observed in the pre-monsoon between the FRAP with β -carotene contents and other antioxidant values except FRAP. It shows carotenoids having antioxidant activities in reducing Fe^{3+} radicals on the UI. Here, there was a significant correlation between IC_{50} with free radical of DPPH, IC_{50} with TPC, and TPC with anthocyanin content in the post monsoon confirm the literature report and may indicate that this compounds are responsible for AOA or at least involved in the reaction of the effect at the post monsoon, which is appropriate with many previous studies aruelles (2021); Arguelles and Sapin 2020; Jesumani et al (2020); Sanger et al (2019); Sivaramakrishnan et al (2017); Farasat et al (2013; 2014); Chew et al. (2008).

Likewise, a lower correlation between β -carotene with DPPH content, IC_{50} , TPC, TFC and Anthocyanin which is only a phenolic compound that is not involved in AOA through this path but there may be several effects involving other active compounds.

Conclusion

In the current study, the assessment of the UI antioxidant activities and properties extract which was harvested from three different locations during per (spring) and post (autumn) monsoon was evaluated. The FRAP content in post monsoons shows a significant difference between all three locations. The highest and lowest FARP content were found in the Tang and Gowatr, respectively. Perhaps, the location of Tang sampling does not yet have extreme environmental pollution compared to Chabahar and Gowatr such as lower tidal ranges that contribute to the sunlight exposure, and disposal of industrial human activities such as fisheries and waste making it into the sea. Based on the obtained results from this study we could claim that macroalgae (*Ulva lactuca*) provides promising antioxidant properties that can be used in various food and pharmaceutical industries.

Abbreviations

UI

Ulva lactuca

AOA

Antioxidant activity

FRAP

Ferric reducing antioxidant power

TFC

Total flavonoid content

TPC

Total phenolic content

ACN

Total anthocyanin content

β C

β -carotene

Declarations

Data Availability Statements

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Author Contributions Statement

Ali Mahdi Abkenar conceived the original idea. Developed the theory and performed the computations. Verified the analytical methods. Carried out the experiment. Discussed the results and wrote the final manuscript.

Conflict of interest

The author declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Aleem AA (1993) The marine algae of Alexandria, Egypt. 139P.
2. Allen LB, Sitonen PH, Thomposon HC (1997) Methods for the determination of arsenic, copper, lead and tin in sucrose, corn syrups and high fructose corn syrups by inductively coupled plasma atomic emission spectrophotometry. J Agric Food Chem 45: 162–165.

3. Antolovich M, Prenzler PD, Patsalides E, Mc Donald S, Robards K (2002). Methods for testing antioxidant activity. *Analyst* 127(1):183-198.
4. Arguelles EDLR (2021) Evaluation of Antioxidant Capacity, Tyrosinase Inhibition, and Antibacterial Activities of Brown Seaweed, *Sargassum ilicifolium* (Turner) C. Agardh 1820 for Cosmeceutical application. *J Fish Environ* 45 (1):64-77.
5. Arguelles EDLR, Monsalud RG, Sapin AB (2019) Chemical composition and *In vitro* antioxidant and antibacterial activities of *Sargassum vulgare* C. Agardh from Lobo, Batangas, Philippines. *J Int Soc Southeast Asian Agri Sci* 25(1): 112-122.
6. Arguelles EDLR, Sapin AB (2020) Bioactive properties of *Sargassum siliquosum* J. Agardh (Fucales, Ochrophyta) and its potential as source of skin-lightening active ingredient for cosmetic application. *J Appli Pharm Sci* 10(7):51-58.
7. Astorg P (1997) Food carotenoids and cancer prevention: an overview of current research. *Trends Food Sci. Technol* 8:406–413.
8. Azam MS, Choi J, Lee MS, Kim HR (2017) Hypopigmenting effects of brown algaederived phytochemicals: A review on molecular mechanisms. *Mar Drugs* 15(10):297. DOI: 10.3390/md15100297.
9. Benita M, Dubinsky Z, Iluz D (2018) *Padina Pavonica*: Morphology and Calcification Functions and Mechanism. *Am J Plant Sci* 9:1156–1168.
10. Bernardini G, Minetti M, Polizzotto G, Biazzo M, Santucci A (2018) Pro-Apoptotic Activity of French Polynesian *Padina Pavonica* Extract on Human Osteosarcoma Cells. *Mar Drugs* 16:504.
11. Billones X.K, Quiao M.A, Roa E.C, Roxas, P.G (2016). Potential Antioxidant Activity of Three *Kappaphycus* spp. Cultivars (Rhodophyta, Solieriaceae) Collected during Two Monsoon Seasons in Kolambugan, Lanao del Norte, Philippines. DOI: 10.48031/msunyear.2016.04.01
12. Biris-Dorhoi ES, Michiu D, Pop CR, Rotar AM, Tofana M, Pop OL, Socaci SA, Farcas AC (2020) Macroalgae-A Sustainable Source of Chemical Compounds with Biological Activities. *Nutrients* 12:3085.
13. Čagalj M, Skroza D, Tabanelli G, Özogul F, Šimat, V (2021) Maximizing the antioxidant capacity of *Padina pavonica* by choosing the right drying and extraction methods. *Processes* 9(4):587.
14. Chewa YL, Lima, YY, Omara M, Khoo KS (2008) Antioxidant activity of three edible seaweeds from two areas in South East Asia. *LWT* 41:1067-1072.
15. Chidambararajan P, Keerthana V, Priyadarshini K, Sakthivel B (2019) In vitro antioxidant and anticancer activity of *Ulva lactuca* l. using molt-3 cell line. *Asian J Pharma Clin Reas* 12(5):75-78.
16. Cotas J, Leandro A, Monteiro P, Pacheco D, Figueirinha A, Gonca Ives AMM, Da Silva GJ, Pereira L (2020) Seaweed Phenolics: From Extraction to Applications. *Mar Drugs* 18:384.
17. Dominguez H, Loret EP (2019) *Ulva lactuca*, A Source of Troubles and Potential Riches. *Mar Drugs* 17:357. Doi: 10.3390/md17060357.

18. Duan X.J, Zhang WW, Li XM, Wang BG (2006) Evaluation of antioxidant property of extract and fractions obtained from a red alga, *Polysiphonia urceolata*. *Food Chem* 95:37-43.
19. Egan S, Harder T, Burke C, Steinberg P, Kjelleberg S, Thomas T (2013) The seaweed holobiont: understanding seaweed-bacteria interactions. *FEMS Microbiol Rev* 37:462–476.
20. Farasat M, Khavari-Nejad RA, Nabavi SMB, Namjooyan F (2014) Antioxidant Activity, Total Phenolics and Flavonoid Contents of some Edible Green Seaweeds from Northern Coasts of the Persian Gulf. *Iranian J Pharma Res*13 (1):163-170.
21. Farasat M, Khavari-Nejad RA, Seyed Nabavi M.B, Namjooyan F (2013) Antioxidant Properties of Some Filamentous Green Algae (*Chaetomorpha* Genus). *Brazilian Arch. Biol. Tech.* 56(6):921-927.
22. Fellah F, Louaileche H, Dehbi-Zebboudj A, Touati N (2017) Seasonal variations in the phenolic compound content and antioxidant activities of three selected species of seaweeds from Tiskerth islet, Bejaia, Algeria. *J mater Environ Sci* 8(12):4451-4456.
23. Fournière M, Bedoux G, Lebonvallet N, Lescchiera R, Goff-Pain CL, Bourgougnon N, Latire T (2021) Poly-and oligosaccharide ulva sp. Fractions from enzyme-assisted extraction modulate the metabolism of extracellular matrix in human skin fibroblasts: Potential in anti-aging dermo-cosmetic applications. *Mar Drugs* 19:156.
24. Garcia-Vaquero M, Rajauria G, Miranda M, Sweeney T, Lopez-Alonso M, O'Doherty J (2021) Seasonal Variation of the Proximate Composition, Mineral Content, Fatty Acid Profiles and Other Phytochemical Constituents of Selected Brown Macroalgae. *Mar Drugs* 19:204. <https://doi.org/10.3390/md19040204>.
25. Granato D, Shahidi F, Wrolstad R, Kilmartin P, Melton LD, Hidalgo FJ, Miyashita K, Camp JV, Alasalvar C, Ismail AB, Elmore S, Birch GG, Charalampopoulos D, Astley SB, Pegg R, Zhou P, Finglas P. (2018). Antioxidant activity, total phenolics and flavonoids contents: Should we ban in vitro screening methods?. *Food Chem* 264:471-475. DOI: 10.1016/j.foodchem.2018.04.012.
26. Handajani F, Prabowo S (2020). Sargassum duplicatum extract reduced arthritis severity score and periarticular tissue matrix metalloproteinase- 1 (MMP-1) expression in *Ajuvan arthritis* exposed to cold stressor. *Sys Rev Pharm* 11:302–307.
27. Heavisides E, Rouger C, Reichel AF, Ulrich C, Wenzel-Storjohann A, Sebens S, Tasdemir D (2018) Seasonal variations in the metabolome and bioactivity profile of *Fucus vesiculosus* extracted by an optimised, pressurised liquid extraction protocol. *Mar Drugs* 16(12):503.
28. Horincar VB, Parfene G, Bahrim G (2011) Evaluation of bioactive compounds in extracts obtained from three Romanian marine algae species. *AC Romanian Biotechnological Letters* 16(6):71-78.
29. Hunter JR (1983). Aspects of the dynamics of the residual circulation of the Arabian Gulf. In *Coastal oceanography* (ed. H.G. Gade et al.), pp. 31-42. NewYork: Plenum Press.
30. Jeloodar N, Farasat M, Tadayoni M (2017) Antioxidant Capacity of Methanolic Extracts of Some Brown, Red and Green Seaweeds of The Persian Gulf. 2nd conferences & Exhibition on Methods to Increase the Shelf-life of Food Products, winter. 2017, Tehran, IRAN. 13p.

31. Jesumani V, Du H, Pei P, Aslam M, Huang N (2020). Comparative study on skin protection activity of polyphenol-rich extract and polysaccharide-rich extract from *Sargassum vachellianum*. PLoS ONE 15(1):e0227308. DOI: 10.1371/journal.pone.0227308.
32. Jose GM, Kurup, GM (2016) *In vitro* antioxidant properties of edible marine algae *Sargassum swartzii*, *Ulva fasciata* and *Chaetomorpha antennina* of Kerala Coast. J Pharma Rep 4(6):100-112.
33. Karkhaneh yousefi M, Seyed Hashtroudi M, Mashinchian Moradi A, Ghasempour A (2020) Seasonal variation of fucoxanthin content in four species of brown seaweeds from Qeshm Island, Persian Gulf and evaluation of their antibacterial and antioxidant activities. Iranian J Fish Sci 19(5): 2394-2408. doi: 10.22092/ijfs.2020.122396.
34. Khairy HM, El-Sheikh MA (2015) Antioxidant activity and mineral composition of three Mediterranean common seaweeds from Abu-Qir Bay, Egypt. Saudi J Biol Sci 22:623-630.
35. Kokabi M, Yousefzadi M, Ali ahmadi A, Feghhi MA, Keshavarz M (2013) Antioxidant Activity of Extracts of Selected Algae from the Persian Gulf, Iranian J Persian Gulf (Mar. Sci.) 4(12):45-50.
36. Ling, ALM, Yasir S, Matanjun P, Abu Bakar, MF (2015) Effect of Different Drying Techniques on the Phytochemical Content and Antioxidant Activity of *Kappaphycus Alvarezii*. J Appl Phycol 27:1717–1723.
37. Liu CC, Zhao GI, Li YN, Ding Z, Liu QG, Li JL (2011) Contribution of phenolics and flavonoids to antioxidant activity of ethanol extract from *Eichhornia crassipes*. Adv Mater Res 156-157: 1372-1377.
38. Lovrić V, Putnik P, Bursać Kovačević, D, Jukić M, Dragović-Uzelac V (2017) The Effect of Microwave-Assisted Extraction on the Phenolic Compounds and Antioxidant Capacity of Blackthorn Flowers. Food Technol Biotechnol 55:243–250.
39. Luo HY, Wang B, Yu CG, Qu YL, Su CL (2010). Evaluation of antioxidant activities of five selected brown seaweeds from China. J Med Plants Res 4(8):2557-2565.
40. MacArtain P, Gill CIR, Brooks M, Campbell R, Rowland IR (2007). Nutritional value of edible seaweeds. Nutr Rev 65:535–543.
41. Mahdi Abkenar A, Yahyavi M, Esmaeili M, Rombenso A (2021) High bioaccumulation factors and ecological risk index of Cd, and Hg in Indian white shrimp, hooded oyster, brown algae and sediment in northern coasts of the Gulf of Oman before and after a monsoon. Reg. Stud. Mar. Sci 41:101552.
42. Mantri V.A, Dineshkumar R, Yadav A, Eswaran K, Shanmugam M, Gajaria T.K (2022) Projections for profitability assessment parameters under short-term, medium-term and long-term evaluation for three farming techniques of *Kappaphycus alvarezii* along eastern coast of India. Aquaculture, 737912.
43. Mau JL, Tsai SY, Tseng YH, Huang SJ (2005) Antioxidant properties of methanolic extracts from *Ganoderma tsugae*. Food chem 93(4), 641-649.
44. Medina-Torres N, Ayora-Talavera T, Espinosa-Andrews H, Sánchez-Contreras, A, Pacheco N (2017) Ultrasound Assisted Extraction for the Recovery of Phenolic Compounds from Vegetable Sources. Agronomy 7(3): 47.

45. Muralidhar AP, Karthireddy S, Chandra P, Kalidas C, Naik RP (2010) Comparative studies on fatty acid composition of three marine macroalgae collected from Mandapam region: south east coast of India. *World Appli Sci* 11(8):958-965.
46. Muzolf-Panek M, Stuper-Szablewska, K (2021) Comprehensive study on the antioxidant capacity and phenolic profiles of black seed and other spices and herbs: effect of solvent and time of extraction. *J Food Meas Charact* 15:4561–4574.
47. NRC, National Research Council (1982) *Diet, Nutr. Cancer*. National Academy Press, Washington, DC.
48. Oyaizu, M (1986) Studies on products of browning reaction prepared from glucosamine. *Jpn J Nutr* 44: 307-314.
49. Pérez M, Falqué E, Domínguez, H (2016) Antimicrobial action of compounds from marine seaweed. *Mar Drugs* 14 (3): 52.
50. Prasedya ES, Martyasari NWR, Apriani R, Mayshara S, Fanani RA, Sunarpi H (2019) Antioxidant activity of *Ulva lactuca* L. from different coastal locations of Lombok Island, Indonesia. *Proceedings of the 2nd International Conference on Bioscience, Biotechnology, and Biometrics 2019 AIP Conf. Proc.* 2199, 020003-1–020003-6; <https://doi.org/10.1063/1.5141281>
51. Reynolds RM, (1993) Physical oceanography of the Gulf, Strait of Hormuz, and the Gulf of Oman: results from the Mitchell Expedition. *Mar Pollut Bull* 27: 35-60. DOI. 10.1016/0025-326X (93)90007-7
52. Sanger G, Rarung LK, Kaseger B.E, Assa JR, Agustin AT (2019) Phenolic content and antioxidant activities of five seaweeds from North Sulawesi, Indonesia. *AACL Bioflux* 12(6):2041-2050.
53. Sanjeeva KKA, Lee JS, Kim WS, Jeon YJ (2017) The potential of brown-algae polysaccharides for the development of anticancer agents: An update on anticancer effects reported for fucoidan and laminaran. *Carbohydr Polym* 177:451–459.
54. Sarani Yaztapeh E, Hosseini Tabatabaei MR, Abkenar AM (2021) The compression of antioxidant activity and β -carotene extracted from three species of native algae of Oman Sea (*Ulva lactuca*, *Sargassum ilicifolium* and *Nizimuddinina zanardini*). *Iranian J Fish Sci* 29(6):53-63.
55. Sarani Yaztapeh, E (2020) Quantitative evaluation of the content of the polyphenolic compounds, anthocyanin & β - caroten of three endemic seaweed specis in chabahar zoone and evaluation of their antioxidant effect. Thesis of Ms.c. Islamic Azad University Zahedan Branch, 92P.
56. Silberfeld T, Bittner L, Fernández-García C, Cruaud C, Rousseau F, de Reviere B, Leliaert F, Payri, CE, De Clerck O (2013) Species Diversity, Phylogeny and Large Scale Biogeographic Patterns of the Genus *Padina* (Phaeophyceae, Dictyotales). *J Phycol* 49:130-142.
57. Sirbu R, Stanciu G, Tomescu A, Ionescu AM, Cadar E (2019) Evaluation of Antioxidant and Antimicrobial Activity in Relation to Total Phenolic Content of Green Algae from Black Sea. *Rev Chim (Bucharest)* 70(4):1197-1203.
58. Sivaramakrishnan T, Swain S, Saravanan K, Kiruba Sankar R, Dam Roy S, Biswas L, Shalini B (2017) In Vitro Antioxidant and Free Radical Scavenging Activity and Chemometric Approach to Reveal Their Variability in Green Macroalgae from South Andaman Coast of India. *Tur J Fish Aqua Sci* 17:639-648.

59. Smith MM, Heemstra P.C (1986). *Smith's Sea Fishes*. Springer-Varlag, Heidelberg, 1047P.
60. Sohrabipour J, Rabii, R (1999) Alist of marine algae of seashores of Persian Gulf and Oman Sea in Hormozgan province. *Iran J Bot* 8:131-162.
61. Thinakaran T, Sivakumar K (2012). Seasonal variation and biochemical studies on certain seaweed from Pamban Coast, Gulf of Mannar biosphere review. *Int J Res Biol Sci* 2:39–44.
62. Trigui N, Gasmi L, Zouari I, Tounsi S (2013) Seasonal variation in phenolic composition, antibacterial and antioxidant activities of *Ulva rigida* (Chlorophyta) and assessment of antiacetylcholinesterase potential. *J Appli Phycol* 25:319–328.
63. USDA, 2010. National nutrient database for standard reference, Release 23, September 2010. Composition of foods raw, processed, prepared.
64. Val A, Platas G, Basilio A, Cabello A, Gorrochategui J, Suay I, Vicente F, Portillo E, Jiménez del Río M, Reina G.G, Peláez F (2001) Screening of antimicrobial activities in red, green and brown macroalgae from *Gran Canaria* (*Canary Islands, Spain*). *Int Microbiol* 4(1):35-40.

Figures




Figure 1

a sampling points on the Oman Sea, Iran.



Figure 2

a green macroalgae (*Ulva*)




Figure 3

comparison of the IC_{50} content in the *Ulva* extract between different locations (left) and monsoon conditions (right) in the northern coasts of the Oman Sea ($P > 0.05$). Mean \pm standard deviation.




Figure 4

comparison of the FRAP content in the UI extract between different locations (left) and monsoon conditions (right) of the northern coasts of the Oman Sea ($P > 0.05$). Mean \pm standard deviation.




Figure 5

comparison of the TPC content in the UI extract between different locations (left) and monsoon conditions (right) of the northern coasts of the Oman Sea ($P > 0.05$). Mean \pm standard deviation.




Figure 6

comparison of the TFC content in the UI extract between different locations (left) and monsoon conditions (right) of the northern coasts of the Oman Sea ($P > 0.05$). Mean \pm standard deviation.




Figure 7

comparison of the ACN in the UI extract between different locations (left) and monsoon conditions (right) of the northern coasts of the Oman Sea ($P > 0.05$). Mean \pm standard deviation.




Figure 8

comparison of the beta C content in the UI extract between different locations (left) and monsoon conditions (right) the northern coasts of the Oman Sea ($P > 0.05$). Mean \pm standard deviation




Figure 9

projections of the variables (a) on to the factor plane defined by principal components (PC1 and PC2); Projection of the scores (b) onto the factor pane in different location.




Figure 10

projections of the variables (a) onto the factor plane defined by principal components (PC1 and PC2); b Projection of the scores (b) onto the factor plane in different location