Remotely monitoring ecosystem respiration from various grasslands along a large-scale east-west transect across northern China

Xuguang Tang (✉ xgtang@niglas.ac.cn)

Southwest University

Yanlian Zhou

Nanjing University

Hengpeng Li

Nanjing Institute of Geography and Limnology Chinese Academy of Sciences

Li Yao

Southwest University

Zhi Ding

Southwest University

Mingguo Ma

Southwest University

Pujia Yu

Southwest University

Research

Keywords: Ecosystem respiration, Grassland, MODIS, Piecewise regression

Posted Date: March 5th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-16165/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Carbon Balance and Management on April 24th, 2020. See the published version at <https://doi.org/10.1186/s13021-020-00141-8>.

Abstract

Background : Grassland ecosystems play an important role in the terrestrial carbon cycles through carbon emission by ecosystem respiration (R_e) and carbon uptake by plant photosynthesis (GPP). Surprisingly, given R_e occupies a large component of annual carbon balance, rather less attention has been paid to developing the estimates of R_e compared to GPP.

Results : Based on 11 flux sites over the diverse grassland ecosystems in northern China, this study examined the amounts of carbon released by R_e as well as the dominant environmental controls across temperate meadow steppe, typical steppe, desert steppe and alpine meadow, respectively. Multi-year mean R_e revealed relatively less CO_2 emitted by the desert steppe in comparison with other grassland ecosystems. Meanwhile, C emissions of all grasslands were mainly controlled by the growing period. Correlation analysis revealed that apart from air and soil temperature, soil water content exerted a strong effect on the variability in R_e , which implied the great potential to derive R_e using relevant remote sensing data. Then, these field-measured R_e data were up-scaled to large areas using time-series MODIS information and remote sensing-based piecewise regression models. These semi-empirical models appeared to work well with a small margin of error (R^2 and RMSE ranged from 0.45 to 0.88 and from 0.21 to 0.69 $\text{g C m}^{-2} \text{d}^{-1}$, respectively).

Conclusions : Generally, the piecewise models from the growth period and dormant season performed better than model developed directly from the entire year. Moreover, the biases between annual mean R_e observations and the remotely-derived products were usually within 20%. Finally, the regional R_e emissions across northern China's grasslands was approximately 100.66 Tg C in 2010, about 1/3 of carbon fixed from the MODIS GPP product. Specially, the desert steppe exhibited the highest ratio, followed by the temperate meadow steppe, typical steppe and alpine meadow. Therefore, this work provides a novel framework to accurately predict the spatio-temporal patterns of R_e over large areas, which can greatly reduce the uncertainties in global carbon estimates and climate projections.

1. Background

Although the terrestrial biosphere absorbs nearly a quarter of anthropogenic CO_2 emissions and plays a critical role in mitigating global climate warming, the land C sequestration potential of different ecosystems remains highly uncertain, leading to large uncertainties in future climate projections (Ballantyne *et al.*, 2015; Green *et al.*, 2019). During the past several decades, substantial advances have allowed for spatially continuous, long-term estimation of terrestrial gross primary productivity (GPP) based on the satellite remote sensing data, climate data and ecosystem models (Running *et al.*, 2004; Zhao *et al.*, 2005; Wu *et al.*, 2011; Zhang *et al.*, 2014). However, net ecosystem C budget depends upon the balance of C fixation through vegetation photosynthesis and C loss from ecosystem respiration (R_e) (Chapin *et al.*, 2006; Schimel *et al.*, 2011), which requires accurate estimates of not only GPP but also R_e across biomes.

Actually, terrestrial R_e from the biosphere to the atmosphere represents a large component of annual carbon budget, which even exceeds the amount of GPP (Milyukova *et al.*, 2002; Messerschmidt *et al.*, 2013). It has excited much interest in evaluating the balance between GPP and R_e on regional to global scales (Tian, 2003; Peylin *et al.*, 2013; Byrne *et al.*, 2018), because small fluctuations in either component caused by natural or human disturbances can ameliorate or exacerbate the buildup of CO_2 in the atmosphere (Wohlfahrt *et al.*, 2008; Tang *et al.*, 2017; Zheng *et al.*, 2018). Particularly, R_e is sensitive to the environmental factors and is highly spatio-temporally heterogeneous across scales (Zhang *et al.*, 2013; Ai *et al.*, 2018), which made it far poorly understood owing to the complicated interactions among physical, chemical, and biological variables in the respiration processes, including autotrophic respiration (R_a) from vegetation itself and heterotrophic respiration (R_h) from diverse soil microbiota (Hinko-najera *et al.*, 2015). Therefore, accurate quantification of the C emissions through R_e is crucial to understand its effect on climate change and global carbon dynamics.

With the development of global FLUXNET community based the eddy covariance (EC) technique across terrestrial ecosystems, it has been possible to continuously monitor the seasonal and interannual variations of carbon fluxes between the biosphere and the atmosphere, which can be further divided into GPP and R_e according to the nighttime based (Reichstein *et al.*, 2005) and daytime based flux-partitioning methods (Lasslop *et al.*, 2010). However, the in situ observations are generally implemented at field scale with low areal coverage ($< 1 \text{ km}^2$) and high cost of constructing and maintaining flux towers (Xiao *et al.*, 2008; Tang *et al.*, 2012; Gu *et al.*, 2018). The availability of spatially continuous data of ecosystem properties and environmental variables important for R_e provides an alternative approach for the large-area estimates. Currently, how to upscale the field-measured data using remote sensing (RS) information is urgently needed for understanding regional and global patterns of ecosystem R_e .

Several studies have recently been conducted to model the spatial distribution of soil respiration (R_s) at alpine grasslands (Huang *et al.*, 2013) and forests (Huang *et al.*, 2014; Wu *et al.*, 2014) using the satellite-based products including land surface temperature (LST) and spectral vegetation index (NDVI or EVI) or leaf area index (LAI). By incorporating the terrain information, Berryman *et al.* (2018) estimated R_s in a typical of the Southern Rocky Mountains with a coefficient of determination (R^2) of 0.45. Jägermeyr *et al.* (2014) firstly developed the models of global R_e according to forested and non-forested biomes. However, the classification system means that parameterization may not take into account the

wide variety of ecosystems on the earth (Gao *et al.*, 2015). Ai *et al.* (2018) also proposed an empirical yet physiologically based model with R^2 and RMSE of 0.55 and $1.67 \text{ g C m}^{-2} \text{ d}^{-1}$ respectively, which is capable of retrieving the patterns in R_e at the global scale. But the model cannot be transferred well to specific ecosystems, and may particularly be inaccurate in hydrologically sensitive areas owing to lack of water index affecting R_e . Nevertheless, previous studies have implied the significant correlations between grassland respiration and vegetation growth status (Huang *et al.*, 2013), as well as the environmental variables such as LST (Wohlfahrt *et al.*, 2008; Ai *et al.*, 2018) from the time-series satellite data.

Grasslands are the dominant landscape in China and account for 40% of the national land area. Geographically, approximately 78% of the grasslands in China exist in the northern temperate and alpine zones, constituting an integral part of the Eurasian grassland ecosystem (Kang *et al.*, 2007; Wei *et al.*, 2019). Due to their large carbon content, grasslands account for one-third of the global terrestrial carbon stock, second only to the forest ecosystem (Schuman *et al.*, 2002; Zhang *et al.*, 2016), and play a key role in China's terrestrial carbon cycle. Zhang *et al.* (2014) found that temperate grasslands in northern China have the potential to sequester carbon, but the capacity of carbon sequestration relies on grassland types and local environmental conditions. Further analysis revealed that the water availability is the dominant environmental factor regulating the annual carbon budget (Chen *et al.*, 2019). Extreme climate events such as drought can significantly reduce the net carbon uptake of grasslands. Moreover, it is predicted that heat waves and droughts will become more frequent in the 21st century (Hua *et al.*, 2017), which may lead to a general decrease in vegetation productivity in these grassland systems of northern China. Although several studies suggested that grasslands might be weak C sinks or near equilibrium (Gilmanov *et al.*, 2006; Zhang *et al.*, 2014), as the main constraint on C budget, a deep understanding of R_e is helpful to project climate change-terrestrial C feedback over different grassland ecosystems. Specially, the goals of this study were: 1) to analyze the differences of carbon released by R_e as well as the dominant environmental variables across the temperate meadow steppe, typical steppe, desert steppe and alpine meadow, respectively; 2) to develop rule-based piecewise regression models to map R_e of diverse grasslands by integrating time-series MODIS products and tower-based observations; and 3) to map the spatial patterns of annual mean R_e for the grassland ecosystems in northern China.

2. Methods And Materials

2.1 Description of the study area

The study was conducted in the grassland ecosystems of northern China, which is characterized by the arid and semi-arid continental monsoon climate with the highest temperature and rainfall period in summer. Following the east-to-west precipitation gradient, the temperate grasslands in northern China alter longitudinally from meadow steppes in the northeast, through typical steppes in the middle, and to desert steppes in the dry northwest. Alpine meadows are the dominant vegetation type in the Tibetan Plateau zone. These grasslands usually start to grow in early May and wither in late September with the peak biomass in July or August, which provide an important resource for livestock production and global carbon sequestration (Zhang *et al.*, 2014; Wei *et al.*, 2019).

During recent years, a series of EC-based flux towers have been installed by the Coordinated Observation and Synthesis in Arid and Semi-arid China (COSAS), as a part of China Flux Observation and Research Network (ChinaFLUX), which can be used to observe the carbon and water exchanges between the atmosphere and these grasslands in the ecologically fragile areas of northern China (Fig.1). In total of 11 flux sites are used in the study, which represent the most prevalent types of grassland ecosystems and a wide range of spatial, ecological, and climatic conditions, including two meadow steppe sites, two typical steppe sites, two desert steppe sites and five alpine meadow sites. Detailed descriptions of these flux sites can be found in the associated literature (Table 1).

2.2 Processing of the EC-based flux data

Both the EC system and the automatic meteorological station were mounted at these grassland sites, which acquired the continuous observations of site-level carbon fluxes (NEE), as well as the relevant climate data, including solar radiation (R_g), air and soil temperatures (T_a and T_s), relative humidity, soil water content (SWC), precipitation (P) and vapor pressure deficit (VPD). Each EC system was comprised of a three-dimensional sonic anemometer (CSAT3, Campbell Scientific, UT, USA) and a Li-7500 open path CO_2 and H_2O gas analyzer (LI-COR Inc., NE, USA). Raw data were continuously recorded at a frequency of 10 Hz on a CR5000 (Campbell Scientific) data logger. The processing procedures including spike detection and despiking, two-dimensional coordinate rotation, time delay removal of H_2O and CO_2 , virtual temperature correction, density effects (WPL correction) and frequency response corrections were completed using the improved EdiRe software package (developed by the University of Edinburgh) to produce a half-hour flux dataset (Mauder *et al.*, 2008). However, owing to instrument malfunctions, power failure, and severe weather conditions, approximately 25% of the one-year observations were lost. Therefore, it was necessary to interpolate these gaps with a standardized gap-filling algorithm. Then, the time-series NEE flux data were partitioned into GPP and R_e separately. In this study, the half-hourly R_e data provided by flux-tower measurements were integrated to the daily time scale, and then averaged over each 8-day period to match the 8-day composite of the MODIS products. The procedures including gap-filling and flux partitioning, were completed using the new R-based package (REddyProc) maintained by the Max Planck Institute for Biogeochemistry (Wutzler *et al.*, 2018).

Currently, there are mainly two methods implemented for flux-partitioning: 1) R_e is estimated from the nighttime temperature and extrapolated to daytime (Reichstein *et al.*, 2005) and 2) the light–response curve is fit to daytime NEE measurements and R_e is estimated from the intercept of the ordinate (Lasslaop *et al.*, 2010), which can avoid the use of potentially problematic nighttime data. The latter approach was chosen for flux partitioning because it uses a hyperbolic light-response curve algorithm, modified to account for the temperature dependency of respiration and the VPD limitation of photosynthesis (Gilmanov *et al.*, 2003). Including the VPD dependency strongly improved the model's ability to reproduce the asymmetric diurnal cycle during periods with high VPD, and enhances the reliability of R_e estimates given that the reduction of GPP by VPD may be otherwise incorrectly attributed to higher R_e . More details can be seen in the associated references.

2.3 MODIS products and processing

R_e is generally comprised of two sources of respiration: R_a from maintenance respiration and growth respiration, and R_h from rhizomicrobial respiration and microbial decomposition of plant residues and other soil organic matter (Hibbard *et al.*, 2005; Gao *et al.*, 2015). Thus, it is strongly affected by plant growth status and climate conditions. This study used the enhanced vegetation index (EVI) and leaf area index (LAI), as well as the land surface water index (LSWI), and mean value of daytime and nighttime temperatures (LST) to represent the vegetation and climate-related variables.

All variables were derived from the time-series MODIS data, which can avoid the complications and difficulties associated with merging disparate data sources. The 8-day land surface reflectance (MOD09A1, V6, with resolution of 500 m), LAI product (MOD15A2, V6, with resolution of 500 m) and LST data (MOD11A2, V6, with resolution of 1 km) were downloaded from the NASA's Earth Observing System Data and Information System (<https://search.earthdata.nasa.gov>). We only used the data described as good quality in the quality layer. These remote sensing-based products were re-sampled to a spatial resolution of 1 km, and the data where corresponded to the geographical location of each flux site were extracted for model development. In addition, the vegetation type map at 1 km resolution in 2010 was obtained from Nanjing Institute of Geography and Limnology, Chinese Academy of Science. The product specifically classified grasslands in China into temperate meadow steppe, typical steppe, desert steppe and alpine meadow, which can meet the needs to estimate R_e across different grassland types in this study. Then, all these data were used to develop models and map the spatio-temporal patterns of R_e in grassland ecosystems in northern China.

2.4 Statistical Analyses

To reveal the dominant environmental factors controlling the variability in R_e on an 8-day time scale over the whole year and different phenological periods (growing season vs dormant season) at these four different grassland ecosystems, the Pearson correlation coefficient (r) was calculated to examine the relationships between site-level R_e observations and these vegetation and climate-related variables. As these grasslands usually start to grow in early May and wither in late September, the growing period and dormant period were defined from WOY 16 to 34 and the rest of one year (WOY 1 to 15, and WOY 35 to 46), respectively. Then, we aimed to develop a rule-based piecewise regression model to capture the seasonal variations in R_e of temperate meadow steppe, typical steppe, desert steppe and alpine meadow in northern China. For each grassland type, the training set and test set of these flux data were instructed in Table 1. The study defined the model developed directly from the entire year as model 1, and the model from the growth period and dormant season as model 2. The model's accuracy was evaluated using two widely-used indicators: R^2 and the root-mean-square error (RMSE). The best model generally had the highest R^2 and lowest RMSE values. Finally, the optimal models were used to map the spatial patterns of annual mean R_e for the grasslands in northern China. All statistical analyses were performed using SPSS 19.0 (IBM, Chicago, IL, USA). In addition, the MODIS Reprojection Tool (MRT) and the Interactive Data Language (IDL) in ENVI 5.3 were used to process over 2000 scenes of MODIS data and large-area estimation.

3. Results

3.1 Differences of annual mean R_e across grasslands

Multi-year mean R_e of the temperate meadow steppe, typical steppe, desert steppe and alpine meadow ecosystems in northern China were exhibited in Fig.2 with apparent differences in magnitude. It revealed that C emissions by R_e were mainly concentrated in the growing season, which was even about twice and seven times of R_e during the dormant period for desert steppe and the other grasslands, respectively. The low R_e of desert steppe throughout the year can be ascribed to relatively sparse vegetation coverage. This study also found that during non-growing season, only small differences in R_e existed among these grassland types. Generally, typical steppe exhibited the strongest R_e of $1.08 \pm 0.23 \text{ g C m}^{-2} \text{ d}^{-1}$, followed by temperate meadow steppe ($1.03 \pm 0.10 \text{ g C m}^{-2} \text{ d}^{-1}$) and alpine meadow ($1.05 \pm 0.48 \text{ g C m}^{-2} \text{ d}^{-1}$), and undoubtedly desert steppe had the weakest R_e . These analyses again emphasized the importance to quantify the patterns of grassland R_e separately for large-area estimation.

3.2 Seasonal variations in R_e and environmental controls

Seasonal dynamics in R_e as well as the vegetation and climate-related variables across the four grassland ecosystems were illustrated in Fig.3 and Fig.4. It implied that except the desert steppe site (CN-Hzz), R_e of the other three grassland types exhibited an apparent single-peak pattern. As the temperature rose in spring, R_e gradually increased with plant growth, and reached the peak in July or August. However, the peak periods came earlier in typical steppe (CN-Nmg) at about week of year–WOY 23, followed by temperate meadow steppe (CN-Cng, WOY 25) and alpine meadow (CN-Ham, WOY 30). Natural rainfall generally occurred in summer with large fluctuations around the year. Contrastingly, relatively smooth SWC can reflect the true water availability. As a proxy of vegetation response to environmental variables, EVI and LAI exhibited consistent trends as R_e . Several periods of the remote-derived LSWI in winter were quite large due to the snow cover on and under the grasslands.

Table 2 revealed that almost all variables were strongly correlated to R_e without considering the phenology information. However, during the dominant growth period for respiration in temperate meadow steppe, typical steppe and alpine meadow, only temperature (T_a and T_s) and SWC strongly and positively affected the variability in R_e . In the desert steppe, R_e was found to be exerted a strong effect by SWC across growing and dormant seasons, which highlighted the water condition rather than temperature as the most important controlling factor in the extremely dry ecosystems. All the remotely-sensed vegetation indexes and climate-related LST and LSWI exhibited strong correlations, implying great potential to quantify the variability in R_e using RS technique.

3.3 Model development for quantitative remote sensing

By integrating site-level R_e observations and time-series MODIS products, the study developed the optimal models for capturing the variability in R_e of different grasslands using the stepwise multiple regression method (Table 3). Mainly two models were proposed with good performances. Generally, the model 2 established from the growth period and dormant season separately had relatively higher accuracy in estimating R_e than that directly from the whole year (model 1). Besides the desert steppe, R^2 and RMSE of the other grasslands varied from 0.77 to 0.88 and from 0.41 to 0.67 g C m⁻² d⁻¹, respectively. The model with only LAI as an explanatory variable also got good estimates for the desert steppe. Fig.5 further certified that the model promised well to estimate 8-day R_e and captured the broad trend of seasonal patterns, especially for the R_{e_model} 2, whereas the R_{e_model} 1 caused individual abnormal R_e estimates in wintertime. In spite of good accuracy, the RS-based models remained lacking of skills to capture abrupt changes in R_e during the summertime growing season.

Meanwhile, this study evaluated the model's performance on annual mean R_e across these four grasslands. Fig.6 revealed that the remotely-derived products slightly underestimated R_e at the temperate meadow steppe and desert steppe sites, but overestimated R_e at the typical steppe and alpine meadow sites. The R_{e_model1} had superior R_e estimates with mean bias of 9.6% at the typical steppe site, while R_{e_model2} only had an underestimation of 6.2% at the temperate meadow steppe site. Both models had similar accuracy at the desert steppe and alpine meadow sites.

3.4 Spatial patterns of satellite-derived R_e

The study mapped the spatial distribution of annual mean R_e for grasslands in northern China using the proposed regression models (Fig.7), which were extrapolated from the 8-day R_e estimates throughout the year of 2010. Generally, our results showed a declining trend from the southeast to the northwest direction. The highest R_e usually occurred in the temperate meadow steppe under good hydrothermal conditions. However, the desert steppe and the Tibetan alpine meadow emitted relatively fewer CO₂ through respiration. The total regional R_e estimate in the northern China's grasslands during 2010 was about 100.66 Tg C, which was approximately 1/3 of the carbon uptake through plant photosynthesis with 332.08 Tg C from the MODIS GPP product. Specially, the ratio of R_e to GPP exhibited the highest value in the desert steppe (88.6%), followed by the temperate meadow steppe (36.9%) and typical steppe (21.5%), with the lowest ratio in the alpine meadow of 18.0%.

4. Discussion

Along the east–west precipitation gradient in northern China, a wide variety of grasslands constitute the dominant landscape in servicing the ecological environment and socio-economics of the region and in supporting diverse species of plants and animals (Kang *et al.*, 2007). The meadow steppes usually occur in the most moist and fertile sites among the four grassland ecosystem types, typically in areas with annual precipitation of about 450 mm. The typical steppes are developed under a semi-arid climate in the temperate zone with annual precipitation of around 350 mm and the plant species are characteristically drought-tolerant. The desert steppes are the most arid grassland type, occurring in areas with annual precipitation between 150 and 250 mm under the influence of continental climate. Alpine steppes are found between 2300 and 5300 m a.s.l. in southwestern China. The common construction plant species are cold- and drought-tolerant grasses and small shrubs. Arid and semiarid ecosystems are often overlooked in this regard because of their spatially sparse vegetation and fragile environmental conditions (Lioubimtseva and Henebry, 2009; Ahlström *et al.*, 2015). In fact, the grassland ecosystems are not only economically important, but recent work has shown water-limited systems to drive the interannual variability in the global C cycle significantly (Soussana *et al.*, 2004; Poulter *et al.*, 2014). Therefore, a deep understanding of the C sequestration potential as well as environmental controls over the northern China's grasslands is vital for reducing the uncertainty in future climate projections given the large area of grasslands.

Accurate quantification of the main C fluxes across terrestrial ecosystems is crucial to our understanding of global carbon balance (Tang *et al.*, 2012; Baldocchi, 2014). Till now, there are fewer successful models of R_e compared to GPP particularly using the RS data directly (Olofsson *et al.*, 2008; Jägermeyr *et al.*, 2014). Reichstein *et al.* (2003) found that soil water and temperature were good predictors for R_s , and adding LAI as a proxy for productivity further improved the accuracy. This study was based on closed-chamber data from forest and shrubland sites across Europe and North America, but it was suggested that the variables could easily be acquired from RS data. Anderson *et al.* (2008) used a model which calculated soil moisture from microwave sensing, soil temperature from thermal imaging, and LAI, showed good agreement with tower flux data over pasture land in Oklahoma. However, model development over such a small area is unlikely to generate a model which is applicable to other ecosystems or climates, and more validation work is also needed. In spite of this, subsequent studies have suggested that the strong relationships between R_s and R_e and GPP (Huang *et al.*, 2014), LST (Olofsson *et al.*, 2008; Huang *et al.*, 2013; Ai *et al.*, 2018) from the time-series RS data.

Changes in temperature and water availability, and thus climate change, are likely to be significant drivers of the future C balance of land systems and their feedbacks to climate change. Grasslands in the arid and semi-arid regions are ecologically fragile and sensitive to climate change and human disturbances, especially to the changes in precipitation (Hu *et al.*, 2010). The precipitation in arid and semi-arid regions is highly variable both temporally and spatially. Fluctuations in carbon budget have been found to be closely associated with interannual and intra-annual variations in precipitation in arid and semi-arid ecosystems, and persistent drought has caused a general decrease in vegetation productivity in the grassland systems of northern China (Zhang *et al.*, 2014). These work also demonstrated that water condition is an important environmental indicator for the estimates of R_e in the grassland ecosystems. Soil moisture affects R_e processes in various ways, including the growth and development of both aboveground vegetation and roots, growth and activity of microbial populations, in addition to gas transport throughout soils (Moyano *et al.*, 2013; Yan *et al.*, 2018). Green *et al.* (2019) also emphasize that the capacity of continents to act as a future carbon sink critically depends on the response of C fluxes to soil moisture. The study used the LSWI derived from the shortwave infrared (SWIR) and the near infrared (NIR) bands of MODIS data to represent land surface water condition, and got reasonably good performance. Actually, LSWI is different from soil moisture (SWC) because most information of LSWI reflects the total amount of liquid water in vegetation (Chandrasekar *et al.*, 2010). An interesting avenue of future work would be to consider employing the soil moisture data from the Advanced Microwave Scanning Radiometer-Earth Observing System (AMSE-E) (Al-Yaari *et al.*, 2014). The problem with many of these studies, however, is that they are too narrow for comparison. They consider one particular site in one particular ecosystem, or attempt to create a global model often focus on a narrow range of only four or five ecosystem types (Jägermeyr *et al.*, 2014; Ai *et al.*, 2018). Grasslands and their huge variety of types are almost never included as a separate category in remote sensing models of carbon flux, and as such are certain to be over or under-estimated.

Our results suggested that simple models relying entirely on spatial data have the potential to estimate R_e over the diverse grassland ecosystems. This result provides a framework for the development of R_e models aimed to obtain spatial pattern in R_e . Meanwhile, the models incorporating the phenology information generally provided better estimations. In this study, we simply separated the growing season and dormant season as these grasslands usually start to grow in early May and wither in late September. To acquire more specific phenological information of each pixel, the global land surface phenology metrics at yearly intervals (MCD12Q2) can be used for auxiliary analysis. But the models would be complicated for large-scale extrapolation owing to strong spatial heterogeneity. In addition, there is a need for more long-term studies in order to monitor the interannual variations of R_e as well as the underlying mechanisms. The study calculated the spatial distribution of grassland R_e in 2010 because we only have a period of classification data with detailed grassland types. The total of carbon emissions of R_e across the northern China's grasslands was accumulated to 100.66 Tg C during 2010, which was approximately 1/3 of the regional carbon uptake through plant photosynthesis with 332.08 Tg C from the MODIS GPP product. Particularly, the carbon use efficiency was the highest in the alpine meadow, followed by the typical steppe, meadow steppe and desert steppe. However, this approach generally need large amounts of R_e data to constrain parameters. The regression coefficients obtained at this study would not work well for other grassland sites with different climate, soil, and vegetation. Although we tried to establish the robust models for the temperate meadow steppe, typical steppe, desert steppe and alpine meadow using 11 flux sites, the representativeness of these limited sites presumably affected the general applicability of our predictive model. Thus, the current Coordinated Observation and Synthesis in Arid and Semi-arid China should be augmented by building more sites across a full range of grassland types.

5. Conclusion

On the basis of 24 EC-based years of flux measurements over 11 grasslands sites under a wide range of geographic, weather and ecological conditions, the study was an attempt to upscale site-level R_e data to the northern China's grasslands including temperate meadow steppe, meadow steppe, desert steppe and alpine meadow, using the satellite-based RS data. The results demonstrated that the rule-based piecewise regression models can successfully estimate the seasonal variations in R_e and provide a new framework to map the regional patterns of large-scale R_e . The reduction of uncertainties in R_e is crucial for projecting climate change impacts on terrestrial carbon cycling and future atmospheric CO₂ concentrations. This work also offers an opportunity to further understand the environmental drivers controlling the variability in R_e . Specially, not only temperate but also soil water content, had a strong correlation with grassland R_e which should not be neglected when developing the RS

models for arid and semiarid ecosystems. Meanwhile, the models incorporating the phenology information generally performed better. The spatial patterns of R_e across the northern China's grasslands exhibited a distinctly declining trend from the southeast to the northwest, with a regional estimate of approximately 100.66 Tg C during 2010. With more periods of grassland classification information, future studies can even evaluate long-term land-use change and its impact on the large-scale R_e of grasslands.

Abbreviations

R_e : ecosystem respiration

GPP: gross primary production

MODIS: Moderate-resolution Imaging Spectroradiometer

R_a : autotrophic respiration

R_h : heterotrophic respiration

EC: eddy covariance

RS: remote sensing

R_s : soil respiration

LST: land surface temperature

NDVI: normalized difference vegetation index

EVI: enhanced vegetation index

LAI: leaf area index

R^2 : coefficient of determination

RMSE: root-mean-square error

r : correlation coefficient

ChinaFLUX: China Flux Observation and Research Network

NEE: net ecosystem carbon flux

R_g : solar radiation

T_a : air temperature

T_s : soil temperature

SWC: soil water content

P: precipitation

VPD: vapor pressure deficit

LSWI: land surface water index

Declarations

Availability of data and materials

The data used in this article are available upon request.

Competing interests

The authors declare no conflict of interests.

Funding

The study was jointly supported by the National Natural Science Foundation of China (Grant No. 41830648 and 41401221), the National Science and Technology Major Project (Grant No. 21-Y20A06-9001-17/18) and Chongqing Basic and Frontier Research Program (Grant No. cstc2018jcyjAX0056).

Authors' contributions

PY conceived the research; XT and ZD contributed to data analysis and wrote the manuscript; YZ and HL provided data support; LY and MM improved the manuscript.

Acknowledgments

All scientists and technicians maintaining the flux site management, data collection and long-term observations are greatly appreciated. The eddy covariance data and meteorological data are acquired from the Coordinated Observation and Synthesis in Arid and Semi-arid China (COSAS), a part of ChinaFLUX. We are also grateful to the Distributed Active Archive Center of the Oak Ridge National Laboratory, and the Earth Observing System Data for making these MODIS products available.

References

- Ai, J., Jia, G., Epstein, H.E., Wang, H., Zhang, A., Hu, Y., 2018. MODIS-based estimates of global terrestrial ecosystem respiration. *J. Geophys. Res. Biogeosciences*. 123, 326–352.
- Anderson, M.C., Norman, J.M., Kustas, W.P., Houborg, R., Starks, P.J., Agam, N., 2008. A thermal-based remote sensing technique for routine mapping of land-surface carbon, water and energy fluxes from field to regional scales. *Remote Sens. Environ.* 112, 4227–4241.
- Al-Yaari, A., Wigneron, J.P., Ducharne, A., Kerr, Y., de Rosnay, P., de Jeu, R., Govind, A., Al Bitar, A., Albergel, C., Muñoz-Sabater, J., Richaume, P., Mialon, A., 2014. Global-scale evaluation of two satellite-based passive microwave soil moisture datasets (SMOS and AMSR-E) with respect to Land Data Assimilation System estimates. *Remote Sens. Environ.* 149, 181–195.
- Baldocchi, D., 2014. Measuring fluxes of trace gases and energy between ecosystems and the atmosphere - the state and future of the eddy covariance method. *Glob. Chang. Biol.* 20, 3600–3609.
- Ballantyne, A.P., Andres, R., Houghton, R., Stocker, B.D., Wanninkhof, R., Anderegg, W., Cooper, L.A., DeGrandpre, M., Tans, P.P., Miller, J.B., Alden, C., White, J.W.C., 2015. Audit of the global carbon budget: Estimate errors and their impact on uptake uncertainty. *Biogeosciences*. 12, 2565–2584.
- Byrne, B., Wunch, D., Jones, D.B.A., Strong, K., Deng, F., Baker, I., Köhler, P., Frankenberg, C., Joiner, J., Arora, V.K., Badawy, B., Harper, A.B., Warneke, T., Petri, C., Kivi, R., Roehl, C.M., 2018. Evaluating GPP and respiration estimates over northern midlatitude ecosystems using solar-induced fluorescence and atmospheric CO₂ measurements. *J. Geophys. Res. Biogeosciences*. 123, 2976–2997.
- Chandrasekar, K., Sesha Sai, M.V.R., Roy, P.S., Dwevedi, R.S., 2010. Land Surface Water Index (LSWI) response to rainfall and NDVI using the MODIS vegetation index product. *Int. J. Remote Sens.* 31, 3987–4005.
- Chapin, F.S., Woodwell, G.M., Randerson, J.T., Rastetter, E.B., Lovett, G.M., Baldocchi, D.D., Clark, D.A., Harmon, M.E., Schimel, D.S., Valentini, R., Wirth, C., Aber, J.D., Cole, J.J., Goulden, M.L., Harden, J.W., Heimann, M., Howarth, R.W., Matson, P.A., McGuire, A.D., Melillo, J.M., Mooney, H.A., Neff, J.C., Houghton, R.A., Pace, M.L., Ryan, M.G., Running, S.W., Sala, O.E., Schlesinger, W.H., Schulze, E.D., 2006. Reconciling carbon-cycle concepts, terminology, and methods. *Ecosystems*. 9(7), 1041–1050.
- Chen, S., Chen, J., Lin, G., Zhang, W., Miao, H., Wei, L., Huang, J., Han, X., 2009. Energy balance and partition in Inner Mongolia steppe ecosystems with different land use types. *Agric. For. Meteorol.* 149, 1800–1809.
- Chen, Y., Gu, H., Wang, M., Gu, Q., Ding, Z., Ma, M., Liu, R., Tang, X. 2019. Contrasting performance of the remotely-derived GPP products over different climate zones across China. *Remote Sens.* 11(16), 1855.
- Dong, G., Guo, J., Chen, J., Sun, G., Gao, S., Hu, L., Wang, Y., 2011. Effects of spring drought on carbon sequestration, evapotranspiration and water use efficiency in the songnen meadow steppe in northeast China. *Ecohydrology*. 4 (2), 211–224.
- Gilmanov, T.G., Johnson, D.A., Saliendra, N.Z., 2003. Growing season CO₂ fluxes in a sagebrush-steppe ecosystem in Idaho: Bowen ratio/energy balance measurements and modeling. *Basic Appl. Ecol.* 4, 167–183.

- Gilmanov, T.G., Svejcar, T.J., Johnson, D.A., Angell, R.F., Saliendra, N.Z., Wylie, B.K., 2006. Long-term dynamics of production, respiration, and net CO₂ exchange in two sagebrush-steppe ecosystems. *Rangel. Ecol. Manag.* 59, 585–599.
- Green, J.K., Seneviratne, S.I., Berg, A.M., Findell, K.L., Hagemann, S., Lawrence, D.M., Gentine, P., 2019. Large influence of soil moisture on long-term terrestrial carbon uptake. *Nature*. 565, 476–479.
- Gu, Q., Wei, J., Luo, S., Ma, M., Tang, X., 2018. Potential and environmental control of carbon sequestration in major ecosystems across arid and semi-arid regions in China. *Sci. Total Environ.* 645, 796–805.
- Hibbard, K.A., Law, B.E., Reichstein, M., Sulzman, J., 2005. An analysis of soil respiration across northern hemisphere temperate ecosystems. *Biogeochemistry*. 73(1), 29–70.
- Hinko-najera, N., Fest, B., Livesley, S.J., Arndt, S.K., 2015. Agricultural and Forest Meteorology Reduced throughfall decreases autotrophic respiration, but not heterotrophic respiration in a dry temperate broadleaved evergreen forest. *Agric. For. Meteorol.* 200, 66–77.
- Hu, Z., Yu, G., Fan, J., Zhong, H., Wang, S., Li, S., 2010. Precipitation-use efficiency along a 4500-km grassland transect. *Global Ecol. Biogeogr.* 19(6), 842–851.
- Hu, Z., Yu, G., Fu, Y., Sun, X., Li, Y., Shi, P., Wang, Y., Zheng, Z., 2008. Effects of vegetation control on ecosystem water use efficiency within and among four grassland ecosystems in China. *Glob. Chang. Biol.* 14(7), 1609–1619.
- Hua, T., Wang, X., Zhang, C., Lang, L., Li, H., 2017. Responses of vegetation activity to drought in northern China. *Land Degrad. Dev.* 28(7), 1913–1921.
- Huang, N., Gu, L., Niu, Z., 2014. Estimating soil respiration using spatial data products: A case study in a deciduous broadleaf forest in the Midwest USA. *J. Geophys. Res.* 119, 6393–6408.
- Huang, N., He, J.S., Niu, Z., 2013. Estimating the spatial pattern of soil respiration in Tibetan alpine grasslands using Landsat TM images and MODIS data. *Ecol. Indic.* 26, 117–125.
- Jägermeyr, J., Gerten, D., Lucht, W., Hostert, P., Migliavacca, M., Nemani, R., 2014. A high-resolution approach to estimating ecosystem respiration at continental scales using operational satellite data. *Glob. Chang. Biol.* 20(4), 1191–1210.
- Kang, L., Han, X., Zhang, Z., Sun, O.J., 2007. Grassland ecosystems in China: Review of current knowledge and research advancement. *Philos. Trans. R. Soc. B Biol. Sci.* 362, 997–1008.
- Kato, T., Tang, Y., Gu, S., Hirota, M., Du, M., Li, Y., Zhao, X., 2006. Temperature and biomass influences on interannual changes in CO₂ exchange in an alpine meadow on the Qinghai-Tibetan Plateau. *Glob. Chang. Biol.* 12(7), 1285–1298.
- Lasslop, G., Reichstein, M., Papale, D., Richardson, A., Arneth, A., Barr, A., Stoy, P., Wohlfahrt, G., 2010. Separation of net ecosystem exchange into assimilation and respiration using a light response curve approach: Critical issues and global evaluation. *Glob. Chang. Biol.* 16, 187–208.
- Lian, J., Huang, M., 2016. Comparison of three remote sensing based models to estimate evapotranspiration in an oasis-desert region. *Agric. Water Manag.* 165, 153–162.
- Liu, S.M., Xu, Z.W., Wang, W.Z., Jia, Z.Z., Zhu, M.J., Bai, J., Wang, J.M., 2011. A comparison of eddy-covariance and large aperture scintillometer measurements with respect to the energy balance closure problem. *Hydrol. Earth Syst. Sci.* 15, 1291–1306.
- Mauder, M., Foken, T., Clement, R., Elbers, J.A., Eugster, W., Grünwald, T., Heusinkveld, B., Kolle, O., 2008. Quality control of CarboEurope flux data - Part 2: Inter-comparison of eddy-covariance software. *Biogeosciences*. 5, 451–462.
- Messerschmidt, J., Parazoo, N., Wunch, D., Deutscher, N.M., Roehl, C., Warneke, T., Wennberg, P.O., 2013. Evaluation of seasonal atmosphere-biosphere exchange estimations with TCCON measurements. *Atmos. Chem. Phys.* 13(10), 5103–5115.
- Milyukova, I.M., Kolle, O., Varlagin, A. V., Vygodskaya, N.N., Schulze, E.D., Lloyd, J., 2002. Carbon balance of a southern taiga spruce stand in European Russia. *Tellus, Ser. B Chem. Phys. Meteorol.* 54(5), 429–442.
- Moyano, F.E., Manzoni, S., Chenu, C., 2013. Responses of soil heterotrophic respiration to moisture availability: An exploration of processes and models. *Soil Biol. Biochem.* 59, 72–85.
- Olofsson, P., Lagergren, F., Lindroth, A., Lindström, J., Klemedtsson, L., Kutsch, W., Eklundh, L., 2008. Towards operational remote sensing of forest carbon balance across Northern Europe. *Biogeosciences*. 5, 817–832.

- Peylin, P., Law, R.M., Gurney, K.R., Chevallier, F., Jacobson, A.R., Maki, T., Niwa, Y., Patra, P.K., Peters, W., Rayner, P.J., Rödenbeck, C., Van Der Laan-Luijkx, I.T., Zhang, X., 2013. Global atmospheric carbon budget: Results from an ensemble of atmospheric CO₂ inversions. *Biogeosciences*. 10, 6699–6720.
- Poulter, B., Frank, D., Ciais, P., Myneni, R.B., Andela, N., Bi, J., Broquet, G., Canadell, J.G., Chevallier, F., Liu, Y., Running, S.W., Sitch, S., Van der Werf, G.R., 2014. Contribution of semi-arid ecosystems to interannual variability of the global carbon cycle. *Nature*. 509, 600–603.
- Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N., Gilmanov, T., Granier, A., Grünwald, T., Havránková, K., Ilvesniemi, H., Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F., Ourcival, J.M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakir, D., Valentini, R., 2005. On the separation of net ecosystem exchange into assimilation and ecosystem respiration: Review and improved algorithm. *Glob. Chang. Biol.* 11, 1424–1439.
- Reichstein, M., Rey, A., Freibauer, A., Tenhunen, J., Valentini, R., Banza, J., Casals, P., Cheng, Y., Grünzweig, J.M., Irvine, J., Joffre, R., Law, B.E., Loustau, D., Miglietta, F., Oechel, W., Ourcival, J.-M., Pereira, J.S., Peressotti, A., Ponti, F., Qi, Y., Rambal, S., Rayment, M., Romanya, J., Rossi, F., Tedeschi, V., Tirone, G., Xu, M., Yakir, D., 2003. Modeling temporal and large-scale spatial variability of soil respiration from soil water availability, temperature and vegetation productivity indices. *Global Biogeochem. Cycles*. 17, 4.
- Running, S.W., Nemani, R.R., Heinsch, F.A., Zhao, M., Reeves, M., Hashimoto, H., 2004. A continuous satellite-derived measure of global terrestrial primary production. *Bioscience*. 54, 547–560.
- Schimel, D.S., House, J.I., Hibbard, K.A., Bousquet, P., Ciais, P., Peylin, P., Braswell, B.H., Apps, M.J., Baker, D., Bondeau, A., Canadell, J., Churkina, G., Cramer, W., Denning, A.S., Field, C.B., Friedlingstein, P., Goodale, C., Heimann, M., Houghton, R.A., Melillo, J.M., Moore, B., Murdiyarso, D., Noble, I., Pacala, S.W., Prentice, I.C., Raupach, M.R., Rayner, P.J., Scholes, R.J., Steffen, W.L., Wirth, C., 2001. Recent patterns and mechanisms of carbon exchange by terrestrial ecosystems. *Nature*. 414(6860): 169.
- Shang, L., Zhang, Y., Lü, S., Wang, S., 2015. Energy exchange of an alpine grassland on the eastern Qinghai-Tibetan Plateau. *Sci. Bull.* 60(4), 435–446.
- Soussana, J.-F., Loiseau, P., Vuichard, N., Ceschia, E., Balesdent, J., Chevallier, T., Arrouays, D., 2004. Carbon cycling and sequestration opportunities in temperate grasslands. *Soil Use Manag.* 20, 219–230.
- Tang, X., Li, H., Ma, M., Yao, L., Peichl, M., Arain, A., Xu, X., Goulden, M., 2017. How do disturbances and climate effects on carbon and water fluxes differ between multi-aged and even-aged coniferous forests? *Sci. Total Environ.* 599, 1583–1597.
- Tang, X., Wang, Z., Liu, D., Song, K., Jia, M., Dong, Z., Munger, J.W., Hollinger, D.Y., Bolstad, P. V., Goldstein, A.H., Desai, A.R., Dragoni, D., Liu, X., 2012. Estimating the net ecosystem exchange for the major forests in the northern United States by integrating MODIS and AmeriFlux data. *Agric. For. Meteorol.* 156, 75–84.
- Tian, H., 2003. Regional carbon dynamics in monsoon Asia and its implications for the global carbon cycle. *Glob. Planet. Change*. 37, 201–217.
- Schuman, G., Janzen, H., Herrick, J., 2002. Soil carbon dynamics and potential carbon sequestration by rangelands. *Environ. Pollut.* 2002, 116(3), 391–396.
- Shao, P., Zeng, X., Sakaguchi, K., Monson, R.K., Zeng, X., 2013. Terrestrial carbon cycle: Climate relations in eight CMIP5 earth system models. *J. Clim.* 26 (22), 8744–8764.
- Wang, Z., Xiao, X., Yan, X., 2010. Modeling gross primary production of maize cropland and degraded grassland in northeastern China. *Agric. For. Meteorol.* 150(9), 1160–1167.
- Wei, J., Chen, Y., Gu, Q., Jiang, C., Ma, M., Song, L., Tang, X., 2019. Potential of the remotely-derived products in monitoring ecosystem water use efficiency across grasslands in Northern China. *Int. J. Remote Sens.* 40, 6203–6223.
- Wohlfahrt, G., Anderson-Dunn, M., Bahn, M., Balzarolo, M., Berninger, F., Campbell, C., Carrara, A., Cescatti, A., Christensen, T., Dore, S., Eugster, W., Friborg, T., Furger, M., Gianelle, D., Gimeno, C., Hargreaves, K., Hari, P., Haslwanter, A., Johansson, T., Marcolla, B., Milford, C., Nagy, Z., Nemitz, E., Rogiers, N., Sanz, M.J., Siegwolf, R.T.W., Susiluoto, S., Sutton, M., Tuba, Z., Ugoni, F., Valentini, R., Zorer, R., Cernusca, A., 2008. Biotic, abiotic, and management controls on the net ecosystem CO₂ exchange of European mountain grassland ecosystems. *Ecosystems*. 11, 1338–1351.
- Wu, C., Chen, J. M., Huang, N., 2011. Predicting gross primary production from the enhanced vegetation index and photosynthetically active radiation: Evaluation and calibration. *Remote Sens. Environ.* 115, 3424–3435.

Wu, C., Gaumont-Guay, D., Andrew Black, T., Jassal, R.S., Xu, S., Chen, J.M., Gonsamo, A., 2014. Soil respiration mapped by exclusively use of MODIS data for forest landscapes of Saskatchewan, Canada. *ISPRS J. Photogramm. Remote Sens.* 94, 80–90.

Wutzler, T., Lucas-Moffat, A., Migliavacca, M., Knauer, J., Sickel, K., Šigut, L., Menzer, O., Reichstein, M., 2018. Basic and extensible post-processing of eddy covariance flux data with REddyProc. *Biogeosciences*. 15, 5015–5030.

Xiao, J., Zhuang, Q., Baldocchi, D.D., Law, B.E., Richardson, A.D., Chen, J., Oren, R., Starr, G., Noormets, A., Ma, S., Verma, S.B., Wharton, S., Wofsy, S.C., Bolstad, P. V., Burns, S.P., Cook, D.R., Curtis, P.S., Drake, B.G., Falk, M., Fischer, M.L., Foster, D.R., Gu, L., Hadley, J.L., Hollinger, D.Y., Katul, G.G., Litvak, M., Martin, T.A., Matamala, R., McNulty, S., Meyers, T.P., Monson, R.K., Munger, J.W., Oechel, W.C., Paw U, K.T., Schmid, H.P., Scott, R.L., Sun, G., Suyker, A.E., Torn, M.S., 2008. Estimation of net ecosystem carbon exchange for the conterminous United States by combining MODIS and AmeriFlux data. *Agric. For. Meteorol.* 148, 1827–1847.

Yan, Z., Bond-Lamberty, B., Todd-Brown, K.E., Bailey, V.L., Li, S., Liu, C., Liu, C., 2018. A moisture function of soil heterotrophic respiration that incorporates microscale processes. *Nat. Commun.* 9, 2562.

Zhang, L., Guo, H., Jia, G., Wylie, B., Gilmanov, T., Howard, D., Ji, L., Xiao, J., Li, J., Yuan, W., Zhao, T., Chen, S., Zhou, G., Kato, T., 2014. Net ecosystem productivity of temperate grasslands in northern China: An upscaling study. *Agric. For. Meteorol.* 184, 71–81.

Zhang, L., Zhou, G., Ji, Y., Bai, Y. 2016. Spatiotemporal dynamic simulation of grassland carbon storage in China. *Sci. China Earth Sci.* 59(10), 1946–1958.

Zhang, Q., Lei, H. M., Yang, D. W., 2013. Seasonal variations in soil respiration, heterotrophic respiration and autotrophic respiration of a wheat and maize rotation cropland in the North China Plain. *Agric. For. Meteorol.* 180, 34–43.

Zhao, M., Heinsch, F.A., Nemani, R.R., Running, S.W., 2005. Improvements of the MODIS terrestrial gross and net primary production global data set. *Remote Sens. Environ.* 95, 164–176.

Zheng, C., Tang, X., Gu, Q., Wang, T., Wei, J., Song, L., Ma, M., 2018. Climatic anomaly and its impact on vegetation phenology, carbon sequestration and water-use efficiency at a humid temperate forest. *J. Hydrol.* 565, 150–159.

Tables

Table 1 Characteristics of the flux sites over grassland ecosystems in northern China

Table 2 Pearson correlation analysis between 8-day R_g and the controlling environmental factors across the four grassland ecosystem types.

Grassland type	Period	R_g (W/m ²)	T_a (°C)	T_s (°C)	VPD (h Pa)	P (mm)	SWC (%)	EVI	LAI	LSWI	LST (°C)
Temperate meadow steppe	Growing	-0.115	0.538**	0.613**	-0.287*	0.252*	0.351**	0.745**	0.692**	0.704**	0.371**
	Dormant	0.411**	0.707**	0.714**	0.685**	0.231*	0.625**	0.378**	0.635**	-0.269*	0.689**
	Entire year	0.570**	0.749**	0.786**	0.520**	0.459**	0.669**	0.869**	0.851**	0.093	0.721**
Typical steppe	Growing	0.073	0.760**	0.707**	-0.149	0.568**	0.495**	0.721**	0.644**	0.748**	0.669**
	Dormant	0.506**	0.554**	0.553**	0.579**	0.269*	0.571**	0.377**	0.451**	-0.235	0.532**
	Entire year	0.644**	0.776**	0.803**	0.581**	0.700**	0.644**	0.861**	0.794**	-0.011	0.759**
Desert steppe	Growing	0.138	0.171	0.261	0.193	0.197	0.538**	0.499**	0.621**	0.338*	0.110
	Dormant	0.086	0.175	0.225	0.056	-0.119	0.611**	0.296*	0.052	-0.262	0.175
	Entire year	0.370**	0.408**	0.449**	0.421**	0.101	0.475**	0.542**	0.653**	-0.198	0.392**
Alpine meadow	Growing	-0.224*	0.535**	0.531**	0.039	0.142	0.397**	0.764**	0.382**	0.739**	0.281*
	Dormant	0.275*	0.652**	0.628**	0.379**	0.445**	0.630**	0.577**	0.566**	-0.206*	0.561**
	Entire year	0.265*	0.629**	0.647**	0.325**	0.484**	0.565**	0.837**	0.548**	0.326**	0.531**

Note: ** and * mean that correlations are significant at the 0.01 level and 0.05 level, respectively. R_g , T_a , T_s , VPD, P and SWC are observed at the flux tower sites listed in Table 1. Time-series EVI, LAI, LSWI and LST data are derived from the corresponding MODIS products. All these data are at an 8-day interval.

Code	Site name	Location	Temperature (°C)	Precipitation (mm)	Elevation (m)	Vegetation type	Height (m)	Soil type	Years	Reference
CN-Cng	Changling	44.5934° N 123.5092° E	5	400	171	Temperate meadow steppe	0.75	Alkali-saline soil	2007~2010	Dong <i>et al.</i> 2011
CN-Nmg	Xilinhot [#]	43.5544° N 116.6714° E	2	350	1250	Typical steppe	0.4	Chestnut soil	2004	Chen <i>et al.</i> , 2009
CN-Du2	Duolun fenced grassland	42.0467° N 116.2836° E	1.7	350	1350	Typical steppe	0.45	Chestnut soil	2006~2008	Chen <i>et al.</i> , 2009
CN-Hzz	Huazhaizi	38.76519° N 100.3186° E	7.5	157	1726	Desert steppe	0.15	Loam	2014~2017	Lian and Huang, 2016
CN-Sw2	Siziwang	41.7781° N 111.8971° E	3.5	283	1438	Desert steppe	0.15	Loam	2011	Shao <i>et al.</i> , 2013
CN-Ham	Haibei	37.6127° N 101.3128° E	-1.7	561	3216	Alpine meadow	0.3	Clay loam	2002~2003	Kato <i>et al.</i> , 2006
CN-Ar	A'rou	38.0473° N 100.4643° E	0.9	403.1	3033	Alpine meadow	0.45	Chestnut soil	2013~2015	Liu <i>et al.</i> , 2011
CN-Ds	Dashalong	38.8399° N 98.9606° E	-3.4	319	3739	Alpine meadow	0.15	Chestnut soil	2014~2015	This study
CN-Dan	Dangxiong	30.4978° N 91.0664° E	1.3	480	4333	Alpine meadow	0.1	Sandy loam	2004~2005	Hu <i>et al.</i> , 2008
CN-Ty	Tongyu degraded grassland [#]	44.5673° N 122.9170° E	5.5	345.4	151	Temperate meadow steppe	0.1	Sand soil	2008*~2009*	Wang <i>et al.</i> , 2010
CN-Mq	Maqu	33.8872° N 102.1407° E	1.9	593	3443	Alpine meadow	0.25	Silt clay loam	2009*	Shang <i>et al.</i> , 2015

Note: [#] represents vegetation type in degraded condition. * means that only part time of the year's flux and climate data are measured at this site, generally during the growing season. Among the eleven eddy covariance flux sites, CN-Cng (2010), CN-Ar (2014), CN-Du2 (2008) and CN-Sw2 (2011) are used for validation as a proxy for temperate meadow steppe, Alpine meadow, typical steppe and desert steppe, respectively.

Table 3 Regression analysis of 8-day R_g and the associated MODIS-derived products across the four grassland ecosystems using the stepwise multiple regression method.

Grassland type	Period	Regression models					Training set		Test set	
		Constant	EVI	LAI	LSWI	LST	R^2	RMSE	R^2	RMSE
Temperate meadow steppe	Growing	1.088	4.078	□	3.743	□	0.57	0.72	0.77	0.45
	Dormant	0.064	2.362	□	0.268	0.014	0.61	0.13		
	Entire year	-0.619	8.487	□	□	0.018	0.80	0.51	0.71	0.63
Typical steppe	Growing	0.561	□	□	6.058	0.096	0.72	0.60	0.88	0.41
	Dormant	0.044	3.463	□	0.659	0.021	0.41	0.26		
	Entire year	-0.083	5.633	□	1.419	0.040	0.84	0.49	0.81	0.50
Desert steppe	Growing	0.213	□	0.758	□	□	0.39	0.32	0.50	0.21
	Dormant	0.122	2.035	□	□	-0.003	0.11	0.17		
	Entire year	0.190	□	0.775	□	□	0.43	0.25	0.45	0.27
Alpine meadow	Growing	-0.321	7.198	-0.208	2.854	□	0.73	0.98	0.85	0.67
	Dormant	-0.264	3.826	0.897	0.804	0.019	0.65	0.20		
	Entire year	-0.520	6.947	□	1.318	0.020	0.81	0.67	0.83	0.69

Note: Time-series EVI, LAI, LSWI and LST data are derived from the corresponding MODIS products. All data are at an 8-day interval. The training set and test set of these flux sites are instructed in Table 1. When using the stepwise multiple regression method to develop models, only the best is exhibited here. The model's performance was evaluated using the site-level flux measurement.

Figures

Figure 1

Spatial location and distribution of the grassland flux sites used in the present study. The base map is derived from MODIS land cover product (MCD12Q1 C5, 500 m resolution) based on the International Geosphere Biosphere Programme (IGBP) global vegetation classification scheme in 2010. The detailed descriptions of these flux sites including CN-Cng, CN-Nmg, CN-Du2, CN-Hzz, CN-Sw2, CN-Ham, CN-Ar, CN-Ds, CN-Dan, CN-Ty and CN-Mq can be seen in Table 1. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 2

Comparisons of multi-year mean R_e during the growing period, dormant period and the entire year across grasslands in northern China. Error bars represent one standard error.

Figure 3

Seasonal trends of (a-d) ecosystem respiration (R_e), (e-h) air temperature (T_a) and soil temperature (T_s), (i-l) solar radiation (R_g) and vapor pressure deficit (VPD), (m-p) precipitation (P) and soil water content (SWC) in an 8-day interval at the representative grassland flux sites for selected years including CN-Cng 2008 (Temperate meadow steppe), CN-Nmg 2004 (Typical steppe), CN-Hzz 2014 (Desert steppe) and CN-Ham 2003 (Alpine meadow).

Figure 4

The seasonal dynamics of 8-day EVI, LAI, LSWI and LST at the representative grassland flux sites for selected years including CN-Cng 2008 (Temperate meadow steppe), CN-Nmg 2004 (Typical steppe), CN-Hzz 2014 (Desert steppe) and CN-Ham 2003 (Alpine meadow).

Figure 6

Comparisons of annual mean Re observations and the remotely-derived products across the temperate meadow steppe, typical steppe, desert steppe and alpine meadow sites.

Figure 7

Spatial patterns of grassland types (a) and annual mean Re (b) over the northern China's grasslands in 2010. The unit of Re is $\text{g C m}^{-2} \text{d}^{-1}$. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.