Fairness evaluation for the elder population in China

Guan Wang (✉ wangguanhz@163.com)

Hangzhou Medical College <https://orcid.org/0000-0001-6238-8002>

Qing Shen

Hangzhou Medical College

Research article

Keywords: Non-communicable diseases, the elderly, socioeconomic status, health financing, kakwani index

Posted Date: March 5th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-16166/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction: Population aging and the increasing burden of non-communicable diseases are increasingly a strain on health systems. The World Health Organization (WHO) uses fairness of health financing as one of the criteria for assessing health system performance. The Chinese government has undertaken a series of health reforms to reduce the pace of disease transition towards non-communicable diseases (NCDs), as well as protecting people from catastrophic health expenditures. The aim of this study was to assess the fairness of health financing among the elderly with different health conditions in China.

Methods: Fairness of health financing analysis was used to evaluate how the burden of health financing is distributed according to the household ability to pay (ATP). The data source was the WHO Study on global AGEing and adult health (SAGE) Wave 1, a national weighted data collected from adults older than 50 years. Chi-square and partial proportional odds model test were applied to assess the distribution of socioeconomic and health behavior factors among different chronic conditions. Kakwani indexes and curves were obtained from concentration index/ curves and Gini coefficient/ATP for progressivity analysis in healthcare financing. Dominance tests were applied for comparing different ways of health financing among health conditions.

Results: More than half of the elderly had at least one chronic condition, and around 20% people suffered from multi-morbidity. Several socioeconomic as well as health behavior factors were associated with developing NCDs. Out-of-pocket payment dominated other health finance sources in all three chronic conditions with a regressive pattern. Even though mandatory insurance had covered more than 70% of the elder population in China, due to the relative lower proportion reimbursement for chronic diseases, people still had to pay a lot for seeking healthcare.

Conclusion: When reimbursement for chronic diseases is relatively low, high mandatory insurance coverage does not ensure fairness of health financing. The Chinese health system should be developed further in order to meet the needs of elderly with different chronic conditions. **Key words :** Non-communicable diseases, the elderly, socioeconomic status, health financing, kakwani index

Background

Aging is a natural process, while population aging is a process which refers to the proportion of the elderly in a population increases all the time(1). Thanks to the relative stable politic environment and fast economic growth in China, the life expectancy has been increased strikingly in the past three decades, from 63.9 years in 1975 to about 76.5 years in 2017(2). The percentage of population with 65 plus years had reached to 7.7% by 2005, which had been ascribed as an “aged society” (3). Since aging population have higher risks in developing chronic diseases, and some other co-morbidities, interventions as well as efficient healthcare delivery specifically for the elderly are getting more concerns(4)(5).

Nowadays, China is undergoing fast industrialization and urbanization. Due to the fast socioeconomic and demographic transition, non-communicable diseases (NCDs) have become the main causes for death and disability in China(6). Non-communicable diseases (NCDs), which also called chronic diseases, is a type of disease which tend to be in a long duration and are the results of combined genetic, lifestyle or environmental factors(7). Cardiovascular diseases, cancers, chronic respiratory diseases, and diabetes are known as the common NCDs(8). NCDs increase dramatically in Chinese society, which is estimated above 260 million and has already put a big pressure in the field of public health(8)(9). As the prevalence of chronic diseases increases with age, multi-morbidity, which is defined as an individual suffers any two or more chronic or acute diseases, has become more common with the elderly(10). Multi-morbidity has brought several challenges to the health system, for more complex diagnosis and treatment, limited beds for overnight stays and polypharmacy prescription(11).

Chinese healthcare system had several transitions since 1949. The most well-known phrase was between 1949–1965, in which the national health advanced dramatically. By conducting a centralized three-tier delivery system, nearly the whole Chinese population were covered by a low and affordable health cost(12). However, since the 1977, along with the reform and opening-up policy, the health inequality was getting obvious. Due to lack of a well-developed health system, larger proportion of Chinese population could not afford to health care, catastrophic expenditure on healthcare occurred a lot(13). It arouse much concern to tackle with the insufficient health system in China(12). Up to 1990's, Chinese government attempted to take some measures to rein the accelerating healthcare costs, such as built-up the urban employee-based basic medical insurance scheme (UEBMI) for urban employees(13). During the post-SARS period in 2003, New Cooperative Medical System(NCMS) has been developed in response to the deterioration in access to the health service in rural areas and attempted to reduce the inequity between urban and rural areas(14). Later in 2007, urban resident-based basic medical insurance scheme (URBMI) has been developed for people living in urban areas without having UEBMI, such as students, children and the elderly without pensions(12)(15). Since then, a multi-level health care financing system has built up gradually for urban and rural population, attempting for healthcare funding equitable distributed(12)(13)(16).

Along with the healthcare reforms for achieving universal health coverage in China, fairness in health system financing has become a major concern for policy-makers as one of the three intrinsic goals in health system performance provided by WHO framework(16)(17).Health system financing mainly addresses two challenges(18). One is about financial risk pooling, in which households should not be impoverished for healthcare payments. The other is ensuring richer people contribute more than the poor when health financing(18). Fairer financing has been regarded as higher level of health and greater equality in health distribution(18). Within a fairer health system financing, poorer people could get access to the prompt and efficient medical treatment especially under the change of disease spectrum, where NCDs make poorer people in more vulnerable situation for more health expenses(12). In China, it mainly has four channels for healthcare financing: government-raised revenue; public health insurance; private health insurance as well as out-of-pocket payment(OOP)(16)(17). Government-raised financing draws revenues from general taxes for health system. Similar to most of other countries, tax is a stable and

important channel of getting fund for health resources in China(16). Public health insurances are constituted of UEBMI, URBMI and NCMS these three types, while due to the different per-capital funds and target population, those service packages are varied from each other(17). Private health insurance is still at initial stage comparing to many OECD countries, which accounts for only around 3% of the whole healthcare financing amount in China(16). OOP payment is to purchase health services directly from healthcare providers, which is more likely to generate catastrophic expenditure(18). Based on the four channels above, it has turned an urgent need to examine whether the health reforms have achieved goals in promoting universal coverage of health service as well as fairness financing(18)(19).

Though fairness of health financing has been assessed in many countries, seldom researches looked on whether is it fair for the elderly with different chronic conditions in respect of health financing. Thus, the aim of this study was to evaluate the health system performance in terms of the fairness of health financing system in China. More specifically, the main objectives were: Described distribution of chronic conditions among various socioeconomic status and risk factors and fairness analysis of Chinese healthcare financing system among the elderly with different chronic conditions.

Methods

Data collection

The data source for this study was the WHO Global AGEing and Adult Health (SAGE), a longitudinal study of health and ageing in China, Ghana, India, Mexico, Russia and South Africa. The aim of SAGE was to provide reliable data on aging people and their health status in low- and middle-income countries (20). The present study used data from Wave 1 (2007–2010) in China. In the SAGE study, the elderly population was defined as aged 50 years and above. This age cut-off was based on the WHO definition of younger and older adults in Low- and middle-income countries (20). Data were collected from Guangdong, Hubei, Jilin, Shanxi, Zhejiang, Shanghai, Shandong and Yunnan with considering geographical and economic effects. All the eligible households were included in conducting household interviews, and individual interviews were from the eligible respondents in the household roster. A proxy questionnaires were used when the individual respondents had cognitive limitations, who may find it hard to complete the interview. All these questionnaires were piloted as part of the pretest in 2005, then translated into Chinese. At last, 10278 household surveys and 15050 individual surveys had been collected. Post-stratification weights were used in both household and individual level analysis to adjust for province, locality, age and sex distributions.

Variables selection

Variables selection from the individual questionnaire

The results were only for the people who had completed interviews (interview were conducted including body measurement, performance tests and blood sample) or partial interview. NCDs selection was based on the self-reported responses to presence of one or more following diseases: arthritis, stroke, angina,

diabetes, chronic lung disease, asthma, cataract, and hypertension. Hypertension was identified as systolic blood pressure was ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg.

Social-demographic variables included age, sex, marriage status and educational level. People above 50 years were selected. Risk factors and health behavior variables contained body mass index (BMI), tobacco using, alcohol consumption, fruit/ vegetables intake as well as physical activity variables. BMI was derived from weight in kilogram (Kg) divided height in meter (m). China's center for disease control guidelines set BMI < 18.5 (kg/m^2) to underweight, $18.5\text{--}23.9$ (kg/m^2) to normal, $24\text{--}27.9$ (kg/m^2) to overweight and > 28 (kg/m^2) to obesity. The fruit/vegetables variable was defined as adequate with intake of fruit and vegetables ≥ 5 servings daily(20). Physical activity was measured by using WHO Global Physical Questionnaires(21).

Variables selection from the household questionnaire

Household asset items were chosen for deriving wealth quintiles. Under household expenditure module, variables like general tax, mandatory health insurance, voluntary health insurance as well as out of pocket (OOP) were picked up for information of healthcare financing. General taxes were consisted of property tax, vehicle tax, income tax and non-health related insurance. Out-of-pocket payments were made up of inpatient care, outpatient care, long term care cost, doctor's consulting fee, drug purchases, alternative or traditional medicine, diagnostic tests, dental care, ambulance fees and other costs with excluding the insurance reimbursement. The time frame for these four payments was set into 12 month. In this Chinese SAGE survey, designers classified UEBMI, URBMI and NCMS as mandatory health insurance.

The WHO SAGE study was approved by the Ethics Review Committee, World Health Organization, Geneva, Switzerland; the Ethics Committee, Shanghai Municipal center for Disease control and Prevention, Shanghai, China. All procedures were approval covered in this study. All participants got informed consent forms before started.

Data analysis

Wealth status was derived from household assets and Principal Component Analysis (PCA) was used to produce the wealth quintile(22). Factor test was a measurement to look at the adequacy of sampling, with using Bartlett's Test and KaiserMeyer-Olkin(KMO)(22)(23). At last, component scores were sort into five categories to get the wealth quintile and ranked into poorest to riches with the value of KMO = 0.871, p-value = 0.000, and the value of correlation matrix = 0.062. Chi-square test was applied for association analysis between socioeconomic risk factors and different health status. Partial proportional odds model was employed for comparing the contributions of each factors to different chronic conditions(24).

Progressivity analysis was chosen for examining the vertical equity on healthcare payments(25). Measuring progressivity was to compare shares of general tax, mandatory health insurance, voluntary health insurance as well as out of pocket payment(OOP) contributed by cumulative wealth of households

ranked by the share of ability to pay(ATP)(25). The fairness of healthcare expenditure was analyzed by using Kakwani index, which was one of the most widely used in measuring inequity in healthcare payment field(18). Kakwani index was formulated as the difference between concentration index for healthcare payments and Gini coefficient(18). Concentration index was known as the measurement of socioeconomic-related health inequality, and Gini coefficient was an indicator for measuring the degree of wealth inequity in a country(26). If health finance was in a vertical equity situation, poorer people would contribute less for the share of health payments than their share of ATP, which was also called progressive system. Whereas it was regressive system if sharing of health payments were greater than ATP sharing(17)(25). As recommended by the World Bank, household assets were measured to reflect the ATP since assets indices could reduce the currency fluctuation comparing to income or expenditure, which have been used more in developing countries(12)(25).

Ordinary least squares(OLS) regression was employed in estimation of CI and Gini coefficient in this study. It compared healthcare payment variables with household fractional ranking according to ATP distribution(25)(26).

$$2\sigma_r^2\left(\frac{z_i}{\gamma}\right) = \alpha + \beta R_i + \varepsilon_i$$

where z_i was a health payment variable for household i and γ was the mean. R_i was the household fractional ranking according to ATP distribution, σ_r^2 was the sample variance of the fractional rank. β in OLS was the estimation for CI or Gini coefficient.

The Kakwani index for healthcare payment was defined as the difference between concentration index for healthcare payments and Gini coefficient(18)(25).

$$\pi k = C - G$$

where C was the concentration index (CI) for healthcare payments, ranging from -1 to 1 . G referred to the Gini coefficient, with a range of 0 to 1 . Values increased as inequality increased. The value of πk was from -2 to 1 . The positive value indicated a progressive system in healthcare spending, to the graph, the concentration curve lied outside of the Lorenz curve, which meant the richer households paid more according to the ATP than poor households. While a negative value indicated a regressive spending, as the concentration curve lied inside of the Lorenz curve(18)(25).

As KI was the difference between CI and Gini coefficient, KI value could be computed by the following OLS regression(17)(25):

$$2\sigma_r^2\left[\frac{h_i}{\eta} - \frac{y_i}{\mu}\right] = \alpha + \theta R_i + \varepsilon_i$$

where h_i was a health payment variable for household i and n is the mean, y_i was the ATP variable and μ was its mean. R_i was the household ranking in ATP distribution and σ_r^2 was a sample variance of fractional ranking. In this formula, θ was estimated for KI value.

Sampling weights were included for all the analysis. Statistical software STATA 14.0 was applied in this study.

Results

Population Health

Table 1

Characteristics of respondents stratified by NCDs, adults aged 50 + China (2007–2010), SAGE Wave1

Characteristics	China (n = 11654)			
	No CD	Single morbidity	Multi-morbidity	P-value
	n = 4880	n = 2961	n = 2257	
Overall rates	49.86%	29.79%	20.35%	
Sex				P = 0.00**
male	2461(54.48)	1361 (28.94)	939 (16.58)	
Female	2419 (45.23)	1601 (30.65)	1318 (24.12)	
Age				P = 0.00**
50–59	2704 (60.76)	1270 (28.98)	509 (10.27)	
60–69	1338 (44.89)	950 (31.08)	763 (24.04)	
70–79	671 (34.27)	612 (30.08)	781 (35.65)	
80+	167 (33.97)	130 (27.76)	204 (38.27)	
Marital status				P = 0.00**
unmarried	47 (54.86)	20 (25.91)	14 (19.23)	
married/cohabit	4189 (50.94)	2489 (29.93)	1779 (19.13)	
widow/separate	644 (42.62)	453 (29.21)	564 (28.16)	
Residence				P = 0.00**
urban	1671 (39.75)	1321 (30.50)	1404 (29.75)	
rural	3209 (56.27)	1641 (29.35)	853 (14.38)	
Education				P = 0.00**
no education	1194 (46.76)	754 (29.92)	632 (23.32)	
less & primary school	2006 (52.40)	1133 (29.55)	784 (18.06)	
secondary & high school	1549 (50.44)	948 (29.59)	714 (19.97)	
college& university	131 (35.59)	127 (33.75)	127 (30.66)	
Wealth				P = 0.00**
1 poorest	1396 (59.36)	629 (27.22)	328 (13.42)	
Actual numbers of observations; Proportions in brackets sum across rows and include survey sampling weights. Person χ^2 tests were undertaken for comparisons. *P-value < 0.05,**P-value < 0.01				

Characteristics	China (n = 11654)		
2	1132 (52.00)	682 (30.71)	418 (17.28)
3	933 (48.56)	594 (29.23)	522 (22.20)
4	716 (39.61)	556 (31.31)	565 (29.08)
5 richest	641 (45.31)	453 (30.25)	397 (24.44)
Household insurance	P = 0.04*		
None of insurance	555 (46.41)	359 (28.20)	334 (25.39)
Mandatory health insurance	3846 (49.94)	2303 (29.72)	1748 (20.34)
Voluntary health insurance	253 (51.67)	152 (29.96)	101 (18.37)
Both of insurance	226 (52.50)	148 (33.08)	74 (14.41)
Body Mass Index	P = 0.00**		
underweight	266 (55.13)	131 (28.77)	83 (16.10)
normal	3314 (54.70)	1734 (27.63)	1224 (17.67)
overweight	1147 (42.93)	914 (33.41)	744 (23.66)
obese	153 (29.68)	183 (35.07)	206 (35.25)
Smoking	P = 0.00**		
Non-smoker	3052 (46.12)	1981 (30.74)	1611 (23.14)
Smoker	1828 (56.13)	980 (28.21)	646 (15.66)
Alcohol drinking	P = 0.00**		
No	3246 (47.61)	2047 (30.27)	1661 (22.13)
Yes	1634 (54.13)	915 (28.90)	596 (16.97)
Fruit/veg intake	P = 0.71		
Inadequate	735 (49.80)	443 (28.99)	362 (21.22)
adequate	4145 (49.86)	2519 (29.91)	1895 (20.23)
Physical activity	P = 0.00**		
Low	1388 (42.79)	945 (30.69)	859 (26.52)
Moderate	1175 (44.25)	802 (30.44)	726 (25.31)

Actual numbers of observations; Proportions in brackets sum across rows and include survey sampling weights. Person χ^2 tests were undertaken for comparisons. *P-value < 0.05,**P-value < 0.01

Characteristics	China (n = 11654)		
High	2317 (57.03)	1215 (28.92)	672 (14.04)
Actual numbers of observations; Proportions in brackets sum across rows and include survey sampling weights. Person χ^2 tests were undertaken for comparisons. *P-value < 0.05;**P-value < 0.01			

Table 2

Results of multiple partial proportional odds model, adults 50 + years China (2007–2010), SAGE Wave1

Comparisons						
Co-variable(s)	No CD vs. (Single morbidity & Multi-morbidity)			No CD & Single morbidity vs. (Multi-morbidity)		
	OR ₁	95%CI	P-value	OR ₂	95%CI	P-value
Sex						
Male	1	Reference		1	Reference	
Female**	1.29	(1.13,1.48)	0.00	1.29	(1.13,1.48)	0.00
Age						
50–59	1	Reference		1	Reference	
60–69**	1.85	(1.65,2.08)	0.00	2.64	(2.21,3.15)	0.00
70–79**	2.69	(2.29,3.17)	0.00	4.09	(3.47,4.83)	0.00
80+**	2.81	(2.13,3.72)	0.00	4.86	(3.50,6.74)	0.00
Marital status						
Unmarried	1	Reference		1	Reference	
Married/cohabit	0.85	(0.46,1.56)	0.59	0.85	(0.46,1.56)	0.59
Widow/separate	0.81	(0.44,1.52)	0.51	0.81	(0.44,1.52)	0.51
Residence						
City	1	Reference		1	Reference	
Rural**	0.63	(0.51,0.78)	0.00	0.49	(0.40,0.60)	0.00
Education						
No education	1	Reference		1	Reference	
Less & primary school	0.99	(0.87,1.13)	0.96	0.99	(0.87,1.13)	0.96
Secondary & high school	1.02	(0.87,1.20)	0.76	1.02	(0.87,1.20)	0.76
College & university	1.03	(0.84,1.27)	0.76	1.03	(0.84,1.27)	0.76

Note: Survey sampling weights applied; OR = odds ratio; 95%CI = 95% confidence Interval; *p-value < 0.05;

**p-value < 0.01

Comparisons						
Wealth						
1 Poorest	1	Reference			Reference	
2**	1.25	(1.08,1.44)	0.00	1.25	(1.08,1.44)	0.00
3	1.21	(0.98,1.50)	0.07	1.21	(0.98,1.50)	0.07
4	1.34	(1.03,1.75)	0.03	1.14	(0.87,1.48)	0.33
5 Richest	1.04	(0.81,1.34)	0.75	1.04	(0.81,1.34)	0.75
Household insurance						
None insurance	1	Reference		1	Reference	
Mandatory insurance	1.12	(0.97,1.29)	0.11	1.12	(0.97,1.29)	0.11
Voluntary insurance	0.96	(0.75,1.23)	0.77	0.96	(0.75,1.23)	0.77
Both of insurance	0.98	(0.74,1.30)	0.88	0.98	(0.74,1.30)	0.88
BMI						
Underweight	1	Reference		1	Reference	
Normal*	1.23	(1.02,1.49)	0.03	1.23	(1.02,1.49)	0.03
Overweight**	1.81	(1.51,2.18)	0.00	1.81	(1.51,2.18)	0.00
Obese**	3.17	(2.53,3.97)	0.00	3.17	(2.53,3.97)	0.00
Smoking						
Non-smoker	1	Reference		1	Reference	
Smoker	0.92	(0.81,1.05)	0.21	0.92	(0.81,1.05)	0.21
Alcohol consumption						
No	1	Reference		1	Reference	
Yes	1.09	(0.97,1.22)	0.13	1.09	(0.97,1.22)	0.13
Fruit/vegetables intake						
Inadequate	1	Reference		1	Reference	

Note: Survey sampling weights applied; OR = odds ratio; 95%CI = 95% confidence interval; *p-value < 0.05;

**p-value < 0.01

Comparisons						
Adequate	1.02	(0.90,1.16)	0.78	1.02	(0.90,1.16)	0.78
Physical activities						
Low	1	Reference		1	Reference	
Moderate	0.95	(0.80,1.12)	0.54	0.95	(0.80,1.12)	0.54
High**	0.73	(0.65,0.82)	0.00	0.64	(0.54,0.76)	0.00
Note: Survey sampling weights applied; OR = odds ratio; 95%CI = 95% confidence Interval; *p-value < 0.05;						
**p-value < 0.01						

Table 1 described the distribution of socioeconomic and behavior risk factors among the elder population in China. In this study, 237 people were excluded for not completing the survey, 1265 people were then excluded for analysis because their ages were under 50 years. After missing check for all selected variables, 10099 people were derived from 15050 of Chinese sample for future analysis. Among these population, 49.86% did not have any chronic diseases, 29.79% had single NCD morbidity and 20.35% were in multi-morbidity status.

By applying chi-square test, sex, age, marriage status, residence, education, wealth, body weight, smoking, alcohol consumption and physical activities were found statistically significant difference with the selected eight chronic diseases. Through adjusting to the partial proportional odds model, sex, older age, wealthier status and overweight were played as risk factors whereas rural residence and physical activities were considered as protective factors in the selected chronic diseases developing, in which, odds ratios of family wealth showed middle-class had higher risks for chronic diseases and odds ratios of age were found to be increased when people were in multiple chronic conditions.

Fairness in healthcare financing

Table 3

Distribution of household assets and health care payments by wealth quintiles, concentration index (CI), and Kakwani index

Classification of health status	Household wealth quintiles	Ability to pay(ATP)	general tax	mandatory health insurance	voluntary health insurance	OOP
No CD	Q1 Poorest	5.00%	1.05%	10.30%	1.20%	11.64%
	Q2	15.98%	13.07%	13.09%	10.29%	16.79%
	Q3	19.75%	21.41%	13.49%	12.49%	19.39%
	Q4	24.00%	27.69%	22.83%	22.81%	22.53%
	Q5 Richest	35.28%	36.77%	40.29%	53.21%	29.65%
	Gini/CI	0.37	0.46	0.43	0.49	0.25
	P-value	-	0.00**	0.00**	0.00**	0.00**
	Kakwani index	-	0.09	0.06	0.12	-0.12
	P-value	0.00**	0.48	0.49	0.32	0.01**
Single morbidity	Q1 Poorest	3.45%	2.33%	6.48%	1.23%	13.50%
	Q2	13.37%	14.70%	11.49%	14.14%	15.49%
	Q3	18.06%	17.17%	16.95%	15.56%	20.38%
	Q4	28.17%	31.19%	23.37%	33.80%	22.49%
	Q5 Richest	36.96%	34.55%	41.71%	35.28%	28.15%
	Gini/CI	0.33	0.41	0.40	0.41	0.09
	P-value	-	0.00**	0.00**	0.00**	0.03*
	Kakwani index	-	0.08	0.07	0.08	-0.23
	P-value	0.00**	0.54	0.63	0.41	0.00**
Multi-morbidity	Q1 Poorest	2.33%	0.17%	4.26%	0.48%	4.36%
	Q2	9.31%	4.82%	10.02%	12.64%	11.19%
	Q3	17.35%	12.25%	10.94%	14.27%	21.24%
*p-value < 0.05;**p-value < 0.01						

Classification of health status	Household wealth quintiles	Ability to pay(ATP)	general tax	mandatory health insurance	voluntary health insurance	OOP
	Q4	29.56%	25.41%	37.35%	29.39%	27.54%
	Q5 Richest	41.45%	57.34%	37.43%	43.23%	35.67%
	Gini/CI	0.28	0.28	0.32	0.30	0.17
	P-value	-	0.01**	0.00**	0.36	0.00**
	Kakwani index	-	-0.01	0.03	0.01	-0.11
	P-value	0.00**	0.92	0.72	0.98	0.00**
*p-value < 0.05;**p-value < 0.01						

Table 4

Tests of dominance between concentration curves for different sources of health finance

Classification of health status	Sources of health financing	General tax	Mandatory health insurance	Voluntary health insurance	OOP
No CD	General tax		D	D	D
	Mandatory health insurance			None	D
	Voluntary health insurance				D
	OOP				
Single morbidity	General tax		D	D	D
	Mandatory health insurance			None	D
	Voluntary health insurance				D
	OOP				
Multi-morbidity	General tax		D	D	D
	Mandatory health insurance			None	D
	Voluntary health insurance				D
	OOP				
Note: D indicated that concentration curve of row source was dominated by the column source; None indicated failed to reject the null hypothesis that curves were indistinguishable at the 5 percent significance level.					

Table 3 reported the distribution of household assets and health care payments by wealth quintile. In all three health status groups, the concentration indices were all positive. This indicated that the rich contributed more in terms of healthcare payments than the poor. The Gini coefficient and CI values were the highest in the non-chronic disease group, and were lower as higher number of chronic conditions. An exception was OOP payment, for which the CI was the lowest in the group with a single chronic condition. CI values for voluntary health insurance were consistently higher than for other types of healthcare payment. This suggested that voluntary health insurance was more unequally distributed. Comparing to other healthcare payments, OOP payment distributed proportionally to the household wealth.

The results of the dominance tests between concentration curves for the different sources of health financing were presented in Table 4. OOP payment dominated the other three health payments regardless of the number of chronic conditions.

The CI value was limited in that it only indicated whether healthcare payments were equally distributed with respect to household wealth. To further evaluate the fairness of health financing, Kakwani Indexes were calculated with considering the ATP in a household. (Values were represented in Table 3). The KI values for general tax, mandatory health insurance, and voluntary health insurance payments were all positive across different health status, with only one exception. These positive values indicate that general tax, and both types of health insurance were all progressive ways of financing. Furthermore, KI values were always higher in the voluntary health insurance group. While KIs in OOP groups were always in negative regardless of health conditions. The lowest KI in OOP classification was -0.23, which was in single morbidity group and the highest KI was in multi-morbidity group with value of -0.11.

It could be seen from Fig. 2 that OOP payment was in a regressive spending, for the curves always lied inside of Lorenz curves, indicating OOP payment was beyond people's ability to pay. In multi-morbidity group, the distance was shortest between OOP and ATP curves, while presented the longest in single morbidity group, stating people with single NCD morbidity suffered a most unfair financing distribution with OOP payment. General tax curve had intersected with ATP in non-chronic diseases group at nearly 60% cumulative wealth sharing, while almost overlapped with ATPs in the other two groups. Mandatory health insurance curves coincided with ATPs mostly in all three groups, showing it was a progressive spending in most cases. Voluntary health insurance payment was below ATP in non-NCD and single morbidity groups, while intersected with ATP in multi-morbidity group, stating elder population from 28–70% cumulative proportions of wealth level in the group, could not afford for the voluntary insurance.

Discussion

This study used data from WHO SAGE WAVE1 (2007–2010) to evaluate the fairness of health financing for the elderly with chronic conditions in China. More than half of the elderly had some chronic conditions, and 20.35% had multi-morbidity; these results fall within the prevalence range reported by researchers(27)(28). Socioeconomic and several health risk factors were found to play an important role in developing chronic diseases. Previous studies found the rise of NCDs in most developing countries were probably linked to the urbanization(29). Due to profound changes in exercise and diet, the risk of having diabetes and cardiovascular diseases attendant increase(29). People living in rural areas were less likely to develop chronic conditions owing to a better natural environment, healthier lifestyle and less densely population comparing to the urban people(29)(30). However, the finding might also arise because the presence of chronic conditions in this study was self-reported. Due to inequalities in the allocation of health resources, access to prompt and effective healthcare is not as good in rural areas, and chronic conditions may not be diagnosed properly(27)(31). There may similarly be a bias towards under-reporting in a poorer population for lack of accurate diagnosis(32).

Healthcare financing mainly consisted of four types, of which out-of-pocket payment stood out in this study. OOP payment was found to be the most regressive type regardless of health status, even though OOP spending decreased from 57.7% in 2002 to 34.8% in 2011 under health reforms(33)(34). In the single morbidity group, the Kakwani Index had the smallest value (-0.23) value comparing to the rest two

groups, which indicated poorer people spent more for their healthcare treatment according to the ATP sharing.

Mandatory health insurance covered more than 70% of elder population in the study. However, the services for treating chronic diseases included in the insurance package were limited, especially in NCMS and URBMI(35), people still need to pay a lot for healthcare expenditure through Out-Of-Pocket, which might account for the KI was lowest in disease group.

In the multi-morbidity group, tax payment was also somewhat regressive. In other words, tax posed a financial burden on the poor with multimorbidity, although less than OOP. A common finding in both high and middle income countries is that tax revenue is a progressive form of healthcare financing(36). In contrast to most high-income countries, indirect taxes such as value-added tax, sales tax, and excise taxes (e.g. on alcohol and tobacco) constitute the majority of total tax revenue in China (52% in 2010) (16). In this study, tobacco and alcohol consumption were taken as risk factors for having chronic conditions. Thus, higher consumption on tobacco and alcohol could partly explained why people in multi-morbidity status were in a regressive system for tax payment. Regarding this, redistribute of income wealth through tax payment, reforms are called to ensure the equity.

We also found the different values of redistribution for household insurance in three groups. Although both mandatory and voluntary health insurance were all in a progressive pattern in three groups, they did not contribute proportionally comparing to the according ATP sharing. As mandatory health insurance was consisted of UEBMI, URBMI and NCMS types, and the way of financing for different types varied from each other, people could face the health payment beyond their ATP(17)(35). UEBMI-covered population were requested to pay a certain proportion of their salary as premiums (17)(37), while it was a fixed number for premium of URBMI and NCMS from the insured health payment, without considering so much about the ability to pay in most of cases(38)(39). It explained partly that in all three groups, mandatory insurance curve was always higher than ATP sharing at the beginning. As a typical pre-paid payment for healthcare financing, national health insurance should play an important role in preventing people from catastrophic and impoverishing in health expenditures(16)(17). To achieve this objective, Chinese government should take measures to ensure the public health insurance could be afforded in any income groups according to their ability to pay.

Even though the value of voluntary health insurance was positive (0.01) in multi-morbidity, the curve fluctuated the most in this group, and the negative effect could be offset due to the crossed over curves(25), which could find at the third picture in Fig. 3, the curve of voluntary insurance was inside of ATP sharing at middle class households. Not like most OECD countries, voluntary health insurance only takes smaller proportion under health financing in China(36). We found the voluntary health insurance covered around 10% of elder population, and the proportion of insured was much smaller comparing to the mandatory health insurance. Since the service benefits insured by URBMI and NCMS were limited for both outpatient and inpatient care, people still had to pay more than half of whole expenses when seeking healthcare(6)(17)(40), which increased the demand for voluntary insurance specifically for the

elderly. Previous studies also showed that the commercial insurance was highly associated with socioeconomic status as well as personal knowledge on insurance purchase(16)(17). People with chronic conditions were more likely to consume voluntary health insurance for reducing out-of-pocket health expenses. As premiums of private insurance were more related to personal risk factors rather than income, it is very likely to exceed people's ATP(41). These might explain why the commercial insurance was in regressive pattern among the middle-class in multi-morbidity group. Heterogeneity of health risks as well as asymmetric information between insurers and insureds could cause problems since people in higher risks of developing diseases might pay more than lower risks(42)(43)(44). Likewise, moral hazard could also occur when policyholders were lack of honesty(45). Due to the imperfect rules of private health insurance, it still has long way to go for a fairness financing system.

Our studies had several limitations. First of all, the data for analysis was almost 10 years ago. However, because Chinese health system has experience a lot during these years, the findings may vary from updated setting. Secondly, SAGE only took arthritis, stroke, angina, diabetes, chronic lung disease, asthma, hypertension, cataracts, oral health as chronic conditions, while more chronic conditions such as cancers or cardiovascular diseases could be included in future questionnaire design for a comprehensive comparison between non-NCDs and NCDs groups. Another weakness was since the questionnaires do not classify the mandatory health insurance into UEBMI, URBMI and NCMS, fairness of the compulsory insurance could not be looked in detail. Future analysis should be carried on this study.

Conclusions

This study used the data from WHO SAGE WAVE1 (2007–2010) for evaluation the fairness of health financing system for the elderly with chronic conditions in China. More than half of the elderly had some chronic conditions, and 20.35% had multi-morbidity. Different socioeconomic status as well as risk factors accounted for reasons in developing the chronic conditions. Out of pocket payment was in regressive pattern among different chronic conditions, and KI value was lowest in single morbidity group. Even though mandatory insurance covered more than 70% of the elderly, but due to the relative lower proportion reimbursement for chronic diseases as well as fragmented public health insurances, people still have to pay a lot for seeking care and the poorer suffered more from the increased health expenses. A health system which allows majority of people could afford for health service is still a big challenge in China especially under population aging and epidemiological transition background. Because of the inconvenience and disparity schemes across regions, Chinese government has to take steps in merging those three insurances. Health system should be developed in order to meet the needs of elderly with different chronic conditions as well as achieve the universal health coverage.

Abbreviations

BMI: Body mass index; NCDs: Non-communicable diseases; SES: Socioeconomic status; NCMS: New Cooperative Medical System; UEBMI: Urban employee-based basic medical insurance scheme; URBMI:

Urban resident-based basic medical insurance scheme; OOP: Out-of-pocket; CI: Concentration index; KI: Kakwani index; SAGE: Study on global AGEing and adult health; WHO: World health organization

Declarations

Acknowledgements

We are grateful to the respondents of SAGE Wave 1 in China, India, Ghana and South Africa and to the WHO for making the WHO-SAGE dataset publicly available. Support for the SAGE-Wave 1 was provided by the United States National Institute on Aging (NIA) Division of Behavioral and Social Research (BSR) through Interagency Agreements (YA1323–08-CN-0020; Y1-AG-1005–01). This paper is based in part on the MSc dissertation of the first author. We are very grateful to Nawi Ng for introduction to the SAGE dataset, Anni-Maria Pulkki-Brännström for help in conceiving the study and much valuable comments on an early version of this manuscript.

Authors' contributions

GW made a substantial contribution to the conception of the study, analyzed the data, wrote the first draft and reviewed the literature. QS helped in developing the draft. All authors read and approved the final draft.

Available of data and materials

The anonymized datasets are in the public domain:

<http://apps.who.int/healthinfo/systems/surveydata/index.php/catalog/central> SAGE is committed to the public release of study instruments, protocols and meta- and micro-data: access is provided upon completion of the Users Agreement available through WHO's SAGE website:

www.who.int/healthinfo/systems/sage and the WHO archive using the National Data Archive application (<http://apps.who.int/healthinfo/systems/surveydata>). The questionnaires and other materials can be found at: <http://www.who.int/healthinfo/sage/cohorts/en/index2.html>. SAGE is committed to the public release of study instruments, protocols and meta- and micro-data: access is provided upon completion of the Users Agreement available through WHO's SAGE website (www.who.int/healthinfo/systems/sage).

Consent for publication

The data were provided after completion of User's agreement available through the WHO SAGE website. The manuscript does not contain individual personal data, therefore statement of consent is not applicable.

Ethics approval and consent to participate

The WHO-SAGE study was approved by the Ethics Review Committee, World Health Organization, Geneva, Switzerland and the individual ethics committees in each of the SAGE countries. Informed consent was

obtained from all respondents before the interviews were initiated. Confidential records of participants are maintained by SAGE country teams.

Funding

There are no funding sources to declare regarding the development and preparation of this manuscript.

Competing interests

The authors declare that they have no competing interests.

References

1. United Nations. Aging. 2019. Available from: <https://www.un.org/en/sections/issues-depth/ageing/>
2. The World Bank. Life expectancy at birth, total (years)-China.2019. Available from: <https://data.worldbank.org/indicator/SP.DYN.LE00.IN?locations=CN>
3. National Bureau of Statistics of China 2006 [Internet]. Beijing: China Statistical Press Available from: <http://www.stats.gov.cn/tjsj/ndsj/2006/indexeh.htm>
4. Boutayeb A, Boutayeb S. The burden of non-communicable diseases in developing countries. *Int J Equity Health*. 2005;4:2. DOI:10.1186/1475-9276-4-2
5. Shrivastava SRBL, Shrivastava PS, Ramasamy J. Health-care of Elderly: Determinants, Needs and Services. *Int J Prev Med*. 2013 Oct;4(10):1224–5.
6. Tang SL, Ehiri J, Long Q. China's biggest, most neglected health challenge: Non-communicable diseases. *Infect Dis Poverty*. 2013;2:7. DOI:10.1186/2049-9957-2-7
7. Sommer I, Griebler U, Mahlknecht P, et al. Socioeconomic inequalities in non-communicable diseases and their risk factors: an overview of systematic reviews. *BMC Public Health*. 2015;15:914. DOI:10.1186/s12889-015-2227-y
8. World Health Organization. A multi-country analysis of non-communicable disease surveillance data. 2010. Available from: http://www.wpro.who.int/publications/PUB_9789290614746/en/
9. China National Plan for NCD Prevention and Treatment (2012-2015). Chinese Center for Disease Control and Prevention. 2012. Available from: http://www.chinacdc.cn/en/ne/201207/t20120725_64430.html
10. Wallace E, Salisbury C, Guthrie B, et al. Managing patients with multimorbidity in primary care. *BMJ*. 2015; 350:h176. DOI:10.1136/bmj.h176
11. Salisbury C. Multimorbidity: redesigning health care for people who use it. *The Lancet*. 2012 Jul 7;380(9836):7–9.
12. Tang S, Meng Q, Chen L, et al. Tackling the challenges to health equity in China. *The Lancet*. 2008 Oct 31;372(9648):1493–501.

13. Zhe D, Michael R P. Evolution of China's health-care system. *The Lancet* 2008. Nov 15;372(9651):1715–6.
14. Yang W. China's new cooperative medical scheme and equity in access to health care: evidence from a longitudinal household survey. *Int J Equity Health*. 2013;12:20. DOI:10.1186/1475-9276-12-20
15. Chen M, Zhao Y, Si L. Who pays for health care in China? The case of Heilongjiang province. *PLoS ONE*. 2014; 9(10):e108867. DOI: 10.1371/journal.pone.0108867
16. Chen M, Chen W, Zhao Y. New evidence on financing equity in China's health care reform - A case study on Gansu province, China. *BMC Health Serv Res*. 2012; 12: 466. DOI: 10.1186/1472-6963-12-466
17. Chen M, Palmer AJ, Si L. Improving equity in health care financing in China during the progression towards Universal Health Coverage. *BMC Health Serv Res*. 2017;17:852. DOI 10.1186/s12913-017-2798-7
18. Christopher JL M, Felicia K, Philip M, et al. Defining and Measuring Fairness in Financial Contribution to the Health System [Internet]. Geneva: World Health Organization; 2000. Available from: <http://www.who.int/healthinfo/paper24.pdf>
19. Garrett L, Chowdhury AMR, Pablos-Méndez A. All for universal health coverage. *The Lancet*. 2009 Oct;374(9697):1294–9.
20. Kowal P, Chatterji S, Naidoo N, et al. Data resource profile: the World Health Organization Study on global AGEing and adult health (SAGE). *International Journal of Epidemiology*. 2012 Dec; 41(6):1639–49.
21. World Health Organization. GPAQ Analysis Guide. 2004. Available from: http://www.who.int/chp/steps/resources/GPAQ_Analysis_Guide.pdf
22. Howe LD, Galobardes B, Matijasevich A, et al. Measuring socio-economic position for epidemiological studies in low- and middle-income countries: a methods of measurement in epidemiology paper. *Int J Epidemiol*. 2012 Jun;41(3):871–86.
23. Vyas S, Kumaranayake L. Constructing socio-economic status indices: how to use principal components analysis. *Health Policy Plan*. 2006 Nov;21(6):459–68.
24. Richard W. Understand and interpreting generalized ordered logit models. *The Journal of Mathematical Sociology*. 2016 Jan;40:1,7-20.
25. Owen O, Eddy Van D, Adam W, Magnus L. Analyzing Health Equity Using Household Survey Data-A guide to techniques and their implementation. Washington, DC: The world bank. 2008.
26. Erreygers G, Van Ourti T. Measuring socioeconomic inequality in health, health care and health financing by means of rank-dependent indices: a recipe for good practice. *J Health Econ*. 2011;30(4):685–694. DOI:10.1016/j.jhealeco.2011.04.004
27. Wang R, Yan Z, Liang Y, et al. Prevalence and Patterns of Chronic Disease Pairs and Multimorbidity among Older Chinese Adults Living in a Rural Area. *PLoS ONE*. 2015 Sep;10(9). DOI: 10.1371/journal.pone.0138521

28. Kunna R1, San Sebastian M2, Stewart Williams J. Measurement and decomposition of socioeconomic inequality in single and multimorbidity in older adults in China and Ghana: results from the WHO study on global AGEing and adult health (SAGE). *Int J Equity Health*. 2017 May 15;16(1):79. DOI: 10.1186/s12939-017-0578-y
29. Leon AD. Cities, urbanization and health. *International Journal of Epidemiology*. 2008; 37(1): 4–8. DOI: [10.1093/ije/dym271](https://doi.org/10.1093/ije/dym271)
30. Kearns G, Brändström A, Tederbrand LG. The urban penalty and the population history of England, Society, Health and Population During the Demographic Transition. 1988. 213-236.
31. Wang HX, Wang JJ, Wong YS, et al. Epidemiology of multimorbidity in China and implications for the healthcare system: cross-sectional survey among 162,464 community household residents in southern China. *BMC Med* **12**, 188 (2014). DOI:10.1186/s12916-014-0188-0
32. Wei Y, Panos K. The less healthy urban population: income-related health inequality in China. *BMC Public Health*. 2012; 12: 804.
33. Zhang L, Liu N. Health reform and out-of-pocket payments: lessons from China, *Health Policy and Planning*. 2014;29(2):217–226.DOI:10.1093/heapol/czt006
34. National Bureau of Statistics of China. Annual data. 2020. Available from: <http://data.stats.gov.cn/easyquery.htm?cn=C01>
35. Meng Q, Fang H, Liu X, et al. Consolidating the social health insurance schemes in China: towards an equitable and efficient health system. *The Lancet*. 2015 Oct;386(10002):1484–92.
36. O'Donnell O, Van Doorslaer E, Rannan-Eliya RP, et al. Who pays for health care in Asia? *Journal of Health Economics*. 2008 Mar;27(2):460–75.
37. Hu S, Tang S, Liu Y, et al. Reform of how health care is paid for in China: challenges and opportunities. *Lancet* 2008; 372:1846-53. DOI: DOI:10.1016/S0140-6736(08)61368-9
38. Lin W, Liu GG, Chen G. The Urban Resident Basic Medical Insurance: a landmark reform towards universal coverage in China. *Health Economics*. 2009 Jul;18 Suppl 2:S83-96.
39. Liu C, Haan A de, Xiulan Z, Warmerdam W. Addressing vulnerability in an emerging economy: China's New Cooperative Medical Scheme (NCMS). *Canadian Journal of Development Studies*. 2011; 32(4):339-413.
40. Zhao, M., Liu, B., Shan, L. et al. Can integration reduce inequity in healthcare utilization? Evidence and hurdles in China. *BMC Health Serv Res*.2019;654. DOI:10.1186/s12913-019-4480-8
41. Alexander N, Claudia Süßmuth D, Florian T. Private health insurance in China finding the winning formula. *Health international*. 2012:74-82.
42. Bank TW. Public expenditure and the role of government in the Chinese health sector. The World Bank; 2005 May; p: 1–8. Report No: 33234.
43. Cutler MD, Finkelstein A, McGarry K. Preference Heterogeneity and Insurance Markets: Explaining a Puzzle of Insurance. *Am Econ Rev*. 2008 May; 98(2): 157–162. DOI 10.1257/aer.98.2.157

44. Cheung D, Padiou Y. Heterogeneity of the effects of health insurance on household savings: Evidence from rural China. 2013. ffhalshs-00848061f
45. Einav L, Finkelstein A. Moral Hazard in Health Insurance: What We Know and How We Know It. J Eur Econ Assoc. 2018 Aug; 16(4):957-982. DOI: 10.1093/jeea/jvy017

Figures

Figure 1

Conceptual cumulative concentration curve for healthcare payment and household wealth

Figure 2

Derivation of study sample from SAGE WAVE1 Chinese population

Figure 3

Concentration curve of healthcare payments and Lorenz curve sorted by health status