

Antifungal activity and genomic characterization of the isolate Bacillus velezensis CMRP 4489, a biocontrol agent for plant-pathogenic fungi

Julia Pezarini Baptista Universidade Estadual de Londrina Gustavo Manoel Teixeira Universidade Estadual de Londrina Maria Luiza Abreu Jesus Universidade Estadual de Londrina Rosiana Bertê Universidade Estadual de Londrina Allan Higashi Universidade Estadual de Londrina Mirela Mosella Universidade Estadual de Londrina **Daniel Vieira Silva** Universidade Estadual de Londrina João Paulo Oliveira Universidade Estadual de Londrina **Danilo Sipoli Sanches** Universidade Tecnológica Federal do Paraná **Jacques Duílio Brancher** Universidade Estadual de Londrina Maria Isabel Balbi-Peña Universidade Estadual de Londrina Ulisses Pereira Padua Universidade Estadual de Londrina Admilton Gonçalves de Oliveira (\square admilton@uel.br) Universidade Estadual de Londrina

Article

Keywords: Rhizosphere colonization, antifungal activity, secondary metabolites

Posted Date: May 10th, 2022

DOI: https://doi.org/10.21203/rs.3.rs-1619465/v1

License: ©) This work is licensed under a Creative Commons Attribution 4.0 International License. Read Full License

Abstract

The development of bio-based products has increased on account of numerous factors, including environmental awareness, human health concerns, and the more frequent selection of resistant pathogens to synthetic products. Among the studied genera for product development are the species of the *Bacillus* genus due to their elevated production of molecules with antimicrobial activity and resistance to extreme environmental conditions through endospore formation. In this context, the antifungal potential of *Bacillus velezensis* strain CMRP 4489 was investigated using *in silico* predictions of secondary metabolites in its genome and *in vitro* tests against the following phytopathogenic fungi: *Sclerotinia sclerotiorum* (SS), *Macrophomina phaseolina* (MP), and *Botrytis cinerea* (BC). The *in silico* predictions indicated that CMRP 4489 possesses several Biosynthetic Gene Clusters (BGCs) capable of producing molecules with antifungal properties and other non-identified BGCs. The *in vitro* assay results evidenced strong antifungal activity, inhibiting more than 60% of the tested fungi, and the isolate's molecules were stable under diverse physicochemical conditions. Therefore, CMRP 4489 showed to be a potential biocontrol agent against various diseases with agronomical importance and can be used under adverse environmental conditions.

Introduction

The world agrarian sector currently recognizes environmental issues as a positive agenda in favor of more efficient, higher-quality production, capable of limiting the expansion of agricultural frontiers and environmental toxicity ¹. Producers, who in recent decades have invested in the development of specialized and highly technological agriculture for greater productivity, are also considering the recovery of degraded areas, investments in the diversification of their properties, crop rotation, mixed planting, and the use of biological control agents (BCAs) ².

The use of BCAs as a form of management or integrated management to protect crops against pests, weeds, and microbial diseases for more sustainable food production and to cause less impact on the environment corroborates the aspirations of a society that increasingly demands safer and cleaner products and processes that are more ecofriendly ³. In addition to environmental and public health factors, it is important to highlight the emergence of a new economic trend - bioeconomy - focused on bio-based industries and businesses ⁴.

BCAs are widely used in many fields, including agriculture, medicine, and forestry products. Multiple genera have been studied as potential biocontrol agents. However, the *Bacillus* genus deserves mention because of its high capacity to synthesize a vast array of beneficial substances of agronomical and industrial interest, as well as being highly resistant to elevated temperatures due to its ability to form endospores ^{5,6}, thus being used in scale in bio-based industries.

Among the different species of the genus *Bacillus*, *B. velezensis* stands out due to its significant environmental versatility and for producing numerous antifungal compounds, such as lipopeptides

⁷,enzymes ⁸, and volatile compounds ⁹, in addition to acting as a Plant Growth-Promoting Rhizobacteria (PGPR) ¹⁰. Therefore, evaluating a new strain of *B. velezensis* could lead to the discovery of novel bioactive compounds, as well as position this species as a promising biocontrol agent for the bio-industry. In the present study, we aimed to assess the antifungal activity and stability of the extracellular metabolites of *B. velezensis* strain CMRP 4489, its influence in the suppression of *Sclerotinia sclerotiorum* on soybean seeds, and its biosynthetic potential, based on genome mining.

Materials And Methods

Plant-pathogenic fungi

Fungal isolates of *S. sclerotiorum* (SS), *Macrophomina phaseolina* (MP), and *Botrytis cinerea* (BC) were grown in potato dextrose agar (PDA) medium (Neogen Corporation, USA) at 25 °C, with a photoperiod of 12 h/12 h, and stored in the same medium at 4°C. All isolates were deposited in the Microbial Culture Collection of the Plant Pathology Laboratory of the Department of Agronomy of the State University of Londrina.

Biological control agent

The *B. velezensis* strain CMRP 4489 (LABIM40) used in this study was originally isolated as an antagonistic contaminant on a plate with fungal growth at the State University of Londrina (UEL), in Paraná, Brazil. Its complete genome has been sequenced and announced ¹¹. CMRP 4489 was maintained at the Laboratory of Microbial Biotechnology at UEL, and was also deposited as strain CMRP 4489 in the *Coleções Microbiológicas da Rede Paranaense* (CMRP) network of the Federal University of Paraná, in Curitiba, Brazil.

Phenotypic characterization and colony architectureof B. velezensis CMRP 4489

B. velezensis CMRP 4489 was grown in Luria-Bertani broth (LBB) medium (Neogen Corporation, USA) at 28 °C and, after 24 h, Gram staining was performed using a Gram-staining kit for morphology and cell wall visualization. Endospore formation was observed using the Wirtz-Conklin method after 48 h of incubation. Scanning electron microscopy (SEM) was conducted to visualize colony morphology. In this step, a colony grown for 48 h was removed from the agar and fixed in a solution containing 2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M sodium cacodylate buffer (pH 7.2) at 4 °C. The samples were kept in this solution overnight for fixation, after which they were washed three times with 0.1 M sodium cacodylate buffer (pH 7.2) for 10 min, followed by dehydration three times for 10 min in an ethanol series (30, 50, 70, 90, and 100%). The samples were then submitted to critical point drying with CO₂ (BALTEC CPD 030 Critical Point Drier), coated with gold (BALTEC SDC 050 Sputter Coater), and observed with an FEI Quanta 200 scanning electron microscope (SEM) operating at 25.0 kV.

Floating pellicle biofilm and swarm expansion assays

The CMRP 4489 strain was evaluated regarding floating pellicle biofilm formation in 24-well plates. An inoculum was prepared from a culture grown overnight at 28 °C in LBB medium (Neogen Corporation, USA), and the suspension was adjusted according to the 0.5 McFarland scale, rendering approximately 1.5×10^8 colony-forming units / mL (CFU/mL). Afterward, 10 µL of the inoculum were added to a 24-well plate (Sarstedt, USA) containing 2 mL of LBB medium, which was incubated at 28 °C for 24 h. For the swarm expansion assays, the CMRP 4489 inoculum was prepared as described above. After that, 90 mm-diameter Petri dishes containing LBB medium with 0.7% agar (Neogen Corporation, USA) were dried in a biological safety chamber (Filterflux, Brazil) for 30 min and inoculated centrally with 10 µL of the inoculum, dried for another 10 min, and incubated at 28 °C for 24 h.

Antagonistic activity

Using an *in vitro* dual-culture assay, the CMRP 4489 inoculum was subjected to antagonism assays against SS, MP, and BC in PDA medium. To this end, 6 mm agar plugs containing 7-day-old fungal mycelia were placed at the center of 90 mm Petri dishes. Five microliters of the CMRP 4489 inoculum (prepared as described in Subheading - *Floating pellicle biofilm and swarm expansion assays*) were transferred onto the Petri dishes at approximately 25 mm from the fungal mycelia plugs at four equidistant points. The plates were incubated at 25°C with a photoperiod of 12 h/12 h. Antifungal activity was expressed as the percentage of mycelial growth inhibition (MGI), according to the following formula:

where *MGI* (%) is the percentage of mycelial growth inhibition; *C* represents the colony radius of the fungal control plates, and *T* is the radius of the fungal colony in the treatment plates ¹². The experiment was repeated twice with 8 replicates, and the results were submitted to analysis of variance (ANOVA) and the means compared by the Tukey test (p<0.05).

Antifungal metabolite production

The CMRP 4489 strain was activated on Luria-Bertani agar (LB) (Neogen Corporation, USA) and incubated at 28 °C for 24 h. For the preparation of the pre-inoculum, colonies were suspended in saline solution (0.85% sodium chloride, w/v), and the concentration was adjusted according to the 0.5 McFarland scale until reaching approximately 1.5×10^8 CFU/mL. For the preparation of the inoculum, 30 µL of pre-inoculum were inoculated separately in 125 mL Erlenmeyer flasks containing 30 mL of two culture media (CM1 or CM2) and incubated at 28 °C for 24 h at 125 rpm (Orbital shaker - Thoth 6430B, Brazil). Afterward, a 1% aliquot of the final volume (v/v) was transferred to a 1000 mL Erlenmeyer flask containing 400 mL of CM1 or CM2:

CM1 – g/L: tryptone 10.0; yeast extract 5.0; NaCl 5.0; pH 7.1.

CM2 – g/L: glucose 20.0; tryptone 12.4; NaCl 5.0; K_2 HPO₄ · 3H₂O 1.5; MnSO₄ · H₂O, 0.04; FeSO₄ · 7H₂O, 1.67; MgCl₂ · 6H₂O, 1.22; pH 7.1.

All culture media were incubated at 28 °C for 72 h at 150 rpm (Orbital shaker - Thoth 6430B, Brazil). Next, the fermentations were centrifuged for 10 min at 8860 × g at 4 °C (Hitachi, CR21G Himac, Japan) to obtain the cell-free supernatants (CFS). All CFS were sterilized by filtration through a cellulose filter with a pore size of 0.22 µm (Millipore, USA), thus obtaining CFS-CM1 and CFS-CM2.

Antifungal activity of the cell-free supernatants

CFS-CM1 and CFS-CM2 were checked for antifungal activity using the agar well diffusion method against SS, MP, and BC (prepared as described in Subheading - *Antagonistic activity*). For each fungus, 6 mm agar plugs containing 7-day-old fungal mycelia were placed at the center of 90 mm Petri dishes. Four symmetrically distributed wells were made approximately 25 mm from the fungal mycelia plugs. A total of 200 μ L of CFS-CM1 and CFS-CM2 were added to the wells, separately per Petri dish (4 experimental units with 4 wells). Sterile CM1 and CM2 were used as blanks. The plates were incubated at 25 °C with a photoperiod of 12/12 h for 96 h. The experiment was repeated twice. The results were submitted to analysis of variance (ANOVA) and the means compared by the Tukey test (p<0.05). SEM was performed to visualize the antifungal activity of the cell-free supernatants (CFS) of the best culture medium identified.

Ultrastructural damage evaluated by scanning electron microscopy

SEM was carried out to visualize the ultrastructural damage against SS of the CFS obtained from the best culture medium identified. In order to observe the ultrastructural damage, an evaluation was carried out at the inhibition zone-SS mycelial growth interface. The control was treated with sterile culture medium, and the assay was conducted in accordance with the method previously described. Plugs measuring 6 mm in diameter were cut and placed in vials containing 2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M sodium cacodylate buffer (pH 7.2) at 4 °C. The samples were kept in this solution overnight for fixation, after which they were washed three times with 0.1 M sodium cacodylate buffer (pH 7.2) for 10 min. Subsequently, the samples were dehydrated three times in an ethanol series (30, 50, 70, 90, and 100%) for 10 min. The samples were submitted to critical point drying with CO₂ (BALTEC CPD 030 Critical Point Drier), coated with gold (BALTEC SDC 050 Sputter Coater), and visualized with an FEI Quanta 200 scanning electron microscope (SEM) operating at 25.0 kV.

Growth curve and antifungal metabolite production at different incubation times

The inoculum preparation methodology used to produce antifungal metabolites and obtain the CFS was carried out as previously describedusing the best culture medium. The samples were removed at different time intervals (0, 12, 24, 36, 48, 60, and 72 h) to determine antifungal activity and bacterial growth. The antifungal activity of the CFS was assessed using the method previously described. Bacterial growth was evaluated according to cell count (CFU/mL) from 10-fold serial dilutions in sterilized saline solution (0.85% sodium chloride, w/v) plated on MYP agar (Neogen Corporation, USA) and incubated at 28 °C for 24 h. This assay was repeated twice. Pearson's correlation coefficient was used to investigate the correlations between growth and antifungal activity. Statistical significance was considered when p<0.05.

Evaluation of stability of the antifungal metabolites at different temperatures, pH, and light wavelengths

In order to evaluate the stability of the antifungal metabolites at different temperatures, pH, and light wavelengths, aliquots of CFS from the best culture medium were used. The samples were exposed to constant temperatures of 28 °C, 70 °C, and 100 °C for 30 min, and 121 °C for 15 min. Subsequently, the samples treated at 28 °C, 70 °C, and 100 °C were sterilized by filtration through a cellulose filter with a pore size of 0.22 µm (Millipore, USA), and antifungal activity was evaluated according to the method previously described. The experiment was performed three times, with eight replications each, and linear regression analysis (p<0.05) was conducted to represent the data. For the pH stability assay, the samples were adjusted to various pH values in the range from 3.0 to 11.0 using 2.0 M HCl or 2.0 M NaOH and maintained at 4 °C for 24 h. Afterward, the samples were readjusted to pH 7.0¹³, sterilized by filtration through a 0.22 µm cellulose filter (Millipore, USA), and submitted to the antifungal activity assay previously described. The evaluation was carried out in two independent experiments, with 4 replicates each, and was based on descriptive analysis. As for the determination of the effect of ultraviolet radiation (365 nm) and white light, the samples were exposed to the two for 12 h at a distance of 15 cm. The untreated CFS was used as a control. After that, all samples were sterilized by filtration through a 0.22 µm cellulose filter (Millipore, Bedford, MA), and their antifungal activity was assessed according to the method previously described. Six repetitions were performed in two independent experiments. The assay was later evaluated using the Kruskal-Wallis nonparametric test (p<0.05).

B. velezensis CMRP 4489 inoculation on soybean seeds: colonization assays by SEM

Seeds of the soybean cultivar 6461RSF IPRO were disinfected with 2% sodium hypochlorite for 2 min and then washed 3 times with sterile water and dried in a biological safety chamber for 20 min. After disinfection, one hundred soybean seeds were treated, in 125 mL Erlenmeyer flasks, with strain CMRP 4489 cultivated in CM2 and adjusted to 5 x 10⁹ CFU/mL (dose of 200 mL/100 kg). The control seeds were treated with CM2 (200 mL/100 kg) alone. After manually homogenizing the flasks for 2 minutes, 10 seeds were transferred to a 150 mm Petri dish containing a filter paper moistened with sterile deionized water. In order to evaluate cell adhesion on day 0 and compare it with the uninoculated seeds, 3 of both inoculated and uninoculated seeds were fixed for further evaluation by SEM. The remaining seeds were incubated in a humid chamber for 7 days at 23°C with a photoperiod of 12/12 h and, every 24 h, moistened with approximately 5 mL of sterile deionized water. After incubation, the tegument and radicle of 3 seven-day-old germinated seeds were fixed for SEM. The procedures used in the SEM assay were conducted according to the method previously described.

B. velezensis CMRP 4489 inoculation on soybean seeds: suppression of Sclerotinia sclerotiorum

Seeds of the soybean cultivar Monsoy 6410 were disinfected according to the method previously described. After disinfection, five hundred soybean seeds were inoculated with *Sclerotinia sclerotiorum* (SS) for each treatment. To this end, SS was cultivated in PDA culture medium and, after 5 days of growth (100% of growth on 150 mm Petri dishes), the seeds were distributed onto the Petri dishes,

maintaining contact with the fungal culture for 24 h. The treatments were as follows: (T1) control, with five hundred uninfected and untreated soybean seeds; (T2) control, with five hundred soybean seeds infected with SS without treatment; (T3) control, with five hundred soybean seeds treated with sterile CM2 (200 mL/100 kg); (T4) chemical treatment, with five hundred soybean seeds treated with a commercial product composed of 52.50 g/L fluazinam + 350.00 g/L thiophanate-methyl (200 mL/100 kg); (T5) biocontrol agent, with five hundred soybean seeds treated with strain CMRP 4489 cultivated in CM2 and adjusted to 5 x 10⁹ CFU/mL (200 mL/100 kg). All treatments were carried out in plastic bags and homogenized by manual shaking for 2 minutes. A layer of sterile filter paper was placed within 150 mm Petri dishes and moistened with sterile distilled water. Subsequently, 13 Petri dishes per treatment/control received 25 seeds and were incubated in a humid chamber at 23 °C, with a photoperiod of 12/12 h. The percentage of germinated seeds and uninfected seeds (absence of SS growth) was evaluated after 7 days of incubation. For the statistical analysis, the data were subjected to analysis of variance (ANOVA), considering a generalized linear model with binomial distribution. When significant, the treatments were compared using the Tukey test (p<0.05). All analyses were processed using the R software (R Core Team, 2019).

Genome mining: prediction of biosynthetic gene clusters and comparative analysis of genes involved in biofilm formation

The prediction of secondary metabolite clusters in the *B. velezensis* CMRP 4489 genome was carried out using the antiSMASH 6.0 webserver ¹⁴. For the comparative analysis of genes involved in biofilm formation and swarming, a database was created using gene sequences associated with swarming and biofilm formation/regulation, which were selected using data collected on the Subtwiki database ¹⁵ and from studies related to root colonization ^{16–21} by *Bacillus subtilis*. This developed database was used to compare similar sequences to the CMRP 4489 genome using the BLASTN program.

Data availability

The genome analyzed during the current study is available in the DDBJ/EMBL/GenBank under the accession number CP023748 (BioProject PRJNA412668, BioSample SAMN07722662).

Results

Phenotypic characterization and colony architectureof *B. velezensis* CMRP 4489

Gram staining showed that CMRP 4489 is a Gram-positive, rod-shaped bacterium that produces endospores. Its rod-shaped morphology and cell association were evidenced by SEM (Supplementary Figure 1).

Floating pellicle biofilm and swarm expansion

The *in vitro* assays evidenced the filming and swarming capacity of the CMRP 4489 strain. Microplate assays revealed a thick film over the entire air-liquid surface, formed after cell growth. In the swarming motility test, growth over the entire surface of the Petri dish was observed after overnight culture. These results indicate that this strain is capable of rhizospheric colonization, an important factor for the establishment of the plant-bacteria interaction (Supplementary Figure 2).

Dual-culture assay

The CMRP 4489 strain showed excellent antagonistic activity against SS, MP, and BC. In the experimental group, CMRP 4489 formed a very clear inhibition zone between the bacterial suspension and the pathogenic fungi (Figure 1a). The mean inhibitory zone was measured, and the percentage of mycelial growth inhibition (MGI%) was calculated. The CMRP 4489 strain inhibited all the fungi, with 55.8%, 64.0%, and 66.6% inhibition of SS, MP, and BC, respectively (Figure 1b).

Antifungal activity of the cell-free supernatant

CFS-MC1 and CFS-MC2 showed excellent antifungal activity against the pathogenic fungi (Figure 2a). When analyzing the effect of CFS-MC1 and CFS-MC2, it was observed that CFS-MC1 exhibited less antifungal activity, with a significant difference compared to CFS-MC2 (Table 1). The inhibitory values presented by CFS-MC1 reached up to 54.36% for SS, 57.27% for MP, and 36.76% for BC, with significant differences when comparing the three fungi. As for CFS-MC2, the inhibitory values reached 65.31% for SS, 68.31% for MP, and 72.87% for BC, with significant differences when comparing the three fungi (Table 1).

Table 1: Antifungal activity against *Sclerotinia sclerotiorum* (SS), *Macrophomina phaseolina* (MP) and *Botrytis cinerea* (BC) of the cell-free supernatants (CFS) obtained by submerged liquid fermentation of *Bacillus velezensis* CMRP 4489 in two culture media (CM1 and CM2).

Plant-pathogenic fungi	MGI (%)			
	CFS-CM1		CFS-CM2	
Sclerotinia sclerotiorum (SS)	54.36	bA	65.31	сВ
Macrophomina phaseolina (MP)	57.27	aA	68.31	bB
<i>Botrytis cinerea</i> (BC)	36.76	cA	72.87	aB

Numbers followed by the same lowercase letter in the column and uppercase letter in the row did not differ according to the Tukey test (p<0.05).

MGI: mycelial growth inhibition

CFS: cell-free supernatants

CM1: culture media (g / L): tryptone 10.0; yeast extract 5.0; NaCl 5.0; pH 7.1.

CM2: culture media (g / L): glucose 20.0; tryptone 12.4; NaCl 5.0; $K_2HPO_4 \cdot 3H_2O 1.5$; MnSO₄ · H₂O, 0.04; FeSO₄ · 7H₂O, 1.67; MgCl₂ · 6H₂O, 1.22; pH 7.1.

Ultrastructural damage of Sclerotinia sclerotiorum

Considering that the CM2 medium showed better performance in producing the antifungal metabolites of the CMRP 4489 strain, it was selected for all of the following assays. The effect of CFS-MC2 on SS growth was substantial. The mycelia of SS in the control presented hyphae with a typical "net" structure and a smooth surface. In the presence of CFS-MC2, the hyphae lost their smoothness and formed unusual surface bulges, indicating that CFS-MC2 inhibited SS growth by causing deformation of the hyphal structure. The obtained results showed that CFS-MC2 distorted and damaged SS hyphae and that fungal growth was inhibited (Figure 2b).

Growth curve of the CMRP 4489 strain and the production of antifungal metabolites in MC2

The kinetic data on cell growth and antifungal activity are shown in Figure 3. The culture of the *B. velezensis* CMRP 4489 strain was carried out at 28°C in CM2 medium for 72 h. Accordingly, two phases could be distinguished during the fermentation process: a growth phase, until 36 h of incubation, followed by a stationary phase, which did not decline until 72 h of cultivation. No lag phase was observed. The production of the antifungal metabolites was dependent on the bacterial growth phase. The CMRP 4489 strain started producing antifungal metabolites at 12 h for BC and MP during the logarithmic growth phase. The strongest antifungal activity, with approximately 70% of inhibition for all fungi and 7×10^9 CFU mL⁻¹ of viable cells, was obtained at 36 h after incubation, which corresponded to the stationary growth phase, and did not decline over the next 36 h. Therefore, under the culture conditions used in this study, the optimal production time of antifungal metabolites for strain CMRP 4489 was 36 h after inoculation, when the viable bacterial biomass was the highest (Figure 3).

Effect of temperature, pH, and light on the CMRP 4489 supernatant

The supernatant continued active after being subjected to high temperatures. However, its activity was lower after autoclaving, being more evident against BC, with an MGI of 73, 65, 64, and 60%; SS, with an MGI of 65, 63, 63, and 62%; and MP, of 68, 62, 64, and 63% at 28°C, 70°C, 100°C, and 121°C, respectively (Figure 4a and 4b). As for the stability tests regarding pH variation, the supernatant presented high stability, without significant changes in activity against all the tested fungi (Figure 4c). Similar results were found in supernatant stability when exposed to UV 365 nm and visible light (Figure 4d).

Colonization of *B. velezensis* CMRP 4489on soybean seeds

Strain CMRP 4489 was able to attach and multiply on the soybean seed surface. In Figure 5, the first column shows untreated seeds, while the second shows the seeds treated with strain CMRP 4489, adjusted to 5 x 10⁹ CFU mL⁻¹ (200 mL 100 kg⁻¹). The first row shows seeds that were fixated on day 0 of treatment; the second row shows the seeds after 7 days of incubation, and the third row shows the radicles of the seeds after 7 days of incubation. All images have inserts at 50x magnification. The control seeds treated with CM2 on day 0 exhibited an absence of adhered microorganisms on the surface (Figure 5a). Seeds treated with CMRP 4489 on day 0 presented several endospores attached to their surface (Figure 5b). After 7 days of incubation, the control seeds (Figure 5c) and radicles (Figure 5e) were not colonized with any type of microorganism. However, those treated with strain CMRP 4489 after 7 days of incubation showed surfaces fully colonized with a net-shaped biofilm (Figure 5d). In the radicles of the seeds treated with strain CMRP 4489, after 7 days of incubation, significant colonization was observed, with net-shaped biofilm and exopolysaccharides (Figure 5f). These results evidence the high potential of CMRP 4489 to colonize soybean roots. The strain transitioned from the seeds to nearby structures, being the first bacterial colonizer of the root system and potentially protective against fungal soil diseases.

Control of *Sclerotinia sclerotiorum* on soybean seeds by inoculation of *B. velezensis* CMRP 4489

In the evaluation of SS suppression in the *in vitro* infected soybean seeds, the results evidenced the high control potential of the CMRP 4489 strain, in addition to not presenting a negative influence on germination. The tested prototype was prepared with 5×10^9 CFU mL⁻¹ and antifungal extracellular metabolites produced during the submerged liquid fermentation stage (Figure 6). The uninfected and untreated control seeds (T1) showed a germination rate of approximately 100% (Figure 6a), and none of them exhibited SS colonization (Figure 6b). Meanwhile, the untreated control seeds infected with SS (T2) presented a germination rate of almost 15% (Figure 6a); however, approximately 95% of the seeds died since they were fully colonized by SS (Figure 6b). The control seeds infected with the fungus and treated only with sterilized CM2 (T3) showed no significant differences compared to T2 (Figures 6a and 6b). On the other hand, the soybean seeds treated with the commercial product (T4) composed of 52.50 g L⁻¹ of fluazinam and 350.00 g L⁻¹ of thiophanate-methyl had a germination rate of roughly 80% (Figure 6a), and the chemical compound suppressed the development of SS. Treatment T4 revealed only around 16% of seeds killed by SS colonization (Figure 6b). The results showed significant differences when comparing the treated groups to the controls. However, the soybean seeds treated with strain CMRP 4489 presented a germination rate of approximately 95% (Figure 6a), differing significantly from the controls and the

chemical treatment (T4). When evaluating the potential to suppress the development of SS in the *in vitro* infected soybean seeds, the CMRP 4489 strain (T5) showed high control performance, differing significantly from the controls and the chemical treatment (Figure 6b). Strain CMRP 4489 suppressed around 95% of SS colonization. This treatment showed that only approximately 5% of the seeds were killed by SS colonization (Figure 6b). These results are directly related to the extracellular antifungal metabolites produced by the CMRP 4489 strain during submerged liquid fermentation and that were present in the prototype used in the assay, corroborating that, in fermentation processes used specifically to produce antimicrobial metabolites, it is possible to elaborate biological products for seed treatment with high potential for controlling this soil-borne fungal pathogen. Furthermore, due to the high colonization potential of the CMRP 4489 strain, as observed in Subheading 3.8, it probably also contributes to the control of SS.

Genome mining: prediction of secondary metabolite clusters and comparative analysis of genes involved in biofilm formation

The antiSMASH 6.0 webserver found 12 BGCs in the genome of the *B. velezensis* CMRP 4489 strain, seven of which presented significant similarity with previously identified clusters in the Minimum Information about a Biosynthetic Gene cluster (MIBiG) repository that are involved in the synthesis of surfactin (86% similarity to known clusters by antiSMASH), macrolactin (100%), bacillaene (100%), fengycin (100%), difficidin (100%), bacillibactin (100%), and bacilysin (100%). The similarity between the clusters of the CMRP 4489 strain and the Bacillus velezensis FZB42 strain ²²⁻²⁴ is noteworthy and considered a model for PGPR and biocontrol strains. The clusters were also represented in a circular genome image (Figure 7) for comparison with other strains of the *Bacillus subtilis* group. Four out of 13 clusters did not show similarity with any of the clusters in the antiSMASH database. Several genes associated with rhizosphere colonization and biofilm production were found in the CMRP 4489 genome (Supplementary Table 1). Key genes related to the root colonization process were found, including: genes regulating the expression of exopolysaccharides sinR (80% similarity with the genes used for comparison) and *sinL* (97%); the operon responsible for the production of exopolysaccharides *epsA-O* (~99%); operon *sfrAA-AC* (79%, 75%, 87%), related to surfactin synthesis; operon *swrA-C* (84%, 82%, 80%), important for swarming motility and flagella production, and operon *vgxM-sipW-tasA* (~99%), associated with the production of biofilm matrix components. Other genes related to root colonization were also found in the genome of the CMRP 4489 strain (Supplementary Table 1).

Discussion

Pesticide use is the most diffused strategy to avoid crop losses caused by plant diseases. However, some aspects, such as pathogen resistance ^{25,26}, environmental contamination ^{27,28}, and damage to human health ²⁹, caused by the excessive and incorrect use of these chemical defensives, have fostered a growing interest in bio-based products.

The use of microorganisms for biological product development is a promising alternative that has gained significant attention over the past decades. According to Van Lenteren *et al.* (2018) ³⁰, over 200 microbial strains are currently being used as commercial products in Brazil, Australia, Canada, the European Union, Japan, New Zealand, and the United States of America.

In the present study, *B. velezensis* CMRP 4489 showed potential antagonism against the analyzed phytopathogenic fungi, as observed in the dual-culture assay. However, for use as a bioproduct, it is necessary to identify the best fermentation system for mass production. Thus, cultivation conditions are important factors that must also be considered in the production of secondary metabolites ³¹. In addition to bacterial biomass, antimicrobial compounds need to be present in large quantities in the fermented product. Therefore, many studies have been developed to formulate more suitable culture media for secondary metabolite production ^{32,33}. In this study, we evaluated two culture media for liquid fermentation. Among the basal media studied, the MC2 medium, supplemented with micronutrients, provided more adequate conditions for strain CMRP 4489 to produce antifungal compounds. In a study on the optimization of multiple responses of biomass and antifungal metabolites with *Bacillus subtilis* EA-CB0015, Mosquera et al. (2014) ³⁴ demonstrated the importance of formulating more suitable culture media to produce high concentrations of biomass and antifungal molecules for biotechnological application. In some cases, depending on the microorganism studied, a simple medium can also provide adequate levels of production of these compounds, as observed with the MC1 medium, and as reported by Zouari et al. (2016) ³⁵ and Torres et al. (2016) ³⁶ when using LB and Mueller-Hinton culture media, respectively.

Aside from nutrient sources, microbial growth also influences the synthesis of biomolecules since molecules from secondary metabolism are more expressed in the stationary phase, as observed in the present study and in Mezghanni *et al.* (2012) ³⁷. When comparing the growth curve and antifungal activity of the CMRP 4489 strain under the culture conditions used in this study, the optimal production time of antifungal metabolites, in which the highest viable biomass of strain CMRP 4489 was observed, was 36 hours after inoculation.

Considering that CFS-MC2 presented better performance, it was chosen to evaluate the ultrastructural damage caused to SS. The results showed that CFS-MC2 distorted and damaged SS hyphae and that fungal growth was inhibited, corroborating the results obtained by Farzand *et al.* (2019)³⁸ and Zhang L and Sun (2018) ³⁹, who analyzed the effect of lipopeptides produced by *Bacillus* species on ultrastructural changes in fungal hyphae. In another study, the shrinkage of *Lasiodiplodia theobromae* hyphae due to antifungal lipopeptides produced by *B. subtilis* B1 was reported, which is also consistent with our findings ⁴⁰

Among the lipopeptides produced by species of the genus *Bacillus* with fungicidal effect is fengycin. In several reports, fengycin has been shown to cause severe damage to fungal hyphae, resulting in their death ^{41,42}. Considering that in the present study 12 BGCs were identified in the genome of the *B*.

velezensis CMRP 4489 strain and that some of them are involved in the synthesis of lipopeptides, including fengycin, this may explain the possible reason behind the effects observed in the hyphae structure of SS herein. However, further studies with the CMRP 4489 strain are underway to identify the antifungal molecules produced by this strain.

Regarding of BGCs found in the *Bacillus* spp. genome, NRPSs and PKSs were the most common ⁴³ and were also observed in the genome of *B. velezensis* CMRP 4489. These and other BGCs are responsible for the production of several secondary metabolites of biotechnological interest, being characterized according to their chemical composition as peptides (bacteriocins, bacilysin), lipopeptides (surfactin, iturin, fengycin), and polyketides (difficidin, bacillaene, macrolactin), among others ⁴⁴

The CMRP 4489 strain has several clusters that have not yet been identified among those already described, some of which share similarities with bacterial isolates with biotechnological potential, as is the case of bacillibactin, macrolactin, surfactin, and bacilysin, present in the QST713 strain of the commercial product Serenade® ⁴⁵, and in strain FZB42 of the commercial product RhizoVital® ^{46,47}. This indicates that CMRP 4489 also has significant biotech potential and could be part of the composition of a commercial product in the future. Among the clusters that are not shared by reference strains are thiopeptides. These peptides, which are found in the genome of several bacteria, synthesized by ribosomes and modified after translation ⁴⁸, may be related to biofilm formation and antibacterial ⁴⁹ and antifungal ⁵⁰ activity, among others. According to Just Baringo *et al.* (2014) ⁵¹, there are more than one hundred families in this cluster.

Other crucial parameters that were considered in this study were related to the stability of the antifungal effect under adverse conditions. These stability attributes are important for the future use of this strain in bioproducts for plant disease control. The results regarding thermostability showed that there was a decrease in activity as the temperature increased; however, the analyzed compounds did not lose their activity. As for pH stability, there was little variation in activity in the tested pH range. Similarly, Yu *et al.* (2002) ⁵² also observed stability in the compounds produced by a strain of *Bacillus* sp. However, while evaluating a strain of *Bacillus* sp., Zhao *et al.* (2010) ⁵³, reported a significant reduction in activity when the compounds were tested regarding temperature and pH. The photostability tests showed that 12-hour exposure to visible light and UV 365 nm did not interfere in the antifungal activity of the supernatant. This was analogous to the compounds of *B. amyloliquefaciens* strain CNU114001, which were also stable under UV light exposure for 30 minutes ⁵⁴

All tests performed herein were essential for the characterization of the supernatants' antifungal activity, and their effect was evaluated based on morphological changes in SS hyphae, showing the potential of the CMRP 4489 strain as a biological control agent. However, another important factor for use as a biocontrol agent is its ability to be an excellent colonizer. That is why we evaluated the colonization of *B. velezensis* CMRP 4489 on soybean seeds and the genes involved in the formation and regulation of colonization mechanisms.

In order for the colonization and formation of microbial communities to take place in the root structures of a plant, it is necessary that the microorganisms have the ability to move around and form cell clusters around the roots. Several genes directly or indirectly associated with colonization were found in the genome of the CMRP 4489 strain, highlighting the presence of *srfAA-AC*, genes involved in the production of surfactin that reduce the surface tension of the matrix and allow greater mobility, and *swrA-C*, genes involved in regulating the expression of flagellins and in improving the swarming capacity of the strain, which is one of the main mechanisms of radicular colonization ⁵⁵. In this study, swarming was also evaluated and evidenced in the CMRP 4489 strain in the *in vitro* assay.

Molecular data from *in vitro* assays regarding biofilm formation and swarming motility suggest that genes associated with biofilm formation and regulation and motility are functional and may confer the capacity of rhizosphere colonization to the strain ^{56,57}. Electron microscopy shows that seed treatment is a viable strategy for using this strain, considering the colonization capacity of CMRP 4489 in seeds and radicles after germination and its antagonistic activity against phytopathogenic fungi. Moreover, its beneficial effects on seed germination make the direct seed coating strategy a possible alternative for its use.

Conclusions

In the recent years the augmented burden on farmers for increased yield has resulted in excess usage of fertilizers and chemical pesticides. Use of chemical fertilizers and antifungal agents to increase production is not sustainable for obvious reasons. The present study suggested the use of biocontrol agent *B. velezensis* CMRP 4489 like as a new alternative for bioformulation for the being used in scale in bio-based industries. The *in vitro* and soybean seeds data in present study has been complimented with the genomic data of the *B. velezensis* CMRP 4489, with mapped on genome of secondary metabolite clusters and comparative analysis of genes involved in biofilm formation, indicating that the biological control agent properties strain CMRP 4489 are genetic and hence stable.

Declarations

Funding: A.G. de Oliveira was supported by a research fellowship from CNPq, Grant No. 315593/2018-0.

Acknowledgments: We acknowledge the financial support from the Brazilian National Council of Scientific and Technological Development (CNPq), the Araucaria Foundation for Support to Scientific and Technological Development of the State of Paraná (FAPPR), and the Simbiose-Agro Company. The authors would also like to thank the Laboratory for Electron Microscopy and Microanalysis – LMEM/UEL, Osvaldo Capello, for his help in carrying out the SEM, and Eurico de Paula Arruda, who provided English editing of the manuscript.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- Umesha, S., Singh, P. K. & Singh, R. P. Microbial biotechnology and sustainable agriculture. in Biotechnology for Sustainable Agriculture: Emerging Approaches and Strategies 185–205 (Elsevier, 2017).
- 2. Berg, G., Zachow, C., Müller, H., Philipps, J. & Tilcher, R. Next-generation bio-products sowing the seeds of success for sustainable agriculture. *Agronomy* vol. 3 648–656 (2013).
- 3. Barea, J. M. *Future challenges and perspectives for applying microbial biotechnology in sustainable agriculture based on a better understanding of plant-microbiome interactions*. Journal of Soil Science and Plant Nutrition vol. 15 (2015).
- Wohlgemuth, R., Twardowski, T. & Aguilar, A. Bioeconomy moving forward step by step A global journey. New Biotechnology 61, 22–28 (2021).
- 5. Perez, K. J. *et al. Bacillus* spp. isolated from puba as a source of biosurfactants and antimicrobial lipopeptides. Frontiers in Microbiology **8**, (2017).
- 6. Fritze, D. Taxonomy of the Genus *Bacillus* and Related Genera: The Aerobic Endospore-Forming Bacteria. Phytopathology **94**, (2004).
- 7. Meena, K. R., Tandon, T., Sharma, A. & Kanwar, S. S. Lipopeptide antibiotic production by Bacillus velezensis KLP2016. Journal of Applied Pharmaceutical Science **8**, 91–98 (2018).
- 8. Xu, T., Zhu, T. & Li, S. β-1,3 1,4-glucanase gene from *Bacillus velezensis* ZJ20 exerts antifungal effect on plant pathogenic fungi. World Journal of Microbiology and Biotechnology **32**, (2016).
- 9. Lim, S. M. *et al.* Diffusible and volatile antifungal compounds produced by an antagonistic *Bacillus velezensis* G341 against various phytopathogenic fungi. Plant Pathology Journal **33**, 488–498 (2017).
- 10. Fan, B. *et al. Bacillus velezensis* FZB42 in 2018: The Gram-Positive Model Strain for Plant Growth Promotion and Biocontrol. Frontiers in Microbiology **9**, (2018).
- 11. Pezarini Baptista, J. *et al.* Complete Genome Sequence of *Bacillus velezensis* LABIM40, an Effective Antagonist of Fungal Plant Pathogens. *Genome Announcements* (2018).
- Yahyazadeh, M., Omidbaigi, R., Zare, R. & Taheri, H. Effect of some essential oils on mycelial growth of *Penicillium digitatum* Sacc. World Journal of Microbiology and Biotechnology 24, 1445–1450 (2008).
- 13. Ji, S. H. *et al.* Biocontrol activity of *Bacillus amyloliquefaciens* CNU114001 against fungal plant diseases. Mycobiology **41**, 234–242 (2013).
- 14. Weber, T. *et al.* AntiSMASH 3.0-A comprehensive resource for the genome mining of biosynthetic gene clusters. Nucleic Acids Research **43**, W237–W243 (2015).
- 15. Pedreira, T., Elfmann, C. & Stülke, J. The current state of Subti Wiki, the database for the model organism *Bacillus subtilis*. Nucleic Acids Research (2021)
- 16. Nagórska, K. *et al.* Importance of eps genes from *Bacillus subtilis* in biofilm formation and swarming. Journal of Applied Genetics **51**, 369–381 (2010).

- 17. Salvetti, S., Faegri, K., Ghelardi, E., Kolstø, A. B. & Senesi, S. Global gene expression profile for swarming *Bacillus cereus* bacteria. Applied and Environmental Microbiology **77**, 5149–5156 (2011).
- Chu, F. *et al.* A novel regulatory protein governing biofilm formation in *Bacillus subtilis*. Molecular Microbiology **68**, 1117–1127 (2008).
- 19. Kearns, D. B. & Losick, R. Swarming motility in undomesticated *Bacillus subtilis*. Molecular Microbiology **49**, 581–590 (2003).
- Kearns, D. B., Chu, F., Rudner, R. & Losick, R. Genes governing swarming in *Bacillus subtilis* and evidence for a phase variation mechanism controlling surface motility. Molecular Microbiology 52, 357–369 (2004).
- 21. Penha, R. O., Vandenberghe, L. P. S., Faulds, C., Soccol, V. T. & Soccol, C. R. *Bacillus* lipopeptides as powerful pest control agents for a more sustainable and healthy agriculture: recent studies and innovations. *Planta* vol. 251 (2020).
- Koumoutsi, A. *et al.* Structural and Functional Characterization of Gene Clusters Directing Nonribosomal Synthesis of Bioactive Cyclic Lipopeptides in *Bacillus amyloliquefaciens* strain FZB42. in Journal of Bacteriology vol. 186 1084–1096 (2004).
- 23. Wu, L. *et al.* Difficidin and bacilysin from *Bacillus amyloliquefaciens* FZB42 have antibacterial activity against *Xanthomonas oryzae* rice pathogens. Scientific Reports **5**, (2015).
- 24. Chen, X. H. *et al.* Structural and functional characterization of three polyketide synthase gene clusters in *Bacillus amyloliquefaciens* FZB42. Journal of Bacteriology **188**, 4024–4036 (2006).
- 25. Lucas, J. A., Hawkins, N. J. & Fraaije, B. A. The Evolution of Fungicide Resistance. Advances in Applied Microbiology **90**, 29–92 (2015).
- Geep, V. *et al.* Resistencia a fungicidas en *Botrytis cinerea* en el Uruguay. Agrociencia Uruguay 16, 97–107 (2012).
- 27. Arias-Estévez, M. *et al.* The mobility and degradation of pesticides in soils and the pollution of groundwater resources. Agriculture, Ecosystems and Environment vol. 123 247–260 (2008).
- Tosi, S., Costa, C., Vesco, U., Quaglia, G. & Guido, G. A 3-year survey of Italian honeybee-collected pollen reveals widespread contamination by agricultural pesticides. Science of the Total Environment 615, 208–218 (2018).
- 29. Cremonese, C. *et al.* Occupational exposure to pesticides, reproductive hormone levels and sperm quality in young Brazilian men. Reproductive Toxicology **67**, 174–185 (2017).
- 30. van Lenteren, J. C., Bolckmans, K., Köhl, J., Ravensberg, W. J. & Urbaneja, A. Biological control using invertebrates and microorganisms: plenty of new opportunities. BioControl **63**, 39–59 (2018).
- 31. Bertrand, S. *et al.* Metabolite induction via microorganism co-culture: A potential way to enhance chemical diversity for drug discovery. Biotechnology Advances vol. 32 1180–1204 (2014).
- 32. Gu, X. B. *et al.* Optimization of medium constituents for a novel lipopeptide production by *Bacillus subtilis* MO-01 by a response surface method. Process Biochemistry **40**, 3196–3201 (2005).

- Shafi, J. *et al.* Optimization of *Bacillus aerius* strain JS-786 cell dry mass and its antifungal activity against *Botrytis cinerea* using response surface methodology. Archives of Biological Sciences 69, 469–480 (2017).
- Mosquera, S., González-Jaramillo, L. M., Orduz, S. & Villegas-Escobar, V. Multiple response optimization of *Bacillus subtilis* EA-CB0015 culture and identification of antifungal metabolites. Biocatalysis and Agricultural Biotechnology **3**, 378–385 (2014).
- 35. Zouari, I., Jlaiel, L., Tounsi, S. & Trigui, M. Biocontrol activity of the endophytic *Bacillus amyloliquefaciens* strain CEIZ-11 against *Pythium aphanidermatum* and purification of its bioactive compounds. Biological Control **100**, 54–62 (2016).
- 36. Torres, M. J., Brandan, C. P., Petroselli, G., Erra-Balsells, R. & Audisio, M. C. Antagonistic effects of *Bacillus subtilis* subsp. *subtilis* and *B. amyloliquefaciens* against *Macrophomina phaseolina*: SEM study of fungal changes and UV-MALDI-TOF MS analysis of their bioactive compounds. *Microbiological Research* 182, 31–39 (2016).
- Mezghanni, H., Khedher, S. ben, Tounsi, S. & Zouari, N. Medium optimization of antifungal activity production by *Bacillus amyloliquefaciens* using statistical experimental design. Preparative Biochemistry and Biotechnology 42, 267–278 (2012).
- 38. Farzand, A. *et al.* Suppression of *Sclerotinia sclerotiorum* by the Induction of Systemic Resistance and Regulation of Antioxidant Pathways in Tomato Using Fengycin Produced by *Bacillus amyloliquefaciens* FZB42. Biomolecules **9**, 613 (2019).
- 39. Zhang, L. & Sun, C. Fengycins, Cyclic Lipopeptides from marine *Bacillus subtilis* strains, kill the plantpathogenic fungus *Magnaporthe grisea* by inducing reactive oxygen species production and chromatin condensation. Applied and Environmental Microbiology **84**, (2018).
- Sajitha, K. L. & Dev, S. A. Quantification of antifungal lipopeptide gene expression levels in *Bacillus subtilis* B1 during antagonism against sapstain fungus on rubberwood. Biological Control 96, 78–85 (2016).
- Liu, J. *et al.* Functions of lipopeptides bacillomycin d and fengycin in antagonism of *Bacillus amyloliquefaciens* C06 towards *Monilinia fructicola*. Journal of Molecular Microbiology and Biotechnology 20, 43–52 (2011).
- 42. Tao, Y., Bie, X., Lv, F., Zhao, H. & Lu, Z. Antifungal activity and mechanism of fengycin in the presence and absence of commercial surfactin against *Rhizopus stolonifer*. The Journal of Microbiology **49**, 146–150 (2011).
- 43. Grubbs, K. J. *et al.* Large-Scale Bioinformatics Analysis of *Bacillus* Genomes Uncovers Conserved Roles of Natural Products in Bacterial Physiology. *mSystems* **2**, (2017).
- 44. Sansinenea, E. & Ortiz, A. Secondary metabolites of soil *Bacillus* spp. Biotechnology Letters vol. 33 1523–1538 (2011).
- 45. Pandin, C. *et al.* Complete genome sequence of *Bacillus velezensis* QST713: A biocontrol agent that protects *Agaricus bisporus* crops against the green mould disease. Journal of Biotechnology **278**, 10−19 (2018).

- Chen, X. H. *et al.* Genome analysis of *Bacillus amyloliquefaciens* FZB42 reveals its potential for biocontrol of plant pathogens. Journal of Biotechnology **140**, 27–37 (2009).
- 47. Fan, B., Blom, J., Klenk, H. P. & Borriss, R. Bacillus *amyloliquefaciens*, *Bacillus velezensis*, and *Bacillus siamensis* form an "Operational Group B. *amyloliquefaciens*" within the *B. subtilis* species complex. *Frontiers in Microbiology* 8, (2017).
- 48. Bogart, J. W. & Bowers, A. A. Thiopeptide Pyridine Synthase TbtD Catalyzes an Intermolecular Formal Aza-Diels-Alder Reaction. Journal of the American Chemical Society **141**, 1842–1846 (2019).
- 49. Bleich, R., Watrous, J. D., Dorrestein, P. C., Bowers, A. A. & Shank, E. A. Thiopeptide antibiotics stimulate biofilm formation in *Bacillus subtilis*. Proceedings of the National Academy of Sciences of the United States of America **112**, 3086–3091 (2015).
- 50. Mizuhara, N. *et al.* Antifungal thiopeptide cyclothiazomycin B1 exhibits growth inhibition accompanying morphological changes via binding to fungal cell wall chitin. Bioorganic and Medicinal Chemistry **19**, 5300–5310 (2011).
- 51. Just-Baringo, X., Albericio, F. & Álvarez, M. Thiopeptide antibiotics: Retrospective and recent advances. Marine Drugs vol. 12 317–351 (2014).
- 52. Yu, G. Y., Sinclair, J. B., Hartman, G. L. & Bertagnolli, B. L. Production of iturin A by *Bacillus amyloliquefaciens* suppressing *Rhizoctonia solani*. Soil Biology & Biochemistry **34**, 955–963 (2002).
- 53. Zhao, Z. *et al.* Study of the antifungal activity of *Bacillus vallismortis* ZZ185 *in vitro* and identification of its antifungal components. Bioresource Technology **101**, 292–297 (2010).
- 54. Liu, C. H. *et al.* Study of the antifungal activity of *Acinetobacter baumannii* LCH001 *in vitro* and identification of its antifungal components. Applied Microbiology and Biotechnology **76**, 459–466 (2007).
- 55. Ghelardi, E. *et al.* Contribution of surfactin and SwrA to flagellin expression, swimming, and surface motility in *Bacillus subtilis*. Applied and Environmental Microbiology **78**, 6540–6544 (2012).
- 56. Ansari, F. A. & Ahmad, I. Biofilm development, plant growth promoting traits and rhizosphere colonization by *Pseudomonas entomophila* FAP1: a promising PGPR. Advances in Microbiology **08**, 235–251 (2018).
- 57. Gao, S., Wu, H., Yu, X., Qian, L. & Gao, X. Swarming motility plays the major role in migration during tomato root colonization by *Bacillus subtilis* SWR01. Biological Control **98**, 11–17 (2016).



Dual-culture assay of *B. velezensis* CMRP 4489 and *Sclerotinia sclerotiorum* (SS), *Macrophomina phaseolina* (MP), and *Botrytis cinerea* (BC). (**a**) Photography of the dual-culture test with the experimental controls (**b**) Mycelial growth inhibition percentage (MGI%). Means followed by the same letter did not differ according to the Tukey test (p<0.05).



Antifungal effect of the cell-free supernatant (CFS) of *Bacillus velezensis* CMRP 4489. (a) Evaluation of antifungal activity against *Sclerotinia sclerotiorum* (SS), *Macrophomina phaseolina* (MP), and *Botrytis cinerea* (BC) by metabolites produced in CM1 and CM2 media. (b) Scanning electron microscopy images of the antifungal effect of the CFS against *Sclerotinia sclerotiorum*. The control refers to the untreated fungal spot and CFS-MC2 refers to the transition area between fungal growth and the inhibition region.



Kinetics of cellular growth and antifungal activity of the cell-free supernatant (CFS) of *Bacillus velezensis* CMRP 4489. Mycelial Growth Inhibition percentage (MGI%); Colony Forming Units per milliliter (CFU/mL).



Mycelial growth inhibition (MGI%) of *Sclerotinia sclerotiorum* (SS), *Macrophomina phaseolina* (MP) and *Botrytis cinerea* (BC) by the cell-free supernatant (CFS) of *Bacillus velezensis* CMRP4489 after thermal, light, and pH treatments. (a) CFS stability at 28°C (no treatment), 70°C, 100°C, and 121°C (b) Mycelial growth of SS, MP, and BC at different temperatures in the thermostability test. (c) CFS photostability assay in UV light (UV), at wavelength of 365 nm, and visible light (VL). Same letters did not differ according to the Kruskal-Wallis test (p<0.05). (d) CFS stability at different levels of pH (3-11).



Scanning electron microscopy of the colonization of *Bacillus velezensis* CMRP4489 on soybean seeds. The first column shows untreated seeds. The second shows the seeds treated with strain CMRP4489, adjusted to 5×10^9 CFU / mL (200 mL / 100 kg). The first row shows seeds that were fixated on day 0 of treatment; the second row shows the seeds after 7 days of incubation, and the third row shows the radicles of the seeds after 7 days of incubation. All images have inserts at 50x magnification.



Control of *Sclerotinia sclerotiorum* on soybean seeds by inoculation of *Bacillus velezensis* CMRP 4489. Evaluation of the % of (A) germinated seeds, (B) uninfected seeds of *Sclerotinia sclerotiorum*. Bars represent the standard deviation based on 13 replicates per treatment. Letters indicate statistical difference by the Tukey test (p<0.05).

Figure 7

Circular representation and comparison of the *Bacillus velezensis* CMRP 4489 strain genome with other strains from the *Bacillus subtilis* group using the BRIG program. From the inside out: representation of the genome size of *B. velezensis* CMRP 4489, GC Skew, GC Content, positions of the clusters of the *B. velezensis* CMRP4489 strain, and the strains FZB42, NGK1, S141, QST713, Bac57, CC178, DSM7, MT45, YP6, 168, and SRCM1036, respectively.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

SupplementaryMaterialsManuscriptBacillusvelezensisCMRP4489.docx