

Impact of Social and Cultural Factors on Incidence, Transmission and Control of Coronavirus Disease in Iran: A Qualitative Study

Saeed Falla-Aliabadi

Shahid Sadoughi University of Medical Sciences

Ahad Heydari

Kurdistan University of Medical Sciences

Farin Fatemi

Semnan University of Medical Sciences

Nooshin Yoshany

Shahid Sadoughi University of Medical Sciences

Mohammad Hasan Lotfi

Shahid Sadoughi University of Medical Sciences

Alireza Sarsangi

GIS and Remote Sensing Department, University of Tehran, Iran

Fahad Hanna (✉ fahad.hanna@torrens.edu.au)

Torrens University Australia

Research Article

Keywords: Covid-19, social factors, cultural factors, community, governance, health policy

Posted Date: May 26th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1627738/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction: COVID-19 pandemic has had mixed reactions from nations, people and governments about ways to cope with, prevent and control the disease. The current study identifies social, cultural and policy factors affecting the incidence and control of Coronavirus disease in Iran.

Methods: A qualitative study consists of content analysis and views of experienced and knowledgeable subjects specialized in social and cultural health management. Data collection has been implemented through appropriate content specific search as well as semi-structured interviews. Data analysis was complemented by inductive content analysis.

Results: There are distinct social and cultural factors in coping with Coronavirus disease. These consisted of three categories of governance, individual and community related factors. A total of 21 subcategories and 215 primary codes that were extracted from the text of interviews as variables of the study and in relation to the research question. Twelve subdomains of governance including vaccination, political factors, restrictions, education, support services, administrative services, transportation, health and treatment, culturalization, legislation, management and finance impacted the spread and mitigation of the pandemic at various levels.

Conclusion: The management of pandemics requires a comprehensive capacity for identifying and determining social and cultural criteria. A healthy partnership between governments and the community may be required to remove unnecessary obstacles that hinder public health attempt to alleviate the risk. The obtained criteria and indicators from this study may be utilized by policy makers in an attempt to strengthen protocols for mitigating pandemics. Further studies may be warranted to confirm these findings

Introduction

Concurrent with Coronavirus disease pandemic in the world, distinct factors based on the social and cultural characteristics of communities have undoubtedly impacted the rate of spread and the level of mortality of the disease in different parts of the world. In various geographical regions, people have displayed different behaviors in coping with the emergent diseases(1). These behaviors are derived from social, economic and cultural processes and they are effective in the manner of reacting to that phenomenon and managing the consequences of that phenomenon, and, will pave the way for increasing and decreasing the vulnerability rate of communities to that phenomenon (2). Factors such as education level, income level, security status, cultural status, the existent of indigenous and immigrant population can have an impact on increasing the vulnerability of community to emergent events (3, 4). Being cognizant of cultural and social factors in different communities contribute to managers and organizational decision makers to take measures in line with reducing the possible harms based on available resources (5).

Considering the existent of various cultures and different social and economic conditions with adjacent countries, Iran has struggled with disease management and treatment of patients since the early months of Coronavirus disease. The official confirmation of the existent of positive test report form of Corona in Iran occurred in February 2020 leading the number of infections and deaths to culminate by March, June, November 2020, and at the months of April and August 2021 (6–9). Given the mortality rate derived from the disease, various approaches were proposed by policy makers for disease management, that followed by a part of the community accompaniment and occasionally the existent of contradictory behaviors with the official policies of the government were seen by a part of the community. These behaviors have been resulted from multifarious social, cultural and economic factors that it is required to be identified through distinct studies. By identifying these factors and examining their role in the incidence and control of Coronavirus disease, it can be taken action in line with applying management policies related to controlling the disease. These determinants can be compared with social and cultural factors in other parts of the world and can investigate the effect of cultural and social factors on disease management. Furthermore, examining the behaviors of community affected by social and cultural factors can be generalized to other emergent diseases that may transpire in the future.

Ample studies have been implemented on the role of social, economic and cultural factors in the incidence of infectious diseases in various parts of the world (10–14). Concurrent with the coronavirus epidemic, considerable studies have been conducted in the domain of social issues given to the effect of this disease on population health. Other studies have examined the social vulnerability of communities resulted from the Coronavirus epidemic and presented the plan for removing the vulnerability (15–17). Some of these essays used Social Vulnerability Index (SVI) data that encompasses 4 domains of socioeconomic status, household composition and disability, racial / ethnic minority and English language proficiency status, and housing and transportation and compared the vulnerability status of different places or groups (18–21).

Some of the studies have investigated socio-cultural determinants related to Coronavirus via a systematic review. For instance, Chu et al. In a study have identified the social consequences of quarantine in communities during the infectious diseases pandemic period and presented strategies to reduce the negative social consequences of quarantine during the COVID-19 epidemic via systematic review (22). In another study, the authors have examined the intervention rate of factors related to the social health in predicting epidemic consequences in models connected with COVID-19 (23). Other researchers have examined the effects of social distance on the living conditions of various people as children, pregnant women, the elderly and patients during the corona epidemic period (24–30). Moreover in some essays the effect of quarantine and distance on the incidence of social anomalies and behavioral problems has been stated (31–34). In another essay, researchers have examined the social and economic factors of the COVID epidemic in 42 Asian countries empirically. In this study, the role of factors as age, status of background diseases, income, poverty, social capital, migration, social participation in the rate of getting infected and in the epidemic of the disease have been investigated (35). On the other hand, the numbers of essays have been conducted in different countries of the world

along with a qualitative approach in the domain of social conditions and its relationship with the issue of Coronavirus epidemics that examines the incidence of behaviors (30, 36–39).

Therefore, given to the importance of identifying socio-cultural factors affecting the incidence, transmission and management of Coronavirus disease in Iran, it is mandatory to conduct a research by applying qualitative approach. The purpose of the current study is to identify the social and cultural criteria and determinants affecting the epidemic and control of Covid-19.

Materials And Methods

Since qualitative studies are very appropriate for identifying prominent variables related to a phenomenon, this study has been conducted qualitatively to identify the economic and social factors affecting the behavior of individuals during the Coronavirus pandemic period in Iran in 2021 (40).

Study Population

In this study experts, managers and specialists who had knowledge, expertise and experience in multifarious domains, namely: health and treatment, epidemiology, sociology, political science, economics, cultural studies and religious experts. They were interested in participating in this research were selected and interviewed purposefully at the very beginning and also in order to identify other experts and clear-sighted individuals, the snowball sampling method was used. Entrance criteria for interviewees included possessing at least a master's degree in the relevant and above-mentioned domains, living in Iran, inclining to cooperate in the present study and encompassing an administrative background in the above-mentioned domains.

Data collection method

In this study, after purposeful selection of population, face-to-face audio recording by MP3 Player was implemented through the interview guide. At the first place, three unstructured interviews were complemented due to specifying the pivots of the interview and completing the questions of interview guide besides the interviews continued along with 17 other participants in a semi-structured manner until a new code was not found in the final interviews and accordingly data saturation was obtained.

At the outset of the interview, the purpose of the study was explained to the populations. At the end of the interview, participants were requested to introduce other experts if they knew therefore the data can be validated by selecting clear-sighted individuals and spending enough time.

Interviews were implemented between June and August 2021. The interviews prolonged between 20 and 50 minutes and averagely 30 minutes. Interviews were implemented by arrange a previous appointment and in a calm and quiet environment. To ensure the accuracy of the data and the interviewer's domination to a great extent as well, the audio files were listened at many times and if there were ambiguous topics or it was revealed the vagueness in formation of new questions, the interviewees would be contacted again. Eventually, the audio files were implemented in word software.

Data analysis

Data collection and data analysis were implemented concurrently. Content analysis was conducted by Graneheim and Lundman method during 8 steps. Moreover the principles of qualitative content analysis of the recommended Graneheim were applied (41).

In this study, after conducting each interview, the text of the interviews was implemented at the first opportunity and the audio file of the interviews was typed. From the units of concepts and meaning, the units of the abridged meaning and then the primary codes were obtained. After that the codes based on a central concept were embedded within a subclass. Then, the sub-classes were revised at many times and compared with each other according to similarities and differences, and then the classes and the themes were existed, as there was the most homogeneity within the classes and the most heterogeneity between classes.

Data Credibility:

In order to ensure the credibility of information, four criteria including credibility, dependability, transferability and confirmability were determined (42).

Ethical considerations

The time and location of the interviews were coordinated with the interviewees via phone calls. At the outset of the interview, participants were asked read and sign informed consent in relation to recording the interview and the researcher presented participant information sheet thereby ensuring that their recorded dialogue and conversations were confidential and that their first and last names to be anonymous and to be coded and the name of each individuals must not be known after implementing and converting into the text.

All experimental protocols in this study were approved by the Research Ethics Committee of Shahid Sadoughi University of Medical Sciences, Yazd, Iran (**IR.SSU.REC.1400.052**). Moreover, all methods were carried out in accordance with relevant guidelines and regulations.

Results

The age of study population was between 34-65 years with an average age of 43 years. Three participants had MA degrees while 17 possessed PHD degrees. In accordance with the purpose of the study, 215 relevant codes were emitted from the content analysis of the questionnaires that in the subsequent step, these codes were divided into 3 main domains including governance factors, individual factors and the factors related to the community and were classified into 21 subdomains (Appendix 1).

Governance factors

In this study, governance factors were identified as the most considerable and comprehensive component affecting the Coronavirus epidemic. Governance factors encompasses 12 subdomains of vaccination, political factors, restrictions, education, support services, administrative services, transportation, health and treatment, culturalization, legislation, management and finance.

Vaccination

Vaccination, as one of the vital strategies to reduce the spread of Coronavirus epidemic, has struggled with many challenges in Iran. Multifarious factors such as absence of public accessibility to vaccines, the loss of valid vaccines and the lack of public satisfaction to the vaccination have significantly hindered the public health efforts to mitigate the impact of the epidemic. Considering the lack of sufficient vaccines in Iran, the process of vaccination has been categorized and it takes a long time for each individual in the community to be vaccinated, which has resulted into the slowing control of Coronavirus. To this point, participant number 5 commented as follow:

"All of the people had no access to the vaccine at the exact time, and at the very beginning the process of vaccination had started with the older age groups. These conditions have caused many people who did not reach the time for their vaccine turn to be frustrated, and occasionally because they were tired of following the protocol, they didn't incline to follow the limitations. The individuals whose relatives died of Coronavirus were depressed and some objected to these existent conditions."

Political factors

Given the existent of political issues in Iran, and the sour international relations and challenges with other countries, as well as the existence of bank and logistic sanctions, the import of high quality and valid vaccines has been limited. Participant number 15 comment in this regard is noted below:

"For instance, our country cannot receive vaccines or health materials and products from developed countries because of international sanctions, and this has caused the process of vaccination and treatment to be difficult. On the other hand, the government officials promised that the government will produce Iranian vaccine that is not clear how effective it will be according to their statements as we are self-sufficient versus other countries that do not send vaccines to us. Some diplomats recommend holding political and religious demonstrations and other ceremonies as gatherings and elections."

Legislation

Ratification of legal restrictions is another obtained subdomain in this study. The results of this study demonstrate that factors such as entry of travelers from foreign country without testing and quarantining, the weak implementation the commute restrictions, non-obedience of certain groups from the enacted restrictions, absence of responsible organizations action toward health offenders and lack of staff due to controlling the Implementation of restrictions contributed to the increase of the spread of the pandemic. Moreover another predicament of the legislation is that there is no coherent management for

implementation of regulations, and the behavior of the managers indicates dual personality/views toward implementation of regulation.

In this regard, participants number 7 and 20 stated the following, respectively:

"Coronavirus disease is a disease that, given to the essence and nature of the disease, we should be able to prevent from spreading it correctly. We should block it. Blocking should be implemented in a way that martial law must be ordained. Commuting must be limited, in other words the governance must be powerful enough to convince the community that they must stay home. It must be closed everywhere for at least twenty days or one month; when it is blocked for a month or twenty days, the spread of the disease will automatically reduce, and if the governance does not possess this power, this can lead to spreading the disease to a great extent."

"The community is expected to follow the protocols, to the same extent, the government is expected to manage the disease by establishing coherent regulation and maintaining a coherent approach. If a single protocol and word is implemented equally to all of the individuals, the people will obey the regulations to a great extent."

Education

The results of this study demonstrated that in many situations, people may not follow health protocols due to loss of knowledge about the nature of the disease and the ways of its transmission, or, in many cases people abstain from vaccination due to spreading propaganda and lack of adequate knowledge, as per participant number 4 statement below:

"Another factor is related to the knowledge and getting aware of the disease. If we do not have sufficient knowledge and awareness and do not follow the principles of health, this will lead to the spread and transmission of the disease to a great extent, and this can be rooted in beliefs of people. We have knowledge and awareness in some cases but we do not believe, it means that we do not understand the dangers of the disease, in other words we do not know what is threatening us, and we do not feel the danger greatly."

Administrative, public and support services

The findings of this study demonstrated that representing administrative services in a virtual way and the public and support services considered by the government for the people could encourage people to stay at home. This may reduce the volume of traffic and decrease the risk of encountering with the virus in the community. The government should provide financial subsistence and support for people to stay at home thereby reducing the potential violation of restrictions due to economic pressures. On the other hand, providing financial subsistence support packages can have a significant effect on reducing public commuting. In this regard, participants number 2, 5 and 17 stated as follow, respectively:

"In many cases, people throng in banks or organizations and they confront with each other in indoor spaces. Unfortunately, electronic services are not presented to the people and they have to go to these places to meet their administrative and banking needs. On the other hand they have to travel from city to city, and when they travel, they may encounter with the people who are carrier of the disease in a taxi or bus."

"The expense of living in big cities is high. In this situation, the expenses of the conditions related to the Coronavirus disease prevention and treatment are imposed on people as well. Particularly in big cities, it is hard to say that people should not use the subway or bus. These transportation vehicles are also crowded and people do not obey the distance when they sit or stand beside each other."

"Furthermore we should provide the minimum supplies and necessities of the people. When we say stores should be closed, as some people do not possess good financial capability. At least affluent people or the government can meet their supplies and needs. Eventually, people need income. Their stores cannot be closed by us unless they do not have financial problems. If the government can contribute to them, fewer people will leave their homes."

Health and treatment services

The preparedness of the health care system has a vital role in controlling Coronavirus disease. The health and treatment system must be able to prepare for identifying and screening the patients and this system must possess sufficient equipment, resources, personnel and hospital beds in the case of the outset of epidemic. In many cases the health and treatment system in Iran has encountered with the lack of medicine, hospital beds, oxygen and above all health and treatment specialists in management and controlling of the disease at the beginning of the expansion of Covid-19. In this regard, participant number 14 commented:

"The weakness of the health and treatment system can also be effective in the expansion of the disease. If we do not have diagnostic test and do not be able to screen it will cause the expansion of the disease to a great extent and if our health and treatment system cannot be able to treat accurately, it can pave the way for disease epidemic. Obeying from the health protocols in places as restaurant or shop can be effective in incidence of the disease."

Culturalization

Culture of the community is one of the vital factors in controlling or expanding diseases as Covid-19. The culture of the people in Iran is such a way that it is mandatory to companion (visiting and meeting the elders of the family) or to participate in weddings or funeral ceremonies that these behaviors increase the incidence and expansion of this disease in the community. In this regard, participant number 5 commented:

"The first measure that should be taken in the country via schools and the media is to build a culture in the domain of personal care so that people can notify of the prevention ways. At the very beginning there

were involvement of news, media, radio and social media and systemic recommendations were given to the people for disease prevention. It is propagated contradictory statements about the nature of the disease and how to cope with it, especially in social media, and some people believe them though."

Financial Policy

Financial and economic policies of the government are effective in controlling the disease to a great extent. Financial contribution of the government as bank cooperation, deferred debt, strengthening charitable organizations financially and the existence of support organizations as the Relief and Welfare Committee to help the families under coverage of the insurance can effect on preventing the expansion of the disease. In this regard, participant number 3 stated:

"Even for the people who live in slums, the existence of health packages as masks and sanitizers can be very effective. There are people who live in difficulty and want to obey the regulations but they cannot follow the route. Financial contributions as providing loan and livelihood contributions and it must be taken action not just words".

Individual factors

The domain of individual factors was broached as an effective element in relation to Coronavirus epidemic by the interviewers in this study. Individual factors were stratified into 7 subdomains after categorizing and combining common codes.

Understanding the Risk

Understanding the risk of people in community and being aware of the consequences of high-risk behaviors is one of the most vital factors affecting the disease epidemic. On the other hand, the absence of knowledge and awareness of people about the disease, its consequences and control strategies can reduce the understanding the risk of people. In this regard participants Number 1 and 19 commented, respectively:

"Some people as youth or athletes possess the exaggerated courage or fearless as if I am in good condition and I will never get sick and in case of getting infected I will overcome the disease. This wishful thinking and misconception have embedded in his mind and have paved the way for not wearing mask."

"Some people soliloquized that I should call on my friend and relatives and greet them but he did not know that if he has been the carrier of the disease, he would have entangled some other people and if God forbid, the same friends or relatives to be died of the disease, the one will not forgive himself."

Personality factors

The behavior of the people is derived from their personality. People may be dissociable or be possessed high social relationships. A group of people may be parsimonious enough to not buy masks and prevention instruments related to Covid-19. Some people are depressed and tired of following the health

protocols and restrictions. These personality features are effective on the behaviors of the people towards prevention from Covid-19. In this regard participant number 6 commented:

"Even some people said apparently that we wear a mask; but it does not go with us and we like to refuse wearing masks. When we got infected to Coronavirus, it is not a problem as we will be treated. Some individuals are tired of this situation and say that I how late I should obey. Maybe this disease may be existed forever. Of course some individuals got along with this condition and accept the Corona protocols as a part of their life."

life style

Another factor affecting on high-risk behaviors during the corona epidemic is lifestyle. Some people live in thickly populated families and communities. Some people smoke, consume alcohol or use drugs, or some people may be weakened physically due to poor condition of nutrition and also their body may be more vulnerable. In this regard participant number 18 commented:

"Some people look the disease more seriously and maybe obey the protocol to a great extent because of their vulnerabilities, as poor nutrition or living at home with the elderly."

health condition

The results of this study demonstrate that health condition of people can affect their behavior before Coronavirus disease. People with rare diseases and also the people who are exposed to the risk of health follow health protocols more than the people who are in good health condition. In contrast, the youth and athletes who are in a healthy physical condition show high-risk behaviors before the disease. The participant number 12 commented:

"Families with people who are infected with chronic disease as diabetes, cancer and background diseases, family members try to obey the health protocols to a great extent and also it is seen less commuting in these families."

Factors Related to the Community

In this study, a group of emitted codes regarding the factors affecting the epidemic of Covid-19 in the community were classified into subdomains of social and cultural factors and were embedded in the domain of community-related factors.

Social factors

Social behaviors as companionships, parties, traveling and visiting are considered as social behaviors related to community norms. Various ethnicities have their own social rituals and behaviors. Moreover, traditional communities in smaller cities rely on their mores to a great extent and do not incline to stop these mores. In this regard participant number 8 and 13 commented as follows, respectively:

"We are a traditional community. We do not quit a series of habits, behaviors and ceremonies, including weddings and funeral ceremonies easily. These situations can pave the way for a more expansion of the disease."

"At the beginning of the disease epidemic, many people quitted important and traditional ceremonies as visiting in Nowruz holidays, as Corona period had been lasted and perhaps the situation had been normalized for the people, this ceremony was held again, of course this ceremony was hardly implemented in the next year."

Cultural factors

The culture of communities is one of the most vital factors that can affect the behavior of people. In general, some communities are circumspect and in some other communities, due to the existence of prevailed and old culture of high risk-taking, the high-risk behaviors are common among individuals. On the other hand, changing the culture in some communities is very tough and time consuming, for instance, canceling and avoiding parties and some traditional ceremonies during the disease epidemic. In this regard participant number 11 stated:

"It is probable that some ethnicities and cities may have their own series of regulations. In this condition, it is not important how hard you will attempt as the restrictions are before their culture, you cannot really hope to see following them therefore in everywhere we must know their own cultures and impose restrictions on them."

Discussion

According to the results of this study, the obtained components along with positive and negative effects on the incidence, transmission and control of coronavirus are embedded in three main categories of governance, individual and social components. Policy-making, making legislation and community awareness on corona disease are the issues that are considered as the most vital components of governance that are undertaken by the government of Iran and the Ministry of Health, in particular. Implementing inappropriate policies or making non-logical decisions without allowing appropriate time by the government may have led to increased rate of mortality and lack of trust of the people towards their government and the privation of resources. For instance, delays in the arrival of a valid vaccine led to relatively high rate of mortality (700 people in August 2021) (9). A study in India on rate of mortality in first and second peak demonstrated that the rate of mortality in the second peak was lower than the first peak as part of population over 65 years of age being vaccinated in the second peak compared to the first peak of the disease (43). At the same time, contradictions about the arrival of foreign vaccines revealed, and some other broached ineffectiveness of the vaccination or its negative effect on community health as well thereby negatively influencing the acceptance of members of the community to vaccinate. This issue was quite prevalent in some Iranian regions along with the people of Japan and the United States has been discussed in a study between 17 countries of the world (44).

Another issue is legislation of governance aiming to manage and reduce the rate of Corona disease. Lack of attention paying and adherence to the setting of restrictions and regulations by non-experts may pave the way for ineffectiveness of such legislation. Following the high proportion of people who got infected in Iran, lengthy periods of restriction were announced in various cities around the country. These restrictions were accompanied by closing various places and prohibiting public gatherings, however, the absence of serious mitigation, planning and compensation resulted into abundant economic problems for the public. Poor support policies by the government caused people to defy the restrictions, which ultimately contributed to increase rate of infection and mortality. These findings are consistent with the results of a study conducted in Santiago, Chile in 2021. The above study demonstrated that there is a strong and significant relationship between social-economic status and the effects of Covid-19 and capacity of community health (45). Our study also found that the deprivation of various services to violators of health protocols and effective monitoring of operation in community may pave the way for reducing the epidemic of the disease, transmission and ultimately the rate of mortality. This finding is consistent with a study in Sweden by Modig et al (2021) where the government restrictions being “recommended” rather than strictly implemented may have contributed to higher rate of mortality due to Covid-19 infection (46). On the contrary, in various other countries, strict regulations were imposed on the public which resulted into a more efficient way for controlling the spread of the disease (47–49). For instance, the appropriate enforcement and control of the social distance policy in some countries, such as Italy, has successfully reduced the rate of mortality in the second peak of Covid-19 disease in the country (47). Distribution of livelihood packages, loan payments, supporting of low-income families who lost livelihoods, supporting and withholding loan repayment, saving individuals and families from business bankruptcy, price reduction of essential materials such as medicine and main nutrients should be taken on board to encourage the public to follow the pandemic rules. A county-level analysis in the United States demonstrated that regions with lower economic status had been faced with higher rate of mortality in the Coronavirus pandemic (48). Furthermore, a more extreme example of poverty in Peru was considered as one of the most influential factors for the excessive rate of mortality due to Covid-19 disease (49). The issue of lack of support from the government of Iran at the economic level, particularly not considering those with poor economic conditions, has also been discussed in another study (50).

Informing the community of the prevention from the pandemic, treatment, and consequences of Covid-19 requires the all-out participation and involvement of health care providers, non-governmental organizations, donors, celebrities, and religious and faith leaders. Considering the fact that medical providers and frontline healthcare workers would struggle with a long-term pandemic, it is imperative to provide sufficient economic, occupational and psychological support to this cadre. High cost of treatment and prevention have been another significant problem. The relative lack of familiarity with the virus and the use of different treatment protocols, with some being ineffective, has led to community pessimism about the treatment cadre, which occasionally caused people to avoid hospitalization leading to condition being exacerbated. A study conducted Iranian cities demonstrated that lack of cure along with other rumors may have paved the way for fear of this epidemic among the public (51). Assuring the

public about the effectiveness of medical treatment in early and timely hospitalization may require a health partnership between governments and the community as a whole.

Another important factor in this study that highlights the role of community in controlling the disease was the individual, cultural and social characteristics. Many of the emerging data on COVID-19 death seemed to be pointing to the fact that older and unhealthy people were more likely to die from it. This was specifically highlighted in a study conducted in Ardebil (Iran), where one of the most predominant mortality risk factors age(52). A proportion of the public ignored the disease by violating the regulations and/or by claiming to be protected due to being young or athletic. The results of a study in the Ohio and Florida states of US between March 15 and December 5, 2020 demonstrated that higher than expected rate of mortality have occurred at the age group of 20–49 years (53). Fatalism, arrogance, insistence, getting susceptible to depression, escaping from loneliness and getting tired of obeying protocols and constant use of masks are considered as other individual and behavioral factors that led to disobeying or neglecting health protocols. Profession type of individuals was also a significant factor in controlling and preventing Coronavirus in community. Findings from the same (above) US study demonstrated that compared with whites, black people, who are placed in a lower economic status and whose professions could not be telecommuted/virtual, exhibited higher mortality rate from the virus (53). The number of clients, size of indoor or outdoor spaces, population density and the degree of obeying the protocols in the workplace can affect the spread and transmission of the disease. At the beginning of the Corona epidemic in Iran, medical officials, taxi drivers, and bank employees had the highest rate of infection and mortality due to COVID-19, however, the rate of death decreased in these groups concurrent with prioritizing vaccination for these individuals (54).

One of the limitations of this study is that it is qualitative by nature which is subject to information bias and views. The information derived from qualitative research can also be highly subjective. However, we have used all efforts to ensure accuracy of data collected from individuals who simply expressed their views about the pandemic. A more quantitative approach coupled with larger sample size and longitudinal design may provide more accurate and reliable data. Furthermore, these findings are relevant to Iranian communities may not necessarily apply to other countries and communities. However, poverty, cultural practices, behavior and other individual characteristics can be common human features, particularly in the developing world.

Conclusion

Managing the COVID pandemic requires a comprehensive knowledge and understanding of social and cultural determinants within each community to facilitate more comprehensive protocols and approaches to mitigate the spread. It is mandatory to provide health, economic and operational infrastructure and support to further enhance the management and control of the pandemic. Moreover, the results of this study demonstrate that in order to improve the effectiveness of disease prevention, managing and control, a more comprehensive cooperation and coordination between all departmental and intradepartmental organizations as well as various institutions of the country, together with a healthy

partnership with the community is required. Therefore, it is highly recommended that trainings, public and specific regulations, programs and the needed resources must be enforced and compiled for preparing and responding to epidemics relevant to the economic and cultural status of the community. The findings of this study can be utilized by policy makers in the domain of cultural, social and health issues in relation with the management of the Covid-19 pandemic and any future pandemic. Further studies mixed with quantitative data may be required to increase the credibility and generalizability of the determinants of Covid-19 spread and control.

Declarations

Ethics approval and consent to participate:

All experimental protocols in this study were approved by the Research Ethics Committee of Shahid Sadoughi University of Medical Sciences, Yazd, Iran (**IR.SSU.REC.1400.052**). Moreover, all methods were carried out in accordance with relevant guidelines and regulations.

Consent for publication:

All co-authors have consented for the manuscript to be published in BMC Public Health

Availability of data and materials:

All data extracted from this qualitative study is available upon request. The majority of the raw data is in Persian, however, a translated version will be available upon request.

Competing interests:

All authors on this manuscript declare no competing interests

Funding:

This study is part of a larger post-doctoral project funded by Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

Authors' contributions:

SFA and AH has made substantial contributions to the initial conception and design of the study, and has taken responsibility for and coordinated the acquisition of data. SFA worked on the analysis of the data while AH, FH, NY and SFA contributed to results interpretation and writing-up of the manuscript. FF, MHL, NY, AS and FH reviewed the data analysis, writing-up and finalization of the manuscript. All authors read and approved the final manuscript.

Acknowledgment

The authors would like to acknowledge members of the communities that participated in this social research and thank each and every staff member who facilitated the work on ground.

References

1. Colizza V, Barrat A, Barthélemy M, Vespignani A. The role of the airline transportation network in the prediction and predictability of global epidemics. *Proceedings of the National Academy of Sciences*. 2006;103(7):2015-20.
2. Weible CM, Nohrstedt D, Cairney P, Carter DP, Crow DA, Durnová AP, et al. COVID-19 and the policy sciences: initial reactions and perspectives. *Policy sciences*. 2020;53(2):225-41.
3. Calderón-Larrañaga A, Vetrano DL, Rizzuto D, Bellander T, Fratiglioni L, Dekhtyar S. High excess mortality in areas with young and socially vulnerable populations during the COVID-19 outbreak in Stockholm Region, Sweden. *BMJ global health*. 2020;5(10):e003595.
4. Bernard KW. Health and national security: a contemporary collision of cultures. *Biosecurity and bioterrorism: biodefense strategy, practice, and science*. 2013;11(2):157-62.
5. Van Bavel JJ, Baicker K, Boggio PS, Capraro V, Cichocka A, Cikara M, et al. Using social and behavioural science to support COVID-19 pandemic response. *Nature human behaviour*. 2020;4(5):460-71.
6. Ghafari M, Hejazi B, Karshenas A, Dascalu S, Kadvidar A, Khosravi M, et al. Lessons for preparedness and reasons for concern from the early COVID-19 epidemic in Iran. *Epidemics*. 2021;36:100472.
7. Doosti-Irani A, Haghdoost AA, Najafi F, Eybpoosh S, Moradi G, Amiri FB, et al. How can the epidemic curve of COVID-19 in Iran be interpreted? *Journal of research in health sciences*. 2020;20(3):e00491.
8. Shafiekhani S, Khalilabad TH, Rafiei S, Sadeghi V, Jafari AH, Gheibi N. Trend and prediction of COVID-19 outbreak in Iran: SEIR and ANFIS model. *Polish Journal of Medical Physics and Engineering*. 2021;27(3):241-9.
9. Available from: <https://www.worldometers.info/coronavirus/>.
10. Ali H, Dumbuya B, Hynie M, Idahosa P, Keil R, Perkins P. The Social and Political Dimensions of the Ebola Response: Global Inequality, Climate Change, and Infectious Disease. *Climate Change Management* 2016. p. 151-69.
11. Barnett T, Fournié G, Gupta S, Seeley J. Some considerations concerning the challenge of incorporating social variables into epidemiological models of infectious disease transmission. *Global Public Health*. 2015;10(4):438-48.
12. Braveman P. Accumulating knowledge on the social determinants of health and infectious disease. *Public Health Reports*. 2011;126(SUPPL. 3):28-30.
13. Hunter PR, Colón-González FJ, Brainard J, Majuru B, Pedrazzoli D, Abubakar I, et al. Can economic indicators predict infectious disease spread? A cross-country panel analysis of 13 European countries. *Scandinavian Journal of Public Health*. 2020;48(4):351-61.

14. Sands P, El Turabi A, Saynisch PA, Dzau VJ. Assessment of economic vulnerability to infectious disease crises. *The Lancet*. 2016;388(10058):2443-8.
15. Biggs EN, Maloney PM, Rung AL, Peters ES, Robinson WT. The Relationship Between Social Vulnerability and COVID-19 Incidence Among Louisiana Census Tracts. *Frontiers in Public Health*. 2021;8.
16. Coelho FC, Lana RM, Cruz OG, Villela DAM, Bastos LS, Pastore Y Piontti A, et al. Assessing the spread of COVID-19 in Brazil: Mobility, morbidity and social vulnerability. *PLoS ONE*. 2020;15(9 September. (
17. Credit K. Neighbourhood inequity: Exploring the factors underlying racial and ethnic disparities in COVID-19 testing and infection rates using ZIP code data in Chicago and New York. *Regional Science Policy and Practice*. 2020;12(6):1249-71.
18. Coelho FC, Lana RM, Cruz OG, Villela DA, Bastos LS, Pastore y Piontti A, et al. Assessing the spread of COVID-19 in Brazil: Mobility, morbidity and social vulnerability. *PLoS One*:(9)15;2020 .e0238214.
19. Roy A, Kar B, editors. Characterizing the spread of COVID-19 from human mobility patterns and SocioDemographic indicators. *Proceedings of the 3rd ACM SIGSPATIAL International Workshop on Advances in Resilient and Intelligent Cities, ARIC 2020*; 2020.
20. Neelon B, Mutiso F, Mueller NT, Pearce JL, Benjamin-Neelon SE. Spatial and temporal trends in social vulnerability and COVID-19 incidence and death rates in the United States. *medRxiv*. 2020.
21. Garcia OEL, Retamales VA, Suarez OAM, Parajuli P, Hingle S, Robinson R. Application of Social Vulnerability Index to Identify High-risk Population of Contracting COVID-19 Infection: a state-level study. *medRxiv*. 2020.
22. Yen-Hao Chu I, Alam P, Larson HJ, Lin L. Social consequences of mass quarantine during epidemics: A systematic review with implications for the COVID-19 response. *Journal of Travel Medicine*. 2020;27(7.(
23. John-Baptiste A, Moulin MS, Ali S. Are COVID-19 models blind to the social determinants of health? A systematic review protocol. *BMJ Open*. 2021;11(7.(
24. Fedorenko EJ, Kibbey MM, Contrada RJ, Farris SG. Psychosocial predictors of virus and social distancing fears in undergraduate students living in a US COVID-19 “hotspot”. *Cognitive Behaviour Therapy*. 2021;50(3):217.33-
25. Morgan AK. Making COVID-19 prevention etiquette of social distancing a reality for the homeless and slum dwellers in Ghana: lessons for consideration. *Local Environment*. 2020:536-9.
26. Prata- Linhares MM, Cardoso TDSG, Lopes-Jr DS, Zukowsky-Tavares C. Social distancing effects on the teaching systems and teacher education programmes in Brazil: reinventing without distorting teaching. *Journal of Education for Teaching*. 2020;46(4):554-64.
27. Roe L, Proudfoot J, Tay Wee Teck J, Irvine RDG, Frankland S, Baldacchino AM. Isolation, Solitude and Social Distancing for People Who Use Drugs: An Ethnographic Perspective. *Frontiers in Psychiatry*. 2021;11.
28. Vilar-Compte M, Pérez V, Teruel G, Alonso A, Pérez-Escamilla R. Costing of actions to safeguard vulnerable Mexican households with young children from the consequences of COVID-19 social

- distancing measures. *International Journal for Equity in Health*. 2020;19(1.(
29. Zainel AA, Qotba H, Al-Maadeed A, Al-Kohji S, Al Mujalli H, Ali A, et al. Psychological and coping strategies related to home isolation and social distancing in children and adolescents during the covid-19 pandemic: Cross-sectional study. *JMIR Formative Research*. 2021;5(4.(
 30. Anderson E, Brigden A, Davies A, Shepherd E, Ingram J. Pregnant women's experiences of social distancing behavioural guidelines during the Covid-19 pandemic 'lockdown' in the UK, a qualitative interview study. *BMC Public Health*. 2021;21(1.(
 31. Antonelli-Salgado T, Monteiro GMC, Marcon G, Roza TH, Zimmerman A, Hoffmann MS, et al. Loneliness, but not social distancing, is associated with the incidence of suicidal ideation during the COVID-19 outbreak: a longitudinal study. *Journal of Affective Disorders*. 2021;290:52-60.
 32. Esteves CS, Oliveira CRD, Argimon IIDL. Social Distancing: Prevalence of Depressive, Anxiety, and Stress Symptoms Among Brazilian Students During the COVID-19 Pandemic. *Frontiers in Public Health*. 2021;8.
 33. Garnett C, Jackson S, Oldham M, Brown J, Steptoe A, Fancourt D. Factors associated with drinking behaviour during COVID-19 social distancing and lockdown among adults in the UK. *Drug and alcohol dependence*. 2021;219:108461.
 34. Pieh C, Plener PL, Probst T, Dale R, Humer E. Assessment of Mental Health of High School Students during Social Distancing and Remote Schooling during the COVID-19 Pandemic in Austria. *JAMA Network Open*. 2021.
 35. Varkey RS, Joy J, Sarmah G, Panda PK. Socioeconomic determinants of COVID-19 in Asian countries: An empirical analysis. *Journal of Public Affairs*. 2020.
 36. Rocha LP, Rose R, Hoch A, Soares C, Fernandes A, Galvão H, et al. The impact of the covid-19 pandemic on the brazilian immigrant community in the U.S: Results from a qualitative study. *International Journal of Environmental Research and Public Health*. 2021.(7)18;
 37. Tan X, Song Y, Liu T. Resilience, vulnerability and adaptability: A qualitative study of COVID-19 lockdown experiences in two Henan villages, China. *PLoS ONE*. 2021;16(2 February).(
 38. Gupta P, Sengupta A. A qualitative analysis of social scientists' opinions on socioeconomic and demographic implications of the lockdown during COVID-19 in India. *Journal of Public Affairs*. 2021;21(3.(
 39. Halvaiepour Z, Nosratabadi M. Identifying social perceptions of people ignoring COVID-19 warnings: a qualitative study in Iran. *BMC Research Notes*. 2021;14(1.(
 40. Yari A, Ardalan A, Ostadtaghizadeh A, Zarezadeh Y, Boubakran MS, Bidarpoor F, et al. Underlying factors affecting death due to flood in Iran: A qualitative content analysis. *International Journal of Disaster Risk Reduction*. 2019;40:101258.
 41. Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts, procedures and measures to achieve trustworthiness. *Nurse education today*. 2004;24(2):105-12.
 42. Lincoln YS, Lynham SA, Guba EG. Paradigmatic controversies, contradictions, and emerging confluences, revisited. *The Sage handbook of qualitative research*. 2011;4:97-128.

43. Purkayastha S, Kundu R, Bhaduri R, Barker D, Kleinsasser M, Ray D, et al. Estimating the wave 1 and wave 2 infection fatality rates from SARS-CoV-2 in India. *BMC Research Notes*. 2021;14(1).
44. Wong LP, Alias H, Danaee M, Ahmed J, Lachyan A, Cai CZ, et al. COVID-19 vaccination intention and vaccine characteristics influencing vaccination acceptance: a global survey of 17 countries. *Infectious diseases of poverty*. 2021;10(1):1-14.
45. Mena GE, Martinez PP, Mahmud AS, Marquet PA, Buckee CO, Santillana M. Socioeconomic status determines COVID-19 incidence and related mortality in Santiago, Chile. *Science*. 2021;37.(6545)2
46. Modig K, Ahlbom A, Ebeling M. Excess mortality from COVID-19: weekly excess death rates by age and sex for Sweden and its most affected region. *European journal of public health*. 2021;31(1):17-22.
47. Michelozzi P, De'Donato F, De Sario M, Stortichini M, Stafoggia M, Niccioli F, et al. Temporal variations in excess mortality during phase 1 and phase 2 of the COVID-19 epidemic in Italy. *Epidemiologia e prevenzione*. 2020;44(5-6 Suppl 2):236-43.
48. Stokes AC, Lundberg DJ, Elo IT, Hempstead K, Bor J, Preston SH. COVID-19 and excess mortality in the United States: A county-level analysis. *PLoS Medicine*. 2021;18(5).
49. Taylor L. Covid-19: Why Peru suffers from one of the highest excess death rates in the world. *The BMJ*. 2021;372.
50. SoleimanvandiAzar N, Irandoost SF, Ahmadi S, Xosravi T, Ranjbar H, Mansourian M, et al. Explaining the reasons for not maintaining the health guidelines to prevent COVID-19 in high-risk jobs: a qualitative study in Iran. *BMC public health*. 2021;21(1):1-15.
51. Zamanian M, Ahmadi D, Sindarreh S, Aleebrahim F, Vardanjani HM, Faghihi SH, et al. Fear and rumor associated with COVID-19 among Iranian adults, 2020. *Journal of Education and Health Promotion*. 2020;9.
52. Adham D, Habibzadeh S, Ghobadi H, Jajin SA, Abbasi-Ghahramanloo A, Moradi-Asl E. Epidemiological characteristics and mortality risk factors among COVID-19 patients in Ardabil, Northwest of Iran. *BMC Emergency Medicine*. 2021;21(1):67.
53. Quast T, Andel R. Excess Mortality Associated With COVID-19 by Demographic Group: Evidence From Florida and Ohio. *Public Health Reports*. 2021:00333549211041550.
54. Poustchi H, Darvishian M, Mohammadi Z, Shayanrad A, Delavari A, Bahadorimonfared A, et al. SARS-CoV-2 antibody seroprevalence in the general population and high-risk occupational groups across 18 cities in Iran: a population-based cross-sectional study. *The Lancet Infectious Diseases*. 2021;21(4):473-81.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supp.table1.docx](#)