

Mental health, smoking, harm reduction and quit attempts – a population survey in England

Leonie Brose (✉ leonie.brose@kcl.ac.uk)

<https://orcid.org/0000-0001-6503-6854>

Jamie Brown

University College London

Debbie Robson

Institute of Psychiatry, Psychology and Neuroscience, King's College London

Ann McNeill

Institute of Psychiatry, Psychology and Neuroscience, King's College London

Research article

Keywords: Tobacco smoking, smoking cessation, mental health, electronic cigarettes, harm reduction

Posted Date: March 9th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-16310/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on August 14th, 2020. See the published version at <https://doi.org/10.1186/s12889-020-09308-x>.

Abstract

Background Tobacco control strategies have led to an overall decline in smoking; however, a large gap remains between people with and without mental health problems, leading to substantial health inequalities. We aimed to assess mental health status of cigarette smokers and recent ex-smokers in England, and smoking and harm reduction behaviour and quit attempts by mental health status.

Methods Data were collected from 5,637 current and 434 recent ex-smokers in 2016/17 in household surveys of representative samples of adults. We calculated weighted prevalence of different indicators of mental health problem: a) ever diagnosis, b) none, moderate or serious past-month distress, c) past-year treatment. We compared weighted smoking status, cigarette type, dependence, motivation to stop smoking, cutting down, use of nicotine replacement therapy or e-cigarettes, short-term abstinence, and quit attempts according to mental health status.

Results Among smokers and recent ex-smokers, 35.9% ever had a diagnosis, 24.3% had experienced moderate, an additional 9.7% serious, past-month distress; 21.9% had had past-year treatment. Those with an indication of a mental health problem were more highly dependent and more likely to smoke roll-your-own cigarettes but also more likely to be motivated to stop smoking, to cut down, use nicotine replacement therapy or e-cigarettes and to have attempted to quit in the past year.

Conclusions About a third of cigarette smokers in England have mental health problems. They are more dependent on cigarettes but also more likely to engage in harm reduction behaviours, more motivated to stop and more likely to attempt to stop smoking.

Introduction

There are substantial inequalities in morbidity and premature mortality between individuals with mental health problems and those without and much of this inequality is due to smoking (1-3). Tobacco control strategies have led to an overall decline in smoking prevalence in countries such as the UK and US, however, there remains a large gap in smoking prevalence between people with and without mental health problems. Among those with a common mental health disorder in England, smoking prevalence remains around 50% higher (4) and this increases further for more severe mental disorders (5). Figures for the US similarly show an increased burden of smoking among those with mental health problems (6, 7). The relationship is complex but appears to be bidirectional (8-10).

Previous research has consistently shown that smokers with mental health problems on average smoke more heavily (4, 5, 7) and extract more nicotine from each cigarette than those without mental health problems (5). People with schizophrenia, bipolar disorder, depression and anxiety who smoke experience more severe symptoms and require higher doses of some psychotropic medicines than non-smokers (5). Smoking is also associated with poorer treatment outcomes; compared with non-smokers with mental health problems, smokers spend longer time in hospital and less time out of hospital (11). There is some evidence, mainly from the US, that they are as likely to make a quit attempt than other smokers (12) but cessation rates are lower. Successful quitting is associated with improved mental health, specifically reduced anxiety, depression and stress and increased psychological quality of life and positive affect (13); additionally, for specific medications for severe mental illness, quitting may reduce the required dosage (5) and thus associated side effects.

Preventing the uptake of smoking, prompting more quit attempts and improving access to smoking cessation or harm reduction interventions are needed to reduce the excess burden of disease and premature mortality associated with smoking among people with mental health problems. Although research about the effectiveness of smoking cessation interventions among this group is promising, advancing their care and treatment is limited, particularly in England and the rest of the UK, by a lack of population level studies about smoking prevalence, motivation to quit, attempts to quit and harm reduction behaviours.

This study aimed to answer the following questions:

1. What is the mental health status of cigarette smokers and recent ex-smokers in England?
2. What are the characteristics of smoking behaviour among cigarette smokers and recent ex-smokers with and without mental health problems?
3. What are the characteristics of harm reduction behaviour among cigarette smokers with and without mental health problems?
4. What is the frequency of quit attempts in cigarette smokers and recent ex-smokers with and without mental health problems?

Methods

Procedure

The present study uses self-reported data from monthly cross-sectional household surveys of representative samples of the population of adults in England collected as part of the ongoing Smoking and Alcohol Toolkit Study. The methodology for the Toolkit Study has been described in detail by Fidler et al (14) and to date, it has been used as basis for approximately 85 peer-reviewed publications. Briefly, each month (wave), a new sample of approximately 1700 adults aged 16 years of age or older in England completes a face-to-face computer-assisted survey. After stratification by geo-demographic classification of population, small geographical areas containing approximately 300 households are allocated randomly to interviewers who visit households within the locality and conduct computer-assisted face-to-face interviews with one member of a household in those areas until a pre-specified quota tailored to the area is fulfilled. This form of sampling has benefits over conventional quota sampling, because the allocation of small areas to interviewers and reduces the impact of selection bias resulting from the selection of properties (15). Response rates cannot be calculated because of the lack of a definitive gross sample: all units fulfilling the criteria of a given quota are interchangeable within the areas. For the present study, measures of mental health were added to the survey for 24 monthly waves from January 2016 to December 2017.

Measures

All measures were self-reported. Generally, questions were completed in an interview with the interviewer recording the responses; the mental health questions were self-completed by the interviewee on a laptop following familiarisation using similar example questions.

Sociodemographic measures included age (16-24; 25-34; 35-44; 45-54; 55-64; 65 and over), gender and occupational grade (coded AB for higher or intermediate managerial, administrative or professional occupation; C1 for Supervisory or clerical and junior managerial, administrative or professional; C2 for skilled manual workers; D for semi-skilled and unskilled manual workers; E for state pensioners, casual and lowest grade workers, unemployed with state benefits only (16)).

Smoking-related measures included smoking status, assessed using "Which of the following best applies to you? Please note we are referring to cigarettes and other kinds of tobacco that you set light to and NOT electronic or 'heat-not-burn' cigarettes. a) I smoke cigarettes (including hand-rolled) every day; b) I smoke cigarettes (including hand-rolled), but not every day; c) I do not smoke cigarettes at all, but I do smoke tobacco of some kind (e.g. pipe, cigar or shisha); d) I have stopped smoking completely in the last year; e) I stopped smoking completely more than a year ago; f) I have never been a smoker (i.e. smoked for a year or more)." Those who had stopped smoking more than a year ago (response e) or had never been smokers (f) were excluded. Those selecting responses a, b or c were categorised as current smokers; current smokers and those who had stopped in the last year (a, b, c or d) were categorised as past-year smokers. For the present study, 186 smokers who reported currently smoking other tobacco products (c) were excluded. Those who had stopped smoking may have included a small number of respondents who had smoked products other than cigarettes. Current cigarette smokers reported smoking cigarettes (manufactured or hand-rolled), type of cigarette smoked (hand-rolled, manufactured or both), strengths of urges to smoke (17), heaviness of smoking index (cigarettes per day and time to first cigarette (18)), motivation to stop smoking (19).

Harm reduction measures included currently attempting to cut down on the number of cigarettes, current use of nicotine replacement therapy (NRT), current use of e-cigarettes/vaping products (EC), number of attempts to quit smoking in the past year, whether abstinence from smoking of one month or longer was achieved during the past year.

Mental health measures included three indicators of mental health problems:

1. Ever diagnosis: Participants were asked: "Since the age of 16, which of the following, if any, has a doctor or health professional ever told you that you had? Depression; Anxiety; Obsessive Compulsive Disorder; Panic Disorder or a phobia; Post-traumatic Stress Disorder; Psychosis; Personality Disorder; Attention Deficit Hyperactivity Disorder; An Eating Disorder; Alcohol Misuse or Dependence; Drug Use or Dependence; Problem Gambling; None of these; Don't know; Prefer not to say." Question wording was adapted from previous surveys (20, 21), the list of diagnosis based on a report on adult psychiatric morbidity in England (22). Response options excluding the final three were presented in a randomised order. Individual diagnoses were dummy-coded. For analysis, a composite measure of 'Any diagnosis' was derived.
2. Past-month distress: Measured using the K6 screener for mental distress in the past 30 days (23, 24): "During the past 30 days, about how often, if at all, did you feel... nervous; hopeless; restless or fidgety; so depressed that nothing could cheer you up; that everything was an effort; worthless?" The options are presented in a randomised order and for each the respondent indicates one of the following: "All of the time (scored 4); Most of the time (3); Some of the time (2); A little of the time (1); None of the time (0)" these options are randomly presented forward or reverse. Additional response options "Don't know; Prefer not to say" are provided. The K6 has been validated and used in a number of population surveys (23-33). A sum score with a possible range from 0 to 24 was calculated and following common practice, scores of 13 and higher categorised as serious distress (23, 27). Some publications have additionally used a category of moderate distress (scores 5-12) (29) which was also used.
3. Past-year treatment: Respondents who selected any of the responses for ever diagnosis were asked for those they selected: "In the last 12 months, which of the following conditions, if any, have you had any treatment or taken any prescribed medication for?".

Sample selection

In 2016 and 2017, 40,831 adults were surveyed of whom 7,651 were past-year (current and recent-ex) smokers who were asked the mental health questions. Smokers who exclusively smoked tobacco products (pipes, cigars) other than cigarettes were excluded (n=186). Those who did not complete the mental health questions or selected 'don't know' or 'prefer not to say' in response to any of them were also excluded, leaving 6,280 past-year smokers with information on mental health. Finally, those with missing data on any of the other variables included in the present analysis were excluded, leaving 6,071 past-year smokers for the main analyses, of whom 5,637 were current cigarette smokers and 434 recent ex-smokers. Type of cigarette smoked was missing for 164, reducing the sample to 5,907 past-year smokers and 5,506 current cigarette smokers if type of cigarette was included in analysis.

Analysis

To address research question 1 (mental health), weighted proportions were used to describe the prevalence of ever having had any mental health diagnosis since the age of 16, the prevalence of moderate and serious distress in the past month and the prevalence of treatment for mental health problems in the past year overall and by diagnosis. All three indicators were also calculated by age, gender and occupational grade.

To address research question 2 (smoking behaviour by mental health), weighted proportions were calculated and chi-square statistics were used to compare smoking status, type of cigarette smoked, dependence (as measured using urges to smoke and heaviness of smoking) of past-year cigarette smokers with and without any diagnosis, with minimal, moderate and serious past-month distress and with and without past-year treatment. Additionally, for current cigarette smokers, motivation to stop smoking was compared using chi-square statistics and spend on tobacco were compared using ANOVAs and independent samples median tests.

To address research question 3 (harm reduction behaviour), weighted proportions for cutting down, using e-cigarettes, using NRT, using either e-cigarettes or NRT, and having achieved at least one month of abstinence in the past year were calculated for current cigarette smokers and compared using chi-square statistics.

To address research question 4 (quit attempts), the mean number of quit attempts made in the past year was compared among past-year smokers with and without mental health problems. A dichotomised version of this measure (at least one quit attempt versus none) was used to calculate prevalence and 95% confidence intervals of having made at least one quit attempt and as outcome in logistic regressions assessing its association with ever diagnosis, past-month levels of distress and past-year treatment among past-year smokers. Unadjusted models were followed by models adjusting for socio-demographics. Additional models adjusted for motivation to stop smoking and type of cigarette; as this information can only be collected from current cigarette smokers, any quit attempt had been unsuccessful.

The analyses plan was pre-registered on the Open Science Framework (<https://osf.io/9pbv8/>). The following changes were made from the registration: Research question 1 was added to provide a fuller description of the population. The present research questions 2 and 3 were initially presented as a single research question. To increase consistency, smokers of other tobacco products such as pipes or cigars were excluded from all analysis.

Results

Table 1 describes sample characteristics overall and by the three indicators of mental health status.

1. Mental health

Overall, 35.9% reported ever having been diagnosed with at least one mental health problem since the age of 16, 34.0% reported having experienced moderate (24.3%) or serious distress (9.7%) in the past month and 21.9% of the sample had received treatment for a mental health in the past year (Table 1). Among those who had ever been diagnosed with a mental health problem, the most common diagnosis was depression, followed by anxiety (Supplemental Table S1). In the past year, 61.0% of those with a diagnosis had received treatment for this diagnosis, however, this ranged from 7.9% for problem gambling to 64.1% for psychosis (Supplemental Table S1). In general, indicators of mental health problems were more common among younger age groups, women and lower socio-economic groups (Table 2).

2. Smoking

About 7% of the sample reported having stopped smoking completely in the past year. Most participants were daily cigarette smokers, and there was little difference between those with and without mental health problems. Overall, about half smoked roll-your own exclusively (44.1%) or alongside manufactured cigarettes (5.4%) and this was more common among cigarette smokers with mental health problems. The vast majority of smokers and recent ex-smokers experienced at least some urges to smoke although few had high heaviness of smoking ratings. Stronger urges and heavier smoking were more common among those with mental health problems (Table 2).

Among current cigarette smokers, only about one in seven (14.4%) reported motivation to stop smoking within the next three months. This was slightly more common among those with any measure of mental health problems, but this was associated with very small effect sizes (Table 2).

Current cigarette smokers with data on spend available (n=5,421) on average spent £22.61 (SD=17.09) per week on tobacco or cigarettes (median £20). This was lower among those with any diagnosis or distress, but not by treatment (Supplemental Table S2), likely in part due to higher rates smoking roll-your own.

3. Harm reduction

Overall, 46.1% of smokers were cutting down, 19.4% were using e-cigarettes and 8.5% nicotine replacement therapy (Table 2). All three harm reduction behaviours were more common among smokers with a mental health problem, albeit all with very small effect sizes (Table 2). In the last year, 12.9% had achieved at least one month of abstinence. This did not differ by ever diagnosis or past-year treatment but by level of distress; those with minimal distress were least likely and those with moderate distress most likely to have achieved a month of abstinence, again with a very small effect size (Table 2).

4. Quit attempts

Just under a third (32.5%) of those who smoked in the past year had made at least one quit attempt in that time. In unadjusted analysis, prevalence of having made at least one quit attempt was higher among past-year smokers with any diagnosis (36.2%, 95% CI: 34.2-38.1; unadjusted OR=1.36, 95% CI: 1.22–1.53, $p<0.001$) than among those without diagnosis (30.4, 95% CI: 29.0-31.8). Prevalence of having made at least one quit attempt increased with distress, from 29.6% (95% CI: 28.2 – 31.0) among those with no or minimal distress through 37.7% (95% CI: 35.3 – 40.1; unadjusted OR=1.45, 95% CI: 1.28 – 1.64, $p<0.001$) among those with moderate distress to 38.8% (95% CI: 35.0-42.7; unadjusted OR=1.54, 95% CI: 1.28 – 1.84, $p<0.001$) among those with serious distress. Treatment in the past year was also associated with higher prevalence of having made a quit attempt (38.3%; 95% CI: 35.7-40.8; unadjusted OR=1.44, 95% CI: 1.27 – 1.64, $p<0.001$) compared with not having had treatment 30.8% (95% CI: 29.6- 32.1). The strength of these associations remained very similar in adjusted analysis (Table 3, 4, 5). When including only current smokers, smokers with any diagnosis (Table 3), past-month distress (Table 4) or past-year treatment (Table 5) were also more likely to have made a quit attempt than smokers without these indicators of a mental health problem.

Discussion

Different indicators of mental health problems showed that substantial proportions of cigarette smokers and recent ex-smokers in England have mental health problems. Just over a third reported ever having had a diagnosis of a mental health problem and around 60% of them had received treatment in the last year, suggesting an ongoing issue which was also indicated by just over a third experiencing moderate or serious mental distress in the past month. Those with an indication of a mental health problem were more highly dependent and more likely to smoke roll-your-own cigarettes. Cigarette smokers with a mental health problem were however more likely to be motivated to stop smoking and to engage in in harm reduction behaviours such as use of e-cigarettes or NRT. Cigarette smokers and recent ex-smokers with mental health problems were more likely to have made at least one quit attempt in the past year and this remained true when including only current smokers (i.e. where any quit attempts had been unsuccessful). Yet, there was little difference by mental health in having achieved at least short-term abstinence among all past-year smokers, suggesting that quit attempts may be less successful in those with mental health problems.

The prevalence of mental health problems found in this population survey is in line with previous estimates that 30% of smokers in the UK have evidence of mental disorder (5) and an estimate that 27.4% of adults in England have ever been diagnosed with a common mental health problem (generalised anxiety disorder, depressive episode, obsessive compulsive disorder, phobias, panic disorder (34), omitting groups of less common diagnoses included in the present estimate). The level of moderate and severe distress found in past-year smokers in England is similar to those from a household survey in the US which found that in 2015, 25.8% of current cigarette smokers had moderate and 9.4% serious distress in the past month (1997-2015 combined: 22.6% and 7.0%) (12). Both a US survey (12, 28) and the present survey found that those with psychological distress were more likely to have made at least one quit attempt in the past year. The present finding that those with a mental health problem are at least as likely to be motivated to stop smoking is in line with earlier findings which used different measures to measure related constructs (4, 28, 35). Previous research from the US found that current cigarette smokers with mental health problems were not statistically more likely to be using electronic cigarettes, this is in contrast to the present findings (36, 37).

The present study had several limitations. Timing or causality of events cannot be established; it could for example be argued that an unsuccessful quit attempt in the past-year could have increased distress and even increased the need for treatment for a mental health problem. However, associations were very similar for the different indicators of mental health problems, making this direction of causality appear less likely. All measures were self-reported; more comprehensive instruments may have differed in their detection of mental health problems. Because the survey is a household survey, it excludes people who are too unwell to take part or in healthcare institutions, thereby excluding those with the most severe mental health problem. Because respondents were asked to select the single option that best describes their smoking, use of other tobacco products such as pipes, cigars or shisha remained unreported among cigarette smokers. However, a strength is that data came from a long-standing established population survey and a sample representative of the adult population in England.

The present study is the first to provide detailed population-level evidence on motivation to quit smoking, attempts to do so and harm reduction behaviours of smokers in England by mental health status. It is also the first to report that smokers with a mental health problem are more likely to engage in harm reduction behaviours compared to those without.

Some implications for policy and clinical practice can be drawn from the findings. The relatively higher levels of motivation, attempts to stop smoking and harm reduction behaviours such as e-cigarette use among smokers with mental health problems compared to those without should be capitalised on by clinicians to reduce the burden from smoking. Specific challenges such as higher dependence on tobacco need to be addressed by making effective support available. A combination of behavioural support and medication is most effective in the wider population of smokers and has been shown to be effective for smokers with severe mental illness (38, 39). In quit attempts and accompanied by behavioural support, e-cigarettes have been shown to be more effective than NRT (40). Additionally, smaller studies in smokers with severe mental illness who were not interested in quitting found that substantial proportions reduced their smoking by at least 50% when supplied with e-cigarettes (41, 42). Clinicians should ensure that patients (both inpatient and community) have their smoking status identified and recorded in health care records. All smokers irrespective of their mental health status should be advised about the most effective way to quit and either provided with treatment or referred for specialist stop smoking support (43). Ensuring harm reduction strategies are accessible should also be considered. Recently in England, the National Health Service has committed to funding tobacco treatment services for all smokers admitted to hospital, including the option to switch to e-cigarettes for mental health patients (44). Mass media campaigns could specifically focus on benefits of smoking cessation for mental health and the effectiveness of support and be tested to ensure that they have good reach and effectiveness among smokers with mental health problems.

Future research needs to assess differences in access to support, outcomes of quit attempts by mental health status and the effectiveness of different types of support including prescription medication, behavioural support and e-cigarettes to identify how differences in quit rates between those with and without mental health problems can be reduced. Associations for specific mental health problems or groups of problems should also be assessed in addition to a catch-all category of any diagnosis.

Conclusions

About a third of smokers in England have mental health problems. This population is more likely to be young, female and poor. They are more dependent on tobacco smoking but also more likely to engage in harm reduction behaviours, more motivated and more likely to attempt to stop smoking.

Declarations

Ethics approval and consent to participate

Ethics approval for the Toolkit Study was originally granted by the University College London Ethics Committee (ID 0498/001). Addition of the mental health questions to the STS was approved by the University College London Research Ethics Committee (ID: 2808/005). All participants provided informed consent.

Consent for publication

Not applicable

Availability of data and materials

All authors have had access to the data. The datasets analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare no competing interests.

Funding

This work was supported by a Cancer Research UK Fellowship awarded to LB (C52999/A19748). For data collection for the Smoking Toolkit Study, Cancer Research UK is the main contributor (C1417/A22962), but the UK Department of Health, Pfizer, GlaxoSmithKline and Johnson and Johnson have also all contributed funding. JB receives support from CRUK (C1417/A22962).

All authors are members of SPECTRUM, a consortium supported by the UK Prevention Research Partnership (MR/S037519/1), which is funded by the British Heart Foundation, Cancer Research UK, Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Health and Social Care Research and Development Division (Welsh Government), Medical Research Council, National Institute for Health Research, Natural Environment Research Council, Public Health Agency (Northern Ireland), The Health Foundation and the Wellcome Trust. The funders had no role on the study design, analysis or interpretation of the data, writing of the report or decision to submit the paper for publication.

Professor McNeill is a National Institute for Health Research (NIHR) Senior Investigator. The views expressed in this article are those of the authors and not necessarily those of the NIHR, or the Department of Health and Social Care.

Authors' contributions

All authors contributed to formulating the research question and designing the study. LB conducted the data analysis and drafted the article, all authors contributed to writing it.

Acknowledgements

We thank all survey participants. We thank Professor Robert West for his contribution to the analysis plan.

References

1. Tam J, Warner KE, Meza R. Smoking and the Reduced Life Expectancy of Individuals With Serious Mental Illness. *Am J Prev Med.* 2016;51(6):958-66.
2. Wahlbeck K, Westman J, Nordentoft M, Gissler M, Laursen TM. Outcomes of Nordic mental health systems: life expectancy of patients with mental disorders. *The British journal of psychiatry : the journal of mental science.* 2011;199(6):453-8.
3. Lawrence D, Hancock KJ, Kisely S. The gap in life expectancy from preventable physical illness in psychiatric patients in Western Australia: retrospective analysis of population based registers. *BMJ (Clinical research ed.)* 2013;346:f2539.
4. Richardson S, McNeill A, Brose LS. Smoking and quitting behaviours by mental health conditions in Great Britain (1993-2014). *Addictive behaviors.* 2019;90:14-9.

5. Royal College of Physicians, Royal College of Psychiatrists. Smoking and mental health. London: RCP; 2013.
6. Smith PH, Chhipa M, Bystrick J, Roy J, Goodwin RD, McKee SA. Cigarette smoking among those with mental disorders in the US population: 2012-2013 update. *Tobacco control*. 2018.
7. Cook BL, Wayne GF, Kafali EN, Liu Z, Shu C, Flores M. Trends in smoking among adults with mental illness and association between mental health treatment and smoking cessation. *JAMA*. 2014;311(2):172-82.
8. Breslau N, Peterson EL, Schultz LR, Chilcoat HD, Andreski P. Major depression and stages of smoking. A longitudinal investigation. *Arch Gen Psychiatry*. 1998;55(2):161-6.
9. Johnson JG, Cohen P, Pine DS, Klein DF, Kasen S, Brook JS. Association between cigarette smoking and anxiety disorders during adolescence and early adulthood. *JAMA*. 2000;284(18):2348-51.
10. Lipari RN, Van Horn S. Smoking and Mental Illness Among Adults in the United States. The CBHSQ Report. Rockville (MD)2013. p. 1-
11. Prochaska JJ, Hall SE, Delucchi K, Hall SM. Efficacy of initiating tobacco dependence treatment in inpatient psychiatry: a randomized controlled trial. *American journal of public health*. 2014;104(8):1557-65.
12. Kulik MC, Glantz SA. Softening Among U.S. Smokers With Psychological Distress: More Quit Attempts and Lower Consumption as Smoking Drops. *Am J Prev Med*. 2017;53(6):810-7.
13. Taylor G, McNeill A, Girling A, Farley A, Lindson-Hawley N, Aveyard P. Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ (Clinical research ed)*. 2014;348:g1151.
14. Fidler JA, Shahab L, West O, Jarvis MJ, McEwen A, Stapleton JA, et al. 'The smoking toolkit study': a national study of smoking and smoking cessation in England. *BMC public health*. 2011;11:479.
15. de Vaus DA. *Surveys in social research*. 5th ed. Crows Nest, Australia: Allen & Unwin; 2002.
16. National Readership Survey. National Readership Survey social-grades. www.nrs.co.uk/nrs-print/lifestyle-and-classification-data/social-grade. 2016.
17. Fidler JA, Shahab L, West R. Strength of urges to smoke as a measure of severity of cigarette dependence: comparison with the Fagerstrom Test for Nicotine Dependence and its components. *Addiction (Abingdon, England)*. 2011;106(3):631-8.
18. Kozlowski LT, Porter CQ, Orleans CT, Pope MA, Heatherton T. Predicting smoking cessation with self-reported measures of nicotine dependence: FTQ, FTND, and HSI. *Drug and alcohol dependence*. 1994;34(3):211-6.
19. Kotz D, Brown J, West R. Predictive validity of the Motivation To Stop Scale (MTSS): a single-item measure of motivation to stop smoking. *Drug and alcohol dependence*. 2013;128(1-2):15-9.
20. Morris CD, Burns EK, Waxmonsky JA, Levinson AH. Smoking cessation behaviors among persons with psychiatric diagnoses: Results from a population-level state survey. *Drug and alcohol dependence*. 2014;136:63-8.
21. The NHS Information Centre. Adult psychiatric morbidity in England, 2007 - Results of a household survey. Appendices and Glossary. Leeds: The NHS Information Centre; 2009.
22. The NHS Information Centre for health and social care. Adult psychiatric morbidity in England, 2007. Results of a household survey. Leeds: The NHS Information Centre; 2009.
23. Kessler RC, Barker PR, Colpe LJ, Epstein JF, Gfroerer JC, Hiripi E, et al. Screening for serious mental illness in the general population. *Arch Gen Psychiatry*. 2003;60(2):184-9.
24. Kessler RC, Green JG, Gruber MJ, Sampson NA, Bromet E, Cuitan M, et al. Screening for serious mental illness in the general population with the K6 screening scale: results from the WHO World Mental Health (WMH) survey initiative. *International journal of methods in psychiatric research*. 2010;19 Suppl 1:4-22.
25. Aldworth J, Colpe LJ, Gfroerer JC, Novak SP, Chromy JR, Barker PR, et al. The National Survey on Drug Use and Health Mental Health Surveillance Study: calibration analysis. *International journal of methods in psychiatric research*. 2010;19 Suppl 1:61-87.

26. Furukawa TA, Kessler RC, Slade T, Andrews G. The performance of the K6 and K10 screening scales for psychological distress in the Australian National Survey of Mental Health and Well-Being. *Psychological medicine*. 2003;33(2):357-62.
27. Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand SL, et al. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychological medicine*. 2002;32(6):959-76.
28. Lawrence D, Mitrou F, Zubrick SR. Non-specific psychological distress, smoking status and smoking cessation: United States National Health Interview Survey 2005. *BMC public health*. 2011;11:256.
29. Prochaska JJ, Sung HY, Max W, Shi Y, Ong M. Validity study of the K6 scale as a measure of moderate mental distress based on mental health treatment need and utilization. *International journal of methods in psychiatric research*. 2012;21(2):88-97.
30. Rush B, Castel S, Brands B, Toneatto T, Veldhuizen S. Validation and comparison of diagnostic accuracy of four screening tools for mental disorders in people seeking treatment for substance use disorders. *Journal of substance abuse treatment*. 2013;44(4):375-83.
31. Streck JM, Weinberger AH, Pacek LR, Gbedemah M, Goodwin RD. Cigarette smoking quit rates among persons with serious psychological distress in the United States from 2008-2016: Are mental health disparities in cigarette use increasing? *Nicotine Tob Res*. 2018.
32. Cornelius BL, Groothoff JW, van der Klink JJ, Brouwer S. The performance of the K10, K6 and GHQ-12 to screen for present state DSM-IV disorders among disability claimants. *BMC public health*. 2013;13:128.
33. Swartz JA, Lurigio AJ. Screening for serious mental illness in populations with co-occurring substance use disorders: Performance of the K6 scale. *Journal of substance abuse treatment*. 2006;31(3):287-96.
34. McManus S, Bebbington P, Jenkins R, Brugha T (eds). *Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014*. Leeds: NHS Digital; 2016.
35. Siru R, Hulse GK, Tait RJ. Assessing motivation to quit smoking in people with mental illness: a review. *Addiction (Abingdon, England)*. 2009;104(5):719-33.
36. Spears CA, Jones DM, Weaver SR, Pechacek TF, Eriksen MP. Use of Electronic Nicotine Delivery Systems among Adults with Mental Health Conditions, 2015. *Int J Environ Res Public Health*. 2016;14(1).
37. Cummins SE, Zhu SH, Tedeschi GJ, Gamst AC, Myers MG. Use of e-cigarettes by individuals with mental health conditions. *Tobacco control*. 2014;23 Suppl 3:iii48-53.
38. Gilbody S, Peckham E, Bailey D, Arundel C, Heron P, Crosland S, et al. Smoking cessation for people with severe mental illness (SCIMITAR+): a pragmatic randomised controlled trial. *The lancet Psychiatry*. 2019;6(5):379-90.
39. Peckham E, Brabyn S, Cook L, Tew G, Gilbody S. Smoking cessation in severe mental ill health: what works? an updated systematic review and meta-analysis. *Bmc Psychiatry*. 2017;17(1):252.
40. Hajek P, Phillips-Waller A, Przulj D, Pesola F, Myers Smith K, Bisal N, et al. A Randomized Trial of E-Cigarettes versus Nicotine-Replacement Therapy. *N Engl J Med*. 2019.
41. Caponnetto P, Auditore R, Russo C, Cappello GC, Polosa R. Impact of an electronic cigarette on smoking reduction and cessation in schizophrenic smokers: a prospective 12-month pilot study. *Int J Environ Res Public Health*. 2013;10(2):446-61.
42. Hickling LM, Perez-Iglesias R, McNeill A, Dawkins L, Moxham J, Ruffell T, et al. A pre-post pilot study of electronic cigarettes to reduce smoking in people with severe mental illness. *Psychological medicine*. 2019;49(6):1033-40.
43. National Institute for Health and Care Excellence. Stop smoking interventions and services. <https://www.nice.org.uk/guidance/ng92>. 2018.
44. NHS England. The NHS Long Term Plan. <https://www.longtermplannhs.uk/>. 2019;accessed 20 Aug 2019.

Tables

Table 1. Weighted prevalence of ever diagnoses, past-month distress and past-year treatment in past-year smokers (unweighted n=6071) and the subset of current smokers (unweighted n=5637)

	Past-year cigarette smokers, %	Current cigarette smokers, %
Ever diagnosis	35.9	35.6
Past-month distress		
None/minimal	66.0	66.3
Moderate	24.3	24.0
Serious	9.7	9.7
Past-year treatment	21.9	21.6

Table 2. Sample description (weighted data) by indicator of mental health problem in past-year smokers (unweighted n=6079) and the subset of current smokers (unweighted n=5637)

	All	Ever diagnosis			Past-month distress				Past-year treatment			
		No	Yes	χ^2 , p, Effect size	Minimal	Moderate	Serious	χ^2 , p, Effect size	No	Yes	χ^2 , p, Effect size	
Past-year cigarette smokers												
Age, %	16-24	18.8	18.4	19.6	$\chi^2=87.890$, p<0.001, V=0.118	15.1	25.9	26.1	$\chi^2=207.65$, p<0.001, V=0.128	18.7	19.2	$\chi^2=74.83$, p<0.001, V=0.109
	25-34	22.5	21.7	24.1		21.8	23.7	24.6		22.2	23.7	
	35-44	18.2	17.1	20.0		17.6	19.2	19.2		17.3	21.3	
	45-54	18.3	17.7	19.5		19.3	15.7	18.1		17.6	20.9	
	55-64	11.9	12.4	11.2		13.4	9.0	9.6		12.4	10.2	
	≥65	10.2	12.8	5.7		12.8	6.5	2.3		11.8	4.7	
Gender, % ¹	Men	52.4	57.6	43.0	$\chi^2=124.73$, p<0.001, V=0.141	55.5	48.2	41.2	$\chi^2=58.32$, p<0.001, V=0.096	56.2	38.7	$\chi^2=132.23$, p<0.001, V=0.145
	Women	47.6	42.4	57.0		44.5	51.8	58.8		43.8	61.3	
Occupational grade, %	AB	14.9	16.0	13.1	$\chi^2=209.54$, p<0.001, V=0.182	16.3	13.2	10.4	$\chi^2=235.35$, p<0.001, V=0.136	15.8	11.7	$\chi^2=253.20$, p<0.001, V=0.200
	C1	24.5	25.1	23.5		24.9	25.2	20.4		25.4	21.5	
	C2	25.7	27.8	21.9		27.7	23.0	18.6		27.3	20.0	
	D	20.2	21.2	18.3		20.7	18.7	19.9		20.5	18.8	
	E	14.7	9.9	23.2		10.4	19.9	30.7		11.0	27.9	
Smoking, %	Daily cigs	81.6	82.0	81.0	$\chi^2=3.58$, p=0.167, V=0.024	82.7	78.8	82.1	$\chi^2=11.44$, p=0.022, V=0.030	81.6	81.8	$\chi^2=8.60$, p=0.014, V=0.037
	Non-daily cigs	10.9	11.0	10.7		10.3	12.6	10.8		11.3	9.3	
	Stopped last year	7.5	7.0	8.3		7.0	8.7	7.2		7.1	8.8	
Type of cigarette, % ²	Manufactured	50.4	56.2	40.2	$\chi^2=146.15$, p<0.001, V=0.154	55.3	43.6	34.1	$\chi^2=131.67$, p<0.001, V=0.103	53.6	39.3	$\chi^2=87.92$, p<0.001, V=0.120
	Roll-your-own	44.1	38.8	53.7		40.0	50.1	57.8		41.5	53.6	
	Mix	5.4	5.0	6.1		4.7	6.3	8.1		4.9	7.2	
Urges to smoke, %	None	16.9	18.9	13.5	$\chi^2=117.41$, p<0.001, V=0.136	18.4	14.8	12.6	$\chi^2=179.27$, p<0.001, V=0.12	18.2	12.3	$\chi^2=111.32$, p<0.001, V=0.133
	1	16.3	17.4	14.4		17.7	14.4	11.7		17.0	13.8	
	2	43.3	43.6	42.6		44.1	43.6	36.7		43.7	41.9	
	3	17.3	15.8	20.0		15.6	19.2	24.0		16.2	21.1	
	4	4.5	3.4	6.4		3.2	5.6	10.4		3.6	7.5	
	5	1.7	0.9	3.0		1.0	2.3	4.6		1.2	3.4	
HSL, %	Low (<4)	87.5	90.4	82.4	$\chi^2=84.35$, p<0.001, V=0.116	89.8	85.2	77.7	$\chi^2=82.78$, p<0.001, V=0.114	89.4	80.7	$\chi^2=75.31$, p<0.001, V=0.109
	High (4+)	12.5	9.6	17.6		10.2	14.8	22.3		10.6	19.3	
≥1 quit attempt, % ³	Yes	32.5	30.4	36.2	$\chi^2=22.00$, p<0.001, V=0.059	29.6	37.7	38.8	$\chi^2=46.28$, p<0.001, V=0.086	30.8	38.3	$\chi^2=27.05$, p<0.001, V=0.065
Abstinence ≥1 month, %	Yes	12.9	12.3	13.8	$\chi^2=3.01$, p=0.083, V=0.022	11.6	16.1	13.4	$\chi^2=20.39$, p<0.001, V=0.057	12.5	14.1	$\chi^2=2.40$, p=0.122, V=0.019
Current cigarette smokers												
MTSS %	In ≤3 months	14.4	13.3	16.3	$\chi^2=9.79$, p=0.002, V=0.041	13.0	17.7	15.6	$\chi^2=19.30$, p<0.001, V=0.057	13.6	17.3	$\chi^2=11.15$, p=0.001, V=0.044
Cutting down, %	Yes	46.1	43.1	51.4	$\chi^2=37.12$, p<0.001, V=0.080	42.8	52.2	53.3	$\chi^2=49.39$, p<0.001, V=0.092	44.5	52.0	$\chi^2=22.57$, p<0.001, V=0.062
Current NRT, %	Yes	8.5	7.5	10.5	$\chi^2=15.60$, p<0.001, V=0.052	7.6	10.0	11.2	$\chi^2=13.33$, p=0.001, V=0.048	7.6	11.8	$\chi^2=22.26$, p<0.001, V=0.062
Current EC, %	Yes	19.4	17.1	23.5	$\chi^2=34.78$, p<0.001, V=0.077	16.6	25.5	23.0	$\chi^2=57.95$, p<0.001, V=0.100	18.1	23.9	$\chi^2=21.10$, p<0.001, V=0.060
Current NRT or EC, %	Yes	25.4	22.6	30.5	$\chi^2=44.64$, p<0.001, V=0.087	22.3	31.6	30.6	$\chi^2=56.02$, p<0.001, V=0.098	23.5	32.4	$\chi^2=41.13$, p<0.001, V=0.084

MTSS: Motivation to stop smoking

Past-year cigarette smokers unweighted n=6071; Current cigarette smokers, unweighted n=5637

¹ The provided third option was not used by anyone in the included sample

² Type of cigarette missing unweighted n=164

³ The median number of quit attempts was 0 for all groups; the mean number of quit attempts overall was 0.59 (SD=3.10), this was similar for those with (M=0.67; SD=3.00) and without diagnosis (M=0.55, SD=3.15; F(1,6316)=2.13, p=0.14), with (M=0.71, SD=2.82) and without past-year treatment (M=0.56, SD=3.17; F(1,6316)=2.42, p=0.112) and differed slightly across levels of distress (minimal: M=0.51,

Table 3. Adjusted associations with having made at least 1 quit attempt in the past year by presence or absence of any diagnosis (unweighted data)

		n=6071 past-year cigarette smokers				n=5506 current smokers			
		AOR	95% CI	95% CI	p	AOR	95% CI	95% CI	p
Any diagnosis	No	1				1			
	Yes	1.32	1.17	1.48	<0.001	1.25	1.09	1.43	0.001
Age	16-24	1				1			
	25-34	1.26	1.06	1.50	0.008	1.19	0.97	1.44	0.090
	35-44	1.21	1.01	1.44	0.042	1.12	0.91	1.37	0.296
	45-54	0.92	0.77	1.11	0.383	0.85	0.69	1.04	0.113
	55-64	0.84	0.69	1.02	0.081	0.79	0.63	0.98	0.033
	65 and over	0.58	0.47	0.72	<0.001	0.59	0.46	0.75	<0.001
Gender	Men	1				1			
	Women	1.04	0.93	1.16	0.504	1.03	0.91	1.17	0.66
Occupational grade	AB	1							
	C1	0.77	0.64	0.91	0.003	0.82	0.67	1.00	0.052
	C2	0.77	0.64	0.92	0.004	0.84	0.68	1.04	0.115
	D	0.62	0.51	0.76	<0.001	0.72	0.55	0.86	0.001
	E	0.63	0.52	0.77	<0.001	0.72	0.57	0.90	0.004
Motivation to stop smoking	Not in next 3 months					1			
	In next 3 months					5.30	4.51	6.22	<0.001
Type of cigarette	Manufactured					1			
	Roll your own / Mix					1.14	1.00	1.29	0.054

Table 4. Adjusted associations with having made at least 1 quit attempt in the past year by level of distress (unweighted data)

		n=6071 past-year cigarette smokers				n=5506 current smokers			
		AOR	95% CI	95% CI	p	AOR	95% CI	95% CI	p
Distress	None/minimal	1				1			
	Moderate	1.41	1.24	1.60	<0.001	1.35	1.17	1.57	<0.001
	Serious	1.51	1.25	1.81	<0.001	1.51	1.22	1.87	<0.001
Age	16-24	1				1			
	25-34	1.32	1.11	1.57	0.002	1.23	1.01	1.50	0.037
	35-44	1.26	1.05	1.51	0.012	1.16	0.95	1.43	0.152
	45-54	0.98	0.82	1.18	0.862	0.89	0.73	1.10	0.297
	55-64	0.90	0.74	1.09	0.290	0.84	0.67	1.05	0.116
	65 and over	0.62	0.50	0.77	<0.001	0.63	0.49	0.80	<0.001
Gender	Men	1				1			
	Women	1.05	0.94	1.17	0.417	1.03	0.91	1.17	0.648
Occupational grade	AB	1				1			
	C1	0.76	0.64	0.90	0.002	0.81	0.66	0.99	0.044
	C2	0.76	0.64	0.92	0.004	0.85	0.69	1.05	0.121
	D	0.62	0.51	0.75	<0.001	0.68	0.54	0.85	0.001
	E	0.61	0.50	0.75	<0.001	0.69	0.54	0.87	0.001
Motivation to stop smoking	Not in next 3 months					1			
	In next 3 months					5.24	4.46	6.16	<0.001
Type of cigarette	Manufactured					1			
	Roll your own / Mix					1.13	0.99	1.29	0.62

Table 5. Adjusted associations with having made at least 1 quit attempt in the past year by past-year treatment for any mental health problem (unweighted data).

		n=6071 past-year cigarette smokers				n=5506 current smokers			
		AOR	95% CI	95% CI	p	AOR	95% CI	95% CI	p
Any treatment	No	1				1			
	Yes	1.41	1.24	1.61	<0.001	1.37	1.18	1.60	<0.001
Age	16-24	1				1			
	25-34	1.26	1.06	1.50	0.007	1.19	0.98	1.45	0.084
	35-44	1.20	1.00	1.44	0.046	1.11	0.90	1.37	0.317
	45-54	0.92	0.77	1.10	0.355	0.84	0.68	1.03	0.100
	55-64	0.84	0.70	1.02	0.087	0.79	0.63	0.99	0.036
	65 and over	0.58	0.47	0.72	<0.001	0.59	0.46	0.75	<0.001
Gender	Men	1				1			
	Women	1.04	0.93	1.16	0.531	1.02	0.90	1.16	0.723
Occupational grade	AB	1				1			
	C1	0.77	0.64	0.91	0.003	0.82	0.67	1.00	0.051
	C2	0.76	0.64	0.92	0.004	0.84	0.68	1.04	0.106
	D	0.62	0.51	0.75	<0.001	0.68	0.54	0.85	0.001
	E	0.62	0.51	0.76	<0.001	0.70	0.55	0.88	0.002
Motivation to stop smoking	Not in next 3 months					1			
	In next 3 months					5.29	4.50	6.21	<0.001
Type of cigarette	Manufactured					1			
	Roll your own / Mix					1.14	1.00	1.30	0.051

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementalmaterial.docx](#)