

# Hypoxia-inducible Factor-1 $\alpha$ protects against Intervertebral Disc Degeneration through Antagonizing Mitochondrial Oxidative Stress.

**Wen Yang**

Department of Orthopaedics, Qilu Hospital of Shandong University

**Chunwang Jia**

Department of Orthopaedics, Qilu Hospital of Shandong University

**Long Liu**

Department of Pathology, Qilu Hospital, Cheeloo College of Medicine, Shandong University

**Yu Fu**

Department of Orthopaedics, The First Affiliated Hospital of Shandong First Medical University

**Yawei Wu**

Department of Orthopedics, Caoxian People's Hospital

**Zhicheng Liu**

The First Clinical Medical School of Shandong University

**Ruixuan Yu**

Department of Orthopaedics, Qilu Hospital of Shandong University

**Xiaojie Ma**

Department of Rheumatology and immunology, Affiliated Hospital of Shandong University of traditional Chinese Medicine

**Ao Gong**

Department of Orthopedics, Affiliated Hospital of Shandong University of traditional Chinese Medicine

**Fangming Liu**

Department of Rehabilitation Medicine, The First Affiliated Hospital of Shandong First Medical University

**Yanni Xia**

Department of Operating Room, Qilu Hospital, Cheeloo College of Medicine, Shandong University

**Yong Hou**

Department of Orthopaedics, Qilu Hospital of Shandong University

**Yuhua Li**

Department of Orthopaedics, Qilu Hospital of Shandong University

**Lei Zhang** (✉ [qygkzl1818@163.com](mailto:qygkzl1818@163.com))

Department of Orthopaedics, The First Affiliated Hospital of Shandong First Medical University

## Research Article

**Keywords:** Intervertebral disc degeneration, Hypoxia inducible factor 1 $\alpha$ , Protective role, TNF- $\alpha$ , IVDD model

**Posted Date:** May 13th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1637390/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.  
[Read Full License](#)

---

# Abstract

Intervertebral disc degeneration (IVDD) demonstrates a gradually increased incidence and has developed into a major health problem worldwide. Nucleus pulposus is characterized by the hypoxic and avascular environment, in which hypoxia inducible factor1 $\alpha$  (HIF-1 $\alpha$ ) has an important role through its participation in extracellular matrix synthesis, energy metabolism, cellular adaptation to stresses and genesis. In this study, the effects of HIF-1 $\alpha$  on mouse primary nucleus pulposus cells (MNPCs) exposed to TNF- $\alpha$  were observed and the potential mechanism was explored, and a rabbit IVDD model was established to verify the protective role of HIF-1 $\alpha$  on IVDD. In vitro results demonstrated that HIF-1 $\alpha$  could attenuate the inflammation, apoptosis and mitochondrial dysfunction induced by TNF- $\alpha$  in MNPCs, and promote cellular anabolism, and inhibit cellular catabolism. In vivo results demonstrated that after establishment of IVDD model in rabbit, disc height and IVD extracellular matrix were decreased in a time-dependent manner, and MRI analysis showed a tendency for decreased T2 values in a time-dependent manner, and supplementation of HIF-1 $\alpha$  improved histological and imaginative IVDD while down-regulation of HIF-1 $\alpha$  exacerbated this degeneration. In summary, HIF-1 $\alpha$  protected against IVDD, possibly through reducing ROS production in mitochondria and consequent inhibition of inflammation, metabolism disorders and apoptosis of MNPCs, which provided a potential therapeutic instrument for treatment of IVDD diseases.

## Introduction

As the population ages, intervertebral disc degeneration (IVDD) demonstrates a gradually increased incidence and has developed into a major health problem worldwide [1–3]. In addition to age, the risk factors of IVDD also include inflammatory cytokines, mechanical trauma, genetic susceptibility, lifestyle factors, certain metabolic disorders, and so on [4–9]. IVDD is the main cause of disability because it often causes chronic low back pain (LBP) [10, 11]. Current treatments of IVDD mainly include pharmacological and surgical interventions, aiming for managing symptoms and minimizing disability. However, both of them are costly, often result in complications and have questionable efficacy [12]. Thus, more and more studies have focused on new therapies for IVDD [13, 14].

Nucleus pulposus is characterized by the hypoxic and avascular environment, in which hypoxia inducible factor1 $\alpha$  (HIF-1 $\alpha$ ) has an important role through its participation in extracellular matrix (ECM) synthesis, energy metabolism, cellular adaptation to stresses and genesis [15, 16]. In the late-stage of IVDD, the expression of HIF-1 $\alpha$  is significantly decreased and neovascularization increases the oxygen concentration. A study reports that HIF-1 $\alpha$  can attenuate the apoptosis of nucleus pulposus derived stem cells induced by excessive mechanical load [14]. However, the effects of HIF-1 $\alpha$  on mouse primary nucleus pulposus cells (MNPCs) exposed to inflammatory cytokines and the potential mechanism are still not investigated. In this study, the effects of HIF-1 $\alpha$  on MNPCs exposed to TNF- $\alpha$  were observed and the potential mechanism was explored, and a rabbit IVDD model was established to verify the protective role of HIF-1 $\alpha$  on IVDD.

## Results

# HIF-1 $\alpha$ attenuated TNF- $\alpha$ -induced inflammation in primary mouse nucleus pulposus cells (MNPCs)

Primary mouse nucleus pulposus cells (MNPCs) were isolated and then co-cultured with TNF- $\alpha$ , TNF- $\alpha$  and ML228 or TNF- $\alpha$  and Oltipraz for 24 h to detect the mRNA levels of COX-2 and iNOS and 48 h to detect their protein levels. As shown in Fig. 1a-1b, ML228 (HIF-1 $\alpha$  activator) could reduce the elevated mRNA expression levels of iNOS and COX-2 induced by TNF- $\alpha$  while Oltipraz (HIF-1 $\alpha$  inhibitor) could not. The western blotting results (Fig. 1c-1e) and immunofluorescence (Fig. 1f-1i) were completely consistent with the real-time PCR results. These suggested that HIF-1 $\alpha$  significantly alleviated the TNF- $\alpha$ -induced inflammatory response in MNPCs.

# HIF-1 $\alpha$ promoted cellular anabolism and inhibited catabolism of MNPCs

Figure 2a-2c showed that activation of HIF-1 $\alpha$  could up-regulate the decreased mRNA levels of Col-2 induced by TNF- $\alpha$  and simultaneously down-regulate the increased mRNA levels of MMP-13 and ADAMTS-5 induced by TNF- $\alpha$  while inhibition of HIF-1 $\alpha$  could not. Figure 2d-2h showed that activation of HIF-1 $\alpha$  could up-regulate the decreased protein levels of Col-2 and Aggrecan induced by TNF- $\alpha$  and simultaneously down-regulate the increased protein levels of MMP-13 and ADAMTS-5 induced by TNF- $\alpha$  while inhibition of HIF-1 $\alpha$  could not. Immunofluorescence (Fig. 2i-2j) was also performed to demonstrate the expression level of MMP-13 and the results were the same as the above.

# HIF-1 $\alpha$ alleviated the TNF- $\alpha$ -mediated apoptosis in MNPCs

Figure 3a-3c showed that activation of HIF-1 $\alpha$  could down-regulate the increased mRNA levels of Bax and cleaved Caspase-3 induced by TNF- $\alpha$  but upregulate the mRNA level of Bcl-2, while inhibition of HIF-1 $\alpha$  could not. Figure 3d-3h showed the same results as the above at protein levels by western blotting and the inhibitory effect of HIF-1 $\alpha$  on cleaved Caspase-3 was particularly significant. Moreover, TUNEL staining of cells (Fig. 3i-3j) was performed, which showed that TNF- $\alpha$  promoted cell death while HIF-1 $\alpha$  repressed this disorganization. Additionally, flow cytometry (Fig. 3k) was performed to test Annexin/PI to reflect the apoptosis rate of MNPCs, which indicated that TNF- $\alpha$  exaggerated the apoptosis of MNPCs and this phenomenon could be diminished by HIF-1 $\alpha$ .

# HIF-1 $\alpha$ alleviated TNF- $\alpha$ -induced mitochondrial dysfunction in MNPCs

DCFDA assays were used to evaluate ROS synthesis in mitochondria. As showed in Fig. 4a-4b, activation of HIF-1 $\alpha$  could decrease the elevated ROS synthesis induced by TNF- $\alpha$  while inhibition of HIF-1 $\alpha$  could not. Moreover, JC-1 and Mitotracker assays were performed to detect the membrane potential of mitochondria (Fig. 4c-4f), which showed TNF- $\alpha$  exacerbated the dysfunction of mitochondria in MNPCs,

and activation of HIF-1 $\alpha$  could alleviate the mitochondrial dysfunction while inhibition of HIF-1 $\alpha$  could not.

## **Exogenous supplementation of HIF-1 $\alpha$ exhibited protective effect on degeneration of NP tissue in vivo**

The rabbit IVDD model was established, and HIF-1 $\alpha$  recombinant protein or Oltipraz were locally delivered into the NP tissue. After 14 weeks, X-ray was performed and the IVD height was demonstrated to be reduced, which could be improved by the application of HIF-1 $\alpha$  (Fig. 5a). MRI was performed at 3, 6, 11 and 14 weeks, which showed higher signal intensity of IVD in HIF-1 $\alpha$  supplementation group compared with PBS group and Oltipraz group (Fig. 5b-5c). These results suggested that HIF-1 $\alpha$  could alleviate the degeneration phenotype of IVD. The IVD samples were then collected for histological analysis. As showed in Fig. 5d-5g, HE, Safranin O and Masson staining results indicated that morphological degeneration score such as height of IVD in this IVDD model was alleviated by supplementation of HIF-1 $\alpha$  in comparison to PBS group and Oltipraz group. Besides, Safranin O and Masson staining results showed that HIF-1 $\alpha$  reduced proteoglycan and collagen loss during the process of IVDD while suppression of HIF-1 $\alpha$  expression aggravated that.

## **Materials And Methods**

### **Isolation and culture of mouse primary nucleus pulposus cells (MNPCs)**

In this study, mice were sacrificed by cervical vertebra dislocation and then soaked in 75% ethyl alcohol for 10 minutes to disinfect the entire body. After the dorsal hair had been shaved, the whole spine was separated from back. The disc tissue was separated under microscope, and cut into pieces and placed in culture dishes. The cells were digested with 0.2% collagenase type II (Gibco, USA) at 37°C for 8 hours. The cells were then cultured in DMEM/F12 (HyClone, USA) supplemented with 10% fetal bovine serum (FBS, Gibco, USA), 1% penicillin and streptomycin (P1400, Solarbio, China) under standard incubation conditions (37°C, 5% CO<sub>2</sub>). The culture medium was replaced every 3 days and the cells were passaged when they reached 80–90% confluence. The cells from within five generations were used in all vitro experiments. In subsequent experiments, the control-group and TNF- $\alpha$  (10 ng/ml) group were cultured under standard incubation conditions (37°C, 5% CO<sub>2</sub>), while ML228 (1 $\mu$ M) group and Oltipraz (10 $\mu$ M) group were cultured under hypoxic conditions (37°C, 1% O<sub>2</sub>, 5% CO<sub>2</sub> and 94% N<sub>2</sub>).

### **Western blotting analysis**

Total protein was extracted from MNPCs of each group with the precooled RIPA Lysis Buffer (P0013C, Beyotime Biotechnology) containing 1mM PMSF on ice for 30 minutes. The collected liquid was centrifuged at 12000 rpm for 15 min at 4°C and the supernatant was retained. Protein concentration was detected with a BCA protein assay kit (PC0020, Solarbio). Then to destroy the 3-dimensional protein

structure, the proteins in loading buffer were heated at 100°C for 10 min. Equal amount of protein from each sample was separated by SDS-PAGE on 8%, 10% or 12% SDS-polyacrylamide gels and then transferred to a polyvinylidene difluoride (PVDF) membrane (Millipore, USA). After blocked with QuickBlock™ Blocking Buffer (P0252, Beyotime Biotechnology) for 20 minutes at room temperature, the membranes were incubated with anti-iNOS(1:1000, 18985-1-AP, Proteintech), anti-COX-2(1:1000, 27308-1-AP, Proteintech), anti-Tubulin(1:1000, 10068-1-AP, Proteintech), anti-Aggrecan(1:1000, 13880-1-AP, Proteintech), anti-Col-2(1:1000, 28459-1-AP, Proteintech), anti-ADAMTs-5(1:1000, ab41037, abcam), anti-MMP-13(1:1000, 18165-1-AP, Proteintech), anti-Bcl-2(1:1000, ab196495, Abcam), anti-Bax(1:1000, BM3964, Boster), anti-Caspase-3(1:1000, 19677-1-AP, Proteintech) antibodies at 4°C overnight. The next day, after washing with Tris-buffered saline Tween-20 (TBST), these membranes were incubated with goat anti-rabbit IgG-HRP secondary antibody (1:5000, Jackson ImmunoResearch) at room temperature for 1 h. Bound antibody was visualized using an enhanced chemiluminescence system (Amersham Life Science, Arlington Heights, IL, USA) and the density of protein bands was quantified using ImageJ software.

## Real-time PCR

An RNAfast200 Kit (220011, Fastagen) was used to extract total RNA from the MNPCs of each group according to recommended procedure. Total RNA (1µg) was reverse-transcribed to complementary DNA (cDNA) using HiScript II Q RT SuperMix for qPCR (R222-01, Vazyme). Real-time PCR was carried out with RealStar Fast SYBR qPCR Mix (A301, GenStar). The experiment was repeated three times for each target gene of each group. The nucleotide sequences of the primers are listed in Table. 1. The expression levels of target genes were normalized to Tubulin and was calculated by the  $2^{-\Delta\Delta CT}$  method

## Immunofluorescence staining

The cells were treated as indicated and after 24 h, the cells were fixed with 4% paraformaldehyde for 20 minutes. After permeabilized with 0.2% Triton X-100 for 20 minutes, samples were blocked by BSA at 37°C for 1 hour. Then, the cells were incubated with anti-iNOS (1:500, 18985-1-AP, Proteintech), anti-COX-2(1:500, 27308-1-AP, Proteintech), MMP-13(1:500, 18165-1-AP, Proteintech) antibodies at 4°C overnight. The next day, the cells were incubated with fluorescently labelled goat anti-rabbit IgG (1:100, Abbkine) for 1 hour at 37°C. The nuclei were stained with DAPI. The images were taken using a fluorescence microscope(ZEISS Vert. A1) and analysed with ImageJ software.

## TUNEL staining

To examine the apoptosis of MNPCs in each experimental group, cells were stained with a TMR (red) Tunel Cell Apoptosis Detection Kit (G1502, Servicebio). All the procedures were performed according to the manufacturer's instructions. The images were captured using a fluorescence microscope (ZEISS Vert. A1).

## Flow cytometry

The apoptosis of MNPCs from each group were detected by flow cytometry. Cells were stained with propidium iodide (PI) and Annexin V-FITC for 15 minutes at room temperature in the dark with a FITC Annexin V Apoptosis Detection Kit (E-CK-A211, Elabscience) according to the manufacturer's instructions. Cell apoptosis was detected with a CytoFLEX S flow cytometer (Beckman Coulter, USA) and the data obtained were analysed with CytExpert software.

## **Reactive oxygen species assay**

To detect intracellular reactive oxygen species (ROS), we used an ROS assay kit (S0033, Beyotime Biotechnology). All the procedures were performed according to the manufacturer's instructions. Briefly, after washing twice with sterile PBS, cells were stained with 10  $\mu$ M DCFDA at 37°C for 20 minutes in the dark. Then, the cells were washed with basal culture medium three times. The images were captured using a fluorescence microscope (ZEISS Vert. A1).

## **JC-1 assay**

The mitochondrial membrane potential changes of MNPCs after treatment were detected with a JC-1 assay kit (C2006, Beyotime). Based on the manufacturer's instructions, each group's cells were stained with the JC-1 staining solution at 37°C for 20 minutes protect from light. Then the cells were washed twice with JC-1 staining buffer and the images were observed and captured using a fluorescence microscope (ZEISS Vert. A1).

## **MitoTracker assay**

MitoTracker staining was performed to visualize the mitochondria and detect mitochondrial membrane potential of each group following the instructions of the Mito-Tracker Red CMXRos (C1049B, Beyotime Biotechnology). The cells were incubated with the culture medium containing 20nM Mito-Tracker Red CMXRos for 30 minutes at 37°C in the dark. Then the images were captured using a fluorescence microscope (ZEISS Vert. A1) after changing fresh culture medium.

## **X-Ray and Magnetic Resonance Imaging (MRI)**

The rabbits in each group were performed X-ray at 14 weeks after the initial puncture. Radiographs were captured at a collimator-to-film distance of 66 cm, an exposure of 63 mAs, and a penetration power of 35 kv. MRI was performed for each group at 3, 6, 11 and 14 weeks, and T2-weighted images (repetition time: 3000 ms; echotime: 80 ms; field of view: 200 mm<sup>2</sup>; slice thickness: 1.4 mm) were obtained by MRI using a 1.5 T system (GE) in the coronal plane. The MRI grade of NPs were evaluated as previously reported.

## **Histological staining**

The rabbits were sacrificed at 14 weeks after indicated surgery and the IVD tissues were collected and fixed in 4% paraformaldehyde for 2 days. After decalcification in 10% EDTA (pH 7.2–7.4), the samples were processed, embedded in paraffin and cut into 5- $\mu$ m sections using a microtome. H&E staining was performed to evaluate the morphological changes of nucleus pulposus with a H&E staining Kit (EE0012, Sparkjade) and histological grading of these samples was valued in accordance with the grading scale

based on the morphology of AF and the cellularity of NP. Safranin O staining was performed to detect changes in proteoglycans with a Safranin O staining kit (G1371, Solarbio) according to the manufacturer's recommended procedure. Masson staining was performed to confirm collagen loss of these samples with a Masson's Trichrome Stain Kit (G1340, Solarbio) according to the manufacturer's recommended procedure. The images were captured by a microscope (Leica DMI3000B).

## **Surgery Procedure**

The protocol of this study was approved by the Institutional Animal Care and Use Committee. Twenty-four New Zealand white rabbits (female), ranging from 2.9 to 3.4 kg in body weight (SLC, Hamamatsu, Japan), were used in this study. Rabbits were housed in separate cages under standard conditions with a light-dark cycle (12h–12h) and dry-bulb room temperature at 22–24°C and provided ad libitum access to tap water and food pellets daily. Rabbits were anesthetized by an intramuscular injection of ketamine hydrochloride (25mg/kg; Ketalar®; Daiichi Sankyo, Tokyo, Japan) mixed with xylazine (5mg/kg; Selactar®, Bayer, Tokyo, Japan). Lateral plain radiographs were obtained to determine baseline IVD height values before the treatment. Under single fluoroscopy (C-arm image intensifier, Philips Med System), rabbits were then placed into a lateral prone position, and a nonionic, isotonic contrast agent (CA) (Iotrolan 240, Bayer Global, Leverkusen, Germany) or MIA (305-53-3, Sigma-Aldrich, St. Louis, MO, USA) was injected into the rabbit IVDs from the posterolateral to the center percutaneously using micro syringes with a 31G needle. L1–L2 discs received 10 µl of CA only, L2–L3 discs received MIA 0.01 mg (in 10 µl CA), L3–L4 MIA 0.1 mg (in 10 µl CA), and L4–L5 MIA 1.0 mg (in 10 µl CA). L5–L6 discs were used as non-injection (NI) controls. The use of the lumbar spine lowest levels (L6–L7) was avoided to eliminate possible influences of the lumbosacral junction. Six rabbits were euthanized at 3, 6, 11, and 14 weeks post-injection, and the spinal columns (L1 to L6 vertebra) were harvested. At each time point, three spinal columns were processed for micro-computed tomography (CT) and another three for magnetic resonance imaging (MRI), followed by histological analysis.

## **Statistical analysis**

Analysis of data were performed with GraphPad Prism (GraphPad Software Inc., USA). Comparisons of various groups were performed using analysis of variance (ANOVA) with Tukey's post hoc test. Data were presented as "mean ± SD". Statistical significance was indicated when two-sided  $P < 0.05$ .

## **Discussion**

In vitro results demonstrated that HIF-1α could attenuate the inflammation, apoptosis and mitochondrial dysfunction induced by TNF-α in MNPCs, and promote cellular anabolism, and inhibit cellular catabolism.

TNF-α is an important pro-inflammatory cytokine which can stimulate inflammatory cascade through binding to the TNFR [17–20]. Its level is significantly elevated in the disc tissue and peripheral serum of patients with IVDD [21–23]. TNF-α can promote the production of ROS and aggravate the degradation of ECM, indicating its critical role in the development of IVDD [24, 25]. Additionally, it can also induce inflammatory response in nucleus pulposus, leading to increase of iNOS and COX2 and acceleration of

intervertebral disk destruction [26]. Therefore, MNPCs stimulated with TNF- $\alpha$  were used to investigate the effects of HIF-1 $\alpha$  on protecting against IVDD in this study. Through co-culturing MNPCs with TNF- $\alpha$ , TNF- $\alpha$  and ML228 or TNF- $\alpha$  and Oltipraz, the results showed that activation of HIF-1 $\alpha$  could down-regulate the expression of iNOS and COX-2 induced by TNF- $\alpha$  while inhibition of HIF-1 $\alpha$  could counteract the effect. This suggested that HIF-1 $\alpha$  could possibly attenuate the inflammatory response in IVDD.

Degradation of nucleus pulposus extracellular matrix is an important cause of IVDD [27]. TNF- $\alpha$  can cause IVDD by promoting expression of MMP-13 and ADAMTS-5 to enhance catabolism and inhibiting the synthesis of Aggrecan and Col-2 to reduce anabolism. Studies have found that HIF-1 $\alpha$  is closely related to the synthesis of ECM in chondrocytes [28, 29]. In our study, activation of HIF-1 $\alpha$  could inhibit catabolism through down-regulating the elevated expression of MMP-13 and ADAMTS-5 induced by TNF- $\alpha$  and enhance anabolism through up-regulating the decreased expression of Aggrecan and Col-2 induced by TNF- $\alpha$  while inhibition of HIF-1 $\alpha$  could not.

Cellular senescence and death occur widely in various tissues, including apoptosis, necrosis and autophagy, and the loss of nucleus pulposus cells due to apoptosis is one of the important causes of IVDD [30]. Studies have reported that massive death of NP cells and significant degeneration of IVD were observed in mice after conditional knockout of HIF-1 $\alpha$  gene [31, 32]. It is well known that TNF- $\alpha$  can induce the apoptosis of NP cells through increasing the expression of proapoptotic cytokines such as BAX and C-Caspase-3 and reducing the expression of antiapoptotic cytokines such as Bcl-2 [33–35], which has been considered as a potential target for investigation of IVDD. In our study, activation of HIF-1 $\alpha$  could down-regulate the increased expressions of BAX and C-Caspase-3 induced by TNF- $\alpha$  and up-regulate the decreased expression of Bcl-2 induced by TNF- $\alpha$  while inhibition of HIF-1 $\alpha$  could counteract the effect. Moreover, TUNEL and flow cytometry also showed the attenuated effect of HIF-1 $\alpha$  on the apoptosis of NP cells induced by TNF- $\alpha$ .

Mitochondria, as the center of cellular energy metabolism, is involved in a variety of signaling pathways and regulates cellular function and survival [36]. After mitochondria is exposed to adverse stimulation, it will occur multiple harmful changes such as increased oxidative stress, swelling and deformation, and decreased membrane potential [37, 38]. These changes can lead to inflammasome activation, cell senescence and death, which plays a crucial role in some degenerative diseases [39]. Studies have reported abnormal mitochondrial morphology and dysfunction in aging NP cells, and mitochondrial dysfunction plays a detrimental role in the development of IVDD [40]. TNF- $\alpha$  can cause damage to the mitochondrial structure, such as swelling and deformation in NP cells [41]. HIF-1 $\alpha$  is a transcription factor that responds to the reduction of intracellular oxygen concentration, and can enhance cellular resistance to oxidative stress as an endogenous anti-oxidative stress regulator [42]. In our study, activation of HIF-1 $\alpha$  reversed the enhanced ROS production and impaired membrane potential in mitochondria induced by TNF- $\alpha$  while inhibition of HIF-1 $\alpha$  was not.

In addition to in vitro experiments, we also evaluated radiological and histological changes of rabbit IVDs through modulation of local HIF-1 $\alpha$  activity in IVD for the first time. After establishment of IVDD model in

rabbit, disc height and IVD extracellular matrix were decreased in a time-dependent manner, and MRI analysis showed a tendency for decreased T2 values in a time-dependent manner, which is in line with previous findings. Intriguingly, our results indicated that supplementation of HIF-1 $\alpha$  improved histological and imaginative IVDD while down-regulation of HIF-1 $\alpha$  exacerbated. These results further confirmed the conclusions of in vitro experiments.

In summary, HIF-1 $\alpha$  protected against IVDD, possibly through reducing ROS production in mitochondria and consequent inhibition of inflammation, metabolism disorders and apoptosis of NP cells, which provided a new idea for exploring therapeutic strategies and targeted drugs for IVDD.

The limitations of this study mainly included two aspects. The first was a small sample size for model rabbits, and the other was that the potential mechanism associated with the protective role of HIF-1 $\alpha$  for IVDD was not verified in vivo.

## **Declarations**

### **Ethics Approval**

All animal experiments described in this study were performed in accordance with institutional guidelines and approved by the Ethical Management Committee of Heze Municipal Hospital (2020-KY005-103).

### **Consent for publication**

Not applicable.

### **Availability of data and materials**

The datasets generated and/or analyzed during the current study are available from the corresponding authors on reasonable request.

### **Competing Interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### **Funding**

This work was supported by the Natural Science Foundation of Shandong Province(Grant ZR2020MH087), The Key Research and Development projects of Shandong Province (Grant 2019GSF107057), Academic Promotion Program of Shandong First Medical University(Grant 2019LJ001).

### **Authors' Contributions**

Y. Xia, Y. Hou, Y. Li and L. Zhang contributed to the study conception and design. In vivo experiments were performed by W. Yang and R. Yu. In vitro experiments were performed by C. Jia and Z. Liu. Data collection and analysis were performed by L. Liu and Y. Fu. The first draft of the manuscript was written by Y. Wu, X. Ma, A. Gong and F. Liu and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

Not applicable.

## References

1. Kos N, Gradisnik L, Velnar T. A Brief Review of the Degenerative Intervertebral Disc Disease. *Med Arch*. 2019;73(6):421-424. doi: 10.5455/medarh.2019.73.421-424.
2. Yurube T, Ito M, Ka Y, Kuroda Rkiuchi, Kakutani K. Autophagy and mTOR signaling during intervertebral disc aging and degeneration. *JOR Spine*. 2020;3(1):e1082. doi: 10.1002/jsp2.1082.
3. Saberi M, Zhang X, Mobasher A. Targeting mitochondrial dysfunction with small molecules in intervertebral disc aging and degeneration. *Geroscience*. 2021;43(2):517-537. doi: 10.1007/s11357-021-00341-1.
4. Clouet J, Vinatier C, Merceron C, Pot-Vaucel M, Hamel O, Weiss P, Grimandi G, Guicheux J. The intervertebral disc: from pathophysiology to tissue engineering. *Joint Bone Spine*. 2009;76(6):614-8. doi: 10.1016/j.jbspin.2009.07.002.
5. Hadjipavlou AG, Tzermiadianos MN, Bogduk N, Zindrick MR. The pathophysiology of disc degeneration: a critical review. *J Bone Joint Surg Br*. 2008 Oct;90(10):1261-70. doi: 10.1302/0301-620X.90B10.20910.
6. Urban JP, Roberts S. Degeneration of the intervertebral disc. *Arthritis Res Ther*. 2003;5(3):120-30. doi: 10.1186/ar629.
7. Board D, Stemper BD, Yoganandan N, Pintar FA, Shender B, Paskoff G. Biomechanics of the aging spine. *Biomed Sci Instrum*. 2006;42:1-6..
8. Hanaei S, Abdollahzade S, Khoshnevisan A, Kepler CK, Rezaei N. Genetic aspects of intervertebral disc degeneration. *Rev Neurosci*. 2015;26(5):581-606. doi: 10.1515/revneuro-2014-0077.
9. Papadakis M, Sapkas G, Papadopoulos EC, Katonis P. Pathophysiology and biomechanics of the aging spine. *Open Orthop J*. 2011;5:335-42. doi: 10.2174/1874325001105010335.
10. Choi YS. Pathophysiology of degenerative disc disease. *Asian Spine J*. 2009;3(1):39-44. doi: 10.4184/asj.2009.3.1.39.
11. Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet*. 2017;389(10070):736-747. doi: 10.1016/S0140-6736(16)30970-9.

12. Kloppenburg M, Berenbaum F. Osteoarthritis year in review 2019: epidemiology and therapy. *Osteoarthritis Cartilage*. 2020 Mar;28(3):242-248. doi: 10.1016/j.joca.2020.01.002.
13. Sakai D, Andersson GB. Stem cell therapy for intervertebral disc regeneration: obstacles and solutions. *Nat Rev Rheumatol*. 2015 Apr;11(4):243-56. doi: 10.1038/nrrheum.2015.13.
14. He R, Wang Z, Cui M, Liu S, Wu W, Chen M, Wu Y, Qu Y, Lin H, Chen S, Wang B, Shao Z. HIF1A Alleviates compression-induced apoptosis of nucleus pulposus derived stem cells via upregulating autophagy. *Autophagy*. 2021 Nov;17(11):3338-3360. doi: 10.1080/15548627.2021.1872227.
15. Semenza GL. Hypoxia-inducible factor 1: oxygen homeostasis and disease pathophysiology. *Trends Mol Med*. 2001;7(8):345-50. doi: 10.1016/s1471-4914(01)02090-1.15.
16. Merceron C, Mangiavini L, Robling A, Wilson TL, Giaccia AJ, Shapiro IM, Schipani E, Risbud MV. Loss of HIF-1 $\alpha$  in the notochord results in cell death and complete disappearance of the nucleus pulposus. *PLoS One*. 2014;9(10):e110768. doi: 10.1371/journal.pone.0110768.
17. Risbud MV, Shapiro IM. Role of cytokines in intervertebral disc degeneration: pain and disc content. *Nat Rev Rheumatol*. 2014;10(1):44-56. doi: 10.1038/nrrheum.2013.160.
18. Puimège L, Libert C, Van Hauwermeiren F. Regulation and dysregulation of tumor necrosis factor receptor-1. *Cytokine Growth Factor Rev*. 2014;25(3):285-300. doi: 10.1016/j.cytogfr.2014.03.004.
19. Walczak H. TNF and ubiquitin at the crossroads of gene activation, cell death, inflammation, and cancer. *Immunol Rev*. 2011 Nov;244(1):9-28. doi: 10.1111/j.1600-065X.2011.01066.x.
20. Borghi A, Verstrepen L, Beyaert R. TRAF2 multitasking in TNF receptor-induced signaling to NF- $\kappa$ B, MAP kinases and cell death. *Biochem Pharmacol*. 2016 Sep 15;116:1-10. doi: 10.1016/j.bcp.2016.03.009.
21. Liang H, Yang X, Liu C, Sun Z, Wang X. Effect of NF- $\kappa$ B signaling pathway on the expression of MIF, TNF- $\alpha$ , IL-6 in the regulation of intervertebral disc degeneration. *J Musculoskelet Neuronal Interact*. 2018;18(4):551-556.
22. Zhang QX, Guo D, Wang FC, Ding WY. Necrosulfonamide (NSA) protects intervertebral disc degeneration via necroptosis and apoptosis inhibition. *Eur Rev Med Pharmacol Sci*. 2020;24(5):2683-2691. doi: 10.26355/eurev\_202003\_20538.
23. Wang S, Wei J, Fan Y, Ding H, Tian H, Zhou X, Cheng L. Progranulin Is Positively Associated with Intervertebral Disc Degeneration by Interaction with IL-10 and IL-17 Through TNF Pathways. *Inflammation*. 2018;41(5):1852-1863. doi: 10.1007/s10753-018-0828-1.
24. Zhao Y, Qiu C, Wang W, Peng J, Cheng X, Shanguan Y, Xu M, Li J, Qu R, Chen X, Jia S, Luo D, Liu L, Li P, Guo F, Vasilev K, Liu L, Hayball J, Dong S, Pan X, Li Y, Guo L, Cheng L, Li W. Cortistatin protects against intervertebral disc degeneration through targeting mitochondrial ROS-dependent NLRP3 inflammasome activation. *Theranostics*. 2020;10(15):7015-7033. doi: 10.7150/thno.45359.
25. Liu K, Wei J, Li G, Liu R, Zhao D, Zhang Y, Shi J, Xie Q, Cheng L. Fexofenadine Protects Against Intervertebral Disc Degeneration Through TNF Signaling. *Front Cell Dev Biol*. 2021;9:687024. doi: 10.3389/fcell.2021.687024.

26. Liu Y, Lin J, Wu X, Guo X, Sun H, Yu B, Shen J, Bai J, Chen Z, Yang H, Geng D, Mao H. Aspirin-Mediated Attenuation of Intervertebral Disc Degeneration by Ameliorating Reactive Oxygen Species In Vivo and In Vitro. *Oxid Med Cell Longev*. 2019;2019:7189854. doi: 10.1155/2019/7189854.
27. Liu H, Pan H, Yang H, Wang J, Zhang K, Li X, Wang H, Ding W, Li B, Zheng Z. LIM mineralization protein-1 suppresses TNF- $\alpha$  induced intervertebral disc degeneration by maintaining nucleus pulposus extracellular matrix production and inhibiting matrix metalloproteinases expression. *J Orthop Res*. 2015;33(3):294-303. doi: 10.1002/jor.22732.
28. Wang P, Meng Q, Wang W, Zhang S, Xiong X, Qin S, Zhang J, Li A, Liu Z. Icariin inhibits the inflammation through down-regulating NF- $\kappa$ B/HIF-2 $\alpha$  signal pathways in chondrocytes. *Biosci Rep*. 2020;40(11):BSR20203107. doi: 10.1042/BSR20203107
29. Wang P, Zhu P, Liu R, Meng Q, Li S. Baicalin promotes extracellular matrix synthesis in chondrocytes via the activation of hypoxia-inducible factor-1 $\alpha$ . *Exp Ther Med*. 2020;20(6):226. doi: 10.3892/etm.2020.9356.
30. Chen D, Xia D, Pan Z, Xu D, Zhou Y, Wu Y, Cai N, Tang Q, Wang C, Yan M, Zhang JJ, Zhou K, Wang Q, Feng Y, Wang X, Xu H, Zhang X, Tian N. Metformin protects against apoptosis and senescence in nucleus pulposus cells and ameliorates disc degeneration in vivo. *Cell Death Dis*. 2016;7(10):e2441. doi: 10.1038/cddis.2016.334.
31. Meng X, Zhuang L, Wang J, Liu Z, Wang Y, Xiao D, Zhang X. Hypoxia-inducible factor (HIF)-1 $\alpha$  knockout accelerates intervertebral disc degeneration in mice. *Int J Clin Exp Pathol*. 2018;11(2):548-557.
32. Silagi ES, Schoepflin ZR, Seifert EL, Merceron C, Schipani E, Shapiro IM, Risbud MV. Bicarbonate Recycling by HIF-1-Dependent Carbonic Anhydrase Isoforms 9 and 12 Is Critical in Maintaining Intracellular pH and Viability of Nucleus Pulposus Cells. *J Bone Miner Res*. 2018;33(2):338-355. doi: 10.1002/jbmr.3293
33. Zhu H, Sun B, Shen Q. TNF- $\alpha$  induces apoptosis of human nucleus pulposus cells via activating the TRIM14/NF- $\kappa$ B signalling pathway. *Artif Cells Nanomed Biotechnol*. 2019;47(1):3004-3012. doi: 10.1080/21691401.2019.1643733.
34. Zhang G, Liao Y, Yang H, Tao J, Ma L, Zuo X. Iridogenin reduces the expression of caspase-3 and matrix metalloproteinases, thus suppressing apoptosis and extracellular matrix degradation in TNF- $\alpha$ -stimulated nucleus pulposus cells. *Chem Biol Interact*. 2021;349:109681. doi: 10.1016/j.cbi.2021.109681.
35. Lin XL, Zheng ZY, Zhang QS, Zhang Z, An YZ. Expression of miR-195 and its target gene Bcl-2 in human intervertebral disc degeneration and their effects on nucleus pulposus cell apoptosis. *J Orthop Surg Res*. 2021;16(1):412. doi: 10.1186/s13018-021-02538-8.
36. Abate M, Festa A, Falco M, Lombardi A, Luce A, Grimaldi A, Zappavigna S, Sperlongano P, Irace C, Caraglia M, Misso G. Mitochondria as playmakers of apoptosis, autophagy and senescence. *Semin Cell Dev Biol*. 2020;98:139-153. doi: 10.1016/j.semcdb.2019.05.022.

37. Jang S, Chapa-Dubocq XR, Parodi-Rullán RM, Fossati S, Javadov S. Beta-Amyloid Instigates Dysfunction of Mitochondria in Cardiac Cells. *Cells*. 2022;11(3):373. doi: 10.3390/cells11030373
38. Marei WFA, Smits A, Mohey-Elsaeed O, Pintelon I, Ginneberge D, Bols PEJ, Moerloose K, Leroy JLMR. Differential effects of high fat diet-induced obesity on oocyte mitochondrial functions in inbred and outbred mice. *Sci Rep*. 2020;10(1):9806. doi: 10.1038/s41598-020-66702-6
39. Sun K, Jing X, Guo J, Yao X, Guo F. Mitophagy in degenerative joint diseases. *Autophagy*. 2021;17(9):2082-2092. doi: 10.1080/15548627.2020.1822097.
40. Xu X, Wang D, Zheng C, Gao B, Fan J, Cheng P, Liu B, Yang L, Luo Z. Progerin accumulation in nucleus pulposus cells impairs mitochondrial function and induces intervertebral disc degeneration and therapeutic effects of sulforaphane. *Theranostics*. 2019 Apr 12;9(8):2252-2267. doi: 10.7150/thno.30658.
41. Zhang Z, Xu T, Chen J, Shao Z, Wang K, Yan Y, Wu C, Lin J, Wang H, Gao W, Zhang X, Wang X. Parkin-mediated mitophagy as a potential therapeutic target for intervertebral disc degeneration. *Cell Death Dis*. 2018 Sep 24;9(10):980. doi: 10.1038/s41419-018-1024-9.
42. Loor G, Schumacker PT. Role of hypoxia-inducible factor in cell survival during myocardial ischemia-reperfusion. *Cell Death Differ*. 2008 Apr;15(4):686-90. doi: 10.1038/cdd.2008.13.

## Table

Table 1 is available in the Supplementary Files section

## Figures

### Figure 1

**HIF-1 $\alpha$  attenuated TNF- $\alpha$ -induced inflammation in primary mouse nucleus pulposus cells (MNPCs).** (a-b) The mRNA levels of iNOS and COX-2 after cells were treated as above, were detected using real-time PCR. (c-e) The protein levels of iNOS and COX-2 were detected using western blotting and (f-i) immunofluorescence staining. The scale bar was 50  $\mu$ m. The values were mean of at least three independent experiments. \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001 and \*\*\*\*P < 0.0001.

### Figure 2

**HIF-1 $\alpha$  promoted cellular anabolism and inhibited catabolism of MNPCs.** (a-c) The expression levels of Col-2, MMP-13 and ADAMTS-5 were detected using real-time PCR. (d-h) The protein levels of Aggrecan, Col-2, MMP-13 and ADAMTS-5 were assayed using western blotting. (i-j) The expression levels of MMP-

13 were detected by immunofluorescence staining. The scale bar was 50  $\mu\text{m}$ . The values were mean of at least three independent experiments. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  and \*\*\*\* $P < 0.0001$ .

### Figure 3

**HIF-1 $\alpha$  alleviated the TNF- $\alpha$ -mediated apoptosis in MNPCs.** (a-c) The expression levels of Bcl-2, BAX and Caspase-3 were detected using real-time PCR. (d-h) The protein levels of Bcl-2, BAX and C-Caspase-3 were assayed using western blotting. (i-j) TUNEL staining was performed to examine the apoptosis. (k) Flow cytometry was performed to detect the apoptosis. The scale bar was 100  $\mu\text{m}$ . The values were mean of at least three independent experiments. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  and \*\*\*\* $P < 0.0001$ .

### Figure 4

**HIF-1 $\alpha$  alleviated TNF- $\alpha$ -induced mitochondrial dysfunction in MNPCs.** (a-b) ROS levels of MNPCs were detected with DCFDA. (c-d) JC-1 and (e-f) MitoTracker were performed to detect the mitochondrial membrane potential of MNPCs. The scale bar was 50  $\mu\text{m}$ . The values were mean of at least three independent experiments. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  and \*\*\*\* $P < 0.0001$ .

### Figure 5

**Exogenous supplementation of HIF-1 $\alpha$  exhibited protective effect on regeneration of NP tissue in vivo.** (a) X-ray and (b-c) MRI to assess the degree of IVDD in rabbits from sham group, PBS group, HIF-1 $\alpha$  group and Oltipraz group. (d-g) HE, Safranin O and Masson staining indicated morphological degeneration score such as height of IVD in this IVDD model was alleviated by supplementation of HIF-1 $\alpha$  in comparison to the PBS group and Oltipraz group. Safranin O and Masson staining showed that HIF-1 $\alpha$  reduced proteoglycan and collagen loss during the process of IVDD but suppression of HIF-1 $\alpha$  expression aggravated that. The histological scores of each indicated group were calculated according to the grading scale previously published. (h) Photos of the surgical procedure in vivo. The scale bar is 500 or 100  $\mu\text{m}$ . The values are mean of at least three independent experiments. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  and \*\*\*\* $P < 0.0001$ .

### Figure 6

**Schematic depicting a proposed model for the role of HIF-1 $\alpha$  in IVDD based on this study.**

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table1.tif](#)
- [GelsandBlots.zip](#)
- [IVDDmodel.mp4](#)