

Supplementation of *Lactobacillus reuteri* isolated from red jungle fowl along with mannanoligosaccharide improves growth performance, immune response and gut health in broiler Chicken

Sudhir Kumar Jaiswal

ICAR-Central Avian Research Institute

Simmi Tomar

ICAR-Central Avian Research Institute

Vishesh Kumar Saxena

ICAR-Central Avian Research Institute

Balraj S

ICAR-Central Avian Research Institute

Sandeep Uniyal (✉ sandeepuniyalvet18@gmail.com)

ICAR-Indian Veterinary research Institute

Research Article

Keywords: lactobacillus reuteri, mannoilgosaccharide, broiler, body weight gain, immunity, genes

Posted Date: May 18th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1645541/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

The present study was undertaken to investigate the effect of supplementation of *Lactobacillus reuteri* isolated from the intestine of red jungle fowl along with mannanoligosaccharide (MOS) on growth performance, intestinal microbial count, immunity and expression of growth and immune related genes in broiler chicken. *Lactobacillus reuteri* was isolated from in the GIT tract of red jungle fowl and was utilized for growth bioassay in broiler. For this total 360 CARIBRO-Vishal broiler chicks were weighed individually and randomly allocated to nine treatment groups, each having five replicates with eight chicks in each following complete randomized block design (CRD). The experiment was conducted for 6 weeks duration. The nine treatment groups were control fed basal diet (T1), basal diet + Antibiotic growth promoter, bacitracin methylene disalicylate (BMD) @20mg/kg feed (T2), basal diet + commercial probiotic @ 0.1 g/kg feed (T3), basal diet + Lab isolated *Lactobacillus reuteri* @ 1×10^6 CFU/g of fermented feed (T4), basal diet + Lab isolated *Lactobacillus reuteri* @ 1×10^7 CFU/g of fermented feed (T5), basal diet + Lab isolated *Lactobacillus reuteri* @ 1×10^8 CFU/g of fermented feed (T6), basal diet + Lab isolated *Lactobacillus reuteri* @ 1×10^6 CFU/g of fermented + 0.1% MOS (T7), basal diet + Lab isolated *Lactobacillus reuteri* @ 1×10^7 CFU/g of fermented + 0.1% MOS (T8), basal diet + Lab isolated *Lactobacillus reuteri* @ 1×10^8 CFU/g of fermented feed + 0.1% MOS (T9). 20% of daily basal ration for broiler chicken was autoclaved and inoculated with 15% of Lactobacillus isolate broth culture having a viable count of 10^6 , 10^7 , and 10^8 CFU/ml and fermented at 37°C for 24 h before adding to daily ration afresh and was mixed well. Results of the present study revealed T9 group supplemented with Lab isolated *Lactobacillus reuteri* at the dose of 1×10^8 CFU/g along with 0.1% MOS significantly ($P < 0.05$) improves body weight, body weight gain, immune response both humoral and cell mediated without effecting the feed intake and feed conversion ratio (FCR). Also the pathogenic bacteria count (Salmonella and E.coli) was significantly ($P < 0.05$) lower in the GIT of T9 group as compared to other groups. The significantly ($p < 0.05$) higher relative expression of growth related genes, IGF-1 and IGF-1R and immune related gene, IL-6 whereas IL-10 and TLR-4 expression were significantly ($P < 0.05$) down regulated in T9 group (Lab isolated *Lactobacillus reuteri* @ 1×10^8 CFU/g of fermented feed + 0.1% MOS). So, it can be concluded from the present study that lactobacillus *reuteri* isolated for the GIT of the red jungle fowl along with MOS is effective in improving the growth performance, immune response and gut health of commercial CARIBRO-Vishal broiler chicken.

Introduction

The chicken gastrointestinal tract (GIT) is rich in microbial biodiversity of more than 500 phylotypes or over 1 million bacterial genes, which equates to 40–50 times of number in the chicken nuclear genome [1]. This microflora has a role in nutrition, detoxification of certain compounds, growth performance, and protection against colonization of pathogens and influences health and well-being of host animals [2]. When these live microorganisms administered in adequate amounts confer a health benefit on the host and with this, the concept of probiotic has been evolved [3]. Lilly and Stillwell coined the term 'probiotics' in 1965, derived from Greek word "pro bios", meaning "for life". Probiotics as per World Health Organization (WHO) and Food and Agriculture Organization (FAO) are "live microorganisms, which when administered in adequate amounts confer a health benefit on the host". Probiotics have numerous advantages as they improves

digestibility and utilization of nutrients, increases growth rate and productivity, inhibits disease producing organisms, prevents diarrhoea due to bacterial infections, reduces stress after vaccination, antibiotic therapy and transportation and stimulates the immune responses [4–6]. Antibiotic growth promoter (AGP) have been used at sub-therapeutic doses in poultry diets to prevent diseases and to promote growth performance for many decades but since the ban imposed by European union in 2006 on the use of these AGP in farm animals due to environmental and public health risk associated with emergences of antibiotic resistance [7]. Therefore, alternatives to the use of AGP must be found to promote growth or production at or near the genetic potential of the modern day poultry. In recent year's use of probiotics that enrich certain bacterial population in the digestive system are considered as alternatives to antibiotic growth promoters in poultry nutrition [8]. Domestic birds raised under commercial conditions are vulnerable to a number of pathogens [2]. The World Health Organization (WHO) has now urged egg and meat producers to use environment friendly alternative methods to control diseases. Phasing out of antibiotic growth promoters from poultry diets in Europe and recent moves toward reduction or removal of these compounds in other parts of the world will likely change the microbial profile of the GIT environment in commercial poultry. Hence every effort would be made to improve the gut efficiency through natural microflora and fauna for better nutrient utilization. Assuming that the Red Jungle Fowl being raised in natural habitats could represent the most natural GIT environment and may harbouring certain uncharacterized strains of microbes imparting better immunity and adaptability to these birds. It is the primary progenitor of the domestic chicken and native to southern Asia, particularly the jungles of India. Keeping above facts in view, the present study has been designed to effect of supplementation of *Lactobacillus reuteri* isolated from the intestine of red jungle fowl along with mannanoligosaccharide (MOS) on growth performance, intestinal microbial count, immunity and expression of growth and immune related genes in broiler chicken raised under intensive system for high economic return.

Material And Methods

Isolation of Lactobacillus

Lactobacillus reuteri was isolated from gut sample of four Red Jungle fowl housed under uniform management and feeding conditions. The purified cultures were stored as 50% glycerol stock mixture (250 µl of broth culture in 1.75 ml of 50% glycerol+50% MRS broth medium) at -80°C until further use [9]. The isolates were sub-cultured at least 2 times before all of the assays. 1 ml from each isolate culture was used for CFU enumeration. Titration was done by serial dilution, plating and counting on MRS agar plates. After that the aliquots was adjusted to 10^6 , 10^7 and 10^8 CFU/ml by using sterile PBS. Broiler (starter and finisher) ration used for feeding the birds in present study were fermented with *Lactobacillus reuteri* for this 20% of daily ration was autoclaved and daily inoculated with 15% of *Lactobacillus reuteri* broth culture having viable count of 10^6 , 10^7 and 10^8 CFU/ml and fermented at 37°C for 24 hours before added to daily ration afresh and mixed well.

Experimental birds and housing management

360 day old CARIBRO-Vishal broiler birds were procured from the ICAR-CARI, Izatnagar hatchery. Each bird was weighed on arrival and randomly assigned to nine groups using completely randomized design (CRD). Each dietary treatment had five replicates having eight broiler birds in each replicate. All the experimental groups of birds are reared in the battery brooder fitted with waterer and feeder from day one to 42nd day of age. Twenty-three-hour light was provided throughout the experimental period. Fresh and clean water were offered *ad libitum* throughout the experimental period. Proper ventilation was maintained in the shed. Daily feed intake, weekly body weight and Mortality, if any was recorded throughout the experimental feeding. The birds are vaccinated for common diseases following standard vaccination schedule.

Experimental feeding of birds

All the experimental birds fed basal diet composed of maize, soybean meal, de-oiled rice bran and fish meal was formulated to target the requirement of the essential nutrients for broiler chickens as per ICAR, 2013. T1 fed basal diet (BD), T2 BD+ Antibiotic BMD (Bacitracin Methylene Di-salicylate) @ 20mg/kg diet, T3 BD+ Commercial multi-strain probiotic (0.1g/kg feed) T4 BD+ Lab isolated *Lactobacillus reuteri* (1×10^6 CFU/g fermented feed) T5 BD+ Lab isolated *Lactobacillus reuteri* (1×10^7 CFU/g fermented feed) T6 BD+ Lab isolated *Lactobacillus reuteri* (1×10^8 CFU/g fermented feed) T7 BD+ Lab isolated *Lactobacillus reuteri* (1×10^6 CFU/g fermented feed)+MOS (0.1g /kg feed) T8 BD+ Lab isolated *Lactobacillus reuteri* (1×10^7 CFU/g fermented feed)+MOS (0.1g /kg feed) T9 BD+ Lab isolated *Lactobacillus reuteri* (1×10^8 CFU/g fermented feed)+MOS (0.1g /kg feed). Weighed amount of each test diet used during the starting period (0-3wks) and finishing period (4-6wks) were offered daily in five replicates of eight chicks each to ensure *ad libitum* feeding at all the time, but with care to avoid spillage and wastage of feed.

Growth performance and feed intake

Body weights of the birds were recorded at 0th day, 1st, 2nd, 3rd, 4th, 5th and 6th week in the morning before offering the feed to birds. Body weights were taken by using platform digital balance. To measure weekly feed intake of the birds weighed quantity of respective diet was offered *ad libitum* daily to quadruplicate groups of each dietary regimen in the morning and the residue was weighed at the end of every week.

Immune response

Immune responses of the experimental birds as affected by different dietary dose of lab isolated *Lactobacillus reuteri* treatments were evaluated in terms of humoral and cell mediated immune response.

Humoral immune response

Humoral immunity was evaluated as antibody titre (HA) against 1% sheep red blood cells (SRBC) suspension. The microtitre haaemagglutinin (HA) procedure [10] was followed to measure total HA antibody titre in chickens on 5th and 10th day of post inoculation

In-Vivo Cell mediated immune response (CMI)

The CMI response to phytohaemagglutinin-P (PHA-P) mitogen was evaluated by the method of [11]. PHA-P (1mg/ml of PBS) was injected intra-dermally in the left foot web of 8 birds/groups. Right foot web of the same birds received 0.1 ml sterile PBS and thus served as control. The skin thickness of foot webs (Right and Left) of injected birds of each group was measured by a micrometer at 0 and 24 hours after injection of mitogen. The foot web swelling was calculated by subtracting the skin thickness at 24 hours from that of 0 hours of injection of both feet. Foot web index (FWI) was calculated by subtracting the difference in thickness at 0 and 24 hrs of mitogen injected foot web with the difference in thickness of control foot web at 0 and 24 hrs.

GI tract microflora population

GIT contents were collected separately from crop, ileum and caecum at the end of experiment from broilers for bacterial enumeration viz Lactobacilli, Salmonella and E. coli. GIT contents were diluted 10-fold with buffered peptone water and vortexed for 2 min; 100 microliters of supernatant was smeared onto appropriate selective media in duplicate plates and incubated under optimum time-temperature combinations. After incubation period bacterial counts were recorded as CFU/ml.

Gene expression analysis

Growth and immunity related gene

At 6th week of age eight birds per group were sacrificed for collection of pectoral muscle and Ileum tissue sample. Total RNA was isolated from the tissue sample by standard TRIZOL method. The purity of RNA was checked before the preparation of first- strand cDNA. Prepared cDNA was stored at -20 °C and used for gene expression studies. Expression of IL-6, IL-10, TLR-4, IGF-1 and IGF-1R gene was quantified with using specific primer pairs (Table 1) for gene of interest (GOI) in Real-Time PCR. GAPDH was used as a reference gene

Statistical analysis

The data generated in the above experiment were statistically analyzed using IBM SPSS version 20 computer package. For comparison of groups, generalized linear model ANOVA procedure and Duncan's multiple range tests were used [12]. The fold expression of gene was calculated by using the $2^{-\Delta\Delta Ct}$ method and analyzed by one way ANOVA

Results

Growth performance

The data pertaining to growth performance of experimental birds as influenced by various dietary treatments has been presented in Table 2. Body weight at 3rd week of age was observe as numerically higher in all dietary treatments groups as compared to control group but highest body weight was recorded in T9 (1×10^8 CFU *Lactobacillus reuteri* + 0.1% MOS) group. The cumulative body weight at 6 week of age

was significantly ($P < 0.05$) higher in T9 followed by statistically similar T8, T2, T7, T6 and T3 whereas significantly lower body weight was recorded in T1. No significant difference ($P > 0.05$) was observed in average body weight gain among the nine different dietary treatment groups in 0-3 and 4-6 week of age. The body weight gain (0-6 week) was significantly ($P < 0.05$) higher in T9 followed by statistically similar T8, T7, T6, T3 and T2 whereas significantly ($P < 0.05$) lower body weight was recorded in T1 followed by statistically similar T8, T7, T6, T3 and T2 group.

Feed Intake and Feed conversion ratio (FCR)

The data pertaining to feed intake (g) and FCR is presented in Table 3. The results indicate that no significant ($P > 0.05$) difference was observed in feed intake among the different dietary supplemented groups during 0-3, 4-6 and 0-6 week of age. The result of FCR showed that there were no significant ($P > 0.05$) differences between the different dietary groups and control group at 0-3, 4-6 and 0-6 week of age. The T9 group (1×10^8 CFU *Lactobacillus reuteri* + 0.1% MOS) showed numerically better FCR than the other groups whereas, the control group showed highest feed conversion ratio than others dietary treatment groups at all times

Intestinal microbial count

The least squares analysis of variance for \log_{10} CFU/g of *Lactobacillus*, *Salmonella* and *E. coli* in crop, ileum and caecum of Broiler fed with different treatments is presented in Table 4. . Enumeration of *Lactobacillus* spp. *Salmonella* spp. and *E. coli* was conducted at 42th day of feeding trial in Broiler. The *Lactobacillus* count in crop was significantly ($P < 0.05$) higher in T9 (1×10^8 CFU *Lactobacillus reuteri* + 0.1% MOS) group and significantly lower in T1 group whereas other treatments resulted in intermediate *Lactobacillus* count in crop at 6th week of age. The *Lactobacillus* count in ileum was significantly ($P < 0.05$) higher in T9 followed by statistically similar T8 and lower in T1 and T2 whereas other treatments resulted in intermediate *Lactobacillus* count in ileum at 6th week of age. In caecum also T9 showed higher ($P < 0.05$) *Lactobacillus* count than control and other dietary treatment groups.

Significantly ($P < 0.05$) lower *Salmonella* count was observed in the crop and ileum of the broilers in T9 group as compared to all the treatment groups at 6th week of age. *Salmonella* count in caecum was significantly ($P < 0.05$) lower in T9 followed by statistically similar in T8, T7, and highest *Salmonella* count was observed in T1 group at 6th week of age. The *E. coli* count in crop was significantly ($P < 0.05$) lower in T9 group followed by statistically similar in T8, T3 and T6 groups whereas significantly ($P < 0.05$) higher in T1 followed by statistically similar T7, T8, T3, T4, T5, T6 and T8 at 6 week of age. In ileum *E. coli* count was significantly ($P < 0.05$) lower in all dietary treatments except T4 and T1. Similar finding was also observed in caecum with significantly ($P < 0.05$) lower *E. coli* count in T9 group followed by statistically similar T8, T7, T6, T5, T4, T3 and higher count observed in T1 group. The results obtained in this work revealed that 1×10^8 CFU of *Lactobacillus reuteri* along with 0.1% MOS significantly increase *Lactobacillus* count and reduce *Salmonella* and *E. coli* count in crop, ileum, and caecum.

Immune response

The results of immune response (humoral and cell mediated immune response) as affected by different treatments are presented in Table 5. The antibody titre (humoral immunity) was significantly ($P<0.05$) lower in T1 group followed by statistically similar T4 and T5, whereas significantly ($P<0.05$) higher antibody titre was obtained in T9 group which did not differ significantly from treatment T8, T2, T7, T3, and T6 groups. The antibody titre, it is revealed that dietary inclusion of lab isolated Lactobacillus along with MOS significantly ($P<0.05$) enhanced the antibody titre values of broilers, more prominently in group T9 and T8. The cell mediated immunity was significantly higher in T9 group followed by statistically similar T8 group, whereas significantly ($P<0.05$) lower cell mediated immunity was obtained in control (T1) group which did not differ significantly from treatment T4, T3, T5, T2, T6 and T7 group.

Gene expression

Growth related gene

The results pertaining to the gene expression of IGF-1 and IGF-1R are illustrated in Fig 1 and 2. The mRNA levels of IGF-1 and IGF-1R gene were quantified by real-time PCR in breast muscle tissue at the end of 6th week. The relative fold expression of IGF-1 gene were significantly ($P<0.05$) up regulated in all dietary treatment groups compare to control (T1). The greater up regulation was observed in T9 group followed by statistically similar T8 and T2 group. Similar pattern was also observed in relative expression of IGF-1R gene, where higher relative expression was recorded in T9 and T2 group compared to other dietary treatment and control.

Immune related gene

The results of the gene expression analysis of IL-6, IL-10 and TLR-4 are illustrated in Fig 3, 4 and 5. The mRNA levels of IL-6, IL-10 and TLR-4 genes were quantified by real-time PCR in ileum tissue at the end of 6th week. The significantly ($P<0.05$) lower relative fold expression of IL-6 gene was observed in treatment group T9 followed by statistically similar T2 and T8 group, whereas higher expression was observed in treatment group T1 followed by statistically similar T4 and T5 groups. The other dietary treatments yielded intermediate fold expression of IL-6 in broiler chicken. The relative fold expression of IL-10 gene were significantly ($P<0.05$) up regulated in T9 group followed by statistically similar T8, T7, T6, T3 and T2 groups whereas lower expression was observed in treatment control (T1) and other dietary treatments yielded intermediate fold expression of IL-10 in broiler chicken. The significantly ($P<0.05$) lower relative fold expression of TLR-4 gene was observed in treatment group T9 followed by statistically similar T2 and T3 group, whereas higher expression was observed in treatment group T1 and other dietary treatments yielded intermediate fold expression of TLR-4 in broiler chicken.

Discussion

Growth performance

The beneficial effect of any supplement is primarily judged through examination of the response in terms of growth of birds for which it is intended to be offered as a part of the diet. Accordingly, growth response

of broiler chicken to dietary supplement was an important part of the study. The response criteria of feeding trial like growth performance viz., body weight, body weight gain, feed intake, feed conversion ratio. From this study it can be concluded that dietary supplementation of lab isolated *Lactobacillus reuteri* @ 1×10^8 CFU along with MOS (0.1%) increases body weight and body weight gain at 6th week of age with no effect on feed intake and feed conversion ratio. This finding is in agreement with several reports demonstrating that probiotic supplemented to the birds improve the body weight, body weight gain and feed conversion ratio [13-18]. However, contrary to our observation body weight, daily weight gain and FCR of broilers were not influenced by the dietary supplementation with probiotics [19-22]. The improvement in the body weight, daily weight gain, feed consumption and feed conversion ratio in the present study may be due to the increase in the digestion and absorption of the nutrients by increasing the secretion and activity of the digestive enzymes processes and reduction of the population of pathogenic bacteria by decreasing the intestinal P^H due to presence of the *Lactobacillus reuteri* bacteria and also maintain the balance between beneficial and harmful bacteria which is required for sound gut health and optimum growth performance [6].

Intestinal microbial count

Feeding of lab isolated *Lactobacillus reuteri* 1×10^8 CFU/g along with MOS (0.1%) to broiler chicken for 6 weeks significantly ($P < 0.05$) increases Lactobacillus count in different parts of GIT (crop, ileum and caecum) with concomitant decrease in pathogenic organisms such as E. coli and Salmonella count. The results were in accordance with [23-25] who reported that reduction of E. coli and Salmonella count in intestine upon Lactobacillus feeding. The mechanism of the antimicrobial action may be due to the potential of Lactobacillus isolates to produce fatty acids, lactic acids, hydrogen peroxide diacetyl, acetoin and the small, heat-stable inhibitory peptides called bacteriocins [26,27] Other proposed mechanisms of pathogen inhibition by the probiotic micro-organisms include competition for nutrients, and by modifying the structure and function of the intestinal epithelium. Prebiotics such as MOS can also flush out pathogens from GIT serving as receptor sites for their adherence [28].

Immune response

Probiotics as an alternative to antibiotics are considered to improve the health status and immunity in poultry birds. It has been reported that probiotic supplementation influenced the production of B-lymphocytes, T lymphocytes, anti-pro inflammatory cytokines and interleukins production in broiler chicken. In present study dietary supplementation of lab isolated *Lactobacillus reuteri* @ 1×10^8 CFU along with MOS (0.1%) enhanced the immune response (humoral and cell mediated immune response) of broiler chicken compared to other levels of *Lactobacillus reuteri*. The results of the present study are in agreement with the finding of [14-17, 29-33] who reported that supplementation of Lactobacillus probiotic enhance immune response by greater lymphocytes cell proliferation, increased production of interleukins and lowering the production of pro-inflammatory cytokines.

Gene expression

Growth related gene

In this study we have found that higher relative expression of both IGF-and IGF-1R gene in birds fed 1×10^8 CFU/g *Lactobacillus reuteri* along with 0.1 % MOS (T9 group) may be due to stimulation of glucose and amino acid uptake, protein synthesis and inhibition of protein degradation by lab isolated *Lactobacillus reuteri* [34]. On the similar lines [35] reported that expression of insulin like growth factor 1 (IGF- 1) and insulin like growth factor receptor 1 (IGF-1R) increased in *Aspergillus awamori*, FOS and combined *Aspergillus awamori* and FOS (symbiotic) fed group. The well documented biological role of IGF-1 is, but not limited to, increased glucose and amino acid uptake, and enhanced DNA and protein synthesis. The IGF I released into the circulation binds to its specific receptor called IGF-1 which stimulates cell proliferation [36]. [37] noted that the IGF-1 is one of the main growth factors that stimulate protein synthesis in muscle tissue. Also, IGFs are important positive modulators of body and muscle growth in mammals and chickens. IGF-1 is a primary mediator of the effects of growth hormone (GH). Growth hormone, made in the anterior pituitary gland, is released into the blood stream and then stimulates the liver to produce IGF-1 which in turn stimulates systemic body growth, and has growth-promoting effects on almost every cell in the body, especially skeletal muscle, cartilage, bone, liver, kidney, nerves, skin, hematopoietic cell, and lungs. In addition to the insulin like effects, IGF-1 can also regulate cell growth and development, especially in nerve cells, as well as cellular DNA synthesis [38]. The IGF-I released into the circulation binds to its specific receptor called by IGF type1 receptor and finally stimulates cell proliferation [36]. Thus, the better growth performance of birds in this study under 1×10^8 CFU/g *Lactobacillus reuteri* along with 0.1 % MOS (T9) can be due to higher corresponding IGF-1 levels. This association of growth performance and IGF-1 expression was reported by many authors. [39] stated that the changes in the plasma IGF-1 levels in the body alter the nutritional condition of young chicken and cause changes in body weight gain. Thus, from the current study the inclusion level of 1×10^8 CFU/g *Lactobacillus reuteri* can be regarded as the requirement level of the chicken.

Immune related gene

In Chicken IL-6, secreted by T cells and macrophages, acts as a pro-inflammatory cytokine in association of the production of acute phase proteins. The IL-6 upregulation in chicken has been associated with Salmonella and Eimeria infection [40-42]. Therefore, a down regulation of IL-6 in the present study favours an anti-inflammatory response and shows that *Lactobacillus reuteri* had an anti-inflammatory effect in the gut. The result of present study are in agreement with the finding of [43, 44] who reported the down regulation of IL-6 in the chicken gut when their diet was supplemented with probiotic. The probiotic bacteria could act in stabilizing intestinal inflammation by balancing the intestinal microflora, maintaining mucosal barrier, modulating and improving the intestinal mucosal immune system, especially by keeping the balance of pro-inflammatory and anti-inflammatory cytokines and production of intestinal IgA [45-48]. The relative fold expression of IL-10 gene were significantly ($P < 0.05$) up regulated in T9 Group (lab isolated lactobacillus 1×10^8 CFU/g along with MOS 0.1 %). The IL-10 act as anti-inflammatory cytokine that controls the nature and extent of inflammatory responses to various microbial infections [49] and is involved in intestinal immunity and homeostasis [50]. Several reports have shown the regulation of IL-10 by probiotic

bacteria [51, 52]. Similar to the results of the present study [53] observed significant increase in IL-10 expression following the *Lactobacillus* supplementation in broiler chicken. However, in contrast to our study [54] reported no significant changes in IL-10 expression in broiler chicken due to the *Lactobacillus* supplementation. The TLR-4 is the principal receptor for lipopolysaccharide, which is a major component of the outer membrane of gram-negative bacteria [55]. The decline in the expression of TLR-4 expression in T9 group in the present study means lesser receptor binding sites for the gram negative bacteria which results in the expulsion of pathogens like salmonella and *E. coli* from the gut. The down regulation of TLR-4 expression would be expected in the intestine of broilers fed the probiotic-supplemented diet because dietary inclusion of the probiotics decreased the population of gram-negative bacteria such as Coliform in the rectum of broilers [44].

Conclusion

In the present study supplementation of *Lactobacillus reuteri* improved growth performance (body weight gain), immune response (humoral and cell mediated immune response), reduces pathogenic gut bacteria (*Salmonella* and *E. Coli*) and upregulated expression of growth related genes (IGF-1 and IGF-1R) and immune related gene, IL-6 whereas expression of IL-10 and TLR-4 gene was down regulated in broiler. Thus, the Red jungle fowl specific lab isolate *Lactobacillus reuteri* may be used in commercial broiler production for enhancing growth and immunity.

Declarations

Acknowledgment

The authors are thankful to ICAR-Central Avian Research Institute, Iztnagar, India for all the infrastructure and financial support.

Funding

Funding for conducting the research was provided by ICAR-Central Avian Research Institute, Iztnagar India

Conflict of Interest Statement

The authors declare that they have no conflict of interest

Authors Contributions

All authors contributed equally to the work

Data availability

All the data is provided with the manuscript

Statement of Animal Ethics

The experiment was approved by Institutional Animal Ethics Committee and conducted under guidelines prescribed by the Committee for the Purpose of Control and Supervision of Experiments on Animals, Government of India.

Consent to participate

All authors give their consent to participate in the publication of this manuscript

Consent for publication

The authors give their consent for publication

References

1. Shang Y, Kumar S, Oakley B, Kim WK (2018) Chicken gut microbiota: importance and detection technology. *Front Vet Sci* 5:254.
2. Paul SS, Chatterjee RN, Raju MVLN, Prakas, B, Rao SVR, Yadav SP, Kannan A (2021) Gut microbial composition differs extensively among Indian native chicken breeds originated in different geographical locations and a commercial broiler line, but breed-specific, as well as across-breed core microbiomes, are found. *Microorganisms* 9: 391.
3. Al-Khalaifah HS (2018). Benefits of probiotics and/or prebiotics for antibiotic-reduced poultry. *Poult Sci* 97(11): 3807–3815.
4. Alayande KA, Aiyegoro OA, Ateba CN (2020) Probiotics in animal husbandry: Applicability and associated risk factors. *Sustainability* 12(3):1087.
5. Jha R, Das R, Oak S, Mishra P (2020) Probiotics (direct-fed microbials) in poultry nutrition and their effects on nutrient utilization, growth and laying performance, and gut health: a systematic review. *Animals* 10(10):1863.
6. Yaqoob MU, Wang G, Wang M (2022) An updated review on probiotics as an alternative of antibiotics in poultry. *Anim Biosci* DOI: 10.5713/ab.21.0485.
7. Castanon JIR (2007) History of the use of antibiotic as growth promoters in European poultry feeds. *Poult Sci* 86(11):2466–2471.
8. Rivera-Pérez W, Barquero-Calvo E, Chaves AJ (2021) Effect of the use of probiotic *Bacillus subtilis* (QST 713) as a growth promoter in broilers: An alternative to bacitracin methylene disalicylate. *Poult Sci* 100(9): 101372.
9. Garriga M, Pascual M, Monfort JM Hugas M (1998) Selection of lactobacilli for chicken probiotic adjuncts. *J Appl Microbiol* 84: 125–132.
10. Siegel PB, Gross WB (1980). Production and persistence of antibodies in chickens to sheep erythrocytes. 1. Directional selection. *Poult Sci* 59(1): 1–5.
11. Cheng S, Lamont SJ (1988) Genetic analysis of immunocompetence measures in a white leghorn chicken line. *Poult Sci* 67(7): 989–995.

12. Snedecor GW, Cochran WG (1994) Statistical method, 6th ed. The Iowa State University Press, Ames, Iowa, USA.
13. Ramlucken U, Ramchuran SO, Moonsamy G, Lalloo R, Thantsha MS, Van Rensburg C J (2020) A novel *Bacillus* based multi-strain probiotic improves growth performance and intestinal properties of *Clostridium perfringens* challenged broilers. *Poult Sci* 99(1):331–341.
14. Rehman A, Arif M, Sajjad N, Al-Ghadi MQ, Alagawany M, Abd El-Hack ME, Swelum AA (2020). Dietary effect of probiotics and prebiotics on broiler performance, carcass, and immunity. *Poult Sci* 99(12): 6946–6953.
15. Zhang B, Zhang H, Yu Y, Zhang R, Wu Y, Yue M, Yang C (2021) Effects of *Bacillus Coagulans* on growth performance, antioxidant capacity, immunity function, and gut health in broilers *Poult Sci* 100(6):101168.
16. Sun Y, Zhang Y, Liu M, Li J, Lai W, Geng S, Zhang L (2022) Effects of dietary *Bacillus amyloliquefaciens* CECT 5940 supplementation on growth performance, antioxidant status, immunity, and digestive enzyme activity of broilers fed corn-wheat-soybean meal diets. *Poult Sci* 101(2): 101585.
17. Chen X, Ishfaq M, Wang J (2022) Effects of *Lactobacillus salivarius* supplementation on the growth performance, liver function, meat quality, immune responses and *Salmonella Pullorum* infection resistance of broilers challenged with Aflatoxin B1. *Poult Sci* 101(3):101651.
18. Qiu K, Wang X, Zhang H, Wang J, Qi G, Wu S (2022). Dietary Supplementation of a New Probiotic Compound Improves the Growth Performance and Health of Broilers by Altering the Composition of Cecal Microflora. *Biology*, 11(5):633.
19. Ramarao S, Reddy M, Raju M, Panda A (2004) Growth, nutrient utilization and immunocompetence in broiler chicken fed probiotic, gut acidifier and antibacterial compounds. *Ind J Poult Sci* 39(2):125–130.
20. Ergün A Yalcin S, Sacakli P (2000) The usage of probiotic and zinc bacitracin in broiler rations. *Ankara Üniversitesi Veteriner Fakültesi Dergisi* 47(3): 271–280.
21. Arslan C (2004) Effect of dietary probiotic supplementation on growth performance in the rock partridge (*Alectoris graeca*). *Turk J Vet Anim Sci* 28(5): 887–891.
22. Gyawali I, Zeng Y, Zhou J, Li J, Wu T, Jiang Q, Zhu C (2022). Effect of Novel *Lactobacillus paracaesi* microcapsule on growth performance, gut health and microbiome community of broiler chickens. *Poult Sci* 101912.
23. Idoui T, Boudjerda J, Karam NE, Leghouchis E (2012) Identification of probiotics properties of *Lactobacillus* strains from poultry crop, Book of Proceedings, 2nd Mediterranean Summit of WPSA. pp. 545–548.
24. Vineetha PG, Tomar S, Saxena VK, Susan C, Sandeep S, Adil K, Mukesh K (.2016) Screening of *Lactobacillus* isolates from gastrointestinal tract of guinea fowl for probiotic qualities using in vitro tests to select species-specific probiotic candidates. *Br Poult Sci* 57(4): 474–482.
25. Ashraf M, Arshad M, Siddique M and Muhammad G (2009) In vitro screening of locally isolated *Lactobacillus* species for probiotic properties. *Pak Vet J* 29(4): 186–190.

26. Soomro AH, Masud, T Anwaar K (2002) Role of lactic acid bacteria (LAB) in food preservation and Human Health: A Review. *Pak J Nutr* 1: 20–24.
27. Simova ED, Beshkova DB, Dimitrov ZP (2009) Characterization and antimicrobial spectrum of bacteriocins produced by lactic acid bacteria isolated from traditional Bulgarian dairy products. *J Appl Microbiol* 106: 692–701.
28. Vineetha PG, Tomar S, Saxena VK, Kapgate M, Suvarna A, Adil K (2017) Effect of laboratory isolated *Lactobacillus plantarum* LGFCP 4 from gastrointestinal tract of guinea fowl on growth performance, carcass traits, intestinal histomorphometry and gastrointestinal microflora population in broiler chicken. *J Anim Physiol Anim Nutr* 101(5): 362–370.
29. Haghighi HR, Gong J, Gyles C L, Hayes MA, Zhou H, Sanei B, Sharif S (2006) Probiotics stimulate production of natural antibodies in chickens. *Clin Vaccine Immuno* 13(9): 975–980.
30. Khaksefidi A Ghoorchi T (2006) Effect of probiotic on performance and immunocompetence in broiler chicks. *J Poult Sci* 43(3): 296–300.
31. Li SP, Zhao XJ, Wang JY (2009) Synergy of *Astragalus polysaccharides* and probiotics (*Lactobacillus* and *Bacillus cereus*) on immunity and intestinal microbiota in chicks. *Poult Sci* 88: 519–525.
32. Stringfellow K, Caldwell D, Lee J, Mohnl M, Beltran R, Schatzmayr G Farnell M (2011) Evaluation of probiotic administration on the immune response of coccidiosis-vaccinated broilers. *Poult Sci* 90(8): 1652–1658.
33. Mahmoud S, Shukry M, Saad M (2013) Lymphocytic proliferation and interleukin-2 production in chickens supplemented with growth promoters. *Res Opi Anim Vet Sci* 3(3): 68–72.
34. Duclos MJ, Chevalier B, Goddard Cand JS. (1993) Regulation of amino acid transport and protein metabolism in myotubes derived from chicken muscle satellite cells by insulinlike growth factor-I. *J Cell Physiol* 157(3): 650–657.
35. Saleh AA, Amber K, El-Magd MA, Atta MS, Mohammed AA, Ragab MM, El-Kader HA (2014) Integrative Effects of Feeding *Aspergillus awamori* and fructooligosaccharide on Growth Performance and Digestibility in Broilers: Promotion Muscle Protein Metabolism. *Bio Med Res Int* 21(2): 41–50.
36. Okumura J, Kita K (1999) Recent Advance in the Relationship between Endocrine Status and Nutrition in Chickens. *Asian-Australas J Anim Sci* 12: 1135–1141.
37. Guobin, C, Xiangping L., Jing L (2011) Temporal and spatial expression of the pax-7 gene during chicken embryo and postnatal development. *J Anim Vet Adv* 10(1): 1785–1788
38. Yakar S, Rosen C.J, Beamer WG, Ackert-Bicknell C.L., Wu Y, Liu JL, LeRoith D (2002) Circulating levels of IGF-1 directly regulate bone growth and density. *J Clin Invest* 110(6): 771–781.
39. Kita K, Kato S, Yaman MA, Okumura J, Yokota H (2002) Dietary L-carnitine increases plasma insulin-like growth factor-I concentration in chicks fed a diet with adequate dietary protein level. *Br. Poult Sci* 43(1): 117–121.
40. Kaiser M.G, Cheeseman JH, Kaiser P, Lamont SJ (2006) Cytokine expression in chicken peripheral blood mononuclear cells after in vitro exposure to *Salmonella enterica* serovar Enteritidis. *Poult Sci* 85(11): 1907–1911.

41. Lynagh GR, Bailey M Kaiser P (2000) Interleukin-6 produced during both murine and avian *Eimeria* infections. *Vet Immunol Immunopathol* 76: 89–102.
42. Wigley P, Kaiser P (2003) Avian cytokines in health and disease. *Revista Brasileira de Ciência Avícola*. 5(1): 1–14.
43. Chichlowski M, Croom WJ, Edens FW, McBride BW, Qiu R, Chiang CC, Daniel LR, Havenstein GB, Koc. MD (2007) Microarchitecture and spatial relationship between bacteria and ileal, cecal, and colonic epithelium in chicks fed a direct-fed microbial, PrimaLac, and salinomycin. *Poult Sci* 86: 1121–1132.
44. Lei Y, Zhang KY, Ding XM, Bai SP, Jung SC (2009) Effect of probiotics on growth performance, development of small intestinal tract and microbial populations in broilers. *Agric Sci Technol* 3: 24–31.
45. Isolauri E, Kirjavainen PV, Salminen S (2002) Probiotics: a role in the treatment of intestinal infection and inflammation. *Gut* 50(3): 54–59.
46. Hua MC, Lin TY, Lai MW, Kong MS, Chang HJ Chen CC (2010) Probiotic Bio-Three induces Th1 and anti-inflammatory effects in PBMC and dendritic cells. *World J Gastroenterol* 16(28): 3529.
47. Ashraf R, Shah NP (2014) Immune system stimulation by probiotic microorganisms. *Crit Rev Food Sci Nutr*. 54: 938–956.
48. Zagato E, Mileti E, Massimiliano L, Fasano F, Budelli A, Penna G, Rescigno M (2014) *Lactobacillus paracasei* CBA L74 Metabolic products and fermented milk for infant formula have anti-inflammatory activity on dendritic cells in vitro and protective effects against colitis and an enteric pathogen in vivo. *PloS one*. 9(2):87615.
49. Couper KN, Blount DG, Riley EM (2008) IL-10: the master regulator of immunity to infection. *J Immuno* 180(9): 5771–5777.
50. Manzanillo P, Eidenschenk C, Ouyang W (2015) Deciphering the crosstalk among IL- 1 and IL-10 family cytokines in intestinal immunity. *Trends Immunol* 36(8): 471–478.
51. Christensen HR, Frøkiær H, Pestka JJ (2002) *Lactobacilli* differentially modulate expression of cytokines and maturation surface markers in murine dendritic cells. *J Immuno* 168(1): 171–178.
52. Chen ML, Pittet MJ, Gorelik L, Flavell RA, Weissleder R, Von Boehmer H, Khazaie K (2005) Regulatory T cells suppress tumor-specific CD8 T cell cytotoxicity through TGF- β signals in vivo. *Proc Nat Acad Sci* 102(2): 419–424.
53. Chen CY, Tsen HY, Lin CL, Yu B and Chen CS (2012) Oral administration of a combination of select lactic acid bacteria strains to reduce the *Salmonella* invasion and inflammation of broiler chicks. *Poult Sci* 91(9): 2139–2147.
54. Haghighi HR, Gong J, Gyles CL, Hayes MA, Zhou H, Sanei B, Sharif S (2006) Probiotics stimulate production of natural antibodies in chickens. *Clin Vaccine Immuno* 13(9): 975–980.
55. Kannaki TR, Reddy MR, Shanmugam M, Verma PC, Sharma RP (2010) Chicken toll-like receptors and their role in immunity. *Worlds Poult Sci J* 66(4): 727–738.

Tables

Table 1. Details of the primers used for real-time PCR analysis of gene in muscle and ileum tissue of broiler

S.No.	Gene	Primer	Annealing temperature (°C)	Product size (bp)	Accession no.
1	GAPDH	F: 5'-CCGTCCTCTCTGGCAAAGTCC-3' R: 5'-AGCCCCAGCCTTCTCCATG-3'	58	264	NM204305
2	IL-6	F: 5' GAAATCCCTCCTCGCCAATCTGA -3' R: 5'TGAAACGGAACAACACTGCCATCT-3'	58	219	JN639847
3	IL-10	F: 5'- TGCGGGAGCTGAGGGTGAAGTTTG-3' R: 5'-CGCGGGGCTGGGCTGAGAG-3'	52	215	AJ621614
4	TLR-4	F: 5'- GTGCCACATCCATACAATAGAAGA-3' R: 5'-ATGGCCCAGATTCAGCTCCT-3'	56	256	JQ280467
5	IGF-1	F: 5' GGTGCTGAGCTGGTTGATGC-3' R: 5'CGTACAGAGCGTGCAGATTTAGGT-3'	58	127	JN942578
6	IGF-1R	F: 5'TAGATCCCTCTACCCTCCAA-3' R: 5'TCTGAAGATCCACTGAGGTACAG-3'	57	291	Heck <i>et al.</i> , 2003

Table 2: Body weight and Body weight gain in broilers fed on different dietary treatments.

Treatment	Body Weight (g)			Body Weight gain (g)		
	Day old	3 week	6 week	0-3 week	4-6 week	0-6 week
T1	39.85	533.55	1554.67 ^a	493.70	676.71	1514.82 ^a
T2	41.45	555.88	1631.86 ^{bc}	515.43	722.96	1591.41 ^{bc}
T3	40.00	543.35	1576.21 ^{abc}	503.36	689.34	1536.21 ^{abc}
T4	38.34	539.59	1561.25 ^{ab}	499.25	675.37	1520.91 ^{ab}
T5	40.14	546.40	1570.11 ^{ab}	506.26	681.07	1529.97 ^{ab}
T6	38.10	553.15	1586.91 ^{abc}	513.05	695.50	1546.81 ^{abc}
T7	39.05	555.43	1599.73 ^{abc}	515.37	706.91	1559.68 ^{abc}
T8	41.19	557.01	1631.69 ^{bc}	516.82	717.53	1591.50 ^{bc}
T9	40.17	562.05	1646.16 ^c	521.88	724.32	1605.99 ^c
P-value	0.230	0.130	0.031	0.166	0.051	0.033

Means with different superscript within column differ significantly (P<0.05).

Table 3: Feed intake and feed conversion ratio (FCR) in broilers fed on different dietary treatments.

Treatment	FCR			Feed Intake (g)		
	0-3 Week	4-6 Week	0-6 Week	0-3 Week	4-6 Week	0-6 Week
T1	1.65	3.23	1.98	813.73	2182.44	2946.17
T2	1.61	2.97	1.87	831.42	2133.75	2965.17
T3	1.62	3.09	1.92	816.24	2132.63	2948.86
T4	1.68	3.19	1.97	838.02	2145.94	2983.96
T5	1.63	3.17	1.95	824.72	2151.21	2975.93
T6	1.59	3.12	1.93	817.77	2169.46	2987.23
T7	1.60	3.06	1.91	822.46	2157.32	2979.79
T8	1.60	3.02	1.88	826.84	2163.77	2990.61
T9	1.56	2.97	1.85	814.30	2148.09	2982.40
P-value	0.609	0.354	0.452	0.987	0.996	1.00

Table 4: Mean log₁₀ value of Lactobacillus spp. Salmonella spp. and E.coli isolated from broiler chicken intestine.

Treatments	Lactobacillus spp			Salmonella spp			E.coli		
	Crop	Ilium	Caeca	Crop	Ilium	Caeca	Crop	Ilium	Caeca
T1	8.38 ^a	8.61 ^a	8.40 ^a	7.31 ^c	7.14 ^c	7.16 ^b	7.35 ^b	7.17 ^c	7.23 ^c
T2	8.58 ^b	8.69 ^{ab}	8.35 ^a	6.94 ^{ab}	6.99 ^{ab}	6.92 ^a	7.19 ^a	7.00 ^a	6.98 ^a
T3	8.84 ^d	8.87	8.71 ^b	6.92 ^a	7.07 ^{abc}	7.13 ^b	7.26 ^{ab}	7.02 ^{ab}	7.10 ^{abc}
T4	8.69 ^c	8.75 ^{bc}	8.61 ^b	7.28 ^c	7.11 ^{bc}	7.12 ^b	7.33 ^b	7.14 ^{bc}	7.21 ^{bc}
T5	8.81 ^d	8.77 ^{bcd}	8.65 ^b	7.25 ^c	7.10 ^{bc}	7.10 ^b	7.31 ^b	7.13 ^{abc}	7.17 ^{bc}
T6	8.84 ^d	8.80 ^{cd}	8.70 ^b	7.23 ^c	7.06 ^{abc}	7.07 ^b	7.25 ^{ab}	7.13 ^{abc}	7.14 ^{bc}
T7	8.75 ^{cd}	8.86 ^d	8.64 ^b	7.26 ^c	7.05 ^{abc}	6.95 ^a	7.31 ^b	7.12 ^{abc}	7.12 ^{abc}
T8	8.83 ^d	8.97 ^e	8.70 ^b	7.03 ^b	7.02 ^{ab}	6.94 ^a	7.27 ^{ab}	7.10 ^{abc}	7.10 ^{abc}
T9	9.03 ^e	9.03 ^e	8.93 ^c	6.91 ^a	6.95 ^a	6.92 ^a	7.20 ^a	7.03 ^{ab}	7.07 ^{ab}
P-value	0.000	0.000	0.000	0.000	0.024	0.00	0.014	0.052	0.023

Means with different superscript within column differ significantly (P<0.05).

Table 5. Humoral and Cell mediated immune response of broilers under different dietary treatments

Treatments	HA Titre	CMI (mm)
T1	6.25 ^a ±0.40	0.46 ± 0.07 ^a
T2	8.38 ^{cd} ±0.37	0.52 ± 0.04 ^a
T3	8.00 ^{bcd} ±0.67	0.51 ± 0.05 ^a
T4	6.38 ^a ±0.37	0.46 ± 0.03 ^a
T5	6.63 ^{ab} ±0.44	0.51 ± 0.05 ^a
T6	7.75 ^{bc} ±0.55	0.56 ± 0.04 ^{ab}
T7	8.00 ^{bcd} ±0.41	0.59 ± 0.04 ^{ab}
T8	8.75 ^{cd} ±0.32	0.72 ± 0.10 ^b
T9	9.25 ^d ±0.43	0.75 ± 0.10 ^b
P-Value	0.000	0.012

Means with different superscript within column differ significantly (P<0.05).

Figures




Figure 1

Relative fold expression of IGF-1 gene in muscle tissue of broiler chicken at 6 wk of age under different dietary treatments. Different superscripts indicates significant expression at the level of 95% ($p \leq 0.05$)




Figure 2

Relative fold expression of IGF-1R gene in Muscle tissue of Broiler chicken at 6 wks of age under different dietary treatments. Different superscripts indicates significant expression at the level of 95% ($p \leq 0.05$)




Figure 3

Relative fold expression of IL-6 gene in Ileum tissue of broiler chicken at 6 wks of age under different dietary treatments. Different superscripts indicates significant expression at the level of 95% ($p \leq 0.05$)




Figure 4

Relative fold expression of IL-10 gene in Ileum tissue of Broiler chicken at 6 wks of age under different dietary treatments. Different superscripts indicates significant expression at the level of 95% ($p \leq 0.05$)




Figure 5

Relative fold expression of TLR-4 gene in Ileum tissue of Broiler chicken at 6 wks of age under different dietary treatments. Means with different superscripts differ significantly ($p \leq 0.05$)