

Evaluating the Knowledge about Older Patients Quiz and the Kogan's Attitudes towards Old People Scale for Nursing Students' Gerontological Learning Experiences in Ghana.

Diana Abudu-Birresborn (✉ diana.abudu@mail.utoronto.ca)

University of Toronto

Sarah Brennenstuhl

University of Toronto

Martine Puts

University of Toronto

Lynn McCleary

Brock University

Vida Yakong

University for Development Studies

Charlene H Chu

University of Toronto

Lisa Cranley

University of Toronto

Research Article

Keywords: knowledge, attitudes, older adults, nursing students, instrument evaluation, Ghana, Africa

Posted Date: May 20th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1654500/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at BMC Nursing on February 21st, 2023. See the published version at <https://doi.org/10.1186/s12912-023-01195-y>.

Abstract

Background: Understanding nursing students' knowledge about and attitudes towards older adults' using a context-specific survey instrument may help to identify, and design effective learning and teaching materials to improve the care for older adults. The study aimed to evaluate the items on the Knowledge about Older Patients Quiz and the Kogan's Attitudes towards Old People Scale suitable for the African context.

Methods: A cross-sectional study was conducted using second- and third-year nursing students from two public Nursing Training Institutions in Ghana. Data were collected from December 2019 - March 2020 using the Knowledge about Older Patients Quiz and the Kogan's Attitudes Towards Old People Scale. Item response theory was employed to evaluate the difficulty level and discrimination indices of the Knowledge about Older Patients Quiz. Corrected item-to-total correlation analysis was conducted for the Kogan's Attitudes towards Old People scale. The internal consistency for both scales was examined.

Results: Of the 170 participants, 169 returned completed surveys. The mean age of participants was 21years (SD = 3.7), and (54%) were female. Of the 30-items of the Knowledge about Older Patients Quiz, seven items were very difficult for most of the students to choose the correct response, and one was easy, as most of the students chose the correct response. Although, 22 items demonstrated appropriate difficulty level, discrimination indices, were used to select the final 15- items that discriminated moderately between upper and lower 25% performing students. The Kuder-Richardson-20 reliability was 0.30. Considering the Kogan's Attitudes towards Old People scale, 10-items were removed following negative and low corrected item-to-total correlation and a high Alpha coefficient if items were deleted. The final 22-items had a Cronbach alpha coefficient of 0.65.

Conclusion: Evaluation of the scales demonstrated essential content validity and moderate internal consistency for the context of our study. Further research should focus on ongoing context specific refinement of the survey instruments to measure nursing students' knowledge about and attitudes towards the care of older adults in the African context.

Background

The population of persons 60 years and older is growing fast in the African region compared to other regions with associated increase in health care utilization by older adults (1). Ghana, like other countries, is experiencing increasing health care utilization by older adults due to multiple complex chronic conditions such as cardiovascular diseases, stroke, diabetes and hypertension (2) resulting in older adults presenting with multiple geriatric syndromes including delirium, incontinence, frailty and falls (3). These complex chronic conditions coupled with complications requires nurses with sufficient gerontological nursing knowledge to provide adequate and quality nursing care (4).

Evidence suggests that nursing students show little interests in working with older adults for reasons such as heavy workload and insufficient knowledge in gerontology (5). Nursing students' ability to demonstrate their knowledge of gerontological nursing content through their responses to knowledge items or test can provide insight to their competencies in caring for older adults (6). It is necessary to examine nursing students' knowledge and attitudes towards older adults to identify opportunities and gaps to facilitate nursing students' preparation in the care of older adults in Africa. However, there are no validated instruments to examine nursing students' knowledge and attitudes towards older adults in the African context.

In Ghana, there is no specialized gerontological nursing care for older adult patients, due to the lack of post-basic specialty programs in gerontology (7). In addition, there are no public long term care facilities or nursing homes for older adults. Older adults are cared for by generalist nurses in emergency departments and medical-surgical wards along with the general population. It is imperative to enhance nursing students' theoretical knowledge and skills to ensure quality care for older adults. Nursing students' with high theoretical knowledge and positive attitudes with or without the preference to working with older adults are likely to demonstrate adequate competencies in providing care for older adults patients (5, 8). Various instruments have been developed and validated for use in assessing nursing students' preparation, such as knowledge and attitudes scales.

The most widely used survey instruments are the Facts on Ageing Quiz (FAQ) and the Kogan's Attitudes towards Old People (KAOP) scale (9, 10). The Knowledge about Older Patients Quiz (KOP-Q) (11) was recently developed and validated to measure knowledge in acute care settings in the Netherlands. Up until now, the KOP-Q has been used to measure nurses' knowledge about older patients in general hospitals in the Netherlands and USA (4). To the best of our knowledge, the KOP-Q has not been used among nursing students in Ghana. Another commonly used is the Kogan's Attitudes towards Old People (KAOP) scale (9) which has been widely used with a reported Cronbach alpha of 0.70 among nursing students in Asia, (12, 13), middle East (14, 15) and a correlation coefficient of 0.71 in Africa (16). However, Faronbi, Adebowale (16) who used it in Nigeria used only nine items of the scale. The authors did not provide a rationale for the use of the nine items. A probable assumption could be that the nine items were context specific and contributed strongly to the overall coefficient of their survey instrument.

In the current study, the KOP-Q and KAOP were adapted for use with Ghanaian nursing students and did not require translation since English is the official language of instruction. However, questions about how suitable these survey instruments are to the African context remain. The purpose of this paper was to evaluate the suitability of the KOP-Q and the KAOP scale for the African context focusing on item difficulty and discrimination for the KOP-Q and corrected item-to-total correlation for the KAOP scale. This study is part of a larger mixed method research project that examined the learning experiences of nursing students to care for older adults in Ghana (17).

Methods

Study Design and Setting

A cross-sectional survey was conducted using a convenience sample (18) of nursing students from two Public Nursing and Midwifery Training Colleges attached to Teaching hospitals. These colleges were selected to represent the northern and southern sectors of Ghana. The northern sector is primarily Muslim dominated and share ethnic, and cultural traditions which are slightly different from the ethnic and cultural traditions of the Christian dominated southern sectors of Ghana. Second- and third-year nursing students were invited to participate in the study because they had more clinical experience than first-year students which was relevant for answering the questionnaires.

Sample

Using the sample size calculation based on our main study (17), 170 nursing students were invited to participate. The 170 nursing students met the Sahin rule of sample size estimate of at least 150 to determine item parameters accurately in unidimensional dichotomous scales (19).

Procedure

Data were collected using survey questionnaires between December 2019 - March 2020. An information session by the principal investigator (DAB) was conducted in the second- and third-year classrooms. An envelope that contained the questionnaires, information sheet and consent form, and invitation to participate in the second phase of our study and a token of appreciation was given to participants who volunteered. The questionnaire had four sections. Sections A focused on the sociodemographic data, Section B included the KOP-Q, Section C included the KOAP scale and Section D was the Self-efficacy to Care for Older Adults' (GES-COA) scale. Sections A, B and C were the focus of this paper. Completed questionnaires were either dropped in a sealed box that was placed at the reception desk at the school or handed to the principal investigator in a sealed envelope onsite.

Measures

The Knowledge about Older Patients Quiz (KOP-Q) (4, 11), is a unidimensional scale that consists of 30 true or false items to measure nursing students' knowledge in the care of older adults. The KOP-Q examines nurses/nursing students' knowledge on the normal ageing process, depression and delirium in older adults, geriatric syndromes, nutrition and fluid imbalance, polypharmacy, referrals, and family caregivers' support. A correct response scored is +1, and an incorrect response scores zero. The scores are summed for a total score. Face validity of 0.91 and construct validity of 0.70 have been reported (11, 20) in the Netherlands and USA.

The Kogan's Attitude Towards Old People Scale (KAOP) (9) consists of 32- item statements. The items of the KAOP scale focused on issues such as the housing arrangements of older adults, the environment and personal appearance, interpersonal relationships with other age groups and dependence of older adults. Item responses are rated on a 5-point Likert scale, ranging from 1 strongly disagree, to 5 strongly

agree. Half of the statements are negative statements and the other half are positive statements regarding attitudes toward older individuals (21). Negative statements were reverse coded, so that there is one total positive score summed which ranges between 32-160. Using the neutral response score of 3, the maximum neutral response 96 was determined as the cut-off point. Higher scores above 96 indicated positive attitudes and scores lower than 96 indicated negative attitudes. In addition to the two scales, socio-demographic data of the participants were collected to describe the sample. These included age, gender, years of school (second /third year), living with/having lived with an older adult, religion, and region (southern /northern sector).

Statistical analysis

All analysis were undertake using SPSS (IBM version 26)

Analysis of KOP-Q

Item difficulty and discrimination indices were calculated as follows. First, the participant knowledge scores were summed, and the total knowledge scores calculated. Next, data were categorized in ranks from the lowest to the highest scores. Participants were categorized into quartiles; the lowest 25%, the middle 50% and the highest 25%. The lowest 25% was calculated using the total number of correct answers in the quartile and divided by the number of participants in that quarter (42 participants). The highest 25 % was calculated using the total number of correct answers in the highest 25% divided by the number of participants in the quarter (42 participants). Participants within the lowest and highest, 25% were used for the discrimination indices (22, 23).

Item difficulty is the percentage of participants who answered an item correctly. Item difficulty was calculated using the formula:

$$ID = \frac{\text{number of students who answered an item correctly}}{\text{total number of participants}} \times 100$$

The score ranges between 0-100 % (24). The lower the difficulty level, the harder the items, suggesting fewer students chose the correct response. The higher the difficulty level, the easier the items, indicating majority of the students chose the correct response (22, 23).

Optimal item difficulty levels are somewhat higher than halfway between guessing and students actually knowing the correct response to an item (22). For dichotomous (true/false) response choice questions, optimal difficulty ranges between 30-85%. Items less than 30% are considered too difficult and items more than 85% are too easy and are recommended to be removed or revised (25).

However, for the selection and examination of final items for the psychometric property analysis of the scale, discrimination indices (DI) was considered (26). The discrimination index (DI) was calculated using the formula:

**DI = Proportion of participants with highest 25% score –
proportion of participants with the lowest 25% score (27).**

Various factors such as number of items, unidimensionality of the scale and item difficulty impact the DI threshold of items (27). A discrimination index threshold of ≥ 0.2 is recommended to determine which items to retain or remove (27, 28). Using our content judgement some items with a DI of 0.19 were approximating 0.2 and included for analysis (27). The DI calculations were repeated after items were removed one after the other which enhanced the DI for the items retained. The Kuder Richardson (KR20) reliability for the retained items was calculated. KR20 is the dichotomous equivalent of the Cronbach alpha coefficient and appropriate for use with survey instruments that measures dichotomous items such as true/false or yes/no responses (29). A KR20 of 0.70 and above is considered acceptable (30).

Analysis for the KAOP scale

To evaluate and improve the internal consistency of KAOP, we calculated the corrected item-to-total correlations and Cronbach alpha if an item was deleted. Corrected item-to-total indicates how much each item correlates with the overall scale (31). Negative and very low corrected item-to-total correlation suggest that items do not correlate adequately with the overall scale and should be removed (32). Negative and very low corrected-item-to-total correlations were deleted one after the other; the deletions increased the overall Cronbach alpha of the scale (31, 33).

The extent and pattern of missing scale item values were examined using descriptive statistics, 95.8% of the sample provided complete data at the item level. With such a small proportion of missing data, complete case analysis was employed. The complete case analysis based on the amount of missing data did not impact the statistical efficiency due to the minimum required sample estimates (19).

Results

Characteristics of participants

Of the 170 questionnaires distributed, 169 were returned (99%). Participants had a mean age of 21 years (SD= 3.7) years. Of the 169 participants, 54% were female, more than half (64 %) of the participants were Christians. The majority (83%) of participants lived with or were currently living with an older adult. About half (49.1%) of the participants were in the second year of their nursing program and the other half in their third year.

KOP- Q

Overall, 22 out of the 30 original items had difficulty level between 30-85% indicating optimum level (Supplementary Table 1). Seven items had item difficulty levels between 7.10% -28.40% indicating very difficult items. Item SB02 (*For older people, bed rest is important to enhance recovery*) had a difficulty

level of 92.6% indicating that it was an easy item (Supplementary Table 1). Although eight items demonstrated very difficult or easy levels, four of the items; SB09, SB28, SB19 and SB21 were retained based on their discrimination index levels (28) (Supplementary Table 2).

For the discrimination indices of the original 30-items, 15 items discriminated moderately (0.24-0.50) between the upper 25% performing students and the lower 25% performing students and were retained. The total knowledge scores for the 15-items ranged between 0-15. The KR20 reliability was 0.30, which is considered low (34). The final 15 items are presented in Supplementary Table 3.

KAOP Scale

Of the 32-items, 10-items were removed due to low or negative corrected-item-to-total correlations. The items did not contribute satisfactorily to the internal consistency of the scale based on the Cronbach alpha if an item is deleted (Supplementary Table 4). Further removal of items did not increase the Cronbach alpha coefficient. The final 22-items had a total score range between 22-110, with a Cronbach alpha coefficient of 0.65 and was considered satisfactory (34) (Supplementary Table 5).

Discussion

Preparing nursing students to face the changing demands of care due to increasing older adult patient population in acute care hospitals is crucial in lower- and middle- income countries (LMICs), including Ghana. Effective evaluation of knowledge acquired in the care of older adults with validated survey instruments that are context-specific is essential to facilitate the evaluation and revision of the curriculum and improve nursing students' readiness to care for older adults. This study showed that 15-items of the original 30-items of the KOP-Q were suitable for the Ghanaian context, of which four items demonstrated low difficulty levels.

The four items examined topics such as distinguishing signs of depression from normal ageing processes, fluid and nutritional imbalance in older adults, effects of medication on geriatric syndromes and family caregiving support services. These topics are either not in the current gerontological nursing curriculum and are not taught or are inadequately taught which may have accounted for the low difficulty levels of the items among Ghanaian nursing students (35). The outcome was not surprising for the item on family caregivers support services. In Ghana, family caregiving is considered the traditional responsibility of adult children and other family members which has little to no support services from government and other agencies (36). As a result, the topic of family caregiving support services may be missing from and not taught in the gerontological nursing curriculum in Ghana. The low difficulty levels of the items highlight a broader gap in gerontological nursing education in the Ghanaian context and can place older adults at a greater risk for poorer outcomes including longer length of stay, increased morbidity, and mortality (37, 38). Thus, there is the need to reinforced nursing students gerontological nursing education to enhance their readiness to meet the complex and demanding needs of older adults (5, 39).

A high percentage of nursing students correctly responded to an item that examined bedrest and recovery in older adults demonstrating the item was a very easy item. It is generally known that bedrest enhances recovery in older adults and is likely the reason for the high percentage score of the item. Very high or low difficulty items make it challenging for assessors to determine which students actually know the content and those who do not know the content and resort to guessing (40). Using discrimination indices, the final items that met both difficulty and discriminatory qualities and measure substantive gerontological nursing topics/areas were retained.

Our study also showed that 22 -items of the original 32 items of the KAOP scale would be applicable for our geographical context. Ten items showed redundancy and did not contribute significantly to the internal consistency of the scale. The remaining 22-items demonstrated moderate internal consistency which is acceptable for analysis. This is similar to findings of a study conducted to assess the validity of the KAOP scale in Thailand, Myanmar, and Indonesia where moderate internal consistency of the scale was reported (13). Researchers have maintained that the number of items of a scale contributes to the internal consistency (41) which may have accounted for the internal consistency of the KAOP scale in this study.

The low internal consistency of the KOP-Q was not surprising because knowledge tests such as this are often very broad in content and measuring multiple theoretical, conceptual, and clinical aspects of a topic. This can result in heterogenous and unrelated items and subsequently a low internal consistency (34, 42, 43). The KR20 reliability is influenced by the difficulty levels, the number of items and the variance of the overall scores (41, 44). Increasing the number of items and students content knowledge about the difficult topics through effective teaching and learning designs can refine the KOP-Q for the African context (41, 44, 45).

Strengths and Limitations

This study was an exploratory study with a high response rate. Despite the low KR20 reliability, decisions were made based on the substantive content concepts in the care of older adults in acute care settings, which discriminated well among students with high and low knowledge about care of older adults. Also, there are other limitations to note. The original scale was developed and tested among registered nurses; however, the scale was used among nursing students in this study who may not have had similar education and experience levels as the nurses in the original study. This may have accounted for the low difficulty levels and discrimination indices of the items. The results should be interpreted with caution.

Implications for nursing education, practice, and research

The present study has the potential to inform nursing education by highlighting nursing students' difficulties and gaps in fundamental topics in gerontological nursing education. The study highlights the potential to improve practice through continuing career development programs to bridge the fundamental knowledge gaps in gerontological nursing care in Ghana. The present study highlights the importance of using substantive content and rigorous methods in research to ensure survey instruments are valid,

appropriate, and context-specific to test students' knowledge. Further testing and adaptation of the KOP-Q should be conducted in LMICs to ensure the psychometric properties are consistent using large samples, especially for the African region.

Conclusion

The revised KOP-Q and the KAOP scale demonstrated substantive content and acceptable internal consistency. Future research should consider the development and psychometric testing of instruments in the care of older adults specific to the African context.

Declarations

Ethical approval and consent to participate

Ethics approval was obtained from the University of Toronto Health Sciences Research Ethics Board (#13408) and the Ghana Health Service Ethics Review Committee (GHS-ERC-001/06/19). Principals of the Nursing and Midwifery Training institutions granted permission for recruitment through the Regional Health Directorates (RHD) of the two Colleges. Participants provided written informed consent before data collection. All methods were carried out in accordance with relevant guidelines and regulations.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study are not publicly available because the datasets are part of a larger study which contains confidential material, but are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

Martine Puts is supported by the Canada Research Chair for the Care of Frail Older Adults.

Authors' contributions

DAB: Study conception and design, data collection, analysis and interpretation of results, draft manuscript preparation and revisions. SB: Study conception and design, data collection, analysis and interpretation of results, draft manuscript preparation and revisions.

MP: Study conception and design, data collection, analysis and interpretation of results, draft manuscript preparation and revisions. LM: Study conception and design, data collection, analysis and interpretation of results, draft manuscript preparation and revisions. VY: Study conception and design, data collection, analysis and interpretation of results, draft manuscript preparation and revisions. CHC: Study conception and design, data collection, analysis and interpretation of results, draft manuscript preparation and revisions

LC: Study conception and design, data collection, analysis and interpretation of results, draft manuscript preparation and revisions. All authors read and approved the final draft of the manuscript.

Acknowledgements

The authors acknowledge the faculty of the two public Nursing and Midwifery Training Colleges for their support and granting permission for the recruitment of nursing students for the study. We also acknowledge the nursing students who voluntarily participated in the study.

References

1. Kamiya Y. Overview of demographic ageing in Africa. Population Division, United Nations Department of Economic and Social Affairs. 2016.
2. de-Graft Aikins A, Addo J, Ofei F, Bosu WK, Agyemang C. Ghana's Burden of Chronic Non-Communicable Diseases: Future Directions in Research, Practice and Policy. *Ghana Medical Journal*. 2012;46(2 Suppl):1–3.
3. Inouye SK, Studenski S, Tinetti ME, Kuchel GA. Geriatric syndromes: clinical, research, and policy implications of a core geriatric concept. *Journal of the American Geriatrics Society*. 2007;55(5):780–91.
4. Dikken J, Hoogerdujin JG, Shortridge-Baggett L, Klaassen S, Schuurmans MJ. Measurement of nurses' attitudes and knowledge regarding acute care older patients: Psychometrics of the OPACS-US combined with the KOP-Q. *Geriatric Nursing*. 2017;38(393–397).
5. Abudu-Birresborn D, McCleary L, Puts M, Yakong V, Cranley L. Preparing Nurses and Nursing Students to Care for Older Adults in Lower and Middle-Income Countries. A Scoping Review. *International journal of nursing studies*. 2019;92:121–34.
6. Gibson SK. Social Learning (Cognitive) Theory and Implications for Human Resource Development. *Advances in Developing Human Resources*. 2004;6(2):193–210.
7. Nursing and Midwifery Council. Revised Curriculum For the Training of Nurses and Midwives in Ghana. Ghana, Accra 2016.
8. Bandura A. Social cognitive theory: An agentic perspective. *Annual review of psychology*. 2001;52(1):1–26.
9. Kogan N. Attitudes toward old people: The development of a scale and an examination of correlates. *Journal of Abnormal and Social Psychology*,. 1961(62):44–54.

10. Palmore EB. The facts on aging quiz: part two. *The Gerontologist*. 1981(2):431–7.
11. Dikken J, Hoogerduijn JG, Kruitwagen C, Schuurmans MJ. Content Validity and Psychometric Characteristics of the “Knowledge about Older Patients Quiz” for Nurses Using Item Response Theory. *Journal of the American Geriatrics Society*. 2016;64(11):2378–83.
12. Runkawatt V, Gustafsson C, Engström G. Different Cultures but Similar Positive Attitudes: A Comparison between Thai and Swedish Nursing Students' Attitudes toward Older People. *Educational Gerontology*. 2013;39(2):92–102.
13. Runkawatt V, Kerdchuen K, Tipkanjanaraykha K, Ubolwan K, Tawetanawanich Y, Nasirin C, et al. Kogan's Attitude Toward Old People: evaluation of validity and reliability assessment among nursing students in Thailand, Myanmar, and Indonesia. *Asian Journal for Public Opinion Research*. 2016;3(3):145–55.
14. Celik SS, Kapucu S, Tuna Z, Akkus Y. Views and attitudes of nursing students towards ageing and older patients. *Australian Journal of Advanced Nursing*. 2010;27(4):24–30.
15. Hweidi IM, Al-Obeisat SM. Jordanian nursing students' attitudes toward the elderly. *Nurse Education Today*. 2006;26(1):23–30.
16. Faronbi JO, Adebawale O, Faronbi GO, Musa OO, Ayamolowo SJ. Perception knowledge and attitude of nursing students towards the care of older patients. *International Journal of Africa Nursing Sciences*. 2017;7:37–42.
17. Abudu-Birresborn D. *Nursing Students' Self-Efficacy to Care for Older Adults in Acute Care Settings in Ghana: A Mixed Method Study*: University of Toronto (Canada); 2022.
18. Polit DF, Beck CA. *Nursing Research: Generating and Assessing Evidence for Nursing Practice*. 9th Edition Wolters Kluwer / Lippincott Williams & Wilkins. 2012.
19. Sahin A, Anil D. The effects of test length and sample size on item parameters in item response theory. 2017.
20. Dikken J, Hoogerduijn JG, Klaassen S, Lagerwey MD, Shortridge-Baggett L, Schuurmans MJ. The Knowledge-about-Older-Patients - Quiz (KOP-Q) for nurses: Cross-cultural validation between the Netherlands and United States of America. *Nurse Educ Today*. 2017;55:26–30.
21. Allen M. *The SAGE Encyclopedia of Communication Research Methods*. 2017.
22. Burud I, Nagandla K, Agarwal P. Impact of distractors in item analysis of multiple choice questions. *International Journal of Research in Medical Sciences*. 2019;7(4):1136–9.
23. Taib F, Yusoff MSB. Difficulty index, discrimination index, sensitivity and specificity of long case and multiple choice questions to predict medical students' examination performance. *Journal of Taibah University Medical Sciences*. 2014;9(2):110–4.
24. Office of Educational Assessment. *Understanding Item Analysis*. University of Washington. 2020.
25. Odukoya JA, Adekeye O, Igbinoba AO, Afolabi A. Item analysis of university-wide multiple choice objective examinations: the experience of a Nigerian private university. *Quality & quantity*. 2018;52(3):983–97.

26. Toland MD. Practical Guide to Conducting an Item Response Theory Analysis. *The Journal of Early Adolescence*. 2014;34(1):120–51.
27. Chiavaroli N, Familiari M. When Majority Doesn't Rule: The Use of Discrimination Indices to Improve the Quality of MCQs. *Bioscience Education*. 2011;17(1):1–7.
28. Rush BR, Rankin DC, White BJ. The impact of item-writing flaws and item complexity on examination item difficulty and discrimination value. *BMC Med Educ*. 2016;16(1):250.
29. Mike Allen. Reliability, Kuder-Richardson Formula. In *The SAGE Encyclopedia of Communication Research Methods*. 2017.
30. McGartland Rubio D. Alpha Reliability. In: Kempf-Leonard K, editor. *Encyclopedia of Social Measurement*. New York: Elsevier; 2005. p. 59–63.
31. Lang L, Zhang L, Zhang P, Li Q, Bian J, Guo Y. Evaluating the reliability and validity of SF-8 with a large representative sample of urban Chinese. *Health and Quality of Life Outcomes*. 2018;16(1):55.
32. Wang M, Batt K, Kessler C, Neff A, Iyer NN, Cooper DL, et al. Internal consistency and item-total correlation of patient-reported outcome instruments and hemophilia joint health score v2.1 in US adult people with hemophilia: results from the Pain, Functional Impairment, and Quality of life (P-FiQ) study. *Patient Prefer Adherence*. 2017;11:1831–9.
33. Mun J, Mun K, Kim S-W. Exploration of Korean Students' Scientific Imagination Using the Scientific Imagination Inventory. *International Journal of Science Education*. 2015;37(13):2091–112.
34. Taber KS. The Use of Cronbach's Alpha When Developing and Reporting Research Instruments in Science Education. *Research in Science Education*. 2018;48(6):1273–96.
35. Considine J, Botti M, Thomas S. Design, format, validity and reliability of multiple choice questions for use in nursing research and education. *Collegian*. 2005;12(1):19–24.
36. Awuviry-Newton K, Byles J, Tavener M. Evolution of familial care and support for older people in Ghana: A literature review 2017.
37. Oates LL, Price CI. Clinical assessments and care interventions to promote oral hydration amongst older patients: a narrative systematic review. *BMC Nursing*. 2017;16(1):4.
38. Vanderwee K, Clays E, Bocquaert I, Verhaeghe S, Lardennois M, Gobert M, et al. Malnutrition and nutritional care practices in hospital wards for older people. *J Adv Nurs*. 2011;67(4):736–46.
39. Dahlke SA, Phinney A, Hall WA, Rodney P, Baumbusch J. Orchestrating care: nursing practice with hospitalised older adults. *International Journal of Older People Nursing*. 2015;10(4):252–62.
40. Schreyer Institute for Teaching Excellence. *Guide to Item Analysis*. 2019.
41. Dillman DA, Smyth JD, Melani L. Internet, mail, and mixed-mode surveys: The tailored design method. 2009.
42. Adadan E, Savasci F. An analysis of 16–17-year-old students' understanding of solution chemistry concepts using a two-tier diagnostic instrument. *International Journal of Science Education*. 2012;34(4):513–44.

43. McGahee TW, Ball J. How to read and really use an item analysis. *Nurse educator*. 2009;34(4):166–71.
44. El-Uri FI, Malas N. Analysis of use of a single best answer format in an undergraduate medical examination. *Qatar Med J*. 2013;2013(1):3–6.
45. Gray JR, Grove SK, Sutherland S. *The Practice of Nursing Research: Appraisal, Synthesis, and generation of Evidence*. Eight Edition Elsevier, Inc. 2017.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterial.docx](#)