

Nuclear Lamin A in Rotator Cuff Tear Margin Tenocytes: An Antiapoptotic and Cell Mechanostat Factor.

Stefano Gumina

Universita degli Studi di Roma La Sapienza

Barbara Peruzzi

Ospedale Pediatrico Bambino Gesù: Ospedale Pediatrico Bambino Gesù

Martina Leopizzi

Universita degli Studi di Roma La Sapienza

Natale Porta

Universita degli Studi di Roma La Sapienza

Valeria Di Maio

Universita degli Studi di Roma La Sapienza

Carlo Della Rocca

Universita degli Studi di Roma La Sapienza

vittorio candela (✉ vittorio.candela@yahoo.it)

Sapienza University of Rome: Universita degli Studi di Roma La Sapienza <https://orcid.org/0000-0002-5962-2598>

Research article

Keywords: Nuclear Lamin A, Mechanostat Factor, Antiapoptotic

Posted Date: February 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-165606/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

The network of intermediate filament proteins underlying the inner nuclear membrane forms the nuclear lamina. A and B-type lamins are the major components of the nuclear lamina. Lamins function in many nuclear activities. The role of lamin-A and transcription factors (NF- κ B) as anti-apoptotic is well documented. Recently, lamin-A has also been considered as mechanosensor protein able to maintain nuclear integrity from mechanical insults.

We aimed to verify how lamin-A expression varies in healthy cuff cells and in those with different sized tears where various mechanical stresses are present.

Methods

Forty-three patients with rotator cuff tear (RCT) [23M-20F, mean age (SD): 63.5 (6.1)] were enrolled. Tissue samples excised from the most medial point of tear margins were analyzed for lamin-A expression by immunohistochemistry. Controls were represented by samples obtained by seven healthy supraspinatus tendons excised from patients submitted to reverse shoulder prosthesis implant [3M-4F, mean age (SD): 67.1 (6.3)]. The intensity of staining was graded, and an H-score was assigned. Statistical analysis was performed.

Results

Our study revealed a moderate intensity of lamin-A in the healthy cuff tendons, a higher expression of this protein in the small tears, and a significant decrease of lamin-A with increasing tear size ($p < 0.0001$).

Conclusions

Our study emphasizes the importance of early repair of small RCTs since nuclear stability is maintained, and the cellular function is protected by lamin-A overexpression. High re-tear of massive cuff repair could be due to cellular apoptosis and nuclear modifications induced by lamin-A lack.

Level of Evidence: III

Introduction

In eukaryotic cells, the nuclear envelope separates chromosome and subnuclear structures from the cytoplasm and regulates the trafficking of macromolecules, including proteins, DNAs and RNAs, between nuclear and cytoplasmic compartments. This transport takes place across nuclear pore complexes that are spaced across the nuclear envelope. The latter is constituted by a double-membrane structure consisting of an inner and outer nuclear membrane. Underlying the inner nuclear layer is a network of intermediate filament proteins forming the nuclear lamina (20–50 nm thick) [1]. The lamina structurally

supports the nuclear envelope, determines the overall shape of the interphase nucleus, and provides an anchoring site for nuclear pore complexes [2]. In vertebrates, A-type and B-type lamins are the significant components of the nuclear lamina. Both are type V intermediate filament proteins that are found exclusively in the nucleus. The two subtypes (A- and B-type lamins) are distinguished by their protein sequences, physical properties, and expression profiles [2]. In humans, A-type lamin is encoded by the LMNA gene, while two separate genes (LMNB1 and 2) encode B-type lamins. By alternative splicing, the LMNA gene also produces lamin C isoforms lacking the final two exons found in lamin A [3].

Lamins function in many nuclear activities such as DNA replication [4], chromatin organization [5], transcriptional regulation, nuclear shape maintenance, and signal transduction [6], nuclear stability [7].

Recently, Swift et al. [8] have also attributed to Lamin-A a role as mechanosensor protein, focusing their attention on the correlation between lamin protein levels and the extracellular matrix stiffness and mechanics. A recent biomechanical study [9] conducted on healthy and injured cuff tendons belonged to adult rats. It has been observed that the defect group exhibited reduced stiffness, reduced ultimate load, and reduced area under the curve at ultimate stress compared to intact tendon group. Furthermore, the authors stated that transverse strain increased with increasing axial load in the defect group region but did not change for the intact group. How this is translated into the cuff tendon cell is not yet known. Since the role of A-type lamins in maintaining nuclear structural integrity from mechanical insults and cell viability has been demonstrated [10–11], we have conducted an immunohistochemical study to verify how A-type lamin expression varies in healthy cuff tendon cells and in those with different sized tears where inevitable different mechanical stresses are present.

Material And Methods

Fifty-four consecutive patients [25M-29F, mean age (SD): 65.2 (5.18)] with rotator cuff tear (RCT) were enrolled. The diagnosis was obtained after physical examination, standard X-Ray (true AP and axillary views), and MRI of the involved shoulder. Exclusion criteria were as follow traumatic RCT; anti-inflammatory drugs assumption during the two months before surgery; V-shaped or L-shaped lesions, diabetes, rheumatologic diseases, and prior surgery.

Tissue samples excised from the most medial point of tear margins [12] were analyzed for lamin A expression by immunohistochemistry. Seven normal supraspinatus tendon biopsies, excised in patients [3M-4F, mean age (SD): 67.1 (6.3)] with proximal humerus fractures with intact rotator cuff submitted to reverse shoulder arthroplasty, were used as controls.

The Southern California Orthopaedic Institute (SCOI) [13] classification of complete RCTs was used to classify tendon tears intraoperatively. We considered the lesions belonging to type I as small, those of types II and III as large, and those of type IV as massive.

Immunohistochemistry was conducted on RCT sections by following the manufacturer's instructions with HRP-DAB-based kit (Dako LSAB Kit peroxidase; Dako, Carpinteria, CA). RCT sections were deparaffinized

and rehydrated in graded ethanol. Endogenous peroxidase activity was blocked by 3% hydrogen peroxide. Antigen retrieval was performed in EDTA buffer pH 8.0 (UCS Diagnostic). The sections were then incubated overnight at 4°C with a rabbit anti-lamin A/C antibody (A0249, Abclonal; Woburn, MA) and subsequently stained using Dako LSAB Kit peroxidase.

For each sample, the nuclear expression of A-type lamin in tenocytes were evaluated by a pathologist using two criteria: the percentage of immunoreactive cells (semiquantitative method) and the intensity of staining (qualitative method). The percentage of immunoreactive cells was estimated by a medium-magnification (20x) microscope examination of the entire section. The intensity of staining was graded as: negative (0), low staining (1), moderate staining (2), and high staining (3). The percentage of cells at each staining intensity level was calculated, and an H-score was assigned using the following formula: $[1 \times (\% \text{ cells } 1+) + 2 \times (\% \text{ cells } 2+) + 3 \times (\% \text{ cells } 3+)]$. The final score, ranging from 0 to 300 (H1 = 0–50; H2 = 51–100; H3 = 101–150; H4 > 150), gives more relative weight to higher intensity staining in a sample.

All participants signed an informed consent form in accordance with the Declaration of Helsinki; the study obtained the local ethics committee approval (Prot xt/2020).

Statistical analysis

All analyses were completed using GraphPad Prism 6.0. Contingency table data were analyzed by the Fisher's exact (Chi-square) test of independence. Accordingly, to Chi-square analysis requirements, the score 0 of A-type lamin intensity (negative staining) was excluded by Chi-square analysis since no samples resulted negative. The level of significance was set at $\alpha = 0.05$. SPSS 20 (IBM, Chicago, IL) for Windows was used.

Results

The studied group was finally composed of 43 cases [23M-20F, mean age (SD): 63.5 (6.1)] and seven controls [3M-4F, mean age (SD): 67.1 (6.3)]. RCTs were intraoperatively classified in 21 small, 14 large and 8 massive tears.

A panel of lamin A immunohistochemical staining is shown in Fig. 1.

The intensity of staining and the H-score of immunoreactive cells, assessed by microscope analysis, are shown in Fig. 2. As regarding the nuclear intensity score of lamin A immunoreactive cells, no negative samples were counted. Study of the distribution of the intensity staining scores showed the 100% of moderate-stained samples for control tendon samples (Fig. 2A), 52% of high-stained, 38% of moderate-stained and 10% of low-stained samples for the small RCTs (Fig. 2B), 36% of moderate-stained and 64% of low-stained samples for the large RCTs (Fig. 2C) and 50% of both moderate- and low-stained samples for the massive RCTs (Fig. 2D). Analysis of the H-score distribution showed 29% of H4-score and 71% of H3-score for control tendon samples (Fig. 2E), 33% of H4-score, 19% of H3-score, 24% of H2-score samples and 24% of H1-score for the small RCTs (Fig. 2F), no H4-score samples, 21% of H3-score, 21% of

H2-score samples and 57% of H1-score for the large RCTs (Fig. 2G) and no H4- and H3 score samples, 63% of H2-score samples and 38% of H1-score for the massive RCTs (Fig. 2H).

The Chi-square test showed a statistically significant correlation between the nuclear expression of A-type lamin and the size of the RCTs (Table I).

Discussion

The skeletal anatomy's potential importance on the genesis of rotator cuff tear has been studied for decades and remains a controversial subject. Recently, tendon degeneration (age-related or degeneration induced by genetics and medical conditions) seems to be the most credited theory for cuff rupture [14–15]. The origin of RCT is represented by an area of the tendon within a few mm of its insertion characterized by relative hypovascularization [16]. The inadequate blood supply is only partially improved by vascular anastomoses of the critical zone, near the tendon insertion. Micro vascularization may become worse in many patients with smoking and alcoholic habit [17–18], hypertension [19], and lung and other cardiovascular diseases [20]. Obesity is also considered a risk factor for rotator cuff tear [21] because it contributes to peripheral vascular deficiencies through its associations with an increased production of adipokines (leptin; adiponectin; plasminogen activator inhibitor; tumor necrosis factor- α ; angiotensinogen; interleukins 6, 8, 10, and 18). All these molecules induce oxidative stress, inflammation, thrombosis, and endothelial dysfunction [20]. The consequent release of many reactive oxygen species (ROS) may lead to degeneration of the tendon causing oxidative stress and cell apoptosis [20].

When cuff tear occurs, multiple stimuli, both mechanical and inflammatory, lead to the altered expression of proteins; some of them are probably synthesized as an attempt to tendon healing. In this regard, increasing in periostin on rotator cuff margins was attributed to the attempt of the tendon to change its viscoelastic properties to prevent increased damage [20]. Our study revealed a moderate intensity of type V intermediate filament proteins lamin A in the healthy cuff tendons (control group), a higher expression of this protein in the small tears, and a significant decrease of lamin A presence with increasing tear size. Since it was demonstrated that cells lacking lamin A have elevated ROS level [22–23], it is plausible that the initial increasing in lamin A observed in our specimens belonged to patients with small rotator cuff tear is ascribable to an attempt to face up ROS levels. However, this attempt is destined to wane given the progressive cell depletion observed in large and massive tears.

In an immunohistochemical analysis on rotator cuff tear margins, nuclear transcription factors, as NF- κ B have been observed [24]. Furthermore, activated NF- κ B on the tear edge increases with increasing tear size. The role of lamin A and of NF- κ B as anti-apoptotic is well documented [2, 25–27], such as it is well known that the same factors that regulate NF- κ B activation act as neoangiogenesis inductors [24]. Interactions between lamin and nuclear transcription factors have been demonstrated in vivo and in vitro [27–28]. We hypothesize that the increased expression of lamin A in the nuclei of small tear tendon tenocytes is initially related to the NF- κ B activation. Therefore, lamin A's action as anti-apoptotic would

occur not only through the protective effect that lamin carries out on the cell nucleus [7–8, 10] but also through the regulation of the NF- κ B.

A proteomic analysis conducted by Swift and Discher [29] revealed that, in vitro, nuclei of mesenchymal cells on a soft substrate are wrinkled and relaxed. In contrast, on the stiff substrate, they are flattened by stress fibers and appear taut and smooth. Furthermore, the native fold of lamin A is maintained on a stiff substrate, but the total quantity of lamin A protein is upregulated. Lammerding et al. [23] observed that the viability of fibroblasts lacking lamin A was significantly reduced under mechanical strain, while under unstrained conditions were similar to wild-type fibroblasts. Our study revealed a moderate intensity and a high H scores of lamin A in the normal cuff, Instead, the Chi square analysis documented a statistically significant correlation between the nuclear expression of lamin A and rotator cuff tear size; in particular, the greatest expression of lamin A was found in the small tears with a significant decrease in lamin A presence with increasing tear size. Increased lamin A levels, in response to extracellular tension, prevent distortion of the nucleus by physical stress, thereby identifying lamin A as a mechanostat factor in cells [29]. Andarawis-Puri et al. [12], in a biomechanical study, demonstrated that the maximum principal strains were directly medial to the tear, corresponding to the direction of tear propagation and to the site where we performed the biopsies. Locke et al. [9] observed that in rat rotator cuff tendons strain concentrations develop near attachment defects. In a biomechanical study conducted on cadaveric cuff tendons, Miller et al [30] registered that the largest strain is particularly concentrated medial and posterior to the tear. With increasing tear size, we observed a decrease in lamin A staining and the H score. In massive tears, no 3–4 points H score were found. This may be due to the reduction of the mechanical stress acting on an extended margin of the tear (and not in a concentrated area, as in small tears occur) and the progressive fatty infiltration to which the muscle of the involved tendon undergoes. This analysis seems to be supported by Swift e Discher theory [29], according to which mechanical loading can cause inhomogeneous straining, making it beneficial to have mechanisms of lamin regulation that act at the local level of individual cells.

The “cellular protection” mechanism progressively fails, and programmed cell death develops. However, in massive tears, some functional cells remain, probably due to the permanent high levels of NF κ B.

The relatively low number of studied patients represents a limit of our study. However, the evaluation of lamin A immunohistochemical staining is a novel and expensive methodology.

Conclusions

Our study emphasizes the importance of early repair of small RCTs since nuclear stability is maintained, and the cellular function is protected by lamin A overexpression.

This study provides a possible explanation for the high re-tear/no healing rate of massive rotator cuff repair; in fact, intracellular modifications leading to apoptosis has already developed.

Abbreviations

RCT

Rotator Cuff Tear

SCOI

Southern California Orthopaedic Institute

ROS

Reactive Oxygen Species

Declarations

Conflict of Interest: nothing to declare.

Acknowledgements

We are thankful for the support of the data analysts from the Department of Statistics of Sapienza University.

Ethics approval and consent to participate

This study was approved by Institutional Review Board of Sapienza University. All participants gave the consent to participate.

Consent for publication

Obtained.

Authors' contributions

SG provided the ideas of this study. SG and VC collected the data. SG and VC wrote the manuscript. BP, ML, NP, VDM analyzed the data. CDC and BP helped with the interpretation of data. The authors have read and approved the final manuscript.

Funding

This research was not funded.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no conflict of interest.

References

1. Hutchison CJ. Lamins: building blocks or regulators of gene expression? *Nat Rev Mol Cell Biol.* 2002 Nov;3(11):848-58.
2. Burke B, Stewart CL. The laminopathies: the functional architecture of the nucleus and its contribution to disease. *Annu Rev Genomics Hum Genet.* 2006;7:369-405.
3. Tran JR, Chen H, Zheng X, Zheng Y. Lamin in inflammation and aging. *Curr Opin Cell Biol.* 2016 Jun;40:124-130.
4. Ellis DJ, Jenkins H, Whitfield WG, Hutchison CJ. GST-lamin fusion proteins act as dominant negative mutants in *Xenopus* egg extract and reveal the function of the lamina in DNA replication. *J Cell Sci.* 1997 Oct;110 (Pt 20):2507-18.
5. Boyle S, Gilchrist S, Bridger JM, Mahy NL, Ellis JA, Bickmore WA. The spatial organization of human chromosomes within the nuclei of normal and emerin-mutant cells. *Hum Mol Genet.* 2001 Feb 1;10(3):211-9.
6. Foti R, Gnan S, Cornacchia D, Dileep V, Bulut-Karslioglu A, Diehl S, Bunes A, Klein FA, Huber W, Johnstone E, Loos R, Bertone P, Gilbert DM, Manke T, Jenuwein T, Buonomo SC. Nuclear Architecture Organized by Rif1 Underpins the Replication-Timing Program. *Mol Cell.* 2016 Jan 21;61(2):260-73.
7. De Vos WH, Houben F, Kamps M, Malhas A, Verheyen F, Cox J, Manders EM, Verstraeten VL, van Steensel MA, Marcelis CL, van den Wijngaard A, Vaux DJ, Ramaekers FC, Broers JL. Repetitive disruptions of the nuclear envelope invoke temporary loss of cellular compartmentalization in laminopathies. *Hum Mol Genet.* 2011 Nov 1;20(21):4175-86.
8. Swift J, Ivanovska IL, Buxboim A, Harada T, Dingal PC, Pinter J, Pajerowski JD, Spinler KR, Shin JW, Tewari M, Rehfeldt F, Speicher DW, Discher DE. Nuclear lamin-A scales with tissue stiffness and enhances matrix-directed differentiation. *Science.* 2013 Aug 30;341(6149):1240104.
9. Locke RC, Peloquin JM, Lemmon EA, Szostek A, Elliott DM, Killian ML. Strain Distribution of Intact Rat Rotator Cuff Tendon-to-Bone Attachments and Attachments With Defects. *J Biomech Eng.* 2017 Nov 1;139(11):1110071–6.
10. Lammerding J, Fong LG, Ji JY, Reue K, Stewart CL, Young SG, Lee RT. Lamins A and C but not lamin B1 regulate nuclear mechanics. *J Biol Chem.* 2006 Sep 1;281(35):25768-80.
11. Zuela N, Bar DZ, Gruenbaum Y. Lamins in development, tissue maintenance and stress. *EMBO Rep.* 2012 Dec;13(12):1070-8.
12. Andarawis-Puri N, Ricchetti ET, Soslowsky LJ. Rotator cuff tendon strain correlates with tear propagation. *J Biomech.* 2009 Jan 19;42(2):158-63.
13. Snyder SJ. Arthroscopic classification of rotator cuff lesions and surgical decision making. In: *Shoulder arthroscopy.* 2nd ed. Philadelphia: JB Lippincott; 2002. p. 204. ISBN:0781735017, 9780781735018.
14. Gumina S, Villani C, Arceri V, Fagnani C, Nisticò L, Venditto T, Castagna A, Candela V. Rotator Cuff Degeneration: The Role of Genetics. *J Bone Joint Surg Am.* 2019 Apr 3;101(7):600-5.

15. Kim HM, Dahiya N, Teefey SA, Middleton WD, Stobbs G, Steger-May K, Yamaguchi K, Keener JD. Location and initiation of degenerative rotator cuff tears: an analysis of three hundred and sixty shoulders. *J Bone Joint Surg Am.* 2010 May;92(5):1088-96.
16. Blevins FT, Djurasovic M, Flatow EL, Vogel KG. Biology of the rotator cuff tendon. *Orthop Clin North Am.* 1997 Jan;28(1):1-16.
17. Carbone S, Gumina S, Arceri V, Campagna V, Fagnani C, Postacchini F. The impact of preoperative smoking habit on rotator cuff tear: cigarette smoking influences rotator cuff tear sizes. *J Shoulder Elbow Surg.* 2012 Jan;21(1):56-60.
18. Passaretti D, Candela V, Venditto T, Giannicola G, Gumina S. Association between alcohol consumption and rotator cuff tear. *Acta Orthop.* 2016;87(2):165-8.
19. Gumina S, Arceri V, Carbone S, Albino P, Passaretti D, Campagna V, Fagnani C, Postacchini F. The association between arterial hypertension and rotator cuff tear: the influence on rotator cuff tear sizes. *J Shoulder Elbow Surg.* 2013 Feb;22(2):229-32.
20. Dandona P, Aljada A, Chaudhuri A, Mohanty P, Garg R. Metabolic syndrome: a comprehensive perspective based on interactions between obesity, diabetes, and inflammation. *Circulation.* 2005 Mar 22;111(11):1448-54.
21. Gumina S, Candela V, Passaretti D, Latino G, Venditto T, Mariani L, Santilli V. The association between body fat and rotator cuff tear: the influence on rotator cuff tear sizes. *J Shoulder Elbow Surg.* 2014 Nov;23(11):1669-74.
22. Pekovic V, Gibbs-Seymour I, Markiewicz E, Alzoghaibi F, Benham AM, Edwards R, Wenhert M, von Zglinicki T, Hutchison CJ. Conserved cysteine residues in the mammalian lamin A tail are essential for cellular responses to ROS generation. *Aging Cell.* 2011 Dec;10(6):1067-79.
23. Lammerding J, Schulze PC, Takahashi T, Kozlov S, Sullivan T, Kamm RD, Stewart CL, Lee RT. Lamin A/C deficiency causes defective nuclear mechanics and mechanotransduction. *J Clin Invest.* 2004 Feb;113(3):370-8.
24. Gumina S, Natalizi S, Melaragni F, Leopizzi M, Carbone S, Postacchini F, Milani A, Della Rocca C. The possible role of the transcription factor nuclear factor- κ B on evolution of rotator cuff tear and on mechanisms of cuff tendon healing. *J Shoulder Elbow Surg.* 2013 May;22(5):673-80.
25. Beg AA, Baltimore D. An essential role for NF-kappaB in preventing TNF-alpha-induced cell death. *Science.* 1996 Nov 1;274(5288):782-4.
26. Wang CY, Mayo MW, Baldwin AS Jr. TNF- and cancer therapy-induced apoptosis: potentiation by inhibition of NF-kappaB. 1996 Nov 1;274(5288):784-7.
27. Dreuillet C, Tillit J, Kress M, Ernoult-Lange M. In vivo and in vitro interaction between human transcription factor MOK2 and nuclear lamin A/C. *Nucleic Acids Res.* 2002 Nov 1;30(21):4634-42.
28. Lloyd DJ, Trembath RC, Shackleton S. A novel interaction between lamin A and SREBP1: implications for partial lipodystrophy and other laminopathies. *Hum Mol Genet.* 2002 Apr 1;11(7):769-77.
29. Swift J, Discher DE. The nuclear lamina is mechano-responsive to ECM elasticity in mature tissue. *J Cell Sci.* 2014 Jul 15;127(Pt 14):3005-15.

30. Miller RM, Fujimaki Y, Araki D, Musahl V, Debski RE. Strain distribution due to propagation of tears in the anterior supraspinatus tendon. J Orthop Res. 2014 Oct;32(10):1283-9.

Tables

Table I - Results of Chi-square analysis of lamin A/C intensity staining and H-score in RCT patients and control (CTL) tendons.

Samples	Lamin A/C intensity staining				Lamin A/C H-score				
	1	2	3	<i>p</i> -value	1	2	3	4	<i>p</i> -value
CTL	0	7	0	<0.0001	0	0	5	2	0.0012
Small	2	8	11		5	5	4	7	
Large	9	5	0		8	3	3	0	
Massive	4	4	0		3	5	0	0	

Lamin A/C intensity staining: 1: low-intensity; 2: moderate-intensity; 3: high-intensity.

Lamin A/C H-score: 1: 0-50; 2: 51-100; 3: 101-150; 4: 151-200.

Figures




Figure 1

Immunohistochemical staining of control (CTL) and rotator cuff tear samples for lamin A/C. Nuclei are stained by Hematoxylin dye. Scale bar: 200 μ m (upper panels) and 100 μ m (lower panels).




Figure 2

Distribution of lamin A/C immunohistochemical staining intensity (upper panels) and H-scores (lower panels) in controls (A, E), small (B,F), large (C,G) and massive (D,H) rotator cuff tear specimens.