

# Two Fe mining sub-products and three thiol compounds alleviate Fe deficiency in soybean (*Glycine max* L.) grown in a calcareous soil in greenhouse conditions

Zahra Gheshlaghi

Reza Khorassani (✉ [khorassani@um.ac.ir](mailto:khorassani@um.ac.ir))

Ferdowsi University of Mashhad Faculty of Agriculture <https://orcid.org/0000-0003-3397-3987>

Javier Abadia

---

## Research Article

**Keywords:** Iron chlorosis, Iron fertilization, Iron nutrition, Iron solubilization, Thiols

**Posted Date:** June 3rd, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1656181/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Purpose

The aim of this work was to test the hypothesis that Fe mining sub-products and thiols can alleviate a moderate Fe-deficiency in the Strategy I species soybean (*Glycine max*) grown in a calcareous soil in greenhouse conditions.

## Methods

Combinations of three Fe sources [Fe(III)-EDDHA and two Fe mining sub-products, both of them containing Fe oxides and one FeS<sub>2</sub>], and three thiols (glutathione, dithiothreitol and thiophenol) were applied in solution to the soil, three times in a 55 day period, and different parameters related to Fe deficiency were measured. The thiol-mediated solubilization of Fe from the Fe mining sub-products was assessed by measuring in the solution total Fe and the reducible Fe pool using an Fe(II) chelator.

## Results

Application of Fe-EDDHA, the two Fe mining sub-products and the three thiols relieved the Fe deficiency symptoms to different extents, increased the Fe concentrations and contents throughout the plant and changed the redox state of leaves and roots, as judged from the changes in reduced and oxidized glutathione, ascorbate and antioxidant enzymes. When using Fe(III)-EDDHA, the addition of thiols led to a better leaf greening. However, the addition of thiols did not cause further greening in the case of the Fe mining sub-products, in spite of being able to solubilize Fe from them.

## Conclusion

Application of Fe-mining sub-products, thiols and the combination of Fe(III)-EDDHA and thiols could be used to alleviate moderate Fe deficiency in *G. max* grown in a calcareous soil.

## Introduction

Iron (Fe) is very abundant in the Earth crust, but in aerated, high pH soils this metal occurs mainly in Fe(III) oxy-hydroxide forms that are not readily available for plants. Therefore, many crops growing in calcareous soils are affected by Fe deficiency and the produce yield and quality are reduced (Briat et al. 2015). To acquire Fe, dicotyledonous plants use a reduction strategy (Strategy I), that includes the elicitation of a root plasma membrane Fe(III) reductase enzyme (FCR) and a Fe(II) membrane transporter, as well as the secretion, depending on the plant species, of a number of substances, including protons, phenolic compounds and flavins (Abadía et al. 2011; Robe et al. 2021; Gheshlaghi et al. 2021). Gramineous plants use a Fe chelation strategy (Strategy II), consisting in the root secretion of

phytosiderophores (PS), which bind Fe in the rhizosphere, and subsequently the Fe(III)-PS complex is taken up by specific root transporters (Connorton et al. 2017).

Iron deficiency in crops is usually controlled by using synthetic Fe(III)-chelate fertilizers, which have to be applied every year (El Jendoubi et al. 2011). Other common fertilization methodologies include foliar sprays with Fe compounds (Álvarez-Fernández et al. 2004; Rodríguez-Lucena et al. 2010a,b), although the transport of the added Fe from the foliage to other plant organs can be a limiting step (Rios et al. 2016). The application of Fe salts, alone or in combination with acidic substances and organic matter, is generally poorly effective in calcareous soils.

Sub-products from the Fe mining industry contain Fe in the form of different oxides and FeS<sub>2</sub> (Cornell and Schwertmann 1996), and this metal can be solubilized by leaching with inorganic (e.g., HCl and SO<sub>4</sub>H<sub>2</sub>) or organic acids such as oxalic acid (Blesa et al 1987; Cornell and Schindler 1987; Panias et al. 1996; Ambikadevi and Lalithambika 2000). Compounds secreted by plants, including carboxylates, phenolics, flavins and flavonoids, can mobilize Fe from Fe oxides and oxi-hydroxides via reductive solubilization (Abadía et al. 2011; Sisó-Terraza et al. 2016; Robe et al. 2021; Gheshlaghi et al. 2021).

Thiol compounds play many roles in the physiology of plants. The water-soluble antioxidant glutathione (GSH; L-g-glutamyl-L-cysteinyl-glycine) is abundant in plants and animals and has redox and regulatory functions, playing roles in ROS scavenging in the GSH-ascorbate cycle and as an electron donor for the enzyme glutathione peroxidase. Iron deficiency has been shown to cause increases in the root concentrations of GSH and ascorbate (Asc) in *Cucumis sativus* (Zaharieva et al. 1999) and *Beta vulgaris* (Zaharieva and Abadía 2003; Zaharieva et al. 2004). The synthesis of GSH in roots is regulated by nitric oxide (NO) (Matamoros et al. 1999), and it has been shown that thiol compounds play an essential role in Fe-deficiency tolerance and NO-mediated Fe-deficiency signalling in plants (Ramírez et al. 2008, 2013; Shanmugam et al. 2012; Kaya et al. 2020; Khan et al. 2021).

Iron oxides can be reductively solubilized by thiols, including GSH (Gheshlaghi et al. 2020, and references therein). It has been recently shown that when GSH was supplied to the roots of *Medicago scutellata* growing in a Fe-rich rock sand, GSH was capable of mobilize Fe from the substrate, increasing plant Fe concentrations and relieving Fe-deficiency symptoms (Gheshlaghi et al. 2020). Also, supplying GSH to the foliage of *M. scutellata* increased internal Fe availability and relieved Fe-deficiency symptoms (Gheshlaghi et al. 2019). The application of GSH to the soil to control Fe deficiency in plants has not been tested yet. However, the usefulness of this practice may be compromised by the fact that soil bacteria and fungi can consume and degrade exogenous GSH (Vergauwen et al. 2013), thus limiting to some extent the amount of GSH available for Fe reduction processes.

Soybean (*Glycine max*) is a species considered to be sensitive to Fe deficiency, and in alkaline and calcareous soils yield and seed quality is often reduced (Hansen et al. 2004; Merry et al. 2022). Soybean is a Strategy I species, and under Fe deficiency shows increases in the root FCR activity (Jolley et al. 1992), the capacity to acidify the medium (Zocchi and Cocucci 1990; Rahman et al. 2022), and the

expression of the FCR -*GmFRO2*, Fe(II) transporter -*GmIRT1*- and ATPase -*GmAHA2*- genes (Waters et al. 2018), as well as other genes in the roots (Moran Lauter et al. 2014; Santos et al. 2013, 2016). However, the decreases in the rhizospheric pH are less strong than in other plant species (Zocchi et al. 2007). Upon Fe deficiency, *G. max* secretes carboxylates (Zocchi 2006; Zocchi et al. 2007) and still uncharacterized phenolic compounds (Brown and Ambler, 1973; Zocchi et al. 2007), but not flavins (Waters et al. 2018). *G. max* roots also contain flavonoids (d'Arcy-Lameta 1986; Graham 1991), compounds that have been found to be involved in Fe-deficiency responses in other species (Masaoka et al. 1993; Gheshlaghi et al. 2020). There are large differences between *G. max* genotypes regarding Fe uptake efficiency in calcareous soils (Lin et al. 1997; Jolley and Brown 1987; García-Mina et al. 2013; Vasconcelos and Grusak, 2014; Waters et al. 2018; Raj et al. 2021). Within the plant, the concentrations of the natural chelators nicotianamine (NA) and citrate (Cit), which can bind Fe and participate in internal Fe transport, were highest in *G. max* among six species analyzed (Ariga et al. 2014). In *G. max*, the expression on NA synthase (NAS) increases under Fe deficiency (Atencio et al., 2021), and overexpression of barley NAS decreased chlorosis and increased Fe deposition in the seed (Nozoye et al. 2014; Nozoye 2018). In *G. max* xylem, Fe was found to be transported by Cit (Tiffin et al. 1970) and Cit-loading proteins are overexpressed in Fe-efficient cultivars compared to Fe-inefficient ones (Rogers et al. 2009; Qiu et al. 2017). Also, transgenic overexpression of *AtFRO2* resulted in an upregulation of Cit and malate (Mal) and increases in Fe concentrations (Vasconcelos et al. 2014).

The management of Fe deficiency in *G. max* usually involves the use of Fe-efficient cultivars and synthetic Fe(III) chelates, which often lead to an incomplete recovery (Wiersma 2005; Gamble et al. 2014). Application of Fe to the foliage can increase to some extent leaf Chl (Rodríguez-Lucena 2010a,b) but not always leads to yield increases (Goos and Johnson 2000; Merry et al. 2022). Many years ago, pyrite ( $\text{FeS}_2$ ) was found to be useful for Fe deficiency management (Wallace et al. 1976, 1980), and it has been recently proposed that a pyrite-rich mine coal waste can ameliorate the nutrition of crops in calcareous soils, including *G. max* (Stander et al. 2022).

In this study, we tested the hypothesis that the application of Fe mining sub-products and thiols can increase the availability of Fe and alleviate a moderate Fe deficiency in *G. max* grown in a calcareous soil in greenhouse conditions. An Fe(III)-chelate, two different Fe mining sub-products (both of them containing Fe oxides and one  $\text{FeS}_2$ ) and three thiols, glutathione (GSH), dithiothreitol (DTT; *threo*-1,4-dimercapto-2,3-butanediol) and thiophenol (PhSH; benzenethiol) were applied to the soil, alone or in combination. Glutathione is a natural compound occurring in plants, whereas DTT and PhSH are synthetic low molecular mass reductants with sulfhydryl (-SH) groups. Parameters assessed were leaf chlorophyll levels, leaf and root biomass, plant height, leaf mineral composition, leaf and root GSH and oxidized GSH (GSSG), abscisic and gibberellic acids, Asc, Cit and Mal in leaves, and Asc peroxidase and GSH reductase (GR) in leaf extracts. Results indicate that the application of Fe sub-products and thiol compounds could be appropriate management practices to control Fe deficiency chlorosis in *G. max* grown in calcareous soils.

# Material And Methods

## Plant and soil materials and experimental design

Seeds of soybean (*Glycine max* L.; Katoul cultivar -also known as DPX 3589-, accession #1.2.3.11, from Aliabad-e Katul, Golestan Province, Iran) were obtained from the Seed and Plant Certification and Registration Institute, Karaj, Iran. Seeds were sterilized for 3 min in 1% (v/v) NaClO, washed for 20 min in sterile distilled water (removing any swollen seeds with broken seed coats), followed by three rinses in sterile distilled water.

The soil used was collected from the top layer (2–15 cm) in a non-agricultural land at the Ferdowsi University of Mashhad, Iran (Mashhad, Razavi Khorasan Province, Iran; 36° 18' 3" N, 59° 31' 52" E). The soil had a loamy texture (USDA classification system; 50% sand, 33% silt, and 17% clay). The soil characteristics and methods used are as follows: 19.3% CaCO<sub>3</sub> (Loeppert and Suarez 1996); pH 7.69 (in H<sub>2</sub>O); 0.85 dS m<sup>-1</sup> electrical conductivity in saturated soil paste (Corwin and Rhoades 1982); and 12.21 cmol<sub>c</sub> kg<sup>-1</sup> soil cation exchange capacity; soil organic carbon content (Walkley-Black method; Nelson and Sommers 1996) 22 g kg<sup>-1</sup>, and total Kjeldahl N (Bremner and Mulvaney 1982) 2 g kg<sup>-1</sup>. Extractable P and K concentrations were 8.9 and 227 mg g<sup>-1</sup>, respectively, and 2-Bis[2 (bis(carboxymethyl)amino)ethyl]amino} acetic acid (DTPA)-extractable Fe was 3.5 mg Fe kg<sup>-1</sup>. The soil was thoroughly homogenized, air-dried, sieved (2 mm), and supplemented with urea, triple superphosphate and KNO<sub>3</sub>, at rates of 80, 136.36 and 42.16 mg kg<sup>-1</sup> soil of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O, respectively. Each pot (17, 15.5 and 11 cm in height, and upper and lower diameters, respectively) was filled with 2.5 kg of soil. Seeds were sowed in the pots, the soil was moistened with Type I water, and after plant germination (at day 7) pots were irrigated to maximum water holding capacity with Type I water. After one week, seedlings were thinned to four per pot.

The experiment was carried out in a greenhouse at the Ferdowsi University of Mashhad. Day/night parameters were 25/18°C, 31/60% RH, and 16/8 h light/dark. The experiment was performed with a factorial, completely randomized design with three replications, using a combination of four Fe treatments and three thiol treatments. The Fe treatments used were implemented with zero Fe, Fe(III)-EDDHA (the Fe(III) chelate of ethylenediamine-*N,N'*-bis(2-hydroxyphenyl)acetic acid; thereafter called Fe-chelate) and two sub-products of the Fe mining industry, a mix of Fe oxides and a sulfur (S)-rich Fe ore. The Fe(III)-EDDHA was Helio Trace Fe(s), (Heliopotasse, Mulhouse, France; 6% EDDHA-chelated Fe, including 4.8% *o,o*EDDHA-chelated Fe). The two Fe mining sub-products used were an Iranian mine waste containing Fe oxides (Fe<sub>oxi</sub>; 70.0% Fe; obtained from a mine Company in Southern Iran) and an Fe ore (Fe<sub>ore</sub>; 60.5% Fe; obtained from Esfarayen Industrial Complex, North Khorasan, Iran), and were passed through a No#100 sieve (105 μm mesh size). The thiol products used were reduced GSH, DTT and PhSH (Sigma-Aldrich, Taufkirchen, Germany; CAS numbers 70-18-8, 3483-12-3 and 108-98-5, respectively). Solutions were prepared at a final volume of 100 mL with no thiols, 5 mM GSH, 0.5 mM DTT or 5 mM PhSH in water. Then, 69.4 mg of Fe-EDDHA, 44.46 mg Fe<sub>oxi</sub> or 43.4 mg Fe<sub>ore</sub> were added, and the

suspensions were shaken for 20 min in the dark at 15°C. The mixtures were then slowly applied to each pot at sunset time at three different dates, 21, 28 and 35 days after sowing. Final Fe doses after the three applications were 5, 37.3 and 31.5 mg Fe kg<sup>-1</sup> soil for Fe(III)-EDDHA, Fe<sub>oxi</sub> and Fe<sub>ore</sub>, respectively.

## Sampling, growth parameters and mineral analysis

Leaf chlorophyll was monitored 50 days after sowing with a SPAD-502 device (Minolta, Osaka, Japan). SPAD values shown are means of 24 measurements in young, fully developed leaves. Sixty days after sowing, plants were randomly selected in each pot, extracted from the soil, thoroughly washed with diluted soap, tap water and thrice with Type I water, and then divided in young (upper 3 trifoliates) and developed leaves (lower 4–5 trifoliates), stems and roots, and fresh weight (FW) was measured. Leaf midribs were not removed. Plant height and biomass measurements were taken in plants from 3 pots per treatment in the case of leaves and 2 pots per treatment in the case of roots. Plant tissues were immediately stored at -80°C for biochemical analysis or placed in an oven at 70°C for 48 h to determine dry weight (DW). Dried tissues were ground to pass through a 1 mm stainless sieve and stored in plastic vials. For mineral analysis, tissues from three plants in a given treatment (two in the case of roots) in each pot were pooled. Nitric–perchloric acid digestion was carried out with standard procedures (AOAC 2000) as described in Gheshlaghi et al. (2019). The concentrations of Fe in the final solutions were determined by ICP-OES (Spectro Arcos, Spectro Analytical Instruments, Kleve, Germany), calibrating the device daily with certified standards and including standard solutions and blanks in each sample set. Three technical replications were made. Leaf 1,10-*o*-phenanthroline-extractable Fe was determined using young leaves harvested 55 days after sowing (Katyal and Sharma 1984; Abadía et al. 1984). Washed leaves (1 g) were finely chopped using a Ti scissor, incubated for 24 h in 10 mL of 83 mM (1.5%) 1,10-*o*-phenanthroline in water, pH 3.0, and Fe in the filtered extracts was determined using AAS (PGI 990, PG Instruments Ltd., Lutterworth, U.K.). To determine total S concentration, 1 g of tissue was dried at 70°C, ashed at 600°C in a muffle furnace, and the ashes were dissolved in 10 mL of 3 M HCl. Samples were filtered through Whatman No. 42 paper, and S was determined by turbidimetry using MgSO<sub>4</sub> as a standard (Bardsley and Lancaster 1960).

## Determination of leaf and root reduced glutathione and ascorbate peroxidase and glutathione reductase in leaf extracts

Reduced and oxidized GSH was measured spectrophotometrically in total foliage and root tissues with 2% metaphosphoric acid (w/v) as in Luwe et al. (1993). Since legumes contain hGSH and hGSSGh in varying amounts, the total GSH determined (tGSH) was the sum of GSH and hGSH, and the total GSSG determined (tGSSG) was the sum of GSSG and hGSSGh. Ascorbate peroxidase (APX; EC 1.11.1.11) and

GR (EC 1.6.4.2) activities were measured in leaf extracts as in Nakano and Asada (1981) and Carlberg and Mannervik (1985), respectively. All these methods were described in detail in Gheshlaghi et al. (2019).

## Determination of ascorbate, gibberellic acid and abscisic acid in leaves

Ascorbate (Asc) was determined with  $\alpha,\alpha$ -dipyridyl (Masato 1980), as described in Gheshlaghi et al. (2019). Extraction and determination of gibberellic acid (GA<sub>3</sub>; thereafter called GA) was carried out according to Berríos et al. (2004) with some modifications. Fresh tissue from the total foliage (0.5 g) was ground in liquid N<sub>2</sub> and homogenized in 2 mL of pure methanol in a ball mill (MM 400, Retsch GmbH, Haan, Germany). The homogenate was centrifuged at 20,000 g at 4°C for 15 min and the supernatant was collected. The sample was adjusted to pH 1–2 using 0.1 M HCl, transferred to a separatory funnel, and after adding 10 mL of ethyl acetate it was vigorously shaken for 2 min. Then, 10 mL phosphate buffer (pH 7.4) was added, shaking again for 3 min. The organic colored phase was discarded and the aqueous phase containing GA was collected. A 3 mL sample aliquot was mixed with 3 mL of 100% ethanol and 3 mL of 3.75 M HCl and stirred. After incubation at RT for 30 min, A<sub>254</sub> was measured spectrophotometrically (WPA S2000). Results were expressed as  $\mu\text{g g DW}^{-1}$  using a GA calibration curve. Abscisic acid (ABA) was determined as in Kelen et al. (2004). Fresh leaf tissue (1.5 g FW from the total foliage) was ground with 60 mL of extraction solution (0.25 g of butylated hydroxytoluene and 0.44 g of Asc in 95% methanol) and the extract was maintained in the dark at 4°C for 16 h. The extract was filtered (Whatman filter paper #42), the residue washed three times with extraction solution, and the pooled extract was concentrated using a freeze drier (FD-10V, Pishtaz Engineering Co., Tehran, Iran) for 24 h and at -50°C. Dried extracts were dissolved with a mixture of 0.5 M phosphate buffer (pH 8.5) and ethyl acetate (1:1; v:v). The ethyl acetate phase was filtered (Whatman No. 42 filter paper), dried at -50°C and re-dissolved in 5 mL of 95% methanol. The mixture was vortexed and filtered (0.22  $\mu\text{m}$  micro-sieve). A 20  $\mu\text{L}$  extract aliquot was analyzed for ABA using a HPLC device (Waters Alliance e2695 XC with PDA detection) fitted with a C<sub>18</sub> column (250×4.6 mm i.d.). The mobile phase was 0.2% acetonitrile:100% methanol (50:50; v:v), and the flow rate was 0.8 mL min<sup>-1</sup>. An ABA standard (Sigma-Aldrich) was used for quantification.

## X-ray Diffraction Analysis and Scanning Electron Microscopy of Fe<sub>oxi</sub> and Fe<sub>ore</sub>

The Fe mining sub-products used (size  $\leq 105 \mu\text{m}$ ) were subjected to X-ray diffraction analysis, and spectra from samples and standards analyzed with a XRD GNR Explorer device (G.N.R. Analytical Instruments Group, Theta/Theta XRD Explorer, Italy) with Match! software (Crystal Impact, Bonn, Germany) in the range 20–80 ° 2theta, with 6002 data points and alpha2 and background subtraction. The products were coated with Au for 60 s using a Sputter Coater (SCDOOS-Baltec, City, Switzerland).

SEM observations were carried out on a Philips XL30 EM-EDS device with an accelerating voltage of 20 kV in the Central Laboratory of the Ferdowsi University. The products were dissolved in by HCl and HClO<sub>4</sub>:HF 1:5 (Hlavay et al. 2004; Tighe et al. 2004), and total Fe determined by ICP-OES (Spectro Arcos, Spectro Analytical Instruments).

## Solubilization of Fe<sub>oxi</sub> and Fe<sub>ore</sub> by thiols

The solubilization of Fe<sub>oxi</sub> and Fe<sub>ore</sub> in the absence and presence of thiols was monitored as shown in Fig. S1. Fifty mg of each product were stirred at 2000 rpm and at 25°C in a tube with 10 mL of assay medium (15 mM MES, pH 6.0, 300 µM BPDS) for different times (5, 10, 30, 60, 90, 110 and 120 min), and samples were centrifuged at 10000 xg for 3 min. Total Fe in the supernatant (thereafter called total ICP-Fe) was determined after digestion with HCl and HClO<sub>4</sub>:HF 1:5 (Hlavay et al. 2004; Tighe et al. 2004) using by ICP-OES (Spectro Arcos, Spectro Analytical Instruments). The supernatant was supplemented with GSH, DTT or PhSH (to final concentrations of 5, 0.5 and 5 mM, respectively), and after 30 min samples were centrifuged again at 10000 xg for 3 min. The supernatant was used to measure i) absorbance in the range 400–700 nm (the complex Fe(II)-BPDS<sub>3</sub> has an absorption peak at 550 nm) using a spectrophotometer (UV2601, Rayleigh, Beiqing Road, Haidian District, Beijing, China), and ii) Fe by ICP-OES (Spectro Arcos, Spectro Analytical Instruments) after acidification of the sample (3% NO<sub>3</sub>H) (thereafter called ICP-Fe).

## Determination of leaf carboxylates

Citric (Cit) and malic (Mal) acids were determined enzymatically according to Mollering (1985, 1989), using specific kits (K-CITR 06/18 and K-LMAL-58A/ K-LMAL-116A 08/18; Megazyme, Wicklow, Ireland), and measuring A<sub>340</sub> spectrophotometrically. Results were expressed as mg Cit or Mal g<sup>-1</sup> FW.

## Data analysis

Data were analyzed by two-way ANOVA using SPSS 13.0 software. Significant differences among treatments were calculated using Duncan's multiple range test (P < 0.01). Significant differences between means for the thiol treatments in a given Fe treatment and for the Fe treatments in a given thiol treatment are indicated in the Figures using lower case and capital letters, respectively.

## Results

### Composition of the Fe sub-products used

The Fe<sub>oxi</sub> sub-product (70.0% Fe) was found to contain (using EDX) 82.2% FeO (wüstite), 2.3% Fe<sub>2</sub>O<sub>3</sub> (hematite) and 7.9% Fe<sub>3</sub>O<sub>4</sub> (magnetite) (Figure S2). The Fe<sub>ore</sub> sub-product (60.5% Fe) was found to

contain 36.5% FeS<sub>2</sub> (pyrite), 33.2% hematite, 14.1% magnetite, 4.9% wüstite and 4.9% Fe<sub>0.4</sub>Mg<sub>0.6</sub>O (magnesiowüstite) (Figure S3).

## Changes in leaf SPAD with Fe and thiol treatments

A picture of the plants at day 50 is shown for some treatments in Figure S4. The chlorosis level of the plants growing with no Fe in the absence of thiols was 2 (some chlorosis in the canopy, as in Merry et al. 2022). In the absence of thiols (grey bars in Fig. 1) all three Fe-containing products led to increases in the SPAD values of young leaves when compared to the zero Fe treatment (in the range from 1.1- to 1.4-fold). In plants grown with no Fe added, the SPAD index increased when GSH, DTT and PhSH were applied (in the range from 1.1- to 1.3-fold). In the case of plants treated with Fe-EDDHA, the SPAD also increased with all thiol compounds tested (in the range from 1.1- to 1.2-fold). However, in plants treated with Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the SPAD did not increase significantly with the thiol treatments. When GSH was applied, all three Fe sources led to increases in the SPAD values when compared to the zero Fe treatment. However, when DTT was applied, Fe treatments did not lead to increases in SPAD, and when PhSH was applied, the application of Fe<sub>oxi</sub> and Fe<sub>ore</sub>, but not that of Fe(III)-EDDHA, led to increases in SPAD.

## Changes in biomass and plant height with Fe and thiol treatments

Changes observed in the shoot and root FW with Fe and thiol treatments are shown in Figure S5 and described in detail in the Supplementary materials file.

When no thiols were added, the only shoot DW change found was a decrease for Fe<sub>oxi</sub> (7%; grey bars in Fig. 2A). In plants grown with zero Fe, the DW was not changed when any of the thiols were applied. In plants treated with the Fe-chelate, the DW increased when GSH and PhSH were applied (1.2-fold). In plants treated with Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the DW decreased only with DTT and PhSH (by 20 and 16%, and 28 and 11%, respectively). When GSH, DTT and PhSH were applied with the Fe-chelate, increases in shoot DW were found when compared to the zero Fe treatment. Decreases in the shoot DW were observed when any of the thiols were applied with Fe<sub>oxi</sub> and when DTT and PhSH were applied with Fe<sub>ore</sub>.

When no thiols were added, all Fe sources led to increases in root DW (1.5- to 1.7-fold; grey bars in Fig. 2B). In plants grown with zero Fe, the DW increased when GSH and DTT were applied (1.1- and 1.3-fold, respectively). In plants treated with Fe-EDDHA, the DW decreased with GSH and DTT (by 18 and 12% respectively) and increased with PhSH (1.3-fold). In plants treated with Fe<sub>oxi</sub>, the DW increased with DTT and PhSH (1.1-fold,) and decreased slightly with GSH (by 3%). In plants treated with Fe<sub>ore</sub>, the root DW decreased with GSH and DTT (by 11 and 14%, respectively) and increased with PhSH (1.1-fold). For any thiol treatment, addition of all Fe sources led to increases in root DW.

Regarding plant height, when no thiols were added all Fe sources led to increases in this parameter (1.9- to 2.1-fold; grey bars in Fig. 2C). In plants grown with zero Fe, plant height increased only when GSH and PhSH were applied (1.4- and 1.2-fold, respectively). In plants treated with any of the Fe products, plant height decreased when thiols were applied (by 15–21% for the Fe-chelate, 37–41% for Fe<sub>oxi</sub> and 28–42% for Fe<sub>ore</sub>). For any thiol treatment, supplementation with the Fe-chelate led to increases in plant height. Other significant changes in plant height were decreases when GSH was supplemented with Fe<sub>ore</sub> and increases when DTT was supplemented with Fe<sub>oxi</sub> and Fe<sub>ore</sub>.

## Changes in Fe concentrations and contents in leaves, stems and roots with Fe and thiol treatments

In the absence of thiols, using the Fe-chelate and Fe<sub>oxi</sub> led to increases in the Fe concentrations in young leaves (1.3- and 1.2-fold, respectively; grey bars in Fig. 3A) In young leaves of plants grown with zero Fe, Fe concentrations decreased when thiols were applied (by 13–35%). In plants treated with the three Fe sources, Fe concentrations increased with DTT (in the range from 1.1- to 1.7-fold), and decreased in the cases of the Fe-chelate and Fe<sub>ore</sub> with the other two thiols (by 18–38% and 21–26% respectively), and in the case of Fe<sub>oxi</sub> with PhSH (by 34%). With any thiol treatment, application of the three Fe sources led to increases in the Fe concentration when compared to the zero Fe treatment, with the only exception of the treatment with GSH with Fe<sub>ore</sub>, where there was no effect.

When no thiols were applied, all Fe compounds led to increases in the Fe concentration in developed leaves when compared to the zero Fe control (1.4- to 1.9-fold; grey bars in Fig. 3B). In developed leaves of plants grown with zero Fe, Fe concentrations increased when thiols were applied (in the range from 1.1- to 1.5-fold). In the case of plants treated with Fe(III)-EDDHA, the Fe concentration increased with DTT (1.2-fold), and decreased with PhSH (by 11%). In plants treated with Fe<sub>oxi</sub>, the leaf Fe concentration increased when GSH and PhSH were applied (2.0- and 1.5-fold, respectively). In plants treated with Fe<sub>ore</sub>, the Fe concentration increased with all thiols tested, with PhSH having the largest effect (1.1-, 1.2- and 1.7-fold). For any thiol treatment, all Fe compounds led to increases in the leaf Fe concentration when compared to the zero Fe control.

Regarding the Fe concentrations in stems, when no thiols were added all Fe compounds led to increases in this parameter (1.3- to 1.6-fold; grey bars in Fig. 3C) In stems of plants grown with zero Fe, the Fe concentrations decreased slightly when DTT was applied (by 4%) and increased with PhSH (1.1-fold). In plants treated with Fe(III)-EDDHA, the Fe concentration increased with DTT (1.1-fold), and decreased with GSH and PhSH (by 4% and 21%, respectively). In plants treated with Fe<sub>oxi</sub>, the leaf Fe concentration increased with all thiols (1.2-fold), whereas in plants treated with Fe<sub>ore</sub>, the Fe concentration increased with DTT and PhSH (1.3-fold). For any thiol treatment, all Fe compounds led to increases in the leaf Fe concentration when compared to the zero Fe control.

In the absence of thiols, only the Fe-chelate led to increases in the Fe concentration in roots when compared to the zero Fe control (2.4-fold; grey bars in Fig. 3D). In plants grown with zero Fe, the Fe concentration only increased with GSH and PhSH (1.3- and 1.6-fold, respectively). In the case of plants treated with Fe(III)-EDDHA, the Fe concentration decreased with all thiols (by 10–27%). Conversely, in plants treated with Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the Fe concentration increased markedly with all thiols (from 1.5- to 1.8-fold and 1.4- to 1.7-fold, respectively). When any of the thiols were applied, all Fe treatments led to increases in the root Fe concentration when compared to the zero Fe control.

When no thiols were applied, the only increase in leaf extractable Fe was when using Fe<sub>oxi</sub> (1.2-fold; grey bars in Figure S6). In plants grown with zero Fe, the extractable Fe (with 1,10-*o*-phenanthroline) increased only when GSH was applied (1.3-fold). In the case of plants treated with Fe(III)-EDDHA, the extractable Fe increased only with DTT and PhSH (1.4- and 1.1-fold, respectively). In plants treated with Fe<sub>oxi</sub>, the extractable Fe decreased only with PhSH (23%), and in those treated with Fe<sub>ore</sub>, the extractable Fe increased with GSH and PhSH (1.6- and 1.4-fold, respectively), but decreased with DTT (by 13%). When GSH was applied, decreases and increases in leaf extractable Fe were found with the Fe-chelate and Fe<sub>ore</sub>, respectively. When DTT was used, increases in leaf extractable Fe were found with the Fe-chelate and Fe<sub>ox</sub>, and decreases with Fe<sub>ore</sub>. Finally, when PhSH was used, increases in leaf extractable Fe were found with Fe<sub>ore</sub>, and decreases with Fe<sub>oxi</sub>.

## Changes in iron contents with Fe and thiol treatments

When no thiols were added, all Fe compounds led to increases in the Fe contents in young leaves when compared to the zero Fe control (1.1- to 1.3-fold; grey bars in Fig. 4A). In leaves of plants grown with zero Fe, the Fe contents decreased when all thiols were applied (by 13–35%). In plants treated with Fe(III)-EDDHA, the leaf Fe content increased with DTT (1.1-fold), and decreased with GSH and PhSH (by 18 and 38%, respectively). In plants treated with Fe<sub>oxi</sub>, the leaf Fe content increased when DTT was applied (1.1-fold) and decreased with PhSH (by 34%). In plants treated with Fe<sub>ore</sub>, the leaf Fe content decreased with GSH and PhSH (by 21–26%) and increased with DTT (1.7-fold). For any thiol treatment, all Fe treatments led to increases in the young leaf Fe content when compared to the zero Fe control, with the only exception of Fe<sub>ore</sub> applied in combination with GSH.

In the absence of thiols, all Fe compounds led to increases in the Fe contents in developed leaves when compared to the zero Fe control (1.4- to 1.9-fold; grey bars in Fig. 4B). In plants grown with zero Fe, the leaf Fe contents increased when thiols were applied (from 1.1- to 1.5-fold). In the case of plants treated with the Fe-chelate, the leaf Fe content showed increases and decreases with DTT and PhSH (1.1-fold and 11%, respectively). In plants treated with Fe<sub>oxi</sub>, the leaf Fe contents increased with GSH and PhSH (2.0- and 1.5-fold, respectively). In plants treated with the Fe<sub>ore</sub>, the leaf Fe contents increased with all thiols applied (from 1.1- to 1.7-fold). With any thiol treatments, increases in developed leaf Fe contents were found when using all three Fe compounds.

Regarding the Fe contents in stems, when no thiols were added all Fe compounds led to increases in this parameter when compared to the zero Fe control (1.5- to 2.5-fold; grey bars in Fig. 4C). In stems of plants grown with zero Fe, the Fe contents did not change when thiols were applied. In the case of plants treated with Fe(III)-EDDHA and Fe<sub>oxi</sub>, the Fe content decreased with DTT and PhSH (from 8–22%). In plants treated with Fe<sub>ore</sub>, the Fe content only increased with PhSH (1.2-fold). For any thiol treatment, all Fe compounds led to increases in stem Fe contents when compared to the zero Fe control.

When no thiols were added, all Fe compounds led to increases in the Fe contents in roots when compared to the zero Fe control (1.7- to 3.8-fold; grey bars in Fig. 4D). In roots of plants grown with zero Fe, Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the Fe content increased with all thiol compounds (from 1.2- to 2.0-fold). In the case of plants treated with the Fe-chelate, the root Fe content decreased with all thiols (by 7–33%). With any thiol treatments, all Fe treatments led to increases in the root Fe concentrations when compared to the zero Fe control.

## Changes in S concentrations and contents with Fe and thiol treatments

In the absence of thiols, the Fe-chelate led to decreases in the S concentrations in the young leaves, whereas Fe<sub>oxi</sub> and Fe<sub>ore</sub> led to increases in this parameter (16% and from 1.2- to 1.3-fold, respectively; grey bars in Figure S7A). In plants grown with zero Fe, the leaf S concentrations only decreased very slightly with DTT. In plants treated with the Fe-chelate, the S concentrations increased with all thiols (from 1.2- to 1.6-fold). In plants treated with Fe<sub>oxi</sub>, the S concentrations increased 1.4-fold with GSH and 1.1-fold with DTT. In plants treated with Fe<sub>ore</sub>, the S concentrations increased with GSH and DTT (from 1.1- to 1.2-fold), and decreased somewhat with PhSH (by 9%). When GSH and DTT were applied, all Fe compounds led to increases in S concentrations, and when PhSH was applied, only Fe<sub>oxi</sub> and Fe<sub>ore</sub> led to increases in S concentrations.

Regarding the root S concentrations, when no thiols were applied the Fe-chelate and Fe<sub>ore</sub> led to S concentration decreases and increases, respectively (32% and 1.1-fold, respectively; grey bars in Figure S7C). In roots of plants grown with zero Fe, the S concentrations increased when all thiols were applied (ca. 1.2-fold). In plants treated with the Fe-chelate and Fe<sub>oxi</sub>, the root S concentration also increased with all thiols were applied (from 1.4- to 1.8-fold and from 1.1- to 1.4-fold, respectively). In plants treated with Fe<sub>ore</sub>, the root S concentration increased 1.5-fold with GSH and 1.1-fold with DTT, and decreased with PhSH (by 6%). When GSH was applied, all three Fe compounds led to increases in root S concentration. When DTT was applied, only the Fe-chelate led to increases in root S concentration, and when PhSH was applied, all Fe compounds led to decreases in S concentrations.

When no thiols were applied, the Fe-chelate and Fe<sub>ore</sub> led to young leaf S content decreases and increases, respectively (21% and 1.2-fold, respectively; grey bars in Figure S7B). In leaves of plants grown

with zero Fe, the S contents did not change when thiols were applied. In plants treated with the Fe-chelate, the leaf S content increased with all thiols were applied (in the range 1.6- to 2.1-fold). In plants treated with Fe<sub>oxi</sub>, the leaf S contents only increased with GSH (1.4-fold). In plants treated with Fe<sub>ore</sub>, the leaf S contents decreased with DTT and PhSH (by 29 and 19%, respectively). When GSH was applied, all three Fe compounds led to increases in S content. When DTT was applied, only the Fe-chelate led to increases in leaf S content, and when PhSH was applied, the Fe-chelate and Fe<sub>oxi</sub> led to increases and decreases in leaf S content, respectively.

In the absence of thiols, Fe<sub>oxi</sub> and Fe<sub>ore</sub> led to root S content increases (1.5- and 1.9-fold, respectively; grey bars in Figure S7D). In roots of plants grown with zero Fe, the S contents increased when all thiols were applied (from 1.2- to 1.6-fold). In plants treated with the Fe-chelate and Fe<sub>oxi</sub>, the S contents also increased with all thiols (from 1.3- to 1.8-fold), whereas in plants treated with Fe<sub>ore</sub>, the S concentration increased with GSH and PhSH (from 1.1- to 1.3-fold) and decreased with DTT (by 8%). When any thiol was applied, all three Fe compounds led to increases in root S contents.

## Changes in antioxidant compounds with Fe and thiol treatments

When no thiols were applied, all three Fe sources led to increases in the leaf tGSH concentrations when compared to the zero Fe treatment (2.1- to 2.9-fold; grey bars in Fig. 5A). In plants grown with zero Fe, Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the leaf tGSH concentrations increased with all thiols (from 1.7- to 2.3-fold, 1.4- to 1.6-fold and 1.2- to 2.3-fold, respectively). In plants treated with the Fe-chelate, the leaf tGSH decreased with DTT (by 13%). With any thiol treatment, all three Fe products led to increases in the leaf tGSH.

In the absence of thiols, all three Fe sources led to increases in the leaf tGSSG concentrations when compared to the zero Fe treatment (1.7- to 2.4-fold; grey bars in Fig. 5B). In plants grown with zero Fe, Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the leaf tGSSG concentrations increased with all thiols (from 1.1- to 1.7-fold, 1.5- to 1.7-fold and 1.3- to 2.2-fold, respectively). In the case of plants treated with the Fe-chelate, the leaf tGSSG decreased with GSH (by 16%). With any thiol treatment, all three Fe products led to increases in the leaf tGSSG when compared to the zero Fe treatment.

Regarding the leaf Asc concentrations, when no thiols were added all three Fe sources led to decreases in this parameter when compared to the zero Fe control (30–35%; grey bars in Fig. 5E). In plants grown with zero Fe, the Asc concentrations decreased with DTT (by 9%), and much more markedly with GSH and PhSH (by 37–43%) (Fig. 5E). In plants treated with the Fe-chelate, the Asc concentrations had decreases (19%) and increases (1.1-fold) with GSH and DTT, respectively. In plants treated with Fe<sub>oxi</sub>, the Asc concentrations also decreased with GSH and PhSH (by 10 and 28%, respectively), and increased with DTT (1.2-fold). In the case of plants treated with Fe<sub>ore</sub>, the Asc concentrations also had decreases with GSH (14%) and increases with DTT and PhSH (1.4- and 1.2-fold, respectively). When GSH was added, the

Fe-chelate and Fe<sub>ore</sub> led to decreases in the leaf Asc concentrations when compared to the zero Fe control. When DTT was added, the Fe-chelate and Fe<sub>oxi</sub> led to decreases in the Asc concentrations, and when PhSH was added, Fe<sub>oxi</sub> and Fe<sub>ore</sub> led to decreases and marked increases in the leaf Asc concentrations.

When no thiols were added, all Fe products led to marked decreases in the root tGSH concentrations when compared to the zero Fe treatment (46–61%; grey bars in Fig. 5C). In plants grown with zero Fe and the Fe-chelate, the root tGSH concentrations decreased with all thiols (by 51–64% and 35–43%, respectively). In plants treated with Fe<sub>oxi</sub>, the root tGSH increased only with GSH (1.2-fold). In plants treated with Fe<sub>ore</sub>, the root tGSH increased with GSH and PhSH (1.7- and 1.5-fold, respectively). When any of the thiols were applied, the Fe-chelate led to decreases in the root tGSH. Decreases in the root tGSH were also found with GSH and Fe<sub>oxi</sub>, and increases with GSH and PhSH and Fe<sub>ore</sub>.

In the absence of thiols, all three Fe products led to marked decreases in the root tGSSG concentration when compared to the zero Fe treatment (47–61%; grey bars in Fig. 5D). In plants grown with zero Fe and the Fe-chelate, the root tGSSG concentrations decreased with all thiols (by 55–68% and 44–47%, respectively). In plants treated with Fe<sub>oxi</sub>, the root tGSSG decreased with GSH (by 12%). In plants treated with Fe<sub>ore</sub>, the root tGSSG increased with GSH and PhSH (1.2- and 1.3-fold, respectively) and decreased with DTT (by 16%). When any of the thiols were applied, the Fe-chelate led to decreases in the root tGSSG when compared to the zero Fe control. Other changes in root tGSSG included decreases when GSH was applied with Fe<sub>oxi</sub> and increases when DTT was applied with Fe<sub>oxi</sub> and when PhSH was applied with Fe<sub>ore</sub>.

## Changes in antioxidant enzyme activities with Fe and thiol treatments

When no thiols were added, all Fe sources led to large increases in the GR activity in leaf extracts when compared to the zero Fe treatment (2.4-fold; grey bars in Fig. 6A). In plants grown with zero Fe, the GR activity increased with the three thiol compounds (from 1.7-fold to 2.4-fold). In plants treated with Fe-EDDHA, the GR activity was not changed with any of the thiols. In plants treated with Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the GR activity increased with GSH ((1.1-fold) and decreased with DTT (18–31%). When GSH was added, Fe<sub>oxi</sub> and Fe<sub>ore</sub> led to minor increases in the GR activity, and when DTT was added, the Fe-chelate and Fe<sub>oxi</sub> led to increases in this parameter. Finally, when PhSH was added, Fe<sub>oxi</sub> and Fe<sub>ore</sub> led to small increases in the GR activity when compared to the zero Fe treatment.

In the absence of thiols, all Fe products led to increases in the APX activity in leaf extracts when compared to the zero Fe treatment (1.4- to 1.5-fold; grey bars in Fig. 6B). In plants grown with zero Fe, the APX activity increased with GSH and PhSH (1.4- and 1.7-fold, respectively). In plants treated with Fe-EDDHA, the APX activity increased and decreased with GSH and DTT (1.2-fold and 13%, respectively). In

plants treated with Fe<sub>oxi</sub>, the APX activity also decreased with DTT and PhSH (by 39 and 20%, respectively). In plants treated with Fe<sub>ore</sub>, the APX activity increased with GSH (1.2-fold) and decreased with DTT and PhSH (by 51 and 19%, respectively). In the case of GSH, the Fe-chelate and Fe<sub>ore</sub>, but not Fe<sub>oxi</sub>, led to increases in APX activity when compared to the zero Fe control. When DTT was added, the Fe-chelate led to increases in APX activity, whereas Fe<sub>oxi</sub> and Fe<sub>ore</sub> led to decreases in this parameter. When PhSH was added, all three Fe-compounds led to decreases in the leaf APX activity when compared to the zero Fe treatment.

## Changes in carboxylates with Fe and thiol treatments

In plants grown in the absence of thiols, all Fe products led to marked decreases in the leaf Cit concentration when compared to the zero Fe control (42–72%; grey bars in Fig. 7A). In plants grown with zero Fe, the Cit concentration increased markedly with all thiol compounds (from 2.1- to 2.9-fold). In plants treated with the Fe-chelate, the Cit concentration increased with DTT and PhSH (1.2-fold). In plants treated with Fe<sub>oxi</sub> the leaf Cit concentration increased with the three thiols (from 1.4- to 1.6-fold). In plants treated with Fe<sub>ore</sub>, the leaf Cit concentration increased markedly with GSH and PhSH (2.3- and 1.9-fold, respectively). In all the thiol treatments the three Fe sources led to marked decreases in the leaf Cit concentration when compared to the zero Fe treatment.

When no thiols were added, all Fe products led to increases in the leaf Mal concentration (1.1- to 1.3-fold; grey bars in Fig. 7B). In plants grown with zero Fe, the Mal concentration decreased when any of the thiols were added (by 26–48%). In the case of plants treated with the Fe-chelate, the Mal concentration decreased only with DTT (by 9%). In plants treated with Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the Mal concentration also decreased with all thiols (11–28%). In all the thiol treatments the three Fe sources led to increases in the leaf Mal concentration when compared to the zero Fe treatment.

## Changes in phytohormones with Fe and thiol treatments

When no thiols were added, all three Fe sources led to major increases in the leaf GA concentration when compared to the zero Fe treatment (4.3- to 5.9-fold; grey bars in Fig. 8A). In plants grown with zero Fe, the leaf GA concentrations showed marked increases when GSH and PhSH were applied (2.3- and 1.9-fold, respectively). In plants treated with the Fe-chelate, the leaf GA concentration decreased by half when any thiol was applied (by 47–51%). In the case of plants treated with Fe<sub>oxi</sub> and Fe<sub>ore</sub>, the leaf GA concentration also decreased by half with all thiols applied (by 46–52% and 48–60% respectively). When DTT was added, all three Fe products led to major increases in the leaf GA concentration when compared to the zero Fe treatment. When GSH was added, the Fe-chelate and Fe<sub>ore</sub> led to increases and decreases in the leaf GA concentration, respectively, and when PhSH was added the Fe-chelate and Fe<sub>ore</sub> led to increases in the leaf GA concentration.

Regarding the leaf ABA concentration, in the absence of thiols all three Fe sources led to small decreases in this parameter when compared to the zero Fe controls (16 to 23%; grey bars in Fig. 8B). In plants grown with zero Fe, the leaf ABA concentration decreased by half with GSH and PhSH and less with DTT (by 10%). In plants treated with any Fe product the leaf ABA concentration did not change with thiols, with the exception of the Fe<sub>ore</sub> with PhSH, where leaf ABA decreased by 13%. In plants treated with GSH, 2-fold increases in the leaf ABA concentration were found with all Fe compounds. When DTT was added, Fe<sub>oxi</sub> and Fe<sub>ore</sub> led to small decreases in the leaf ABA concentration, and when PhSH was added, all three Fe products led to major increases in the leaf ABA concentration when compared to the zero Fe control.

## Changes in DTPA-extractable soil Fe with Fe and thiol treatments

In the absence of thiols, only the Fe<sub>oxi</sub> led to increases in DTPA-extractable Fe (1.2-fold; grey lines in Fig. 9). When plants were grown with zero Fe, the DTPA-extractable soil Fe only increased with GSH (1.3-fold). In plants treated with the Fe-chelate, the DTPA-extractable Fe increased with DTT and PhSH (1.4- and 1.1-fold, respectively). In plants treated with Fe<sub>oxi</sub>, the DTPA-extractable Fe increased with GSH and DTT (1.1-fold), and decreased with PhSH (by 23%). In plants treated with Fe<sub>ore</sub>, the DTPA-extractable Fe increased with GSH and PhSH (1.6- and 1.6-fold, respectively), and decreased with DTT (by 13%). When GSH was applied, decreases and increases in DTPA-extractable Fe were found with the Fe-chelate and Fe<sub>ore</sub>. When DTT was applied, increases in DTPA-extractable Fe were observed with Fe-chelate and Fe<sub>oxi</sub>, and decreases with Fe<sub>ore</sub>. When PhSH was applied, decreases in DTPA-extractable Fe were observed with Fe<sub>oxi</sub>, and increases with Fe<sub>ore</sub>.

## Solubilization of Fe oxides by thiols

Samples were homogenized in 15 mM MES, pH 6.0, supplemented with 300 µM BPDS, in the absence of thiols. After centrifugation to remove the bulk of the products, the total Fe (total ICP-Fe) in the still slightly opalescent solution was 8.9 and 175.6 mg Fe L<sup>-1</sup> with the Fe<sub>oxi</sub> and Fe<sub>ore</sub>, respectively. When Fe was measured by ICP without a strong acid digestion (ICP-Fe), the Fe concentration in the solution was two orders of magnitude lower with Fe<sub>oxi</sub> than with Fe<sub>ore</sub> (0.1 and 9.3 mg L<sup>-1</sup>, respectively; grey bars in Fig. 10A). In the absence of thiols there was a measurable Fe(II) chelation by BPDS in both products, being larger at 120 min than at 60 min (grey bars in Fig. 10B,C).

The addition of thiols led to large increases in the ICP Fe in solution (measured after acidification), with the increases being higher for DTT in the case of Fe<sub>oxi</sub> and for GSH in the case of Fe<sub>ore</sub> (Fig. 10A). The addition of thiols also led to large increases in the Fe(II) chelated by BPDS, which were larger when the incubation time increased (Fig. 10B,C, and Fig S8). The Fe(II) values were higher for Fe<sub>oxi</sub> than for Fe<sub>ore</sub>

for all thiol compounds and at all incubation times. The highest and lowest solubilized Fe(II) values were for GSH and PhSH, respectively, both when using Fe<sub>oxi</sub> and Fe<sub>ore</sub>.

In the absence of thiols, the ICP-Fe accounted for approximately 1% and 5% of the total ICP-Fe (for Fe<sub>oxi</sub> and Fe<sub>ore</sub>, respectively), whereas in the presence of thiols, the ICP-Fe accounted for ca. 53% and 64% of the total ICP-Fe (for Fe<sub>oxi</sub> and Fe<sub>ore</sub>, respectively). The Fe chelated by BPDS constituted a larger fraction of the ICP-Fe in the case of Fe<sub>oxi</sub> (ca. 1%, 16%, 16% and 9% in the cases of the no thiol treatment, GSH, DTT and PhSH, respectively) than in the case of Fe<sub>ore</sub> (< 1% in all cases).

## Discussion

### Effects of Fe deficiency in *G. max*

When *G. max* plants were grown with zero Fe added, they showed symptoms of a moderate Fe-deficiency (SPAD values were 33% lower than that observed with the best Fe + thiol treatment; Fig. 1). Plants were small, young leaves were light green and root biomass was low (Fig. 2). These moderate symptoms observed in plants grown in a 19% CaCO<sub>3</sub> soil indicate that the *G. max* Katoul cultivar used is an Fe-efficient (Fe-deficiency tolerant) one. Iron concentrations in developed leaves (ca. 82 µg g<sup>-1</sup> DW) were above the low Fe range for *G. max* (51 µg g<sup>-1</sup> DW; Benton Jones et al. 1991), and the Fe concentration in young leaves was even higher (Fig. 3). Leaf chlorosis in the presence of relatively high leaf Fe concentrations, a phenomenon that has been called the “Fe-chlorosis paradox”, have been observed in previous studies with *G. max* (Santos et al. 2015, 2019; Chen et al. 2020). Leaves of Fe-deficient plants did not appear to show a strong oxidative stress, since the tGSH and tGSSG levels and the APX and GR activities were relatively low (Figs. 5–6), and this was likely associated with the moderate leaf chlorosis observed. However, the root concentrations of tGSH and tGSSG were quite high, in line with the large decrease in root biomass (Fig. 5). On the other hand, the leaf Asc (Fig. 5E), Cit (Fig. 7A) and ABA levels (Fig. 8B) were higher, and GA levels were lower (Fig. 8B), than those in the plants supplied with the Fe-chelate.

When Fe deficiency is severe, oxidative stress in leaves and roots is a common occurrence in many plant species (López-Millán et al. 2013), including *G. max* (Mira et al. 2021; Rahman et al. 2022) and *Medicago* spp. (M’sehli et al. 2009; Gheshlaghi et al. 2019). However, when the Fe deficiency is moderate the oxidative stress is less intense, and for instance in *G. max* lower APX levels occur in Fe-efficient cultivars when compared to Fe-inefficient ones (Santos et al. 2019; Mira et al. 2021). Also, in *M. sativa* APX was low in Fe-deficient plants when compared to the Fe-sufficient controls (Rahman et al. 2021). Citrate concentrations are usually increased throughout the plant with Fe deficiency (Abadía et al. 2002), and for instance in Fe-deficient *M. scutellata* the leaf Cit (and Asc) levels were higher when compared to those in Fe-sufficient controls (Gheshlaghi et al. 2019). However, in a metabolomics study, decreases in Cit and Mal have been observed in strongly Fe-deficient *G. max* leaves (Lima et al. 2014). On the other hand, the low and high leaf GA and ABA levels, respectively, confirm previous results in *G. max* (Chen et al. 2020)

and are likely associated with the decreases in plant size and biomass. In *Oryza sativa*, Fe deficiency has also been shown to decrease biologically active GA in leaves (Wang et al. 2017).

## Effects of applying Fe-EDDHA alone

The Fe-chelate at a  $5 \text{ mg Fe kg}^{-1}$  soil dose was partially efficient in alleviating Fe-deficiency symptoms in *G. max*, leading to modest increases in SPAD and shoot biomass. A heatmap summarizing the changes is included in Fig. 11. However, increases were more marked in root biomass (Fig. 2B), and especially in plant height (Fig. 2C), Fe concentrations (Fig. 3) and Fe contents (Fig. 4) in leaves, stems and roots, with little change in leaf extractable Fe (Fig. S6). These positive but limited effects are in line with previous results with Fe(III)-EDDHA in *G. max* (Wiersma 2005; García-Marco et al. 2006; Gamble et al. 2014). The oxidative stress situation changed markedly with the Fe-chelate treatment when compared to that in the Fe-deficient plants. In the leaves, the application of Fe-chelate led to marked increases in tGSH, tGSSG (Fig. 5A,B) and GR (Fig. 6), as well as decreases in Asc (Fig. 5), suggesting the elicitation of a moderate oxidative stress, likely associated with by the increase in Fe being transported. Conversely, in the roots there were marked decreases in tGSH and tGSSG (Fig. 5C,D), indicating that the application of Fe-chelate relieved the oxidative stress in this plant compartment. A large increase in leaf GA, with decreases in leaf ABA, was also found (Fig. 8), in agreement with a previous study (Chen et al 2020), and in line with the observed increases in plant height and shoot and root biomass (Fig. 2). Leaf Cit concentrations decreased with no changes in Mal concentrations (Fig. 7), suggesting that Cit could be being used for Fe transport (Rellán-Álvarez et al. 2011), in line with previous studies in *M. scutellata* (Gheshlaghi et al. 2019). The reason behind the decreases in leaf S concentrations and contents and root S concentrations with Fe-chelate (Fig. S7) is likely to be related to the shift in use of this element, which is needed for the Fe-deficiency responses (Astolfi et al., 2021).

## Effects of applying Fe<sub>oxi</sub> and Fe<sub>ore</sub> alone

When stirred in buffer, the two Fe sub-products maintain a significant part of the Fe in solution/suspension (< 1 and 6% of the total Fe added in the cases of Fe<sub>oxi</sub> and Fe<sub>ore</sub>, respectively) (Fig. 10). Part of this Fe is likely in small particles that cannot be removed by centrifugation. A small fraction of this Fe in solution/suspension can be measured using ICP without acidification (1% and 6% of the total ICP-Fe when using Fe<sub>ore</sub> and Fe<sub>oxi</sub>, respectively), and an even smaller part is converted, in a time-dependent manner, into chemical forms that are reduced in the presence of BPDS (Fig. 10). These data indicate that Fe<sub>oxi</sub> and Fe<sub>ore</sub> are capable to deliver Fe in chemical forms that are potentially available for plants.

Indeed, when the two Fe sub-products (Fe<sub>oxi</sub> and Fe<sub>ore</sub>; applied at  $31.5$  and  $37.3 \text{ mg Fe kg}^{-1}$  soil doses, respectively) were used alone, they were also efficient in alleviating Fe-deficiency symptoms in *G. max*, as judged by the increases in leaf SPAD (Fig. 1), root biomass and plant height (Fig. 2), with the leaf

regreening being similar with both products and significantly better than that found with the Fe-chelate (see heatmap in Fig. 11). These results indicate that the *G. max* cultivar used was an efficient one, capable of taking up Fe from Fe<sub>oxi</sub> and Fe<sub>ore</sub> using the natural Fe-acquisition toolbox in this plant species, which includes the secretion to the rhizosphere of protons, carboxylates and phenolics (Zocchi et al. 2007). Other changes in Fe chlorosis parameters were in the same direction, but less intense, than those caused by the Fe-chelate, including increases in leaf Fe concentrations and contents, root Fe contents (Figs. 3–4), and leaf tGSH and tGSSG (Fig. 5A,B). This suggests that Fe<sub>oxi</sub> and Fe<sub>ore</sub> also elicit a slight oxidative stress in leaves, but less intense than that observed with the Fe-chelate, likely because of a more controlled Fe supply and trafficking within the plant. In some parameters, such as the shoot biomass and root Fe concentrations, there were little changes when compared to the zero Fe control. Conversely, the decreases in root tGSH and tGSSG (Fig. 5B,C), leaf ABA (Fig. 8) and leaf Cit (Fig. 7) were more marked than those observed with the Fe-chelate. Changes in the rest of parameters, including the large increase in leaf GA (Fig. 8), were quite similar to those found with the Fe-chelate.

The positive results of Fe mining sub-products in alleviating Fe-deficiency are in line with previous studies in *G. max*, and indicate that this species can take up Fe from Fe chemical forms present in these materials using their natural acquisition mechanisms. The pioneering studies of Wallace et al. (1976, 1980) indicated that the application of FeS<sub>2</sub> could be useful for controlling Fe deficiency. Recently, the application of Fe-humic nanofertilizers containing the Fe oxi-hydroxide ferrihydrite have been shown to alleviate Fe deficiency in *G. max* (Cieschi et al. 2019). In *O. sativa*, the application of nanoparticles containing zero-valent Fe have been shown to cause minor increases in leaf chlorophyll, whereas nanoparticles containing Fe<sub>2</sub>O<sub>4</sub> and Fe<sub>2</sub>O<sub>4</sub> were not effective, in spite of the increases in plant Fe concentrations observed with the three nanoparticle types; these applications reduced the concentrations of GA and in some cases that of indole-3-acetic acid (Li et al. 2021). The possibility that FeS<sub>2</sub>-rich mine coal wastes can ameliorate the nutrition of *G. max* and other crops in calcareous soils is currently being explored (Stander et al. 2022).

## Effects of applying thiols alone

In the absence of Fe fertilization, the three thiols used (GSH, DTT and PhSH), alleviated Fe-deficiency symptoms in soybean, as judged by the leaf SPAD values, although growth parameters were not much changed, with the exception of increases in plant height with GSH and PhSH and root DW with DTT (see heatmap in Fig. 11). The Fe concentrations and contents in developed leaves and roots tended to increase and the Fe concentrations and contents in young leaves decreased (Figs. 3–4), suggesting that the increase in leaf chlorophyll was associated to an Fe remobilization from pre-existing Fe pools as well as to a new Fe uptake. The decrease caused by thiols in the Fe concentration and contents in young leaves is remarkable, and suggest remobilization of Fe forms in this plant part. The application of thiols led to increases in leaf tGSH, tGSSG (Fig. 5A,B) and antioxidant enzyme activities (Fig. 6), and decreases in root tGSH and tGSSG (Fig. 5C,D) and leaf Asc (Fig. 5E), again in line with an increase in Fe transport within the plant. Leaf hormone changes were not homogeneous, with GSH and PhSH causing large

increases in GA and decreases in ABA, whereas DTT caused no major changes in leaf hormone concentrations (Fig. 8). All thiols led to large increases and decreases in Cit and Mal concentrations in leaves, respectively, whereas the opposite was found with the Fe products (Fig. 7). The only thiol causing an increase in the soil DTPA-extractable Fe was GSH (Fig. 9). The increases in root S concentrations and contents with the addition of thiols were expected.

The positive results of thiols in alleviating Fe-deficiency are in line with those observed in previous studies. In Fe-deficient Arabidopsis grown in nutrient solution the application of GSH increased the expression levels of Fe uptake- and Fe transport-related genes and the Fe concentrations (Koen et al. 2012), and increased leaf chlorophyll levels, but not Fe concentrations, and preserved cell redox homeostasis (Ramírez et al. 2013). Later, it was shown that supplying GSH to the foliage of soil-grown *M. scutellata* relieved Fe-deficiency symptoms, and this was ascribed to the solubilization of pre-existing Fe pools within the plant via a reductive mechanism (Gheshlaghi et al. 2019). Also, when GSH was applied to the roots of *M. scutellata* growing in a Fe-rich rock sand, Fe was mobilized from the substrate via a reductive solubilization mechanism, increasing plant Fe concentrations and relieving Fe-deficiency symptoms (Gheshlaghi et al. 2020). Additional studies have focused on the use of NaHS, a donor of H<sub>2</sub>S, which is rapidly incorporated into plant thiols such as GSH (Chen et al. 2011). The application of NaHS has been shown to promote chlorophyll synthesis in Fe-deficient *G. max* (Chen et al. 2020), *Zea mays* (Chen et al. 2015) and *Fragaria x ananassa* (Kaya and Ashraf, 2019).

## Effects of applying Fe(III)-EDDHA together with thiols

The application of the Fe(III)-chelate supplemented with GSH and PhSH tended to decrease somewhat the Fe concentrations and contents in roots, stems and young leaves when compared to the Fe-chelate alone (see heatmap in Figure S9). Furthermore, the application of Fe-chelate supplemented with thiols increased leaf SPAD (Fig. 1), although plant height was decreased, and shoot biomass increased with GSH and PhSH when compared to the Fe-chelate alone (Fig. 2). The addition of thiols caused increases in the root tGSH and GSSG (Fig. 5C,D) and normalized the leaf and root S concentrations and contents, which were low when the Fe-chelate was applied alone (Fig. S7). When compared with the use of the Fe-chelate alone, thiols moderated the increases in leaf GA (Fig. 8) and the decreases in Cit (Fig. 7).

## Effects of applying Fe<sub>oxi</sub> and Fe<sub>ore</sub> together with thiols

When Fe<sub>oxi</sub> and Fe<sub>ore</sub> are stirred in water, a small part of the Fe occurring in solution/suspension was converted in the presence of thiols and BPDS, in a time-dependent manner, into chemical forms that can be available for plants (Fig. 10B,C). The amount of Fe forms that are in solution/suspension in both Fe-containing products was much larger in the case of Fe<sub>ore</sub> (Fig. 10A), although the Fe that can be trapped by the Fe(II) chelator *o*-phenatroline was larger with Fe<sub>oxi</sub> than with Fe<sub>ore</sub> (Fig. 10B,C), indicating that in the latter case the Fe put in solution was in non-reducible forms.

Indeed, the application of Fe<sub>oxi</sub> and Fe<sub>ore</sub> supplemented with thiols increased the Fe concentrations and contents in roots, stems and developed leaves (Figs. 3–4). However, the application of Fe<sub>oxi</sub> and Fe<sub>ore</sub> supplemented with thiols did not provide a further significant greening (Fig. 1), possibly because the effects of Fe<sub>oxi</sub> and Fe<sub>ore</sub> were already quite high when used alone. In all cases, when used in combination with Fe-containing products the thiols caused marked decreases in plant height (Fig. 2E), and the application of GSH and PhSH decreased the Fe concentrations and contents in young leaves, as it occurs when the thiols were applied alone (Fig. 3–4). On the other hand, the application of Fe<sub>oxi</sub> and Fe<sub>ore</sub> in combination with thiols decreased markedly the leaf GA (Fig. 8) and increased the leaf tGSH and tGSSG (Fig. 5A,B).

## Concluding Remarks

Results confirm the hypothesis that the application of Fe-mining sub-products, thiols and the combination of Fe(III)-EDDHA and thiols could be used to control a moderate Fe deficiency in *G. max* grown in a calcareous soil, with the different treatments relieving Fe deficiency symptoms to different extents. At the doses used, the two Fe-mining sub-products are more effective than the Fe-chelate, and this is likely due to their ability to release Fe forms that could be solubilized and the metal taken up by the Fe-acquisition toolbox existing in *G. max*.

Thiols are also effective in alleviating Fe deficiency symptoms in *G. max*, confirming for the first time that they are capable to facilitate Fe mobilization and uptake from the Fe oxy-hydroxide forms naturally occurring in a calcareous soil. This has been previously shown to occur in plants grown in nutrient solutions and in a rock sand substrate. Therefore, although the microbiome present in the calcareous soil is likely to consume thiols, at the doses used plants were still able to take advantage of the capacity of these compounds to mobilize Fe.

When applied in combination, the thiols only improved greening in the case of Fe(III)-EDDHA, and this was associated to the mobilization in old leaves and stems of Fe forms elicited by the application of the Fe-chelate. In the case of the Fe mining sub-products the leaf Fe concentrations and contents also increased with the application of thiols, associated with a solubilization of Fe forms from the substrate, but leaves did not regreen further, probably to the already marked effects of the Fe sub-products alone.

The doses used in this study are equivalent to approximately 175, 112 and 109 kg ha<sup>-1</sup> of Fe-chelate, Fe<sub>oxi</sub> and Fe<sub>ore</sub>, respectively, and 387, 19 and 139 kg ha<sup>-1</sup> of GSH, DTT and PhSH, respectively. The results obtained suggest that the use of reasonable doses of Fe mining sub-products and GSH, which are cheap and easily available worldwide, can be an alternative for the alleviation of a moderate Fe deficiency chlorosis in *G. max*. New experiments should be envisaged to test the applicability of these management techniques in field conditions.

## Declarations

# Acknowledgements

Authors acknowledge the support of the Department of Soil Science, Faculty of Agriculture, Ferdowsi University of Mashhad, Iran. This work was part of the research project on *G. max* 2/51555. Support was obtained by the Spanish Ministry of Science and Innovation (Grant PID2020-115856RB-100 funded by MCIN/AEI/10.13039/501100011033) and the Aragón Government (group A09-20R). Authors thank J. Rodríguez-Celma for reading the manuscript and provide useful comments.

## References

1. Abadía J, Monge E, Montañés L, Heras L (1984) Extraction of iron from plant leaves by Fe (II) chelators. *J Plant Nutr* 7:777–784. <https://doi.org/10.1080/01904168409363241>
2. Abadía J, López-Millán A-F, Rombolà A, Abadía A (2002) Organic acids and Fe deficiency: A review. *Plant Soil* 241:75–86. <https://doi.org/10.1023/A:1016093317898>
3. Abadía J, Vázquez S, Rellán-Álvarez R, El-Jendoubi H, Abadía A, Álvarez-Fernández A, López-Millán A-F (2011) Towards a knowledge-based correction of iron chlorosis. *Plant Physiol Biochem* 49:471–482. <https://doi.org/10.1016/j.plaphy.2011.01.026>
4. Álvarez-Fernández A, García-Laviña P, Fidalgo C, Abadía J, Abadía A (2004) Foliar fertilization to control iron chlorosis in pear (*Pyrus communis* L.) trees. *Plant Soil* 263:5–15. <https://doi.org/10.1023/B:PLSO.0000047717.97167.d4>
5. Ambikadevi VR, Lalithambika M (2000) Effect of organic acids on ferric iron removal from iron-stained kaolinite. *Appl Clay Sci* 16:133–145. [https://doi.org/10.1016/S0169-1317\(99\)00038-1](https://doi.org/10.1016/S0169-1317(99)00038-1)
6. AOAC (2000) Official Methods of Analysis Association of Analytical Chemists, p.334. Washington DC
7. Ariga T, Hazama K, Yanagisawa S, Yoneyama T (2014) Chemical forms of iron in xylem sap from graminaceous and non-graminaceous plants. *Soil Sci Plant Nutr* 60:460–469. <https://doi.org/10.1080/00380768.2014.922406>
8. Astolfi S, Celletti S, Viganì G, Mimmo T, Cesco S (2021) Interaction between sulfur and iron in plants. *Front Plant Sci* 12:670308. <https://doi.org/10.3389/fpls.2021.670308>
9. Atencio L, Salazar J, Moran Lauter AN, Gonzales MD, O'Rourke JA, Graham MA (2021) Characterizing short and long term iron stress responses in iron deficiency tolerant and susceptible soybean (*Glycine max* L. Merr.). *Plant Stress* 2:100012. <https://doi.org/10.1016/j.stress.2021.100012>
10. Bardsley CE, Lancaster JD (1960) Determination of reserve sulfur and soluble sulfates in soils. *Soil Sci Soc Am J* 24:265–268. <https://doi.org/10.2136/sssaj1960.03615995002400040015x>
11. Benton-Jones J, Wolf B, Mills HA (1991) Plant analysis handbook: A practical sampling, preparation, analysis, and interpretation guide. Micro-Macro-Publishing, Athens, GA
12. Berríos J, Illanes A, Aroca G (2004) Spectrophotometric method for determining gibberellic acid in fermentation broths. *Biotechnol Lett* 26:67–70. <https://doi.org/10.1023/B:BILE.0000009463.98203.8b>

13. Blesa MA, Marinovich HA, Baumgartner EC, Maroto AJG (1987) mechanism of dissolution of magnetite by oxalic acid-ferrous ion solutions. *Inorg Chem* 26:3713–3717.  
<https://doi.org/10.1021/ic00269a019>
14. Bremner JM, Mulvaney CS(1982) Nitrogen-total. In: Page, A.L., Miller, R.H., Keeney, D.R. (Eds.), *Methods of soil analysis, Part 2, second ed., Agron. Monogr. 9*, pp. 595–624 ASA-CSSA-SSSA, Madison, Wisconsin
15. Briat JF, Dubos C, Gaymard F (2015) Iron nutrition, biomass production, and plant product quality. *Trends Plant Sci* 20:33–40. <https://doi.org/10.1016/j.tplants.2014.07.005>
16. Carlberg I, Mannervik B (1985) Glutathione reductase. *Meth Enzymol* 113:484–490
17. Brown JC, Ambler JE (1973) “Reductants” released by roots of Fe-deficient soybeans. *Agron J* 65:311–314. <https://doi.org/10.2134/agronj1973.00021962006500020037x>
18. Chen J, Wu FH, Shang YT, Wang WH, Hu WJ, Simon M, Liu X, Shangguan ZP, Zheng HL (2015) Hydrogen sulphide improves adaptation of *Zea mays* seedlings to iron deficiency. *J Exp Bot* 66:6605–6622. <https://doi.org/10.1093/jxb/erv368>
19. Chen J, Zhang NN, Pan Q, Lin X-Y, Shangguan Z, Zhang J-H, Wei G-H (2020) Hydrogen sulphide alleviates iron deficiency by promoting iron availability and plant hormone levels in *Glycine max* seedlings. *BMC Plant Biol* 20:383. <https://doi.org/10.1186/s12870-020-02601-2>
20. Chen S, Jia H, Wang X, Shi C, Wang X, Ma P, Wang J, Ren M, Li J (2020) Hydrogen sulfide positively regulates abscisic acid signaling through persulfidation of SnRK2.6 in guard cells. *Mol Plant* 13:732–744. <https://doi.org/10.1016/j.molp.2020.01.004>
21. Chen J, Wu FH, Wang WH, Zheng CJ, Lin GH, Dong XJ, He JX, Pei ZM, Zheng HL (2011) Hydrogen sulphide enhances photosynthesis through promoting chloroplast biogenesis, photosynthetic enzyme expression, and thiol redox modification in *Spinacia oleracea* seedlings. *J Exp Bot* 62:4481–4493. <https://doi.org/10.1093/jxb/err145>
22. Cieschi MT, Polyakov AY, Lebedev VA, Volkov DS, Pankratov DA, Veligzhanin AA, Perminova IV, Lucena JJ (2019) Eco-friendly iron-humic nanofertilizers synthesis for the prevention of iron chlorosis in soybean (*Glycine max*) grown in calcareous soil. *Front Plant Sci* 10:413.  
<https://doi.org/10.3389/fpls.2019.00413>
23. Connorton JM, Balk J, Rodríguez-Celma J (2017) Iron homeostasis in plants-a brief overview. *Metallomics* 9:813–823. <https://doi.org/10.1039/c7mt00136c>
24. Cornell RM, Schindler PW (1987) Photochemical dissolution of goethite in acid/oxalate solution. *Clays Clay Miner* 35:347–352. <https://doi.org/10.1346/CCMN.1987.0350504>
25. Cornell RM, Schwertmann U(2004) *The iron oxides: structure, properties, reactions, occurrences and uses*, 2nd edition. Wiley-VCH, Verlag GmbH & Co. KGaA, Weinheim, Germany. ISBN: 3-527-30274-3.  
<https://www.wiley.com/en-us/9783527606443>
26. Corwin DL, Rhoades JD (1982) An improved technique for determining soil electrical conductivity-depth relations from above-ground electromagnetic measurements. *Soil Sci Soc Am J* 46:517–520.  
<https://doi.org/10.2136/sssaj1982.03615995004600030014x>

27. d'Arcy-Lameta A (1986) Study of soybean and lentil root exudates. *Plant Soil* 92:113–123. <https://doi.org/10.1007/BF02372272>
28. El-Jendoubi H, Melgar JC, Álvarez-Fernández A, Sanz M, Abadía A, Abadía J (2011) Setting good practices to assess the efficiency of iron fertilizers. *Plant Physiol Biochem* 49:483–488. <https://doi.org/10.1016/j.plaphy.2011.02.013>
29. Gamble AV, Howe JA, Delaney D, van Santen E, Yates R (2014) Iron chelates alleviate iron chlorosis in soybean on high pH soils. *Agron J* 106:1251–1257. <https://doi.org/10.2134/agronj13.0474>
30. García-Marco S, Martínez N, Yunta F, Hernández-Apaolaza L, Lucena JJ(2006) Effectiveness of ethylenediamine-N(*o*-hydroxyphenylacetic)-N'( *p*-hydroxyphenylacetic) acid (*o,p*-EDDHA) to supply iron to plants. *Plant Soil* 279:31–40. <https://doi.org/10.1007/s11104-005-8218-5>
31. García-Mina JM, Bacaicoa E, Fuentes M, Casanova E (2013) Fine regulation of leaf iron use efficiency and iron root uptake under limited iron bioavailability. *Plant Sci* 198:39–45. <https://doi.org/10.1016/j.plantsci.2012.10.001>
32. Gheshlaghi Z, Khorassani R, Abadía J, Kafi M, Fotovat A (2019) Glutathione foliar fertilisation prevents lime-induced iron chlorosis in soil grown *Medicago scutellata*. *J Plant Nutr Soil Sci* 182:607–624. <https://doi.org/10.1002/jpln.201800692>
33. Gheshlaghi Z, Khorassani R, Abadía J, Álvarez-Fernández A, Luis-Villarroya A, Fotovat A, Kafi M (2020) Glutathione supplementation prevents iron deficiency in *Medicago scutellata* grown in rock sand under different levels of bicarbonate. *Plant Soil* 446:43–63. <https://doi.org/10.1007/s11104-019-04314-4>
34. Gheshlaghi Z, Luis-Villarroya A, Álvarez-Fernández A, Khorassani R, Abadía J (2021) Iron deficient *Medicago scutellata* grown in nutrient solution at high pH accumulates and secretes large amounts of flavins. *Plant Sci* 303:110664. <https://doi.org/10.1016/j.plantsci.2020.110664>
35. Goos RJ, Johnson BE (2000) A comparison of three methods for reducing iron-deficiency chlorosis in soybean. *Agron J* 92:1135–1139. <https://doi.org/10.2134/agronj2000.9261135x>
36. Graham TL (1991) Flavonoid and isoflavonoid distribution in developing soybean seedling tissues and in seed and root exudates. *Plant Physiol* 95:594–603. <https://doi.org/10.1104/pp.95.2.594>
37. Hansen NC, Jolley VD, Naeve SL, Goos RJ (2004) Iron deficiency of soybean in the North Central U.S. and associated soil properties. *Soil Sci Plant Nutr* 50:983–987. <https://doi.org/10.1080/00380768.2004.10408564>
38. Hlavay J, Prohaska T, Weisz M, Wenzel WW, Stinger GJ(2004) Determination of trace elements bound to soil and sediment fractions (IUPAC Technical Report). *Pure Appl Chem* 76:415–442. <https://doi.org/10.1351/pac200476020415>
39. Katyal JC, Sharma BD (1984) Some modification in the assay of Fe<sup>2+</sup> in 1–10, *o*-phenanthroline extracts of fresh plant tissues. *Plant Soil* 79:449–450. <https://doi.org/10.1007/BF02184337>
40. Jolley VD, Brown JC (1987) Soybean response to iron-deficiency stress as related to iron supply in the growth medium. *J Plant Nutr* 10:637–651. <https://doi.org/10.1080/01904168709363597>

41. Jolley VD, Stevens WB, Terry RE, Orf JH (1992) Root iron-reduction capacity for genotypic evaluation of iron efficiency in soybean. *J Plant Nutr* 15:1679–1690.  
<https://doi.org/10.1080/01904169209364430>
42. Kaya C, Ashraf M (2019) The mechanism of hydrogen sulfide mitigation of iron deficiency-induced chlorosis in strawberry (*Fragaria × ananassa*) plants. *Protoplasma* 256:371–382.  
<https://doi.org/10.1007/s00709-018-1298-x>
43. Kaya C, Higgs D, Ashraf M, Alyemeni MN, Ahmad P (2020) Integrative roles of nitric oxide and hydrogen sulfide in melatonin-induced tolerance of pepper (*Capsicum annuum* L.) plants to iron deficiency and salt stress alone or in combination. *Physiol Plant* 168:256–277.  
<https://doi.org/10.1111/ppl.12976>
44. Kelen M, Çubuk Demiralay E, Şen S, Özkan G (2004) Separation of abscisic acid, indole-3-acetic acid, gibberellic acid in 99 R (*Vitis berlandieri* × *Vitis rupestris*) and rose oil (*Rosa damascena* Mill.) by reversed phase liquid chromatography. *Turkish J Chem* 28:603–610
45. Khan MN, Siddiqui MH, Al-Solami MA, Basahi RA, Siddiqui ZH, Alamri S (2021) Cysteine and hydrogen sulfide: A complementary association for plant acclimation to abiotic stress. In: Khan MN, Siddiqui MH, Alamri S, Corpas FJ (eds) *Hydrogen Sulfide and Plant Acclimation to Abiotic Stresses, Plant in Challenging Environments*, vol 1. Springer, Cham, pp 187–214. [https://doi.org/10.1007/978-3-030-73678-1\\_11](https://doi.org/10.1007/978-3-030-73678-1_11)
46. Koen E, Szymańska K, Klinguer A, Dobrowolska G, Besson-Bard A, Wendehenne D (2012) Nitric oxide and glutathione impact the expression of iron uptake- and iron transport-related genes as well as the content of metals in *A. thaliana* plants grown under iron deficiency. *Plant Signal Behav* 7:1246–1250. <https://doi.org/10.4161/psb.21548>
47. Li M, Zhang P, Adeel M, Guo Z, Chetwynd AJ, Ma C, Bai T, Hao Y, Rui Y (2021) Physiological impacts of zero valent iron, Fe<sub>3</sub>O<sub>4</sub> and Fe<sub>2</sub>O<sub>3</sub> nanoparticles in rice plants and their potential as Fe fertilizers. *Environ Pollut* 269:116134. <https://doi.org/10.1016/j.envpol.2020.116134>
48. Lima MRM, Diaz SO, Lamego I, Grusak MA, Vasconcelos MW, Gil AM (2014) Nuclear magnetic resonance metabolomics of iron deficiency in soybean leaves. *J Proteome Res* 13:3075–3087.  
<https://doi.org/10.1021/pr500279f>
49. Lin S, Cianzio S, Shoemaker R (1997) Mapping genetic loci for iron deficiency chlorosis in soybean. *Mol Breed* 3:219–229. <https://doi.org/10.1023/A:1009637320805>
50. Loeppert RH, Suarez DL (1996) Carbonate and gypsum. *Methods of soil analysis Part III, Chemical methods*. Soil Science Society of America, Madison, USA, pp 437–474
51. López-Millán A-F, Grusak MA, Abadía A, Abadía J (2013) Iron deficiency in plants: an insight from proteomic approaches. *Front Plant Sci* 4:254. <https://doi.org/10.3389/fpls.2013.00254>
52. Luwe M, Takahama U, Heber U (1993) Role of ascorbate in detoxifying ozone in the apoplast of spinach (*Spinacia oleracea* L.) leaves. *Plant Physiol* 101:969–976.  
<https://doi.org/10.1104/pp.101.3.969>

53. Dell'Orto MW, Donnini M, De Nisi S, Zocchi G, Abdelly C, Gharsalli M (2009) Variability of metabolic responses and antioxidant defence in two lines of *Medicago ciliaris* to Fe deficiency. *Plant Soil* 320:219–230. <https://doi.org/10.1007/s11104-008-9887-7>
54. Masaoka Y, Kojima M, Sugihara S, Yoshihara T, Koshino M, Ichihara A (1993) Dissolution of ferric phosphate by alfalfa (*Medicago sativa* L.) root exudates. *Plant Soil* 155–156:75–78. <https://doi.org/10.1007/BF00024987>
55. Masato O (1980) An improved method for determination of l-ascorbic acid and l-dehydroascorbic acid in blood plasma. *Clin Chim Acta* 103:259–268. [https://doi.org/10.1016/0009-8981\(80\)90144-8](https://doi.org/10.1016/0009-8981(80)90144-8)
56. Matamoros MA, Moran JF, Iturbe-Ormaetxe I, Rubio MC, Becana M (1999) Glutathione and homoglutathione synthesis in legume root nodules. *Plant Physiol* 121:879–888. <https://doi.org/10.1104/pp.121.3.879>
57. Merry R, Dobbels AA, Sadok W, Naeva S, Stupar RM, Lorenz AJ (2022) Iron deficiency in soybean. *Crop Sci* 62:36–52. <https://doi.org/10.1002/csc2.20661>
58. Mira MM, Asmundson B, Renault S, Hill RD, Stasolla C (2021) Suppression of the soybean (*Glycine max*) Phytoglobin GmPgb1 improves tolerance to iron stress. *Acta Physiol Plant* 43:147. <https://doi.org/10.1007/s11738-021-03315-0>
59. Mollering H (1985) L-malate. In: Bergmeyer HU (ed) *Methods of Enzymatic Analysis*, vol VII, 3rd edn. VCH Publishers (UK) Ltd., Cambridge, UK, pp 39–47
60. Mollering H (1989) Citrate. In: Bergmeyer HU (ed) *Methods of Enzymatic Analysis*, vol VII, 3rd edn. VCH Publishers (UK) Ltd., Cambridge, UK, pp 2–12
61. Moran Lauter AN, Peiffer GA, Yin T, Whitham SA, Cook D, Shoemaker RC, Graham MA (2014) Identification of candidate genes involved in early iron deficiency chlorosis signaling in soybean (*Glycine max*) roots and leaves. *BMC Genomics* 15:702. <https://doi.org/10.1186/1471-2164-15-702>
62. Nakano Y, Asada K (1981) Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol* 22:867–880. <https://doi.org/10.1093/oxfordjournals.pcp.a076232>
63. Nelson DW, Sommers LE (1996) Total carbon, organic carbon, and organic matter. In: Sparks DL (ed) *Methods of Soil Analysis, Chemical Methods*, Vol. 5, part 3. Soil Science Society of America Inc., Madison, WI, USA, pp 961–1010
64. Nozoye T (2018) The nicotianamine synthase gene is a useful candidate for improving the nutritional qualities and Fe-deficiency tolerance of various crops. *Front Plant Sci* 9:340. <https://doi.org/10.3389/fpls.2018.00340>
65. Nozoye T, Kim S, Kakei Y, Takahashi M, Nakanishi H, Nishizawa NK (2014) Enhanced levels of nicotianamine promote iron accumulation and tolerance to calcareous soil in soybean. *Biosci Biotechnol Biochem* 78:1677–1684. <https://doi.org/10.1080/09168451.2014.936350>
66. Papias D, Taxiarchou M, Paspaliaris I, Kontopoulos A (1996) Mechanisms of dissolution of iron oxides in aqueous oxalic acid solutions. *Hydrometallurgy* 42:257–265.

[https://doi.org/10.1016/0304-386X\(95\)00104-0](https://doi.org/10.1016/0304-386X(95)00104-0)


67. Qiu W, Dai J, Wang N, Guo X, Zhang X, Zuo Y (2017) Effects of Fe-deficient conditions on Fe uptake and utilization in P-efficient soybean. *Plant Physiol Biochem* 112:1–8.  
<https://doi.org/10.1016/j.plaphy.2016.12.010>
68. Rahman MA, Kabir AH, Song Y, Guo X, Zhang X, Zuo Y (2021) Nitric oxide prevents Fe deficiency-induced photosynthetic disturbance, and oxidative stress in alfalfa by regulating Fe acquisition and antioxidant defense. *Antioxidants* 10:1556. <https://doi.org/10.3390/antiox10101556>
69. Rahman MA, Bagchi R, El-Shehawi AM, Elseehy MM, Anee SA, Lee K-W, Kabir AH (2022) Physiological and molecular characterization of strategy-I responses and expression of Fe-transporters in Fe-deficient soybean. *South Afr J Bot* 147:942–950.  
<https://doi.org/10.1016/j.sajb.2022.03.052>
70. Raj KK, Pandey RN, Singh B, Meena MC, Talukdar A, Chakraborty D (2021) Response of soybean genotypes to iron limiting stress in calcareous vertisol under ambient and elevated CO<sub>2</sub> and temperature conditions. *J Environ Biol* 42:295–301. <https://doi.org/10.22438/JEB/42/2/MRN-1226>
71. Ramírez L, Graziano M, Lamattina L (2008) Decoding plant responses to iron deficiency. *Plant Signal Behav* 3:795–797. <https://doi.org/10.4161/psb.3.10.5874>
72. Ramírez L, Bartoli CG, Lamattina L (2013) Glutathione and ascorbic acid protect Arabidopsis plants against detrimental effects of iron deficiency. *J Exp Bot* 64:3169–3178.  
<https://doi.org/10.1093/jxb/ert153>
73. Rellán-Álvarez R, El-Jendoubi H, Wohlgemut G, Abadía A, Fiehn O, Abadía J, Álvarez-Fernández A (2011) Metabolite profile changes in xylem sap and leaf extracts of strategy I plants in response to iron deficiency and resupply. *Front Plant Sci* 2:66. <https://doi.org/10.3389/fpls.2011.00066>
74. Rios JJ, Carrasco-Gil S, Abadía A, Abadía J (2016) Using Perls staining to trace the iron uptake pathway in leaves of a Prunus rootstock treated with iron foliar fertilizers. *Front Plant Sci* 7:893.  
<https://doi.org/10.3389/fpls.2016.00893>
75. Robe K, Izquierdo E, Vignols F, Rouached H, Dubos C (2021) The coumarins: secondary metabolites playing a primary role in plant nutrition and health. *Trends Plant Sci* 26:248–259.  
<https://doi.org/10.1016/j.tplants.2020.10.008>
76. Rodríguez-Lucena P, Roperio E, Hernández-Apaolaza L, Lucena JJ (2010a) Iron supply to soybean plants through the foliar application of IDHA/Fe<sup>3+</sup>: effect of plant nutritional status and adjuvants. *J Sci Food Agric* 90:2633–2640. <https://doi.org/10.1002/jsfa.4132>
77. Rodríguez-Lucena P, Hernández-Apaolaza L, Lucena JJ (2010b) Comparison of iron chelates and complexes supplied as foliar sprays and in nutrient solution to correct iron chlorosis of soybean. *J Plant Nutr Soil Sci* 173:120–126. <https://doi.org/10.1002/jpln.200800256>
78. Rogers EE, Wu X, Stacey G, Nguyen HT (2009) Two MATE proteins play a role in iron efficiency in soybean. *J Plant Physiol* 166:1453–1459. <https://doi.org/10.1016/j.jplph.2009.02.009>
79. Santos CS, Silva AI, Serrão I, Carvalho AL, Vasconcelos MW (2013) Transcriptomic analysis of iron deficiency related genes in the legumes. *Food Res Int* 54:1162–1171.

<https://doi.org/10.1016/j.foodres.2013.06.024>

80. Santos CS, Roriz M, Carvalho SMP, Vasconcelos MW (2015) Iron partitioning at an early growth stage impacts iron deficiency responses in soybean plants (*Glycine max* L.). *Front Plant Sci* 6:325. <https://doi.org/10.3389/fpls.2015.00325>
81. Santos CS, Serrão I, Vasconcelos MW (2016) Comparative analysis of iron deficiency chlorosis responses in soybean (*Glycine max*) and barrel medic (*Medicago truncatula*). *Rev Ciências Agrárias* 39:538–549. <https://doi.org/10.19084/rca16090>
82. Santos CS, Ozgur R, Uzilday B, Turkan I, Roriz M, Rangel AOSS, Calvalho SMP, Vasconcelos MW (2019) Understanding the role of the antioxidant system and the tetrapyrrole cycle in iron deficiency chlorosis. *Plants* 8:348. <https://doi.org/10.3390/plants8090348>
83. Shanmugam V, Tsednee M, Yeh K-C (2012) Zinc tolerance induced by iron 1 reveals the importance of glutathione in the cross-homeostasis between zinc and iron in *Arabidopsis thaliana*. *Plant J* 69:1006–1017. <https://doi.org/10.1111/j.1365-313X.2011.04850.x>
84. Sisó-Terraza P, Rios JJ, Abadía J, Abadía A, Álvarez-Fernández A (2016) Flavins secreted by roots of iron-deficient *Beta vulgaris* enable mining of ferric oxide via reductive mechanisms. *New Phytol* 209:733–745. <https://doi.org/10.1111/nph.13633>
85. Stander HM, Harrison STL, Broadhurst JL (2022) Using south african sulfide-enriched coal processing waste for amelioration of calcareous soil: A pre-feasibility study. *Min Eng* 180:107457. <https://doi.org/10.1016/j.mineng.2022.107457>
86. Tiffin LO (1970) Translocation of iron citrate and phosphorus in xylem exudate of soybean. *Plant Physiol* 45:280–283. <https://doi.org/10.1104/pp.45.3.280>
87. Tighe M, Lockwood P, Wilson S, Lisle L (2004) Comparison of digestion methods for ICP-OES analysis of a wide range of analytes in heavy metal contaminated soil samples with specific reference to arsenic and antimony. *Commun Soil Sci Plant Anal* 35:1369–1385. <https://doi.org/10.1081/CSS-120037552>
88. Vasconcelos MW, Grusak MA (2014) Morpho-physiological parameters affecting iron deficiency chlorosis in soybean (*Glycine max* L.). *Plant Soil* 374:161–172. <https://doi.org/10.1007/s11104-013-1842-6>
89. Vasconcelos MW, Clemente TE, Grusak MA (2014) Evaluation of constitutive iron reductase (AtFRO2) expression on mineral accumulation and distribution in soybean (*Glycine max* L.). *Front Plant Sci* 5:112. <https://doi.org/10.3389/fpls.2014.00112>
90. Vergauwen B, Verstraete K, Senadheera DB, Dansercoer A, Cvitkovitch DG, Guédon E, Savvides SN (2013) Molecular and structural basis of glutathione import in Gram-positive bacteria via GshT and the cystine ABC importer TcyBC of *Streptococcus mutans*. *Mol Microbiol* 89:288–303. <https://doi.org/10.1111/mmi.12274>
91. Wallace A, Mueller RT, Patel PM, Soufi SM (1976) Use of waste pyrites from mine operations on highly calcareous soil. *Commun Soil Sci Plant Anal* 7:57–60. <https://doi.org/10.1080/00103627609366613>


92. Wallace A, Romney EM, Wallace GA, Mueller RT, Cha JW (1980) Pyrite as a source of iron for plants. *J Plant Nutr* 2:193–195. <https://doi.org/10.1080/01904168009362762>
93. Wang B, Wei H, Xue Z, Zhang WH (2017) Gibberellins regulate iron deficiency-response by influencing iron transport and translocation in rice seedlings (*Oryza sativa*). *Ann Bot* 119:945–956. <https://doi.org/10.1093/aob/mcw250>
94. Waters BM, Amundsen K, Graef G (2018) Gene expression profiling of iron deficiency chlorosis sensitive and tolerant soybean indicates key roles for phenylpropanoids under alkalinity stress. *Front Plant Sci* 9:10. <https://doi.org/10.3389/fpls.2018.00010>
95. Wiersma JV (2005) High rates of Fe-EDDHA and seed iron concentration suggest partial solutions to iron deficiency in soybean. *Agron J* 97:924–934. <https://doi.org/10.2134/agronj2004.0309>
96. Zaharieva TB, Abadía J (2003) Iron deficiency enhances the levels of ascorbate, glutathione, and related enzymes in sugar beet roots. *Protoplasma* 221:269–275. <https://doi.org/10.1007/s00709-002-0051-6>
97. Zaharieva T, Yamashita K, Matsumoto H (1999) Iron deficiency induced changes in ascorbate content and enzyme activities related to ascorbate metabolism in cucumber roots. *Plant Cell Physiol* 40:273–280. <https://doi.org/10.1093/oxfordjournals.pcp.a029538>
98. Zaharieva TB, Gogorcena Y, Abadía J (2004) Dynamics of metabolic responses to iron deficiency in sugar beet roots. *Plant Sci* 166:1045–1050. <https://doi.org/10.1016/j.plantsci.2003.12.017>
99. Zocchi G (2006) Metabolic changes in iron-stressed dicotyledonous plants. In: Barton LL, Abadía J (eds) *Iron Nutrition in Plants and Rhizospheric Microorganisms*. Springer, Dordrecht, The Netherlands, pp 359–370. [https://doi.org/10.1007/1-4020-4743-6\\_18](https://doi.org/10.1007/1-4020-4743-6_18)
100. Zocchi G, Cocucci S (1990) Fe uptake mechanism in Fe-efficient cucumber roots. *Plant Physiol* 92:908–911. <https://doi.org/10.1104/pp.92.4.908>
101. Zocchi G, De Nisi P, Dell’Orto M, Espen L, Gallina PM (2007) Iron deficiency differently affects metabolic responses in soybean roots. *J Exp Bot* 58:993–1000. <https://doi.org/10.1093/jxb/erl259>

## Figures




**Figure 1**

SPAD index in young leaves 50 days after sowing. Treatments were applied to the soil at days 21, 28 and 35, and consisted in 0, 5, 31.8 or 37.3 mg Fe kg<sup>-1</sup> soil (from Fe(III)-EDDHA, Fe<sub>oxi</sub> or Fe<sub>ore</sub>, respectively), supplemented with no thiols, 5 mM GSH, 0.5 mM DTT or 5 mM PhSH. Values shown are means ± SE (n = 24). Letters above the columns indicate significant differences at  $p \leq 0.01$  for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).




**Figure 2**

Plant biomass and height at the end of the experiment (at day 60). Treatments are as described in Figure 1. Shoot DW (**A**), root DW (**B**) (in g plant<sup>-1</sup>), and plant height (**C**; in cm). Values shown are means  $\pm$  SE (n = 9 for FW and 12 for height). Letters above the columns indicate significant differences at  $p \leq 0.01$  for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).



**Figure 3**

Iron concentrations at the end of the experiment (at day 60) in young leaves (A), developed leaves (B), stems (C) and roots (D) (in  $\mu\text{g g}^{-1}\text{DW}$ ). Treatments are as described in Figure 1. Values shown are means  $\pm$  SE ( $n = 3$ ). Letters above the columns indicate significant differences at  $p \leq 0.01$  for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).




**Figure 4**

Iron contents at the end of the experiment (at day 60) in young (A) and developed (B) leaves, stems (C) and root (D), (in µg plant<sup>-1</sup>). Treatments are as described in Figure 1. Values shown are means ± SE (n =

3). Letters above the columns indicate significant differences at  $p \leq 0.01$  for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).


**Figure 5**

Antioxidant compounds at the end of the experiment (at day 60) (in  $\text{nmol g}^{-1}$  FW). Treatments are as described in Figure 1. Reduced glutathione (tGSH; in  $\text{nmol g}^{-1}$  FW) in leaves (A) and roots (C), oxidized glutathione (tGSSG; in  $\text{nmol g}^{-1}$  FW) in leaves (B) and roots (D) and ascorbate (Asc;  $\mu\text{mol g}^{-1}$  FW) in leaves (E). Values shown are means  $\pm$  SE ( $n = 3$ ). Letters above the columns indicate significant differences at  $p \leq 0.01$  for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).




**Figure 6**

Antioxidant enzymes in leaves at the end of the experiment (at day 60). Treatments are as described in Figure 1. Glutathione reductase (GR; A, in  $\text{U mg}^{-1}$  protein) and ascorbate peroxidase activity (APX; B, in  $\text{U mg}^{-1}$  protein). Values shown are means  $\pm$  SE ( $n = 3$ ). Letters above the columns indicate significant differences at  $p \leq 0.01$  for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).



**Figure 7**


Leaf carboxylates at the end of the experiment (at day 60). Treatments are as described in Figure 1. Leaf citrate (A) and malate (B) (in mg g<sup>-1</sup> DW). Values shown are means ± SE (n = 3). Letters above the columns indicate significant differences at  $p \leq 0.01$  for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).



**Figure 8**


Gibberellic and abscisic acid in leaves at the end of the experiment (at day 60). Treatments are as described in Figure 1. Gibbelleric acid (GA, in µg g<sup>-1</sup> DW) (A) and abscisic acid (ABA; in µg g<sup>-1</sup> DW) (B). Values shown are means ± SE (n = 3). Letters above the columns indicate significant differences at  $p \leq$

0.01 for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).



**Figure 9**

DTPA-extractable soil Fe concentration at the end of the experiment (at day 60) (in  $\text{mg kg}^{-1}$  soil). Treatments are as described in Figure 1. Values shown are means  $\pm$  SE ( $n = 3$ ). Letters above the columns indicate significant differences at  $p \leq 0.01$  for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).



**Figure 10**

Iron measured by ICP-OES (without acid digestion; ICP-Fe) and Fe(II) in the solution after incubation of Fe<sub>oxi</sub> and Fe<sub>ore</sub> with BPDS (in mg L<sup>-1</sup>). Total ICP-Fe (after acid digestion) in the solution was 8.9 and 175.6 mg Fe L<sup>-1</sup> with the Fe<sub>oxi</sub> and Fe<sub>ore</sub>, respectively. ICP-Fe after 120 min (A), and Fe(II) measured spectrophotometrically after 60 (A) and 120 min (B). Fifty mg of product was stirred with 10 mL of buffer in 300 μM BPDS for different times. Values shown are means ± SE (n = 3). Letters above the columns indicate significant differences at *p* ≤ 0.01 for the thiol treatments in a given Fe treatment (in lower case) and for the Fe treatments in a given thiol treatment (in capitals).




Figure 11

Heatmap summarizing the changes observed with the Fe-containing products or thiols were applied alone.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterials.v.102.docx](#)