Delayed Pancreatic Fistula: An Unaccustomed Complication Following Pancreaticoduodenectomy.

Suvendu Sekhar Jena (✉ suvu1078@gmail.com)

Sir Ganga Ram Hospital <https://orcid.org/0000-0002-3072-0418>

Dibyasingh Meher

VMMC and Safdarjung Hospital: Vardhman Mahavir Medical College and Safdarjung Hospital

Rahul Ranjan

VMMC and Safdarjung Hospital: Vardhman Mahavir Medical College and Safdarjung Hospital

Case report

Keywords: Post-operative pancreatic fistula, Occult pancreatic fistula, overt pancreatic fistula
Pancreaticoduodenectomy, Percutaneous drainage, Endoscopic stenting

Posted Date: February 2nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-167066/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Post-operative pancreatic fistula is a morbid complication after pancreaticoduodenectomy. Though most of them present in the immediate post-operative period, few case reports have mentioned it even 7 years after index surgery. Here, we report a delayed presentation of pancreatic fistula 6 months after surgery.

Case report

A 57 year old female underwent Whipple's pancreaticoduodenectomy for pancreatic head adenocarcinoma and was discharged with an uneventful post-operative recovery. She presented after 6 months with complaints of abdominal pain and distension which upon evaluation found to be a pancreatic enzyme rich multiloculated collection. It was managed with per-cutaneous drain placement.

Discussion

Pancreatic fistula remained a major cause of morbidity and mortality even after 100 years of its existence. It can be overt fistula which manifest in the immediate post-operative period or occult fistula which manifests long after primary surgery. Various causes of delayed fistula are anastomotic site stricture, previous chemotherapy, infection. The management options available are percutaneous drainage, endoscopic stenting of anastomotic stricture or redoing the anastomosis.

Conclusion

Pancreatic fistula can have a delayed presentation which can be diagnosed and managed with regular follow up.

Introduction

After its inception by Whipple in 1914, pancreaticoduodenectomy (PD) became preferred choice of surgery in both benign and malignant diseases of periampullary region.^[1] Though operative mortality (3–5%) is decreased but morbidity still remained high which around 30–65%.^[2] Dealing with pancreatic stump is most important challenge and mostly the cause of postoperative complication.^[3] Pancreatic fistula is one of major complication after PD which is around 2–28%.⁴ There are lots of literature described about post-operative pancreatic fistula (POPF), but none of them have been mentioned about the timing of presentation apart from few case reports about delayed pancreatic fistula according to our knowledge. Here we present another case of pancreatic fistula presented 6 months after PD.

Case Presentation

Fifty seven year old female, without any co-morbidity, with past history of cholecystectomy, presented with history of progressively worsening jaundice. It was associated with itching, occasional fevers with

chills and rigor. She was evaluated with ultrasound abdomen which showed mild dilatation of the intrahepatic biliary tree and the common hepatic duct with a dilated common bile duct measuring 9 mm and main pancreatic duct is dilated and measuring 4.1 mm. Computed tomography (CT) angiography of abdomen showed periampullary mild heterogeneous enhancing mass of size (2.0 x 1.9 x 1.7) cm mainly involving the lower end of the common bile duct causing upstream dilatation of the biliary radicals and extrahepatic duct, also causes mild dilatation of the main pancreatic duct, associated with some pancreatitis involving the head and uncinate process with stranding around the distal SMV along the right side without any vascular invasion. There was narrowing of celiac artery with collaterals between celiac artery and superior mesenteric artery over the head of pancreas. **(Figure – 1)** In view of features of cholangitis, she was planned for endoscopic stenting of common bile duct but percutaneous transhepatic drainage was done in view of repeated failed cannulation. After optimization, she underwent Whipple's pancreaticoduodenectomy with division of median arcuate ligament. Intra-operatively, there was a pancreatic head mass of size (3 x 1) cm with a large aorto-caval lymph node which on frozen section was negative for metastatic deposit, pancreas was soft with a duct diameter of 3mm **(Figure – 2)** The histopathological report showed moderately differentiated adenocarcinoma of the head of pancreas with lymphovascular invasion and peri-neural invasion with 5 positive out of 42 lymph nodes with a pathological stage of pT1cpN2pMx. The drain fluid amylase sent on post-operative day 3 (POD) from both drains were 8 U/L and 9 U/L. Both the drains were removed and she was discharged on POD 12. She received adjuvant chemotherapy in the form of 2 cycles of gemcitabine and capecitabine. Post 2nd cycle chemotherapy she developed pain abdomen which was initially of low intensity but gradually worsened. She was admitted to a local hospital with hypotension and a total leukocyte count of 26,000/cu mm. After optimization, she presented here and on evaluation with CT abdomen there was a large multiloculated collection with enhancing wall in the upper and lower abdomen with haziness around the PJ site underneath the anterior abdominal wall with a distended stomach. **(Figure – 3 and 4)** Feeding was started after using nasojejunal tube. CT guided percutaneous drain was placed and drained fluid on analysis had an amylase of 1131 U/L and lipase of 3003 U/L. Somatostatin analogue was started due to raised drain fluid enzyme levels. Drain fluid culture was positive for Klebsiella pneumonia and antibiotics were changed as per sensitivity. Gradually drain output reduced and subsequent CT abdomen showed no significant collection. **(Figure – 5)** She was discharged with per-cutaneous drain in situ which was removed on outpatient basis. Now she is doing well in subsequent follow up 15 months post surgery.

Discussion And Conclusion

PD continues to be a difficult surgery even 100 years after its inception by Dr Walter Kausch and its modification by Dr. Allen Oldfather Whipple. With improvement of surgical techniques, though the mortality rate dropped, the morbidity still remained on higher side. Delayed gastric emptying, wound infection, post-pancreatectomy hemorrhage, postoperative pancreatic fistula (POPF), intra-abdominal abscess are amongst the common and serious complications following PD. Pancreatic anastomosis leak causing POPF still remains the leading cause of morbidity following PD even in specialized centers.^[5, 6, 7] The main hurdle to manage POPF was lack of standardized and universally accepted definition until

2004 when 37 pancreatic surgeons from 15 countries formed International Study Group of Pancreatic Fistula (ISGPF) and purposed a definition for POPF. According to ISGPF, POPF was defined as measurable fluid output on or after postoperative day 3, with amylase content higher than three times the upper normal serum level. [8] It was revised in 2016 and divided POPF into biochemical fistula and clinically relevant fistula. [9]

The risk factors of POPF following PD can be various like etiologies for surgery, pancreatic parenchymal texture, duct diameter, surgical techniques, blood loss, high BMI. In regards to etiologies, PD done for pancreatic adenocarcinoma has a low risk for fistula formation in comparison to ampullary, duodenal, distal cholangiocarcinoma. [10,11,12,13] Risk of POPF increased many fold in patients with hard pancreas and small pancreatic duct diameter < (2-3) mm. [10,14,15]

Depending upon timing of its presentation, Veillette et al divided POPF into overt and occult fistula. Overt fistulas are defined by their evidence within first post-operative week and causes major morbidity and mortality. Occult fistulas are those which are not manifested in the initial post-operative period but subsequently resulted in intra-abdominal collection, abscess formation, hemorrhage and death. In their series, there were 13 occult fistulas. They also found that occult fistulas were responsible for repeated hospital admissions but there was no difference between them in relation to intervention, reoperation or mortality. But the occult fistulas in their study presented within 90 days of surgery. [16]

So, delayed presentation of POPF is a rare presentation which is rarer after PD and is only limited to individual case reports. [17] There are few case reports of delayed POPF after splenectomy. [18,19] Though the cause of delayed presentation of fistula is yet to be established, there are various plausible explanations. Pancreatic duct obstruction, recurrent pancreatitis, ischemia and fibrosis may lead to anastomotic disruption causing delayed POPF. [20] Pancreatojejunal anastomotic site stricture secondary to chemotherapy for pancreatic carcinoma may be another explanation. [21] Smoking which creates a thrombogenic environment along with its vasoconstrictive property may be another factor responsible for delayed POPF. [22] Small amount of initial leakage which often goes unnoticed may be aggravated by infection causing delayed fistula is another theory behind POPF. Gray zone between POPF and post-operative collections was described by Barreto et al. They proposed drains failing to drain properly due to blockage, displaced or misplaced drains and vascular factors may be responsible for delayed presentation of POPF. [23] In this case, though her immediate post-operative drain fluid amylase levels were normal ruling out an overt fistula, she presented with collections after receiving chemotherapy which may be the cause of delayed POPF by causing pancretojejunal anastomotic stricture.

Due to its rarity, the exact time duration of its presentation from index surgery is yet to be elucidated. Veillette et al in their series found all occult fistulas presented within 90 days of index surgery. [16] In a case report by Faraj et al, the duration was 7 years which is by far the longest interval reported in literature. [20] Ito et al reported a gap of 1 year and it was 8 months by Perez et al. In our case, patient presented to us after 6 months of index surgery. [24, 3]

Management of this unique condition may range from percutaneous drainage, endoscopic stenting of anastomotic stricture to redo of the anastomosis. [3, 20, 24] Patient was managed with antibiotics and percutaneous drainage in our case.

A delayed pancreatic fistula after PD is a rare complication and is limited to only individual case reports. This is the fourth case report on delayed POPF in literature to our best of knowledge. A little is known about its etiopathogenesis and few plausible explanations are available regarding the same without any concrete evidence. Few management options are available in the form of percutaneous drainage, stenting of anastomotic stricture and redoing the anastomosis. Though this study provides a different prospective to the existing literature about the delayed presentation of pancreatic fistula, a case series involving large number of patients is required to establish its etiopathogenesis and management.

Abbreviations

PD- Pancreaticoduodenectomy.

POPF- Post operative Pancreatic Fistula.

CT- Computed Tomography.

POD – Post- operative day.

ISGPF - International Study Group of Pancreatic Fistula.

Declarations

Ethical Committee and consent to participate - Ethical committee clearance was not needed as per the local ethical committee regulations. Consent to participate in the study was taken from the patient.

Consent for publication - Consent for publications was taken from the patient.

Availability of data and materials - All the data and material is available with the author.

Funding – The research received no specific grant from any funding agency in public, commercial or not-for-profit sector.

Author Contribution - Dr Suvendu Sekhar Jena - Data collection and analysis, Writing Case report and Discussion. Dr Dibyasingh Meher – Writing abstract, introduction and article name. Dr Rahul Ranjan – Study designing and Final proofing.

Acknowledgements – Not applicable.

References

1. Conzo G, Gambardella C, Tartaglia E, Sciascia V, Mauriello C, Napolitano S et al. Pancreatic fistula following pancreatoduodenectomy. Evaluation of different surgical approaches in the management of pancreatic stump. Literature review. *Int J Surg*. 2015; 21 Suppl 1:S4-9.
2. Machado NO. Pancreatic fistula after pancreatectomy: definitions, risk factors, preventive measures, and management-review. *Int J Surg Oncol*. 2012;602478.
3. Perez NP Jr, Forcione DG, Ferrone CR. Late pancreatic fistula after pancreaticoduodenectomy: A case report and review of the literature. *Case Rep Pancreat Cancer*. 2016;2:65-70.
4. Haddad LB, Scatton O, Randone B, Andraus W, Massault PP, Dousset B et al. Pancreatic fistula after pancreaticoduodenectomy: the conservative treatment of choice. *HPB*. 2009;11:203-9.
5. Rumstadt B, Schwab M, Korth P, Samman M, Trede M. Hemorrhage after pancreatoduodenectomy. *Ann Surg*. 1998 Feb;227(2):236-41.
6. Cullen JJ, Sarr MG, Ilstrup DM. Pancreatic anastomotic leak after pancreaticoduodenectomy: incidence, significance, and management. *American Journal of Surgery*. 1994;168: 295–8.
7. Strasberg SM, Drebin JA, Soper NJ. Evolution and current status of the whipple procedure: an update for gastroenterologists. *Gastroenterology*. 1997;113: 983–94.
8. Bassi C, Butturini G, Molinari E, Mascetta G, Salvia R, Falconi M et al. Pancreatic fistula rate after pancreatic resection. The importance of definitions. *Dig Surg*. 2004;21:54-9.
9. Bassi C, Dervenis C, Butturini G, et al. Postoperative pancreatic fistula: an international study group (ISGPF) definition. *Surgery*. 2005;138:8–13.
10. Callery MP, Pratt WB, Kent TS, et al. A prospectively validated clinical risk score accurately predicts pancreatic fistula after pancreatoduodenectomy. *J Am Coll Surg*. 2013;216:1–14.
11. Vin Y, Sima CS, Getrajdman GI, Brown KT, Covey A, Brennan MF et al. Management and outcomes of postpancreatectomy fistula, leak, and abscess: results of 908 patients resected at a single institution between 2000 and 2005. *J Am Coll Surg*. 2008;207:490-8.
12. Schmidt CM, Choi J, Powell ES, Yiannoutsos CT, Zyromski NJ, Nakeeb A et al. Pancreatic fistula following pancreaticoduodenectomy: clinical predictors and patient outcomes. *HPB Surg*. 2009;2009:404520.
13. van Berge Henegouwen MI, De Wit LT, Van Gulik TM, Obertop H, Gouma DJ. Incidence, risk factors, and treatment of pancreatic leakage after pancreaticoduodenectomy: drainage versus resection of the pancreatic remnant. *J Am Coll Surg*. 1997;185:18–24.
14. Muscari F, Suc B, Kirzin S, Hay JM, Fournanier G, Fingerhut A et al. Risk factors for mortality and intraabdominal complications after pancreatoduodenectomy: multivariate analysis in 300 patients. *Surgery*. 2006;139:591–8.
15. Yeo CJ, Cameron JL, Lillemoe KD, Sauter PK, Coleman J, Sohn TA et al. Does prophylactic octreotide decrease the rates of pancreatic fistula and other complications after pancreaticoduodenectomy? Results of a prospective randomized placebo-controlled trial. *Ann Surg*. 2000;232:419–429.

16. Veillette G, Dominguez I, Ferrone C, Thayer SP, McGrath D, Warshaw AL et al. Implications and management of pancreatic fistulas following pancreaticoduodenectomy: the Massachusetts General Hospital experience. *Arch Surg.* 2008;143:476–81.
17. Casadei R, Bassi F, Calculli L, Zanini N, Greco VM, Minni F. Report of three cases of chronic pancreatic fistulas treated with prolamine as a sclerosing substance following pancreatic resection. *JOP.* 2006; 7:41-6.
18. Jin SG, Chen ZY, Yan LN, Zeng Y. Delayed internal pancreatic fistula with pancreatic pleural effusion postsplenectomy. *World J Gastroenterol* 2010; 16:4494-6.
19. Yagmurdur MC, Gür G, Aytakin C, Karakayali H, Boyacioglu S, Bilgin N. Atypical presentation of chronic pancreatic fistula: a case successfully managed with long-acting somatostatin. *Turk J Gastroenterol* 2003; 14:200-3.
20. Faraj W, Abou Zahr Z, Mukherji D, Zaghal A, Khalife M. Pancreatic anastomosis disruption seven years postpancreaticoduodenectomy. *Case Rep Med.* 2010;2010:436739.
21. Cioffi JL, McDuffie LA, Roch AM, Zyromski NJ, Ceppa EP, Schmidt CM et al. Pancreaticojejunostomy stricture after pancreatoduodenectomy: outcomes after operative revision. *J Gastrointest Surg.* 2016;20:293–9.
22. Mathur AK, Ranney DN, Patel SP, Lee DS, Bednar F, Lynch RJ et al. The effect of smoking on biliary complications following liver transplantation. *Transpl Int.* 2011;24:58–66.
23. Barreto G, D'Souza MA, Shukla PJ, Shrikhande SV. The gray zone between postpancreaticoduodenectomy collections and pancreatic fistula. *Pancreas* 2008;37:422-5.
24. Ito Y, Irino T, Egawa T, Hayashi S, Nagashima A. Delayed pancreatic fistula after pancreaticoduodenectomy. A case report. *JOP.* 2010;12:410-2.

Figures

Figure 1

Computed tomography of abdomen showing pancreatic head mass (Red arrow), multiple collaterals around the pancreatic head (White arrow).

Figure 2

Whipple's Pancreaticoduodenectomy specimen.

Figure 3

CT abdomen showing a collection with enhancing wall (White arrow).

Figure 4

Coronal section of CT abdomen showing multiloculated collection (Double white arrow) with a distended stomach (Single white arrow).

Figure 5

CT abdomen post drainage showing no significant collection.