

Anxiety, Perceived Stress and Coping Strategies in Nursing Students: A Cross-Sectional Correlational Descriptive Study

María Dolores Onieva-Zafra

Department of Nursing, Physiotherapy and Occupational Therapy, Nursing Faculty of Ciudad Real. Universidad de Castilla-La Mancha, Spain

Juan José Fernández-Muñoz

Department of Psychology, Universidad Rey Juan Carlos I, Alcorcón, Madrid, Spain

Elia Fernández-Martínez (✉ elia.fernandez@denf.uhu.es)

Universidad de Huelva - Campus El Carmen <https://orcid.org/0000-0002-7700-999X>

Francisco José García-Sánchez

Department of Nursing, Physiotherapy and Occupational Therapy, Nursing Faculty of Ciudad Real. Universidad de Castilla-La Mancha, Spain

Ana Abreu-Sánchez

Department of Nursing. University of Huelva, Spain

María Laura Parra-Fernández

Department of Nursing, Physiotherapy and Occupational Therapy, Nursing Faculty of Ciudad Real. Universidad de Castilla-La Mancha, Spain

Research article

Keywords: Anxiety, coping behavior, nursing students, perceived stress

Posted Date: August 14th, 2020

DOI: <https://doi.org/10.21203/rs.2.10722/v3>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on October 19th, 2020. See the published version at <https://doi.org/10.1186/s12909-020-02294-z>.

Abstract

Background: Clinical training is recognized as being a stressful experience for many nursing students. The levels of stress and anxiety may increase or decrease during their educational training depending on coping behavior strategies and other factors. The objective of this was to investigate the relationship between anxiety, perceived stress and coping strategies used by nursing students during their clinical training.

Methods: A cross-sectional correlational descriptive study. The sample consisted of 190 nursing students enrolled in the Nursing Faculty of Ciudad Real University in Spain. Participants provided data on background characteristics and completed the following instruments: Perceived Stress Scale, State-Trait Anxiety Inventory and Coping Behavior Inventory. Relationships between scores were examined using Spearman's rho.

Results: The mean age of participants was 20.71 ± 3.89 years (range 18-46 years). One half of the students (47.92%) indicated a moderate level of stress with a mean of Perceived Stress Scale score of $22.78 (\pm 8.54)$. Veteran nursing students perceived higher levels of stress than novice students. The results showed a significant correlation for perceived stress and state anxiety ($r = 0.463, p < .000$) and also for trait anxiety ($r = 0.718, p < .000$). There was also a significant relationship between the total amount of perceived stress and the following domains of the coping behavior inventory: problem solving ($r = -.452, p < .01$), self-criticism ($r = .408, p < .01$), wishful thinking ($r = .459, p < .01$), social support ($r = -.220, p < .01$), cognitive restructuring ($r = -.375, p < .01$), and social withdrawal ($r = .388, p < .01$). In the current study, the coping strategy most frequently used by students was problem-solving, followed by social support and cognitive restructuring.

Conclusions: Nursing students in our study presented a moderate level of stress and the correlation with anxiety was significant. Nursing teachers and clinical preceptors/mentors should be encouraged to develop programs that could help prepare nursing students to cope with the challenges they are about to face during their clinical placements.

Background

Research on stress levels in health professionals is an issue of current interest that merits concern [1]. This is not only due to the many causes of intrinsic stress referred by healthcare professionals, but also because it is worth considering the negative and chronic effects that stress can have in the same over time [2].

In nursing, the subject of stress has received much attention in the literature, and continues to be the topic of many studies [3, 4]. According to several studies, the practical part of a nurse's education is much more stressful than the academic part and the perceived lack of knowledge and skills are considered to be one of the common stressors for many students (Yildiz et al., 2014). Furthermore, the first experience in

clinical practice includes stressors such as fear of making mistakes, having to handle emergency situations, irregularities in clinical practice and visiting specialized units [7].

Generally, nursing students do not have the same responsibility in the individual care of patients in clinical practice as registered nurses, however they are exposed to some of the same stressors. Examples of the same include the relationships with other professionals, the notorious ranking that exists in hospitals, difficult situations regarding the treatment of patients and dealing with family members and the way they experience the death of the patients they care for [8]. Furthermore, nursing students coexist with other stressors that are typical considering their role as students, such as those related with their academic program and their role as nursing students [9]. This is because, as opposed to other degree programs, nursing students are in touch with the job market which requires a certain responsibility in the wellbeing of their patients, distancing them, at times from the student campus life and especially, from normal social activities enjoyed by their same-age colleagues.

Low or moderate levels of stress may enhance students' motivation, leading to greater perseverance when studying and achieving future goals [10]. Conversely, high levels of stress can have a negative influence on students, leading to depression and despair, and therefore affecting the health and academic level of students [11]. Stress is unavoidable and, in most cases, it is difficult to overcome, however, a good coping strategy help students to improve their academic results [12].

Coping mechanisms are absolutely necessary when trying to deal with the stress and anxiety that nursing students face on a daily basis. Longitudinal studies have shown that stress levels in nursing students may increase or decrease during their educational training depending on coping behavior strategies. However, as noted by Jimenez et al.[13], these differences regarding stress levels over the course of professional training should be considered with caution, as different programs exist in different countries. Whereas some studies identify the first or second year as the most stressful year for students [14], for others, the third year is the most stressful regarding the clinical aspect (Edwards et al., 2004; Evans and Kelly, 2004). However, other studies do not specify the year, revealing results in which stress increases according to the training or the academic year [19, 20] or decreases as the student becomes more trained [21]. Besides these differences, coping strategies vary according to the characteristics of the individual and the context where the stressors are found. The problem solving strategy has been the most cited in different studies as the best way to cope with stress. On the other hand, emotional based coping strategies were found the least effective strategy [22]. Therefore, the main aim of this study was to investigate the relationship between anxiety, perceived stress and coping strategies used by nursing students during their clinical training. Moreover, we performed a comparative analysis between perceived stress and coping strategies by gender, course and clinical placement was applied, based on the previous literature.

Methods

Study design, setting and participants.

A cross-sectional descriptive and correlational design was used. Students were recruited from graduate levels of the Ciudad Real Nursing Faculty of Castilla-La Mancha University during the 2017/18 academic year. Currently, in Spain the nursing degree lasts four years with a total of 240 credits, under the European Credits Transfer System (ETCS). At the University of Castilla-La-Mancha (UCLM), during the first academic year, nursing students only have academic subjects (theoretical-practical) which are taught at the Faculty, having no contact at this point with the hospital world. After the first year, once the basic core subjects are taught, the student begins clinical placements. These start in the first semester of the second year and end in the fourth year with subject, that are purely care-based and which take place at the hospital (Anex 1). In those three years they alternate clinical placement with academic subjects.

A prior sample size calculation was not performed, rather the sample size was based on the entire population of students, as described previously. Clinical placements are an essential part of the acquisition of competencies and skills for nursing students, constituting an important element of their education.

The eligibility criteria for this study were as follows: all participants were students enrolled in an academic year of the nursing degree course taught at the university (except first year students). Participants must have been present in the classroom when the researcher visited to collect data and informed consent was required for participation. The population of nursing students studying at the university was 340 students, 115 were removed because they did not fulfill the inclusive criteria; thirty-three of them did not participate at the research and finally 192 student agree to participate and they self-administered the questionnaire. The response rate was 85.33%.

Data collection instruments.

Data were collected from the students in the study group using a 20- minute online self-report questionnaire containing the following measures:

-Demographic characteristics: The demographic questionnaire was constructed by the researchers and was based on the recent literature. This included items such as age, gender, relationship status, study year of the nursing degree etc.

-Perceived stress scale (PSS). The PSS-14 was designed for measuring the degree to which daily life situations are evaluated as stressors. The PSS-14 consists of multiple choice questions measuring stressful experiences and responses to stress over the previous four weeks. The European Spanish version PSS (14-item) has demonstrated adequate reliability (internal consistency, $\alpha = .81$) [23]. Our study demonstrated an $\alpha = .87$. The range for total scores on the PSS-14 is from 0 to 56. Stress scores below the 25th percentile (0 to 17) were interpreted as low stress, scores between the 25th and 75th percentile (18 to 28.5) were interpreted as moderate stress and scores above the 75th percentile (28.6 to 56) were interpreted as high stress. On the Stress Survey, each item's mean stress rating was

calculated. An item with an average of 5 was interpreted as causing “severe stress”, 4 “a lot of stress”, 3 “moderate stress”, 2 “a little stress” and 1 “no stress”.

- The State-Trait Anxiety Inventory (STAI): Internal consistency coefficients for the Spanish version scale have ranged from .86 to .95 (Spielberger et al., 2008). In our sample we obtained an alpha of 0.91 for STAI State and an Alpha of 0.86 for STAI trait. The STAI has 40 items with 20 items allocated to each of the subscales and it is used with a 4-point Likert format (score from 0 to 3). Range of scores for each subscale is 0–60, the higher score indicating greater anxiety. First, the State Anxiety Scale (S-Anxiety) evaluates the current state of anxiety and the Trait Anxiety Scale (T-Anxiety) evaluates relatively stable aspects of “anxiety proneness,” including general states of calmness, confidence, and security [24].

-The Coping Strategy Inventory (CSI) is a self-report questionnaire designed to assess coping thoughts and behaviors in response to a specific stressor [25]. We used the existing Spanish version (Cano-García et al., 2007). The internal consistency coefficients were between .63 and .89 [26] and between .64 and .85 in our sample for each primary subscale. The CSI has 40 items, Each item on the CSI may be scored using a 5-point Likert format (score from 0 to 4), with a range score from 0 to 160. It has 8 primary subscales that are: problem solving, self-criticism, expression of emotions, wishful thinking, social support, cognitive restructuring, problem avoidance, and social withdrawal.

Statistical data Analyses

Descriptive analysis and reliability

Data analyses were carried out using the Statistical Software Package for the Social Sciences (SPSS) version 23.0. Data were examined by calculating the means, standard deviation (\pm SD), absolute and relative frequencies and percentages, in order to generate a descriptive statistical analysis. In order to measure the internal consistency and **homogeneity** of the three questionnaires, the Cronbach's alpha test was performed, accepting a coefficient ≥ 0.70 as an ideal value. The individual analysis of each item was carried out using the homogeneity index which was assessed with the Spearman correlation coefficient. Each item with a coefficient >0.30 was considered useful for evaluating the attribute. Additionally, there were no items that did not fulfil this condition.

Correlations and regression analysis

After checking the non-normal distribution of the total scores of the scales in our sample by the Kolmogorov-Smirnov test, the relevant non parametric tests were used for the comparison of means between groups according to gender, course, clinical placement and previous training in Health Sciences.

Correlations between the scores of the different scales used were assessed using the Spearman's rho correlation coefficient. The accepted confidence interval was 95% and the significance level for all analyses was set at $p < .05$; moreover, with the significant correlation between perceived stress scale and STAI and CSI a hierarchical regression model was applied to assess the independent variables that contributed significantly to the variance in the score on the PSS. These independent variables were

entered into the regression model in 5 steps. Changes in R^2 were reported after each step of the regression model to further determine the association of the additional variables. The significance criterion of the critical F value for entry into the regression equation was set at $p < .05$ being considered significant in all tests.

Results

Descriptive results for perceived stress scale, and dimensions of anxiety and coping scale.

The mean age of participants was 20.71 ± 3.89 years (range 18-46 years). Most students were female (86.5%) and 17.7% of the students had previous training in health sciences. Up to 52.1% of students were undergoing their first clinical placement in the second year.

The mean PSS score was $22.78 (\pm 8.54)$, indicating a moderate level of stress, and the stress scores ranged from 5 to 47 out of a possible 56. In our study, most participants (47.92%) indicated a moderate level of stress. The Anxiety state score was $17.64 (\pm 9.01)$ classified as 'no problem' with a minimum score of 3 and maximum of 54 and Anxiety trait of $20.13 (\pm 8.74)$ classified as 'mild anxiety' with a minimum score of 4 and maximum of 46.

Comparative analysis between PSS, STAI and clinical placement.

Regarding type of clinical placements, no significant differences were found when comparing the mean perceived stress ($p = .352$) using ANOVA. However, significant differences were identified in relation to anxiety state ($p = .002$). When comparing clinical placements two by two, statistically significant differences were identified between Primary Care and Special Services (15.9 ± 8.75 vs 23.77 ± 11.16 , $p = .006$), Geriatrics and Special Services (16.18 ± 7.53 vs 23.77 ± 11.16 , $p = .004$) and Internal Medicine and Special Services (16.14 ± 7.75 vs 23.77 ± 11.16 , $p = .001$). 100% of the students who displayed severe anxiety in the state STAI were in specialized services.

Relationships between gender, academic year and dimension of CSI.

The CSI displays significant differences between gender for the dimensions Expression of emotion, Social support and Problem avoidance, as, in all cases, the mean of these scores was higher for the female sex. However, for the total score on the scale, despite the fact that the mean was higher in women (23.22 ± 8.55) than in men (20 ± 8.04), significant differences were not found between gender ($p = .069$) for the total scale score. Regarding the students' academic year, the dimensions that were found to be statistically significant were wishful thinking and social withdrawal. The total mean score of the test for students in their second year was 21.30 ± 8.65 and the total mean score for students in their third year was 24.40 ± 8.16 , the statistical analysis showed significance ($p = 0.009$) (Table 3).

Correlation and hierarchical regression analysis

A Spearman's rho correlation was used to investigate the relationship between total perceived stress and anxiety (state and trait) and the PSS score with the total score on the CSI and all subscales. The results displayed a significant correlation for the total on the PSS and the state STAI ($r = .463, p < .01$) and for the total PSS and the trait STAI ($r = .718, p < .01$). Regarding the stress perceived and the coping strategies, the results revealed a significant relationship between the total perceived stress and the following domains: problem solving ($r = -.452, p < .01$), self-criticism ($r = .408, p < .01$), wishful thinking ($r = .459, p < .01$), social support ($r = -.220, p < .01$), cognitive restructuring ($r = -.375, p < .01$), and social withdrawal ($r = .388, p < .01$), of the CSI. (Table 4).

Table 5 shows the hierarchical regression analysis developed in this study.

For the first step the adjustment index of the model was significant $F(1, 191) 212.186, p < .01$ and the variable anxiety trait was a significant predictor of perceived stress scale ($B = .726, t = 14.56, p < .01$).

The significant variable included at the second model were: anxiety trait ($B = .624, t = 12.32, p < .01$) and wishful thinking ($B = .266, t = 5.26, p < .01$) and the adjustment was $F(2, 191) 134.82, p < .01$.

For the third model: anxiety trait ($B = .564, t = 11.28, p < .01$), wishful thinking ($B = .263, t = 5.47, p < .01$), cognitive restructuring ($B = -.211, t = -4.53, p < .01$) and the adjustment was: $F(3, 191) 106.01, p < .01$.

For the fourth model anxiety trait ($B = .486, t = 8.98, p < .01$), wishful thinking ($B = .269, t = 5.72, p < .01$), cognitive restructuring ($B = -.215, t = -4.75, p < .01$), anxiety state ($B = .162, t = 3.30, p < .05$) and the adjustment was: $F(4, 191) 86.44, p < .01$.

For the step 5: anxiety trait ($B = .450, t = 8.15, p < .01$), wishful thinking ($B = .268, t = 5.79, p < .01$), cognitive restructuring ($B = -.133, t = -2.41, p < .05$), anxiety state ($B = .170, t = 3.52, p < .01$), problem solving ($B = -.147, t = -2.56, p < .05$) and the adjustment was: $F(5, 191) 72.53, p < .01$. The fifth model explained 66.1% of the variance in perceived stress and it has been the model with the best fit.

Discussion

This study was conducted to assess nursing students' perceived stress levels and its association with anxiety as well as the coping behaviors used to reduce the effect of stress during clinical training.

Overall, 47.92 % of the students were found to experience moderate level of perceived stress and only 25% perceived a high degree of stress. Furthermore, the correlation between perceived stress and anxiety was significant in the present study, i.e. students with high scores on perceived stress had higher scores for anxiety.

The prevalence of stress among nursing students found in the literature is variable, which could be due to the different academic programs available worldwide and the use of different scales for measuring the same [27, 28]. However, stress levels may also become affected because of different perceptions regarding stress across cultures and among different individuals. Lazarus and Folkman defined stress as

“a particular relationship between the person and the environment that is appraised by the person as taxing or exceeding his or her resources and endangering his or her well-being” [29]. Besides these academic and personal differences, according to this definition of stress, it is important to include the effects of the environment, and more concretely in our case, the different clinical placements the students visited throughout their training, on the levels of stress or anxiety presented by the student. The results of other studies suggest that mental health nurses are much more vulnerable to stress and have fewer resources to face the same (John Galvin et al., 2015). Regarding other specialized services such as intensive care or emergency care, our study suggested that the students had higher levels of anxiety and stress during this clinical placement units.

Regarding the year of studies, our findings support previous studies where the students with the most experience displayed higher levels of anxiety, whereas the most inexperienced showed lower levels of stress and anxiety [19, 20]. This may be because they feel that their teachers and other nurses expect more from them as they are more experienced and therefore more knowledgeable students, thus increasing their stress levels. However, this interpretation of data should be linked to the previously cited findings, i.e., considering the clinical placements performed by the students. For example, clinical placements in more specialized services are usually completed during the later years, whereas during the first years of study, training takes place in more general services requiring more basic competencies for care and patient responsibility. Therefore, students who have more extensive training, but who are also required to have a greater level of competencies and skills during patient care are those that are exposed to a greater level of anxiety and stress. However, as suggested by Jimenez et al.[13], it is important to exercise caution in these interpretations as, in this sense, the different training programs, despite being based on the ECTS system, should not be centered only on the number of credits that should be completed each year, but they should insist on coordinating and developing parallel competencies over time, as differences in training programs exist even within the same country.

In the current study the coping strategies most frequently used by students were problem-solving followed by social support and cognitive restructuring. According to Folkman and Lazarus, problem solving is one of the more effective ways to deal with stress as it focuses on behaviors in order to manage or alter the problem [29]. Problem solving has been found to be the most utilized coping strategy in different studies with nursing students [32–34], despite the fact that these studies have used a measurement scale for facing stress that is different to the one used in this study. In terms of the relationship between perceived stress and coping strategies, our findings indicate that among these three domains (problem solving, cognitive restructuring and social support) an inverse correlation exists, indicating that people who suffer less stress, will use these strategies more. Similarly, the positive correlation with the following domains shows how people with greater stress have more anxiety trait and state and use strategies such as wishful thinking, self-criticism, social withdrawal and problem avoidance. Results of this study showed that the greatest predictor of perceived stress were anxiety trait. As for the domains or strategies used to cope with stress, in our study, the use of certain strategies such as problem solving and cognitive restructuring, were found to be considered predictors of less stress while the use of wishful thinking appeared as a predictive factor of greater stress. In other studies, using

other coping tools, they found a positive relationship in terms of protection regarding the mental health of the student, in those students who used an optimistic strategy. In this sense, it may be that our study sample did not understand culturally speaking what wishful thinking meant or that they did not know strategies really framed in this state of optimism or illusion. On the one hand could be a protective factor in terms of mental health but that are not giving truly optimal results in terms of reducing the stress they suffer [35].

In a qualitative study by Lopez V et al., nursing students of a University in Singapore reported that talking about their negative emotions with their peers and positive reframing of their negative circumstances were the most used strategies when facing perceived stress (such strategies would be framed within the domains of social support and/or expression of emotion). However, relationships between students and their clinical educators and nurses and medical staff have been widely reported in the literature [13, 32, 36, 37], as being difficult relationships based on a lack of emotional or social support. Both teachers and mentors should be responsible for the proper implementation of coping strategies as basic tools in the skills to be acquired during their competencies in the clinic. The university faculty should not only be aware of the stress levels of students, but also consider how they manage this stress, i.e. whether they use good and effective tools for coping with the same, as this will be key in their development as a nurse. Getting to know the level of stress and/or anxiety that is experienced by our students is important to determine which negative effects should be changed in their behaviors to improve coping.

Implications for education

In this sense, and in line with other research, the study of coping strategies appears instrumental for the prevention of stress and it is essential that these strategies should begin to be trained within the university facilities. These programs could help prepare nursing students to cope with the challenges they are about to face during their clinical rotations. For example, in Spain no university uses a nurse student peer mentoring or support program. This peer mentoring program, where third year students mentor first year students, was implemented in other foreign universities in order to reduce the anxiety experienced by first year students nurses and to facilitate a smooth transition to clinical practice situations [38]. Other strategies have demonstrated to be effective in the management of stress and anxiety in nursing students, such as the use of biofeedback and mindfulness and meditation interventions [39] or emotional freedom techniques [40].

These findings call for a greater challenge to nurse educators in create a planning strategy to prevent recurrence of stress based in the use of the coping strategies most closely correlated with lower levels of perceived stress.

Limitations

This study has limitations that must be considered when interpreting these results. First, these findings cannot be generalized, as the study was conducted in nursing students of only one faculty of nursing and therefore, the socio-demographic structure of the sample was not necessarily the same as that of other

faculties in Spain. The performance of longitudinal studies conducted over several academic years is recommended as these could show changes in perceived stress over time. Further studies based on qualitative techniques would provide more detail regarding the stressor factors and its relationship with levels of anxiety and coping strategies. However, despite these limitations, the results of this study appear to concur with previous findings on this topic.

Conclusions

In light of these findings, we recommend that the teaching of positive coping strategies should be implemented in the nursing curriculum *prior* to clinical placements. Qualitative research focused on the student's perception on their clinical experience may be helpful for developing an effective clinical teaching strategy in nursing education.

Declarations

Ethics approval and consent to participate

Ethical approval for the study was obtained from the Research Ethics Committee - number C- 121. All procedures were followed in accordance with the Helsinki Declaration. Permission to conduct the study was obtained from the management of the Nursing Faculty. Before data were collected, all students were informed of the purpose of the study and informed written consent was obtained. In addition, all participants were assured that their anonymity and confidentiality would be maintained and that they were entitled to drop out of the study at any time.

Consent for Publication

This manuscript does not include any identifiable details, images, or videos relating to an individual person.

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests.

The authors declare that they have no competing interests.

Funding

This research did not receive any specific grant from funding agencies in the public commercial or not-for-profit sectors.

Author's Contribution

Study conception and design. O-Z MD., F-M E., P-F ML.,

Data collection, statistical expertise, analysis and interpretation of data. O-Z MD., F-M E., P-F ML., F-M JJ, G-S FJ., A-S A.

Manuscript preparation, supervision, administrative support and critical revision of the paper. O-Z MD., F-M E., P-F ML., G-S FJ., A-S A.

All authors read and approved the final manuscript.

Acknowledgements

The authors thank the students who took part in this study and generously granted us their time and provided us details about their experiences in clinical practice.

Abbreviations

ETCS: European Credits Transfer System.

PSS: Perceived stress scale

STAI: The State-Trait Anxiety Inventory

CSI: The Coping Strategy Inventory

References

1. Thimmapuram J, Pargament R, Sibliss K, Grim R, Risques R, Toorens E. Effect of mindfulness meditation on burnout, emotional wellness, and telomere length in health care professionals. *J Community Hosp Intern Med Perspect*. 2017;7:21–7.
2. Duffy E, Avalos G, Dowling M. Secondary traumatic stress among emergency nurses: A cross-sectional study. *Int Emerg Nurs*. 2015;23:53–8.
3. Adriaenssens J, Hamelink A, Bogaert P Van. Predictors of occupational stress and well-being in First-Line Nurse Managers: A cross-sectional survey study. *Int J Nurs Stud*. 2017;73:85–92.
4. Hamaideh SH, Al-omari H, Al-modallal H. Nursing students' perceived stress and coping behaviors in clinical training in Saudi Arabia. *J Ment Heal*. 2016;8237 February.
5. Yildiz U, Ozbas A, Cavdar I, Yildizeli S, Onler E. Assessment of nursing students' stress levels and coping strategies in operating room practice. *Nurse Educ Pract*. 2014;:1–4.
6. Zupiria Gorostidi X, Huitzi Egilegor X, Jose Alberdi Erice M, Jose Uranga Iturriotz M, Eizmendi Garate I, Barandiaran Lasa M, et al. Stress sources in nursing practice. Evolution during nursing training. *Nurse Educ Today*. 2007;27:777–87.

7. Shaban IA, Khater WA, Akhu-Zaheya LM. Undergraduate nursing students' stress sources and coping behaviours during their initial period of clinical training: A Jordanian perspective. *Nurse Educ Pract.* 2012;12:204–9.
8. Watson R, Yanhua C, Ip MYK, Smith GD, Wong TKS, Deary IJ. The structure of stress: Confirmatory factor analysis of a Chinese version of the stressors in Nursing Students Scale (SINS). *YNEDT.* 2013;33:160–5.
9. Priesack A, Alcock J. Well-being and self-efficacy in a sample of undergraduate nurse students: A small survey study. *Nurse Educ today.* 2015;35:e16–20.
10. Gibbons C. Stress, coping and burn-out in nursing students. *Int J Nurs Stud.* 2010;47:1299–309.
11. Riley JM, Collins D, Collins J. Nursing students' commitment and the mediating effect of stress. *Nurse Educ Today.* 2019;76:172–7.
12. Wang W, Xu H, Wang B, Zhu E. The mediating effects of learning motivation on the association between perceived stress and positive-deactivating academic emotions in nursing students undergoing skills training. *J Korean Acad Nurs.* 2019;49:495–504.
13. Jimenez C, Navia-Osorio PM, Diaz CV. Stress and health in novice and experienced nursing students. *J Adv Nurs.* 2010;66:442–55.
14. Admi H, Moshe-Eilon Y, Sharon D, Mann M. Nursing students' stress and satisfaction in clinical practice along different stages: A cross-sectional study. *Nurse Educ Today.* 2018;68:86–92.
15. Suen WQ, Lim S, Wang W, Kowitlawakul Y. Stressors and expectations of undergraduate nursing students during clinical practice in Singapore. *Int J Nurs Pract.* 2016;22:574–83.
16. Edwards H, Smith S, Courtney M, Finlayson K, Chapman H. The impact of clinical placement location on nursing students' competence and preparedness for practice. *Nurse Educ Today.* 2004;24:248–55.
17. Timmins F, Kaliszer M. Aspects of nurse education programmes that frequently cause stress to nursing students – fact-finding sample survey. *Nurse Educ Today.* 2002;22:203–11.
18. Evans W, Kelly B. Pre-registration diploma student nurse stress and coping measures. *Nurse Educ Today.* 2004;24:473–82.
19. Deary IJ, Watson R, Hogston R. A longitudinal cohort study of burnout and attrition in nursing students. *J Adv Nurs.* 2003;43:71–81.
20. Watson R, Deary I, Thompson D, Li G. A study of stress and burnout in nursing students in Hong Kong: a questionnaire survey. *Int J Nurs Stud.* 2008;45:1534–42.
21. Lo R. A longitudinal study of perceived level of stress, coping and self-esteem of undergraduate nursing students: an Australian case study. *J Adv Nurs.* 2002;39:119–26.
22. Labrague LJ, McEnroe-Petitte DM, Gloe D, Thomas L, Papathanasiou I V, Tsaras K. A literature review on stress and coping strategies in nursing students. *Journal of Mental Health.* 2017;26:471–80.
23. Remor E. Psychometric properties of a European Spanish version of the Perceived Stress Scale (PSS). *Span J Psychol.* 2006;9:86–93.

24. Spielberger CD, Gorsuch RL, Lushene RE. Manual for the State-Trait Anxiety Inventory. 1970.
25. Tobin D. User manual for the Coping Strategies Inventory. 2001. Unpubl Manuscr. 2001.
26. Martínez FJCGLRFJG. Adaptación española del Inventario de Estrategias de Afrontamiento. *Actas Esp Psiquiatr*. 2007;35(1):29–39.
27. Gazzaz ZJ, Baig M, Al Alhendi BSM, Al Suliman MMO, Al Alhendi AS, Al-Grad MSH, et al. Perceived stress, reasons for and sources of stress among medical students at Rabigh Medical College, King Abdulaziz University, Jeddah, Saudi Arabia. *BMC Med Educ*. 2018;18:29.
28. McCarthy B, Trace A, O'Donovan M, Brady-Nevin C, Murphy M, O'Shea M, et al. Nursing and midwifery students' stress and coping during their undergraduate education programmes: An integrative review. *Nurse Educ Today*. 2018;61:197–209.
29. Lazarus RS, Folkman S. Stress, appraisal, and coping.
30. Galvin J, Smith AP, Shen C. Stress in U.K. Mental Health Training: A Multi-dimensional Comparison Study. *Br J Educ Soc Behav Sci Orig Res Artic Galvin Smith; BJESBS*. 2015;9:161–75.
31. Tully A. Stress, sources of stress and ways of coping among psychiatric nursing students. *J Psychiatr Ment Health Nurs*. 2004;11:43–7.
32. Al-Zayyat AS, Al-Gamal E. Perceived stress and coping strategies among Jordanian nursing students during clinical practice in psychiatric/mental health courses. *Int J Ment Health Nurs*. 2014;23:326–35.
33. Chen Y-W, Hung C-H. Predictors of Taiwanese baccalaureate nursing students' physio–psycho-social responses during clinical practicum. *Nurse Educ Today*. 2014;34:73–7.
34. Al-Gamal E, Alhosain A, Alsunaye K. Stress and coping strategies among Saudi nursing students during clinical education. *Perspect Psychiatr Care*. 2018;54:198–205.
35. Karaca A, Yildirim N, Cangur S, Acikgoz F, Akkus D. Relationship between mental health of nursing students and coping, self-esteem and social support. *Nurse Educ Today*. 2019;76:44–50.
36. Galvin J, Suominen E, Morgan C, O'Connell E-J, Smith AP. Mental health nursing students' experiences of stress during training: a thematic analysis of qualitative interviews. *J Psychiatr Ment Health Nurs*. 2015;22:773–83.
37. Bagcivan G, Cinar FI, Tosun N, Korkmaz R. Determination of nursing students' expectations for faculty members and the perceived stressors during their education. *Contemp Nurse*. 2015;50:58–71.
38. Hogan R, Fox D, Barratt-See G. Peer to peer mentoring: Outcomes of third-year midwifery students mentoring first-year students. *Women and Birth*. 2017;30:206–13.
39. Ratanasiripong P, Park JF, Ratanasiripong N, Kathalae D. Stress and Anxiety Management in Nursing Students: Biofeedback and Mindfulness Meditation. *J Nurs Educ*. 2015;54:520–4.
40. Patterson SL. The effect of emotional freedom technique on stress and anxiety in nursing students: A pilot study. *Nurse Educ Today*. 2016;40:104–10.

Tables

Table 1. Means, standard deviation and skewness and kurtosis for the scores of participants in perceived stress (PSS), anxiety (STAI) and dimensions of coping (CSI).

Constructs	Dimensions	M	SD	Min	Max	Skewness	Kurtosis
Stress	Perceived stress	22.78	8.53	5	47	.303	.155
Anxiety	Anxiety state	17.63	9.01	3	54	.894	.813
	Anxiety trait	20.13	8.73	4	46	.541	-.126
Coping	Problem solving	15.08	3.43	4	20	-.450	-.072
	Self criticism	6.54	4.56	0	20	.484	-.152
	Expression of emotion	9.88	4.37	0	20	-.127	-.406
	Wishful thinking	9.45	5.15	0	20	.248	-.863
	Social support	13.9	4.61	0	20	-.708	.067
	Cognitive restructuring	12.51	3.78	2	20	-.212	-.188
	Problem avoidance	7.36	1.62	3	13	-.018	.560
	Social Withdrawal	6.28	3.70	0	19	.495	.012

N = 192; Skewness Standard Error = .175; kurtosis Standard Error = .349

Table 2. Comparison of the level of stress and anxiety classified by clinical placement type

	STAI STATE (Mean±SD)	PERCEIVED STRESS (Mean±SD)
Geriatrics	16.18± 7.53	24.35±8.65
Mental health	19.40± 10.72	25.40±6.87
Primary care	15.9 ±8.75	22.33±7.30
Internal medicine	16.14± 7.75	21.13±9.08
Specialized services	23.77± 11.16	23.64±9.00
Mother-child health	18.27 ±7.92	24.09±7.03
TOTAL	17.64±9.01	22.79±8.54

Table 3. Bivariate analysis of the mean score for the dimensions of the CSI compared by academic year and gender

	Gender	(M, SD)	U	p	Year	(M-SD)	U	p
				value				value
Problem solving	Female	15.22±3.42	1828.50	.208	2nd	15.33±3.49	4192.50	.287
	Male	14.23±3.50			3rd	14.83±3.39		
Self-criticism	Female	6.59±4.61	2051.00	.684	2nd	5.98±4.35	3952.50	.091
	Male	6.27±4.30			3rd	7.16±4.74		
Expression of emotion	Female	10.23±4.34	1432.01	.006**	2nd	10.14±4.28	4390.00	.584
	Male	7.61±3.98			3rd	9.60±4.48		
Wishful thinking	Female	9.59±5.25	1911.00	.348	2nd	8.62±5.28	3610.50	.010*
	Male	8.58±4.51			3rd	10.36±4.90		
Social support	Female	14.14±4.65	1594.00	.032*	2nd	14.01±4.65	4465.00	.725
	Male	12.38±4.15			3rd	13.78±4.60		
Cognitive restructuring	Female	12.68±3.87	1703.50	.083	2nd	12.51±3.83	4543.50	.883
	Male	11.42±3.07			3rd	12.51±3.76		
Problem avoidance	Female	7.47±1.56	1647.00	.048*	2nd	7.47±1.71	4270.00	.382
	Male	6.69±1.89			3rd	7.25±1.52		
Social withdrawal	Female	6.27±3.66	2135.50	.932	2nd	5.50±3.30	3509.00	.004*
	Male	6.34±4.03			3rd	7.13±3.94		

∴.05; ** p < .01; U= U Mann Whitney

Table 4. Spearman's correlation coefficient between perceived stress, STAI state and trait and dimension of CSI.

	1	2	3	4	5	6	7	8	9	10
1. Perceived stress										
1. Anxiety state	.463**									
1. Anxiety trait	.718**	.264**								
1. Problem solving	-.452**	.050	-.180*							
1. Self criticism	.408**	.144*	.316**	-.281**						
1. Expression of emotion	-.086	.051	-.063	.269**	.059					
1. Wishful thinking	.459**	.078	.313**	-.142	.534**	.196**				
1. Social support	-.220**	.031	-.088	.439**	-.042	.591**	.085			
1. Cognitive restructuring	-.375**	.003	-.145*	.573**	-.214**	.298**	-.071	.471**		
1. Problem avoidance	.105	.423**	.461**	.053	.168*	.106	.120	.004	-.053	
1. Social Withdrawal	.388**	.138	.289**	-.266**	.396**	-.104	.424**	-.275**	-.115	.091

.05. ** p < .01.

Table 5. Summary of Stepwise Regression Analyses to determine predictors of Perceived stress.

Independent Variables	B	SE B	β	t
Step 1				
Anxiety trait	.709	.049	.726	14.56**
Step 2				
Anxiety trait	.609	.049	.624	12.32**
Wishful thinking	.440	.084	.266	5.26**
Step 3				
Anxiety trait	.550	.049	.564	11.28**
Wishful thinking	.436	.080	.263	5.47**
Cognitive restructuring	-.475	.105	-.211	-4.53**
Step 4				
Anxiety trait	.474	.053	.486	8.98**
Wishful thinking	.445	.078	.269	5.72**
Cognitive restructuring Anxiety state	-.485	.102	-.215	-4.75**
	.153	.046	.162	3.30*
Step 5				
Anxiety trait	.439	.054	.450	8.15**
Wishful thinking	.444	.077	.268	5.79**
Cognitive restructuring Anxiety state	-.300	.124	-.133	-2.41*
Problem solving	.161	.046	.170	3.52**
	-.364	.142	-.147	-2.56*

$R^2 = .528$ for step 1; $R^2 = .588$ for step 2; $R^2 = .628$ for step 3; $R^2 = .649$ for step 4; $R^2 = .661$ for step 5. * $p < .05$; ** $p < .01$

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [STROBEchecklistcrosssectional.doc](#)
- [DATABASE.xls](#)