

*Tg*ROP18 Targets IL20RB for Host-Defense-Related-STAT3 Activation During *T. gondii* Infection

Ling Kong

Southern Medical University

Dan Jiang

Southern Medical University

Cheng He

Southern Medical University

Jing Xia

Southern Medical University

Haixia We

Southern Medical University

Lijuan Zhou

Southern Medical University

Hong-Juan Peng (✉ floriapeng@hotmail.com)

Southern Medical University

Research

Keywords: *Toxoplasma gondii*, ROP18, IL20RB, STAT3, proinflammatory immunity

Posted Date: June 17th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-16783/v3>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at *Parasites & Vectors* on August 7th, 2020.

See the published version at <https://doi.org/10.1186/s13071-020-04251-7>.

Abstract

Background: *Toxoplasma gondii* is an opportunistic protozoan infecting almost one-third of the world's population. *T. gondii* rhoptry protein 18 (*TgROP18*) is a key virulence factor determining the parasite's acute virulence and is secreted into host cells during infection. We previously identified the interaction of *TgROP18* and host cell immune-related receptor protein IL20RB, and observed the activation of STAT3 in human keratinocytes (HaCaT) cells infected by the *rop16* knockout RH strain, though *TgROP16* is regarded as being responsible for host STAT3 activation during *T. gondii* invasion. Therefore, we hypothesize *TgROP18* can activate host STAT3 through binding to IL20RB.

Methods: CRISPR-CAS9 technology was used to generate the ROP16 and ROP18 double knockout RH strain, RH- Δ rop16 Δ rop18. SDS-PAGE and Western blot were used to detect STAT3 activation in different HaCaT cells with high endogenous IL20RB expression treated with *T. gondii* tachyzoites infection, recombinant ROP18, or IL-20. FRET and co-immunoprecipitation (Co-IP) was used to detect the protein-protein interaction.

Results: We observed that *TgROP18* was involved in a synergic activation of the host JAK/STAT3 pathway together with *TgROP16* in human HaCaT cells infected with *T. gondii* or treated with recombinant *TgROP18* protein, stimulating host proinflammatory immune responses such as expression of TNF- α . The effect of recombinant ROP18 on STAT3 phosphorylation was presented in a dose-dependent manner. Additionally, *TgROP18* was identified to target IL20RB on its extracellular domain. When we treated different cell lines with the recombinant ROP18, STAT3 phosphorylation could only be observed in the cells with endogenous IL20RB expression, such as HaCaT cells.

Conclusions: These findings indicate that *TgROP18*-IL20RB interaction upon *T. gondii* invasion was involved in STAT3 activation, which is associated with host cell defense.

1. Introduction

Toxoplasma gondii is an obligate intracellular parasite of the phylum apicomplexan, which resides in the nucleated cells of warm-blooded vertebrates (1, 2). It is reported that approximately one-third of the world's population are chronically infected with *T. gondii* (3, 4). *T. gondii* infection in immunocompetent individuals is usually asymptomatic, whereas, in immunosuppressed individuals or in primary infection during pregnancy, it may cause severe toxoplasmosis and even death (5-7).

The apical complex of the apicomplexan organisms, consisting of conoid and apical organelles, is responsible for host cell penetration and the establishment of intracellular parasitism. The apical organelles, including rhoptries, micronemes and dense granules, are characteristic secretory vesicles which discharge proteins involved in the cell invasion(8). Host cell invasion by *T. gondii* is a rapid but complicated process (9), including cell membrane attachment, protein secretion, moving junction formation and inclusion (10-12). Host cell activities, including innate and adaptive immune responses, nutrient metabolism, cytoskeletal rearrangement, and apoptosis, are manipulated during *Toxoplasma*

tachyzoite invasion (13-16). During invasion, many of the effectors are injected into host cells by *T. gondii*, including rhoptry proteins (ROPs), dense granule proteins (GRAs), and microneme proteins (MIC) (17, 18). These injected ROPs, such as ROP16 and ROP18, have been known to activate host transcription factors, such as signal transducer and activator of transcription (STAT), to inhibit host immune responses (19-21). Moreover, previous study has shown that a large number of rhoptry proteins are injected into cells that are not infected, and result in transcription factor activation comparable to that of infected cells (18). Meanwhile, *T. gondii* modulates host immune responses to maintain a balance between pro-inflammatory and anti-inflammatory reactions to allow its successful parasitism, and thereby to establish a long-term (chronic) infection(22).

TgROP18 is a serine/threonine kinase secreted from the rhoptry during parasite invasion, and is known as a critical contributor to the virulence of *T. gondii* (23). Detectable *TgROP18* is reported to be only secreted by type I and type II strains, but not by type III strains. *TgROP18* can suppress the differentiation of host neural stem cells(24), inhibit the host NF- κ B pathway by promoting p65 degradation in HEK 293T cells (25), and target phosphorylation of IFN-inducible small GTPases (IRGs) in murine cells or ATF6 β degradation in human cells for disarming the host's innate and adaptive immunity (26, 27).

IL20RB is an orphan receptor among the class II cytokine receptors, serving as a common β receptor chain in both type I and type II IL-20 receptors (IL-20R) (28). Type I IL-20R is a heterodimeric receptor that mediates IL-19, IL-20 and IL-24 signaling, and is composed of two subunits, IL20RA and IL20RB; type II IL-20R mediates IL-20 and IL-24 signaling and is composed of IL22RA1 and IL20RB (29). Both types of receptors are expressed in various non-immune tissues, especially in epithelial cells (30). Upon activation, IL20RB phosphorylates JAK1 and activates the JAK/STAT3 pathway; phosphorylated STAT3 then forms homodimers and translocate to the nucleus to initiate transcription of downstream proinflammatory cytokines, including tumor necrotic factor - α (TNF- α) and Monocyte Chemoattractant Protein 1 (MCP-1), by binding to STAT-responsive promoter elements (31, 32). It is well documented that STAT3 is the dominant transcription factor activated by both type I and type II IL20 receptors(31).

It is generally recognized that *TgROP16* is dominantly responsible for host STAT3 activation during *T. gondii* invasion(33, 34), and no study has been conducted to show if ROP18 functions on STAT3 activation. In our study, we were surprised to observe the activation of STAT3 in human keratinocyte (HaCaT) cells infected by the *rop16* knockout RH strain(RH- Δ *rop16*). Combined with the finding of *TgROP18*-human IL20RB interaction in our previous research (35), we hypothesized that *TgROP18* may activate STAT3 through binding with IL-20R to activate the JAK/STAT3 pathway. The HaCaT cell is a type of epithelial cell expressing both type I and type II IL-20 receptors in high levels, therefore, it was used as a cell model in this research.

2. Materials And Methods

2.1 Cell culture and *T. gondii* propagation

Human foreskin fibroblasts (HFF), African green monkey kidney cells transformed by SV40 (COS-7), human embryonic kidney 293T (HEK 293T) cells, and human keratinocytes (HaCaT) cells were bought from American Type Culture Collection (ATCC, USA), and cultured in Dulbecco's Modified Eagles Medium (DMEM, Gibco/Invitrogen, USA) containing 10% (v/v) FBS (Gibco/Invitrogen, USA) and 1% gentamicin (10mg/ml, Invitrogen, USA) at 5% CO₂ and 37°C. For fluorescence resonance energy transfer (FRET) experiments, COS-7 cells were grown on coverslips in 12-well plates (Corning, USA). For pSTAT3 detection, HaCaT or COS-7 cells were seeded on 6-well plates and grown to 100% confluence, and then cultured in serum-free medium overnight before treatment with IL-20 (R&D Systems), recombinant GST-ROP18, or *T. gondii* infection.

The tachyzoites of CEP, CEP-*TgROP18I*, and CEP-*TgROP18II* were a generous gift from Jon Boyle at the University of Pittsburgh. The tachyzoites of *T. gondii* RH strain, *rop16* knockout strain (RH- Δ *rop16*), *rop18* knockout strain (RH- Δ *rop18*), and *rop16 + rop18* double knockout strain (RH- Δ *rop16* Δ *rop18*) were propagated in HFFs with DMEM supplemented with 1% FBS, as well as the CEP and its mutant strains. Before experiments, the tachyzoites cultured in HFFs were purified by passing through a 3 μ m filter after the host cells were ruptured or syringed broken, and then counted.

2.2 Mammalian eukaryotic expression

The cDNA of IL20RB on pDONR was generously gifted by Prof. Wen-Bin Ma at the School of Life Science, Sun Yat-Sen University. The cDNA of IL20RB was amplified using PCR with *Pfu* DNA polymerase (TransGen Biotech, China), to add the HA tag on its C-terminus and incorporate *SaI*I and *SacI*I restriction sites on the 5'- and 3'- ends, respectively. To perform FRET and co-immunoprecipitation (Co-IP) assays, IL20RB-HA, and its extracellular and intracellular domain was then cloned into pEYFP-C1 to form pEYFPC1-IL20RB-HA, pEYFPC1-IL20RB-Extr-HA (Extr, extracellular domain of IL20RB), and pEYFPC1-IL20RB-Cyt-HA (Cyt, intracellular domain of IL20RB). The construction of pCFPN1-ROP18-3 \times FLAG has been described in detail elsewhere (16). All the primers used are shown in Supplementary Data (Table S1).

2.3 CRISPR-CAS9 mediated generation of RH- Δ *rop16* Δ *rop18* double knockout strain

RH- Δ *rop16* was constructed in our previous research. To disrupt the *rop18* gene in the RH- Δ *rop16* strain, a recombinant CRISPR plasmid, pSAG1::CAS9-U6::sgROP18, which contains an sgRNA specifically targeting the site very close to and downstream from the start codon of *rop18* gene, was constructed, using a Q5[®] Site-Directed mutagenesis kit (NEB, USA) following the published protocol(36). A chloramphenicol-resistance gene (chloramphenicol acetyltransferase, CAT) was used as the screening marker. The homologous recombination plasmid (pBlue-5'-3'-ROP18-homo-CAT) was constructed as follows (the procedure is also shown in Figure S1A); A 991 bp fragment and a 965 bp fragment homologous to the upstream and downstream of the gRNA targeted site of the *rop18* gene, were respectively subcloned to the 5' and 3'- end of the CAT gene cassette on pBlue-script SK II (-) plasmid (16). Subsequently, these two recombinant plasmids were co-transfected into RH- Δ *rop16* tachyzoites by

electroporation. The transfected tachyzoites were used to infect HFF cells, and cultured in chloramphenicol free DEMEM complete medium for 48 h. To screen the recombinants, the tachyzoites were cultured in the DEMEM complete medium containing 20 mmol/L of chloramphenicol for 4 passages. Finally, the double-knockout RH- $\Delta rop16\Delta rop18$ tachyzoites were screened through monoclonal screening, and identified by PCR. All the PCR primers used for homologous template amplification, plasmid construction, and recombinant identification are shown in Table S1.

2.4 Detection of STAT3 activation in different *T. gondii* tachyzoites infected or IL-20 treated HaCaT cells

2.4.1 Cell preparation

- A. HaCaT cell infection with RH-WT, RH- $\Delta rop16$, RH- $\Delta rop18$, and RH- $\Delta rop16\Delta rop18$ strains: HaCaT cells were cultured in 6-well plates to 100% confluence. After the cells were washed 3 times with PBS and switched to serum-free medium and starved overnight, RH-WT, RH- $\Delta rop16$, RH- $\Delta rop18$, and RH- $\Delta rop16\Delta rop18$ strains were used to infect the cells by one well for each, at multiplicity of infection (MOI) 10 for 30min; another well was not infected for control.
- B. HaCaT cell treatment with IL20, or infection with RH-WT and RH- $\Delta rop18$ strains: HaCaT cells were cultured in 6-well plates to 100% confluence. After the cells were starved overnight, two wells of cells were treated with 400 ng of IL-20 for 30 min and 24 h, respectively; two wells were infected with RH-WT at MOI 10 for 30min and 24h, respectively, and two wells were infected with RH- $\Delta rop18$ under the same conditions; another well was not infected for control.
- C. HaCaT cell treatment with recombinant GST-ROP18 and GST at different amount: The construction of the prokaryotic expression vector and expression of GST-ROP18 and GST have been described in detail previously(16). The *Escherichia coli* expressed GST-ROP18 and GST proteins were purified using the GST fusion protein purification kit (GenScript Inc., China) following the manufacturer's instructions. The concentration of the purified protein was detected with NanoDrop 2000 (Thermo Scientific, USA). HaCaT cells were cultured in 6-well plates to 100% confluence. The cells were starved overnight, and treated with 0.3, 0.6, 1.2, and 2.4mg of GST and GST-ROP18, separately, one well for each treatment. Another well was treated with 10 ul of elution buffer (20mM glutathione, 50mM Tris-HCl, pH 8.0) for control.
- D. HaCaT cell treatment with IL-20, GST-ROP18 and GST, and detection of the co-localization of ROP18 and IL20RB on HaCaT cell membrane

COS-7, HaCaT and HFF cells were grown in 6-well plates to 100% confluence. The serum starved cells were treated with IL-20 (200ng), recombinant GST (2 μ g), or recombinant GST-ROP18 (2 μ g) for 30min, and elution buffer (50mM Tris-HCl 8.0 with 20mM glutathione, pH 8.0) as a negative control, one well for each treatment.

1. HaCaT cell infection with type III CEP, CEP-*TgROP18I*, and CEP-*TgROP18II* strains

HaCaT cells were grown in 6-well plates to 100% confluence, cells were starved overnight and infected with CEP, CEP-*TgROP18I*, and CEP-*TgROP18II* at MOI 10 for 30min, one well for each infection. Another well was uninfected for control.

2.4.2 Detection of STAT3 activation

The culture medium was aspirated, the cells were then washed with PBS three times, and the unrecruited parasites were washed off. The cells were harvested and lysed. Fifty milligrams of total proteins were subjected to SDS-PAGE and Western blot to detect activation of STAT3 (phosphorylation at Y705 or S727). β -actin was detected as the loading control for the whole cell lysate. Densitometric quantitation of each band in A was applied using Image J. The optical density of Stat3-pY705 and Stat3-pS727 was normalized to the density of the total STAT3, and divided by the uninfected group whose normalized value was set as 1.

2.4.3 Statistical analysis

All experiments were performed at least in triplicate. Data are presented as means \pm SD of three independent experiments. Statistical significance was determined using the Kruskal–Wallis H-test with Bonferroni correction or independent t-test. Values of $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$ were defined as statistically significant.

2.5 Western blot analysis and co-immunoprecipitation

COS-7 cells transfected with recombinant plasmid or HaCaT cells infected with parasites were lysed in lysis buffer (Beyotime Biotechnology) containing protease inhibitor cocktail (Thermo Fisher Scientific) or/and phosphatase inhibitors cocktail (TransGen Biotech, China). The cell lysates were centrifuged at $14,000\times g$ for 5 minutes at 4 °C. The supernatants (50mg of total proteins) were separated by SDS-PAGE and then transferred to PVDF membranes using Trans-Blot SD Semi-Dry Electrophoretic Transfer (Bio-Rad, USA). Then, the PVDF membranes were blocked in PBS or TBS containing 5% Bovine Serum Albumin (BSA, [Sigma-Aldrich](#), German) and 0.05% Tween-20. After blocking, the PVDF membrane was incubated with primary antibodies overnight at 4 °C. Then, membranes were probed with secondary antibodies conjugated with horseradish peroxidase (HRP) for 2 h. The proteins of interest were visualized by luminescence generated using ClarityTM Western ECL Substrate (Bio-Rad, USA) and imaged with ChemiDoc Touch Imaging System (BIO-RAD, USA).

For co-immunoprecipitation (Co-IP), the supernatants of cell lysates were incubated with 4 μ g of anti-HA tag antibody at 4°C for 1 h. Then, Protein A - Agarose (Santa Cruz, USA) was added to the mix and incubated for 12 h with gentle rotation at 4 °C. The agarose beads were collected by centrifugation for 5 min at $1,000\times g$ at 4°C and washed four times with PBS. Then the immunoprecipitates were resuspended in SDS-PAGE sample buffer (Takara, Japan) and boiled for 10min followed by soaking immediately in ice water. The boiled samples were centrifuged at high speed, the supernatant was then loaded for SDS-PAGE, and then subjected to Western blot analysis with the antibodies described below.

2.6 Fluorescence Resonance Energy Transfer (FRET) experiment

COS-7 cells were seeded on coverslips in a 12-well plate, and grown to 80% confluence. The experimental group was co-transfected with pCFPN1-ROP18-3×FLAG and pEYFPC1-IL20RB-HA for 48h. The mono-fluorescence groups were transfected with pCFPN1-ROP18-3×FLAG or pEYFPC1-IL20RB-HA separately for adjustment of the fluorescence base line of donor (eCFP) and acceptor (eYFP). The negative control group was co-transfected with pECFPN1 and pEYFPC1, and the positive control group was transfected with pEYFP-ECFP. The co-localization of ECFP-ROP18 with EYFP-IL20RB was observed under a FluoView FV1000 confocal microscope (Olympus, Japan), and the FRET efficiency and intermolecular distance were calculated by sensitized emission (SE) program. The detection was repeated in different cells three times. The difference between groups was analyzed with the t-test using SPSS software (** $p < 0.01$).

2.7 Reverse Transcription PCR (RT-PCR) and Quantitative Reverse Transcription PCR (qRT-PCR)

The cells were starved overnight before infection and detection. The transcription level of IL20RB in COS-7, HEK293T, HFF, and HaCaT cells was evaluated by RT-PCR. The transcription levels of IL20RB, IL22RA1, and TNF- α in infected cells were analyzed by qRT-PCR. The normal HaCaT cells, and the cells infected with RH-WT or RH- Δ rop18 at MOI 10 for 10, 20 and 30 min, were harvested separately. Transcription level of iNOS, IL20, IL22, and IL10 was detected by qRT-PCR. All the PCR primers are shown in Table S1. The translation level of IL10 was detected by Western blot.

Total RNA isolation from infected or uninfected cells was performed using Trizol[®] Reagent (Ambion[™], USA), and genomic DNAs were immediately removed by TURBO DNAase (Invitrogen, USA). cDNAs were generated with the RevertAid First Strand cDNA Synthesis Kit (Thermo Fisher Scientific, USA). PCR was conducted using *Pfu* DNA polymerase (TransGen Biotech, China), and qRT-PCR was performed using SuperReal PreMix Plus (Tiangen Biotech, China). All of the procedures were performed following the manufacturer's instructions. Gene-specific primers are shown in Table S1. Each experiment was performed in triplicate and the reactions were conducted in a BioRad Applied Biosystems 7500 qPCR System (BioRad laboratories, Inc, USA). The relative transcription level of each gene was normalized to the mRNA of an internal control gene, GAPDH. The transcription levels of the target genes were compared by the change-in-cycling-threshold ($\Delta\Delta$ CT) method.

2.8 Detection of the co-localization of ROP18 and IL20RB on HaCaT cell membrane

HaCaT cells were grown on coverslips in 12-well plates to 100% confluence. The overnight serum starved cells were treated with recombinant GST (1 μ g), or recombinant GST-ROP18 (1 μ g) for 30min, and then fixed in 300 μ l 4% paraformaldehyde. The cells were then permeabilized with 0.5% Triton X-100/PBS for 10 min, then washed with PBS three times, 5 min for each. The cells were blocked with 500 μ l 10% BSA for 1 h at room temperature, followed by incubation with Rabbit anti-IL20RB and Mouse anti-GST monoclonal antibody in 10% BSA overnight at 4 °C. After washing with PBS three times, 5 min for each, the cells were incubated with the secondary antibodies Alexa Fluor 488 Goat Anti-Rabbit IgG and Alexa Fluor 594 Goat Anti-Mouse IgG in 10% BSA for 1 h at room temperature in the dark. The cells were

washed with PBS for three times, 5 min for each. The coverslips were mounted with DAPI mounting oil and observed under a fluorescence microscope (Nikon eclipse Ni) with a 100× objective lens (NA 1.40).

2.9 Antibodies:

The primary antibodies for Western blot: Mouse poly-clonal Ab anti-*TgROP18* (1:1000) were raised in our laboratory; mouse mAb anti-*TgSAG1* (1:1000), rabbit anti-HA tag (1:5000), rabbit mAb anti-stat3 (phosphor Y705, 1:200,000), rabbit mAb anti-stat3 (phosphor S727, 1:5000), and rat mAb anti-IL20RB (1:1000) were purchased from ABcam Pharmatech, Inc, United Kingdom. Mouse mAb anti-GST (1:2000), mouse mAb anti-DDDDk (1:2000), and rabbit polyclonal Ab anti-TNF- α (1:1000) were purchased from ABclonal Biotechnology, USA. Mouse mAb anti-stat3 (1:1000), and rabbit mAb anti-b-actin (1:1000) were purchased from [Cell Signaling Technology](#), USA. Rabbit anti IL-10 monoclonal antibody (1:1000) was purchased from Affbiotech Biosciences, UK.

The secondary antibodies for Western blot: Goat anti-mouse IgG-HRP (1:2000) and goat anti-rabbit IgG-HRP (1:2000) were purchased from Santa Cruz [Biotechnology](#), USA. Goat anti-rat IgG-HRP (1:2000) was purchased from Bioss Inc (China). Alexa Fluor 488 Goat anti-Rabbit IgG (1:2000) was purchased from Invitrogen, and Alexa Fluor 594 Goat anti-Mouse IgG (1:2000) was purchased from Thermo Fisher Scientific.

The antibody for IP or Co-IP: Rabbit anti-HA tag (4 μ g) was purchased from [Cell Signaling Technology](#), USA.

3. Results

3.1 CRISPR/CAS9 mediated generation of RH- Δ rop16 Δ rop18 strain

The CRISPR-Cas9 system was adopted to disrupt the *rop18* gene in the RH- Δ rop16 strain to generate a double-knockout strain RH- Δ rop16 Δ rop18. A guide RNA (gRNA) and CAS9 expression plasmid pSAG1::CAS9-U6::sgROP18, and the pBlue-5'-3'-ROP18-homo-CAT plasmid containing the CAT reading frame flanking the regions homologous to the *rop18* gene were successfully constructed (Figure S1a). Tachyzoites of RH- Δ rop16 strain were co-transfected with these two plasmids. The recombinant tachyzoites were screened with chloramphenicol, and monoclonal recombinant was obtained through limiting dilution. Seven single clones were obtained and verified with PCR by confirming the disruption of *rop18* and integration of the CAT marker (Figure S1b). No. 1–3 double-knockout RH- Δ rop16 Δ rop18 strain single clones were selected and further confirmed by Western blotting (WB) (Figure S1c).

3.2 Detection of STAT3 activation in RH- Δ rop16 infected HaCaT cells

It is reported that ROP16 is the dominant factor activating STAT3 during *T. gondii* infection; a defective STAT3 activation has been observed in the macrophages infected by the RH-*rop16* KO strain (33). Based on this information, we next tried the other possible parasite kinases on the other cell types. HaCaT cells were infected with RH-WT, RH- Δ rop16, RH- Δ rop18 strains, and two RH- Δ rop16 Δ rop18 mono-clones at MOI

10 for 30 min, and were subjected to assessment of STAT3 activation (phosphorylation at Y705 and S727). To our surprise, a significant STAT3-Y705 phosphorylation was observed in HaCaT cells when infected with either RH- $\Delta rop16$ or RH- $\Delta rop18$ when compared with the uninfected cells or RH- $\Delta rop16\Delta rop18$ infected cells, but phosphorylation was still significantly weaker than the RH-WT infected cells (Figure 1). These results implied that host cell STAT3 could be activated by *TgROP18* in addition to *TgROP16*. However, STAT3-S727 phosphorylation which has been reported to facilitate the full activation of STAT3 (37), was not significantly different among all the infected or uninfected groups (Figure 1).

3.3 STAT3 activation induced by recombinant TgROP18 added to HaCaT cells culture in a dose-dependent manner

We further infected HaCaT cells with RH-WT and RH- $\Delta rop18$ strains for 30min and 24h, respectively. Four hundred ng of IL-20 (a ligand of IL20R) was used as a positive control. Then, the STAT3 phosphorylation at Y705 and S727 was detected in the total protein extracted from the different groups of cells. To our surprise, a higher level of STAT3-Y705 phosphorylation was revealed in IL-20 treatment and RH-WT infected groups than in the RH- $\Delta rop18$ infected group at 30min post infection, and no STAT3-Y705 phosphorylation was observed in the control group (untreated cells). Although weak STAT3-Y705 phosphorylation was still observed in these treatment groups at 24 h post infection, the STAT3-Y705 phosphorylation observed in the RH-WT infected group was stronger than that observed in IL-20 treatment or RH- $\Delta rop18$ infection group (Figure 2 a&b). Furthermore, we found that *TgROP18* did not affect the expression of total STAT3 and STAT3-S727 phosphorylation (Figure 2 a&b). These results demonstrated that ROP18 significantly phosphorylated STAT3 at Y705 but not at S727 during *Toxoplasma gondii* infection.

However, previous study elaborated that STAT3 could be activated by overexpressed (intracellular) *TgROP16* but not overexpressed (intracellular) *TgROP18* (33). Given this fact, we speculated that the phosphorylation of STAT3 was induced by extracellular *TgROP18*. Hence, the recombinant GST-ROP18 expressed by *E. coli* was added to the cell culture medium to determine the effect of *TgROP18* on STAT3 activation. The phosphorylation of STAT3 was detected in HaCaT cells which had been incubated with different concentrations of recombinant GST-ROP18 for 30 min. Intriguingly, we found that the STAT3-Y705 phosphorylation was distinctly induced by GST-*TgROP18* in a dose-dependent manner, but the STAT3 expression was not affected by this recombinant protein added to the cell culture (Figure 2 a&d). This result suggested that the STAT3 activation could be mediated by extracellular *TgROP18* without changing the total STAT3 level. To further confirm this phenomenon, HaCaT cells were infected with RH-WT and RH- $\Delta rop18$ tachyzoites at an MOI of 2, 4, 6, 8, and 10 for 30 min, separately. We found that RH-WT induced a much higher level of STAT3-Y705 phosphorylation than RH- $\Delta rop18$ at the same MOI index, and the phosphorylation levels resulted by both of these two strains increased with the MOI index from 2-8 (at MOI 2, 4, 6, 8), before stabilizing between MOI 8 and 10 (Figure 2 e &f).

3.4 TgROP18 interacts with host cell IL20RB at the extracellular domain CFPN1-ROP18

In our previous study, a Bi-molecular fluorescence complementation (BiFc) protein interaction screening was performed to identify the human interactome of *TgROP18*, and the human IL20RB was screened as an interaction protein of *TgROP18* (35). Fluorescence resonance energy transfer (FRET) was conducted to further confirm this interaction. The results indicated that the FRET efficiency and intermolecular distance of CFP-*TgROP18* and IL20RB-YFP (in the pCFPN1-ROP18/ pYFPC1-IL20RB co-transfection group) was not significantly different from that of CFP-YFP (in the positive control group), but was significantly different from that of CFP and YFP (in the negative control group) (Figure 3a and b). This interaction was also verified by Co-IP assay in vitro. FLAG tagged ROP18 and HA tagged IL20RB were detected in the immunoprecipitates with anti-HA antibody, but not in those precipitated with normal IgG (Figure 3c). Because IL20RB is known as a transmembrane glycoprotein, to further investigate which domain of IL20RB is binding with *TgROP18*, Co-IP assay was performed to detect the interaction of *TgROP18* with the extracellular/intracellular domain of IL20RB. The WB results demonstrated that *TgROP18* could interact with the extracellular domain of IL20RB, but not its intracellular (Cyt) domain (Figure 4).

All these data indicated that host cell IL20R could be a receptor of *TgROP18* to activate STAT3 through targeting the extracellular domain of its beta chain IL20RB.

3.5 STAT3 activated by ROP18 in cells expressing IL20RB

Since *TgROP18* - IL20RB interaction occurred in the extracellular domain of IL20RB, we speculated that the IL20RB receptor on the host cell membrane was responsible for the STAT3 activation induced by ROP18, thus, the expression of IL20RB in several types of cells was examined. The mRNAs of IL20RA and IL22RA1, which comprise the alpha chain of IL20R and are required for complete IL20R formation and indispensable for inducing downstream signaling events, were also detected. Among COS-7, HEK293T, HFF, and HaCaT cells, IL20RB was found to be transcribed in higher levels in HaCaT cells, as well as the other components to form the IL20R receptor (Figure S2). We further detected the IL20RB protein in HFF and HaCaT cells with WB, and found IL20RB was highly expressed in HaCaT cells, instead of HFF cells (Figure 5a). Considering IL20RB is a transmembrane glycoprotein, and the binding site with *TgROP18* is the extracellular domain, it possibly mediated the signal transduction of *TgROP18* during invasion when the parasite or the secreted *TgROP18* contacted the cell membrane. To test this, COS-7, HFF and HaCaT cells were selected and treated with recombinant GST-ROP18 for 30 min, with the cells treated with IL-20 as a positive control, and the untreated cells and the cells treated with recombinant GST as negative controls. Our results indicated that STAT3-Y705 phosphorylation was significantly induced by the recombinant ROP18 and IL-20 in HaCaT cells, but not in COS-7 and HFF cells (Figure 5 b&c). Therefore, we concluded that *TgROP18* mediated STAT3 activation in the cells expressing a high level of IL20RB such as HaCaT cells. We further examined the transcription and expression of *tnf-a* and *iNOS*, a downstream gene of the IL20R-JAK/STAT3 pathway regulated by STAT3 activation, using HaCaT cells uninfected or infected with RH-WT or RH- Δ *rop18*. The most significant *tnf-a* and *iNOS* transcription; and *tnf-a* translation was observed at 20 min post infection in both RH-WT and RH- Δ *rop18* infection groups, but RH-WT resulted in a significantly higher level than RH- Δ *rop18* (Figure 5 d-f, Figure S4a). As STAT3 is

the main transcription factor activated by IL-20 receptors and can be activated by IL10 family members, such as IL-10, IL-20, and IL-24, in HaCaT cells (30, 31), we further confirmed that these IL10 family members were not produced by HaCaT cells after *T. gondii* infection (Figure S4 b-e). However, a significant STAT3 activation in HaCaT cells, by the recombinant ROP18 and endogenous ROP18 secreted by *T. gondii* was observed in our study (Figure 1-2, 5-6).

Meanwhile, using qRT-PCR, we also detected the transcription of *il20rb* and one of its alpha chain *il20ra1* in HaCaT cells after infection by RH-WT and RH- Δ *rop18* at 5, 10, 20, and 30 min post infection. Using WB, the translation of *il20rb* in HaCaT cells after treatment with 400ng of IL-20 and infection by RH-WT or RH- Δ *rop18* was detected at 30 min post infection. Our results indicated that both the transcription and translation of IL20RB was not affected by *TgROP18* until after 30 min post infection (Figure S3).

3.6 Both type I and type II *TgROP18* activate host cell STAT3

To explore whether type II ROP18 (*TgROP18II*) can activate host cell STAT3 in the same way as *TgROP18I*, HaCaT cells were infected with the type III strain CEP, and the CEP expressing type I ROP18 (CEP-*TgROP18I*); and type II ROP18 (CEP-*TgROP18II*), for 30 min. The cells were harvested and total protein was extracted followed by detection of STAT3-Y705 phosphorylation. Compared to the CEP (type III) strain, both CEP-*TgROP18I*, and CEP-*TgROP18II* infection showed significant STAT3 activation in HaCaT cells, but no significant difference existed between CEP-*TgROP18I*, and CEP-*TgROP18II* infection (Figure 6 This result indicated that both type I and type II ROP18 could similarly activate STAT3 in HaCaT cells.

4. Discussion

In this study, we demonstrated that host cell STAT3 could be activated by *T. gondii* virulence factor ROP18 (of both type I and Type II stains) to regulate host innate immune responses in HaCaT cells during invasion. The recombinant *TgROP18* and the native *TgROP18* secreted by parasites bound to IL20RB on the cell membrane to activate STAT3 in HaCaT cells. *TgROP18* interacted with host IL20RB and the binding site was on the IL20RB extracellular domain. In addition, *TgROP18* only mediated STAT3 activation in the cells expressing IL20R with a high level (such as HaCaT cells), and ultimately resulted in a promotion of TNF- α and iNOS transcription (Figure7 and Figure S4 a), which represented the activation of STAT3 .

We therefore deduced the whole process as follows. Upon *T. gondii* infection, it first attaches firmly to the cell membrane. After attaching, the host cell manipulation by *T. gondii* initiates less than a minute (38). During this period, *TgROP18* secreted from *T. gondii* rhoptries bind to the host cell IL20RB, which was only expressed in some cell types on the membrane, to induce downstream signaling events. The interaction of ROP18 and IL20RB was further confirmed by FRET and Co-IP. It had been reported that overexpressed ROP18 had no effect on STAT3 activation (19, 33). Therefore, *TgROP18* may function through binding with IL20RB on the cell membrane to activate STAT3 and JAK. This phenomenon was observed in HaCaT cells with high IL20RB expression, but not in COS-7 and HFF cells with very low

IL20RB expression. HaCaT expressed both type I and type II IL-20 receptors with high levels and showed significant STAT3 activation when treated with recombinant GST-ROP18.

It is generally believed that JAK/STAT3 pathway activation is typically presented with STAT3-Y705 phosphorylation, the phosphorylated STAT3 then form homodimers and translocate to the cell nucleus where they induce transcription of target genes such as TNF- α and iNOS (39). Therefore, STAT3-Y705 phosphorylation was used to evaluate the activation of the JAK/STAT3 pathway. Though the host cell STAT3 activation by overexpressed ROP16 but not overexpressed ROP18 is well clarified (33), we still observed that the *Tg*ROP16 defective RH strain showed a significant but relatively lower STAT3 activation compared to the wild type RH strain. Combined with the finding that GST-ROP18 incubated cells showed significant STAT3 activation compared to the GST incubated cells, we can deduce that STAT3 activation by *Tg*ROP18 was due to the binding of *Tg*ROP18 on the extracellular domain of IL20RB. These results also explained why the STAT3 activation mediated by *Tg*ROP18 was considerably more distinct in HaCaT cells at 20-30min post infection, earlier than the point at which most tachyzoites were attaching to, but not yet entering the cells. Though it has been reported that IL20RB can induce not only STAT3, but also STAT1 and STAT5, STAT1 and STAT5 activation needs a much higher concentration of IL-19, IL-20, or IL-24 than STAT3 does (20, 31-34). Moreover, STAT3 is the main transcription factor activated by type I and type II IL-20 receptors and has been verified to be activated by IL-19, IL-20, and IL-24 in HaCaT cells (30, 31). Therefore, we further confirmed that these IL10 family members were not produced by HaCaT cells after *T. gondii* infection (Figure S4 b-e). However, a significant activation effect of ROP18 on STAT3 through the IL20RB during *T. gondii* infection in HaCaT cells was observed in our study. On the other hand, examination of whether STAT1 and STAT5 signaling can be induced by ROP18 through IL20R receptor complex, and its potential functions, requires further research.

In this study, we could not exclude the possibility of IL19, IL-20 or IL-24 binding with IL-20R other than ROP18 *in vivo*. Considering the interaction of *Tg*ROP18 with IL20RB occurred at an early infection stage, and the major cytokines strongly induced by *T. gondii* at this stage are type I cytokines, and not IL-20 subfamily cytokines (22, 40-43), therefore, the existence of a competition between ROP18 and the IL-20 subfamily working on IL-20R is not possible. It is possible that *Tg*ROP18 functions on IL-20R at the early stage of infection, while IL-20 subfamily cytokines work on IL-20R at the relatively later stage of infection.

The activation of the JAK/STAT3 pathway enhances the expression of its downstream proinflammatory factors such as TNF- α , which are closely related to early inflammatory initiation and immune responses. In previous study, TNF- α was found to be up-regulated in *T. gondii* infection (44). Furthermore, TNF- α is not only an activator inducing higher levels of IFN- γ (44) but also an important activator of the NF- κ B pathway (45); both play an important role in anti-*T. gondii* infection. In previous studies, ROP18 mainly exerts their immunosuppressive action in *T. gondii* infection to promote their survival (13, 25, 27, 46). In our study, we revealed that *Tg*ROP18 activated the IL20R-JAK/STAT3 pathway for anti-infection against *T. gondii*, and mainly played a role in early infection. Once the invasion is accomplished, other signaling pathways play important roles in anti-*T. gondii* infection will be soon triggered by ROP18, such as NF- κ B signaling (25, 47). The recent research has clarified that *T. gondii* inhibits host immunity through ROP16-

mediated activation of STAT3(48), which may explain why the *TgROP18* activated IL20R-JAK/STAT3 pathway only caused transiently enhanced production of TNF- α during early infection, since the anti-infection effect was covered by ROP16.

Ultimately, similar STAT3 activation results were observed in CEP-*TgROP18I* and CEP-*TgROP18II* infected HaCaT cells at 30min post infection, which suggested that both type I and type II ROP18 could activate STAT3 in HaCaT cells at the early stage of infection. However, whether they were involved in the same mechanism still needs to be confirmed.

5. Conclusions

This study determined that the immune-related receptor protein IL20RB was targeted by *TgROP18* on its extracellular domain upon *T. gondii* invasion, and led to synergic activation of the host JAK/STAT3 pathway together with *TgROP16*, triggering proinflammatory immune responses against *T. gondii*. Herein, we elucidated a novel molecular mechanism of manipulation of *TgROP18* on host immune responses and provided new insights into the *T. gondii*-host interaction.

Declarations

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported by National Key R&D Program of China (2017YFD0500400), National Natural Science Foundation of China (81772217, 201828006, 81971954, 81572012), Science and Technology Planning Project of Guangdong Province (2018A050506038), and Key project of Guangzhou science research [201904020011], Guangdong Provincial Natural Science Foundation Project (2016A030311025, 2017A030313694) to HJP

Author Contributions

Conceptualization: Ling Kong, Hongjuan Peng; Formal analysis: Ling Kong, Dan Jiang; Funding acquisition: Hongjuan Peng; Investigation: Ling Kong, Cheng He, Jing Xia, Haixia Wei, Lijuan Zhou; Methodology: Ling Kong, Cheng He, Jing Xia, Haixia Wei; Project administration: Hongjuan Peng; Resources: Hongjuan Peng; Supervision: Hongjuan Peng; Validation: Hongjuan Peng; Writing - original draft: Ling Kong, Hongjuan Peng; Writing - review & editing: Ling Kong, Hongjuan Peng. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

We thank Prof. Wen-Bin Ma in the School of Medicine of Sun Yat-Sen University for kindly providing pDONR cloning plasmid.

Conflicts of Interest

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Not applicable

References

1. Laliberte J, Carruthers VB. Host cell manipulation by the human pathogen *Toxoplasma gondii*. Cellular and molecular life sciences. 2008;65(12):1900-15.
2. Webster JP, Dubey JP. Toxoplasmosis of Animals and Humans. BioMed Central; 2010. p. 3:112.
3. Tenter AM, Heckeroth AR, Weiss LM. *Toxoplasma gondii*: from animals to humans. International journal for parasitology. 2000;30(12):1217-58.
4. Montoya JG, Liesenfeld O. Toxoplasmosis. Lancet (London, England). 2004;363(9425):1965-76.
5. Goldstein EJ, Montoya JG, Remington JS. Management of *Toxoplasma gondii* infection during pregnancy. Clinical Infectious Diseases. 2008;47(4):554-66.
6. Dubey J, Jones J. *Toxoplasma gondii* infection in humans and animals in the United States. International journal for parasitology. 2008;38(11):1257-78.
7. Hampton MM. Congenital Toxoplasmosis: A Review. Neonatal network : NN. 2015;34(5):274-8.
8. Blackman MJ, Bannister LH. Apical organelles of Apicomplexa: biology and isolation by subcellular fractionation. Molecular and biochemical parasitology. 2001;117(1):11-25.
9. Morisaki JH, Heuser JE, Sibley LD. Invasion of *Toxoplasma gondii* occurs by active penetration of the host cell. Journal of cell science. 1995;108(6):2457-64.
10. Alexander DL, Mital J, Ward GE, Bradley P, Boothroyd JC. Identification of the moving junction complex of *Toxoplasma gondii*: a collaboration between distinct secretory organelles. PLoS pathogens. 2005;1(2):e17.

11. Carruthers V, Boothroyd JC. Pulling together: an integrated model of *Toxoplasma* cell invasion. *Current opinion in microbiology*. 2007;10(1):83-9.
12. Shen B, Sibley LD. The moving junction, a key portal to host cell invasion by apicomplexan parasites. *Current opinion in microbiology*. 2012;15(4):449-55.
13. Yamamoto M, Takeda K. Inhibition of ATF6 β -dependent host adaptive immune response by a *Toxoplasma* virulence factor ROP18. *Virulence*. 2012;3(1):77-80.
14. Blader IJ, Koshy AA. *Toxoplasma gondii* development of its replicative niche: in its host cell and beyond. *Eukaryotic cell*. 2014;13(8):965-76.
15. Yarovinsky F. Innate immunity to *Toxoplasma gondii* infection. *Nature Reviews Immunology*. 2014;14(2):109-21.
16. He C, Kong L, Zhou L, Xia J, Wei H, Liu M, et al. Host Cell Vimentin Restrains *Toxoplasma gondii* Invasion and Phosphorylation of Vimentin is Partially Regulated by Interaction with TgROP18. *International journal of biological sciences*. 2017;13(9):1126-37.
17. Koshy AA, Fouts AE, Lodoen MB, Alkan O, Blau HM, Boothroyd JC. *Toxoplasma* secreting Cre recombinase for analysis of host-parasite interactions. *Nature methods*. 2010;7(4):307-9.
18. Koshy AA, Dietrich HK, Christian DA, Melehani JH, Shastri AJ, Hunter CA, et al. *Toxoplasma* co-opts host cells it does not invade. *PLoS pathogens*. 2012;8(7):e1002825.
19. Saeij J, Coller S, Boyle J, Jerome M, White M, Boothroyd J. *Toxoplasma* co-opts host gene expression by injection of a polymorphic kinase homologue. *Nature*. 2007;445(7125):324-7.
20. Hunter CA, Sibley LD. Modulation of innate immunity by *Toxoplasma gondii* virulence effectors. *Nature Reviews Microbiology*. 2012;10(11):766-78.
21. Steinfeldt T, Könen-Waisman S, Tong L, Pawlowski N, Lamkemeyer T, Sibley LD, et al. Phosphorylation of mouse immunity-related GTPase (IRG) resistance proteins is an evasion strategy for virulent *Toxoplasma gondii*. *PLoS biology*. 2010;8(12):e1000576.
22. Denkers EY. From cells to signaling cascades: manipulation of innate immunity by *Toxoplasma gondii*. *FEMS Immunology & Medical Microbiology*. 2003;39(3):193-203.
23. Taylor S, Barragan A, Su C, Fux B, Fentress S, Tang K, et al. A secreted serine-threonine kinase determines virulence in the eukaryotic pathogen *Toxoplasma gondii*. *Science*. 2006;314(5806):1776-80.
24. Zhang X, Su R, Cheng Z, Zhu W, Li Y, Wang Y, et al. A mechanistic study of *Toxoplasma gondii* ROP18 inhibiting differentiation of C17. 2 neural stem cells. *Parasites & vectors*. 2017;10(1):585.
25. Du J, An R, Chen L, Shen Y, Chen Y, Cheng L, et al. *Toxoplasma gondii* virulence factor ROP18 inhibits the host NF- κ B pathway by promoting p65 degradation. *Journal of Biological Chemistry*. 2014;289(18):12578-92.
26. Fentress SJ, Behnke MS, Dunay IR, Mashayekhi M, Rommereim LM, Fox BA, et al. Phosphorylation of immunity-related GTPases by a *Toxoplasma gondii*-secreted kinase promotes macrophage survival and virulence. *Cell host & microbe*. 2010;8(6):484-95.

27. Yamamoto M, Ma JS, Mueller C, Kamiyama N, Saiga H, Kubo E, et al. ATF6 β is a host cellular target of the *Toxoplasma gondii* virulence factor ROP18. *Journal of Experimental Medicine*. 2011;jem. 20101660.
28. Foster D, Parrish-Novak J, Fox B, Xu W. Cytokine–receptor pairing: accelerating discovery of cytokine function. *Nature Reviews Drug Discovery*. 2004;3(2):160-70.
29. Logsdon NJ, Deshpande A, Harris BD, Rajashankar KR, Walter MR. Structural basis for receptor sharing and activation by interleukin-20 receptor-2 (IL-20R2) binding cytokines. *Proceedings of the National Academy of Sciences*. 2012;109(31):12704-9.
30. Kunz S, Wolk K, Witte E, Witte K, Doecke WD, Volk HD, et al. Interleukin (IL)-19, IL-20 and IL-24 are produced by and act on keratinocytes and are distinct from classical ILs. *Experimental dermatology*. 2006;15(12):991-1004.
31. Wegenka UM. IL-20: biological functions mediated through two types of receptor complexes. *Cytokine & growth factor reviews*. 2010;21(5):353-63.
32. Blumberg H, Conklin D, Xu W, Grossmann A, Brender T, Carollo S, et al. Interleukin 20: discovery, receptor identification, and role in epidermal function. *Cell*. 2001;104(1):9-19.
33. Yamamoto M, Standley DM, Takashima S, Saiga H, Okuyama M, Kayama H, et al. A single polymorphic amino acid on *Toxoplasma gondii* kinase ROP16 determines the direct and strain-specific activation of Stat3. *Journal of Experimental Medicine*. 2009;206(12):2747-60.
34. Hakimi M-A, Olias P, Sibley LD. *Toxoplasma* effectors targeting host signaling and transcription. *Clinical microbiology reviews*. 2017;30(3):615-45.
35. Xia J, Kong L, Zhou L-J, He C, Peng H-J. Genome-wide BiFC-based proteomic analysis of *Toxoplasma gondii* ROP18's human interactome shows its key role in regulation of cell immunity and apoptosis. *Frontiers in Immunology*. 2018;9:61.
36. Shen B, Brown KM, Lee TD, Sibley LD. Efficient gene disruption in diverse strains of *Toxoplasma gondii* using CRISPR/CAS9. *MBio*. 2014;5(3):e01114-14.
37. Ihle JN. The Stat family in cytokine signaling. *Current opinion in cell biology*. 2001;13(2):211-7.
38. Ong Y-C, Reese ML, Boothroyd JC. *Toxoplasma* rhoptry protein 16 (ROP16) subverts host function by direct tyrosine phosphorylation of STAT6. *Journal of Biological Chemistry*. 2010;285(37):28731-40.
39. Hebenstreit D, Horejs-Hoeck J, Duschl A. JAK/STAT-dependent gene regulation by cytokines. *Drug News Perspect*. 2005;18(4):243-9.
40. Johnson LL. SCID mouse models of acute and relapsing chronic *Toxoplasma gondii* infections. *Infection and immunity*. 1992;60(9):3719-24.
41. Gazzinelli RT, Hieny S, Wynn TA, Wolf S, Sher A. Interleukin 12 is required for the T-lymphocyte-independent induction of interferon gamma by an intracellular parasite and induces resistance in T-cell-deficient hosts. *Proceedings of the National Academy of Sciences*. 1993;90(13):6115-9.
42. Khan IA, Matsuura T, Kasper LH. Interleukin-12 enhances murine survival against acute toxoplasmosis. *Infection and immunity*. 1994;62(5):1639-42.

43. Coutermarsh-Ott SL, Doran JT, Campbell C, Williams TM, Lindsay DS, Allen IC. Caspase-11 Modulates Inflammation and Attenuates *Toxoplasma gondii* Pathogenesis. Mediators of inflammation. 2016;2016:9848263.
44. Hunter CA, Subauste CS, Van Cleave V, Remington JS. Production of gamma interferon by natural killer cells from *Toxoplasma gondii*-infected SCID mice: regulation by interleukin-10, interleukin-12, and tumor necrosis factor alpha. Infection and immunity. 1994;62(7):2818-24.
45. Basak S, Hoffmann A. Crosstalk via the NF- κ B signaling system. Cytokine & growth factor reviews. 2008;19(3):187-97.
46. Cheng L, Chen Y, Chen L, Shen Y, Shen J, An R, et al. Interactions between the ROP18 kinase and host cell proteins that aid in the parasitism of *Toxoplasma gondii*. Acta tropica. 2012;122(3):255-60.
47. Shapira S, Harb OS, Margarit J, Matrajt M, Han J, Hoffmann A, et al. Initiation and termination of NF- κ B signaling by the intracellular protozoan parasite *Toxoplasma gondii*. J Cell Sci. 2005;118(15):3501-8.
48. Chen L, Christian DA, Kochanowsky JA, Phan AT, Clark JT, Wang S, et al. The *Toxoplasma gondii* virulence factor ROP16 acts in cis and trans, and suppresses T cell responses. The Journal of experimental medicine. 2020;217(3).

Figures




Figure 1

TgROP18 functions in STAT3 phosphorylation in HaCaT cells during *T. gondii* infection. HaCaT were infected with indicated parasites and harvested. (a) Total proteins of HaCaT cells were loaded and activation of STAT3 was determined by Western blot using anti-phospho-Stat3-Y705 and anti-phospho-Stat3-S727. (b) Densitometric quantitation of each band in A was applied using Image J. Statistical analysis was conducted with the Kruskal–Wallis H-test with Bonferroni correction, using SPSS 13.2 software (* $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$). Error bars show the standard errors (SD).




Figure 2

Extracellular TgROP18 resulted in STAT3-Y705 phosphorylation in HaCaT cells in dose-dependent manner without affecting the level of total STAT3. (a) HaCaT cells were infected with indicated parasites, the control group was treated with the IL-20 vehicle solution (BSA-PBS), the positive control was stimulated with IL-20. (c) HaCaT cells were treated with indicated amount of recombinant GST and GST-ROP18, and elution buffer was used as control. (d) HaCaT cells were infected with gradient MOI of indicated parasite, or uninfected as control. (a,c,e) The total proteins of HaCaT cells were extracted and subjected to Western blot (WB). (b, d, f) Densitometric quantitation of each band in a, c and e was applied using Image J. Statistical analysis was conducted by Kruskal–Wallis H-test with Bonferroni correction, using SPSS 13.2 software (* $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$). Error bars show the standard errors (SD).




Figure 3

Identification of IL-20RB interacting with TgROP18 with FRET and Co-IP (a) Identification of the interaction using FRET. The COS-7 cells co-transfected with pECFPN1-ROP18 and pEYFPC1-IL20RB (EXP) were used to detect the FRET efficiency between CFPN1-ROP18 and YFPC1-IL20RB. The COS-7 cells co-transfected with pECFPN1 and pEYFPC1 were used for negative control (NC) and the cells transfected with pECFPN1-EYFP as the positive control (PC). The FRET efficiency is shown in the right column where a color matched FRET efficiency scale is indicated for each image. The red end of the scale indicates a higher FRET efficiency. (b) The calculated FRET efficiency (left) and intermolecular distance (right) between ROP18 and IL20RB is shown as a column. (c) The interaction of ROP18 with IL20RB was further confirmed through immunoprecipitation assay (Co-IP) in vitro. Group 1, cells transfected with pEYFP-IL20RB-HA; Group 2, cells transfected with pCFP-ROP18i-3×FLAG; Group 3, cells co-transfected with pCFP-ROP18i-3×FLAG and pEYFP-IL20RB-HA; Group 4, negative control group.





Figure 4

Identification of IL-20RB extracellular domain as the TgROP18-binding domain. The interaction of ROP18 with the Extr/Cyt domain of IL20RB was identified with Co-IP. Lysates of COS-7 cells transiently transfected or co-transfected with the indicated plasmids, were immunoprecipitated with anti-HA. Starting fractions (Input) and immunoprecipitates (IP) were analyzed with WB to detect FLAG-ROP18 and HA-Extr/Cyt domain of IL20RB. Group 1, cells transfected with pEYFPC1-IL20RB-Extr-HA or pEYFPC1-IL20RB-Cyt-HA; Group 2, cells transfected with pCFP-ROP18i-3xFLAG; Group 3, cells co-transfected with pCFP-ROP18i-3xFLAG and pEYFPC1-IL20RB-Extr-HA or pEYFPC1-IL20RB-Cyt-HA; Group 4, negative control group.



indicated times. TNF- α transcription and translation was detected by qRT-PCR and WB. (c&f) Densitometric quantitation of each band in B or E was applied using Image J. Statistical analysis was performed with the Kruskal–Wallis H-test with Bonferroni correction (** $p < 0.01$).




Figure 6

Both TgROP18 and TgROP18 induce STAT3 activation in HaCaT cells. (a) HaCaT were infected with indicated parasites for 30min. Cell lysates were examined for STAT3 activation by WB. (b) Densitometric quantitation of STAT3 phosphorylation was applied with Image J and normalized to which of the total STAT3. Statistical analysis was conducted with the Kruskal–Wallis H-test with Bonferroni correction ($*p < 0.05$), and error bars show the standard errors of the mean for three independent experiments.




Figure 7

Schematic illustration of JAK/STAT3 pathway activated by TgROP18 through targeting IL20RB. Type I IL-20R is shown on the left and Type II IL-20R on the right. TgROP18 is secreted from rhoptries of *T. gondii* and binds to host cell IL20RB on the extracellular domain during invasion. The β -chain of IL20R (IL20RB) associates with and phosphorylates TYK2, while the α -chain (IL20RA or IL22RA1) associates with and phosphorylates JAK1. Activated JAK1 and TYK2 undergo cross-phosphorylation and

autophosphorylation. Thereafter, the tyrosine residues within the intracellular domain of IL-20RA (IL-20RA1) are phosphorylated and create the docking sites for STAT3. Meanwhile, free STAT3 in the cytoplasm are recruited and phosphorylated by JAK1 and TYK2. Phosphorylated STAT3 then forms homodimers and translocates to the nucleus to initiate transcription of STAT3 responsive genes such as TNF- α through STAT-responsive promoter elements. STAT3, signal transducers and activators of transcription 3; JAK1, Janus Kinase 1; TYK2, Tyrosine Kinase 2; Y: tyrosine residue; P, phosphorylation.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.docx](#)
- [supplement2.tif](#)
- [supplement3.tif](#)
- [supplement4.tif](#)
- [supplement5.tif](#)
- [Graphicalabstract.jpg](#)
- [FigS4.tif](#)