

Association of epicardial fat thickness with left ventricular diastolic function parameters in a community population

Wei Ma (✉ kmmawei@sina.com)

Peking University First Hospital <https://orcid.org/0000-0003-1305-3980>

Baowei Zhang

Peking University First Hospital

Ying Yang

Peking University First Hospital

Litong Qi

Peking University First Hospital

Jin Zhou

Peking University First Hospital

Min Li

Peking University First Hospital

Jia Jia

Peking University First Hospital

Yan Zhang

Peking University First Hospital

Huo Yong

Peking University First Hospital

Research article

Keywords: epicardial fat, thickness, left ventricle, diastolic function

Posted Date: April 12th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-171598/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Purpose

We examined the relationship between EFT measured by echocardiography and LV diastolic function parameters in a Beijing community population.

Methods

We included 1004 participants in this study. Echocardiographic parameters including E and A peak velocity, the early diastolic velocities (e') of the septal and lateral of mitral annulus using tissue doppler imaging, E/e' , and EFT, were measured. EFT1 was measured perpendicularly on the right ventricular free wall at end-diastole in the extension line of the aortic root. EFT2 was the maximum thickness measured perpendicularly on the right ventricular free wall at end-diastole. Multivariate linear regression was used to analyze the relationship between EFT and the mean e' and E/e' .

Results

The mean age of the participants was 63.91 ± 9.02 years old (51.4% men). EFT1 and EFT2 were negatively correlated with e' lat, e' sep, and e' mean ($p < 0.05$), and positively correlated with E/e' lat, E/e' sep, and E/e' mean. Multivariate regression analysis showed that EFT1 and EFT2 were independently and negatively correlated with e' mean (EFT1: $\beta = -0.089$ [95% confidence interval = $-0.177, -0.000$, $p = 0.050$]; EFT2: $\beta = -0.078$ [95% confidence interval = $-0.143, -0.012$, $p = 0.020$]). There were no interactions between EFT and any covariates, including age or heart groups, sex, BMI, or presence of hypertension, diabetes, or coronary heart disease in relation to LV diastolic dysfunction.

Conclusions

EFT was negatively and independently correlated with e' mean, suggesting that more attention to this type of adipose fat is required for cardiovascular disease therapy.

Introduction

There has been a progressive increase in the incidence of obesity with the increasing prevalence of unhealthy lifestyles in recent years. An increase in visceral fat is the main manifestation of the abnormal fat distribution in obesity, which is closely related to cardiovascular disease, metabolic syndrome [1]. Therefore, abnormal fat distribution should be taken into account when considering cardiovascular risk assessment.

There are two types of fat around the heart, including epicardial and pericardial fat[2]. Epicardial fat is located between the myocardium and visceral pericardium. Pericardial fat is located on the external surface of the parietal pericardium [2]. Epicardial fat is an important endocrine and paracrine organ producing variety of active substances [2]. Epicardial fat is closely associated with obesity, metabolic syndrome, and heart disease, especially coronary artery disease (CAD) [3], and can predict the cardiovascular disease and mortality in patients with type 2 diabetes [4].

Left ventricular (LV) diastolic dysfunction is one of the main reason of all-cause mortality [5]. Obesity is one of the important risk factors for the heart failure with preserved EF. [6]. Intramyocardial fat was correlated with LV diastolic

dysfunction in HFpEF patients [7]. Further, an increased epicardial fat thickness (EFT) was correlated with LV diastolic function impairment in obesity and coronary heart disease (CHD) patients [8–10]. However, there is still a lack of community population research. Thus, we examined the relationship between EFT measured by echocardiography and LV diastolic function parameters in a Beijing community population.

Methods

Population

All residents who were more than 40 years of age and lived in Shi Jing Shan district of Beijing were invited. Residents were contacted by recruitment advertisements or by telephone, and those volunteering to participate were included. Of the 5593 individuals who were ≥ 40 years of age, 1058 (18.9%) volunteered to participate in this study. The investigation started in July 2004 and ended in June 2005. Participants who had echocardiography were recruited in this study, while participants who had valve disease, evident arrhythmia (including atrial fibrillation), regional wall movement abnormal, or a LV ejection fraction (LVEF) $< 50\%$ were excluded. A final 1004 participants were included in this study. The study was approved by the institutional review board of Peking University First Hospital, and informed consent was obtained from all participants.

Definition of cardiovascular risk factors and disease

Waist circumference (WC) was measured in the standing position using the middle circumference between the lower rib margin and the iliac crest. Body mass index (BMI) was calculated. After a rest period of 15 min, blood pressure (BP) was measured three times, with 5-min intervals between each measurement, at the right upper arm in a sitting position with a mercury sphygmomanometer. Mean systolic BP and diastolic BP were calculated based on the three measurements. Total cholesterol and total triglyceride were analyzed by a fasting blood sample using standard techniques in the Beijing Hypertension League Institute. Participants with cigarette smoking history were identified as smokers. Hypertension was defined as office systolic BP ≥ 140 mmHg and/or diastolic BP ≥ 90 mmHg, or history or the usage of antihypertensive drugs. Diabetes was diagnosed according to each participant's interview and use of hypoglycemic drugs. Participants with a fasting glucose ≥ 7.0 mmol/L and a 2-h glucose ≥ 11.1 mmol/L on the oral glucose tolerance test were also defined as diabetic. Stroke, including intracerebral hemorrhage, cerebral infarction, and transient ischemic attack, was determined based on a history of data collected from hospitalizations and outpatient records, which were confirmed by computed tomography (CT) or magnetic resonance imaging (MRI) scan [11]. A history of old myocardial infarction, percutaneous coronary intervention, and coronary artery bypass grafting were all included in CHD.

Echocardiography

Echocardiography was performed using a 3-MHz transducer in an ultrasound system (Vivid-7; General Electric). According to the guidelines [12], standard images were collected and stored. One experienced echocardiography doctor blinded to the clinical picture of the participants measured the echo parameters in Peking University First Hospital central lab.

LV internal dimensions, LV wall thickness, and LVEF (measured by the Teichholtz method) were measured following the recommendations [12]. LV mass (LVM) was calculated as $LVM = 0.8 \cdot 1.04 \cdot ([PWTd + VSTd + LVIDd]^3 - [LVIDd]^3) + 0.6$ g, where PWTd and SWTd are the posterior and septal wall thickness at end diastole, respectively, and LVIDd is the LV dimension at end diastole [12]. LV mass index (LVMI) was calculated. The biplane method was used to measure the maximum left atrial volume (LAV), and LAV index (LAVI) was calculated. Transmitral inflow velocities were

measured using pulsed doppler at the mitral valve leaflet tips in the apical four-chamber view. E wave velocity, A wave velocity, E/A ratio and the E wave deceleration time (DT) was measured. Tissue doppler imaging was used to measure LV myocardial velocities in the apical four-chamber view, the early diastolic velocities (e') of the septal and lateral mitral annulus were measured. The e' mean was calculated as the mean of the septal e' (e' sep) and lateral e' (e' lat). The E/e' was calculated, and the E/e' mean was calculated as the mean of the septal E/e' (E/e' sep) and lateral E/e' (E/e' lat) regions. The mean doppler values were obtained over three different cardiac cycles.

EFT was identified as the echo-free space between the myocardium and the visceral layer of the pericardium from the parasternal long axis view. EFT1 was measured in the extension line of the aortic root and perpendicularly on the right ventricular free wall. EFT2 was the maximum thickness measured perpendicularly on the right ventricular free wall. Both EFTs were measured at end-diastole. The intra class correlation coefficient was used to evaluate intra-researcher and inter-researcher consistency. Interobserver values for EFT1 and EFT2 were 0.897 (95% confidence interval [CI]: 0.794–0.949, $p < 0.05$) and 0.914 (95% CI: 0.827–0.958, $p < 0.05$), respectively, while intraobserver values for EFT1 and EFT2 were 0.532 (95% CI: 0.218–0.746, $p < 0.05$) and 0.548 (95% CI: 0.240–0.756, $p < 0.05$), respectively.

Statistical analysis

Measurement data are presented as mean \pm standard deviation, while counting data are presented as percentages. Spearman correlation was used to analyze the correlation of EFT with echocardiographic parameters. Multivariate linear regression was used to analyze the relationship between EFT and e' mean and E/e' mean, adjusting for age, heart rate, WC, hypertension, diabetes, and CHD. Subgroup analyses and interaction tests were used to examine the EFT2 and e' mean according to age (< 65 years and ≥ 65 years), sex (male and female), BMI (< 28 kg/m² and ≥ 28 kg/m²), heart rate (< 80 beats per min and ≥ 80 beats per min), hypertension (yes or no), diabetes mellitus (yes or no), and CHD (yes or no). A p -value < 0.05 (two-sided) was considered statistically significant for all tests. All analyses were performed with statistical software (Empower(R), www.empowerstats.com; X&Y solutions, Inc., Boston, MA, USA; R [<http://www.R-project.org>] v3.4.3; SPSS v13.0).

Results

The general characteristics of the participants are shown (Table 1). The mean age was 63.91 ± 9.02 years of age, and 51.4% of participants were male. The prevalence of hypertension was 80.0%, diabetes was 29.4%, CHD was 12%, and stroke was 16.1%. The echocardiography parameters of the participants are shown in Table 2. The e' lat, e' sep, and e' mean were all reduced compared to normal values, while there were no changes in indicators of LV filling pressure (E/e' sep, E/e' lat, and E/e' mean).

Table 1
General characteristics of the study participants

Variables	Values
Sex(male,n%)	516(51.4)
age(years)	63.91 ± 9.02
BMI(kg/m ²)	25.88 ± 3.40
WC(cm)	88.72 ± 9.21
SBP(mmHg)	133.98 ± 17.95
DBP(mmHg)	80.07 ± 10.97
HR(/min)	71.09 ± 10.04
Smoking(n,%)	352(35.0)
Hypertension(n,%)	803(80.0)
Diabetes(n,%)	295(29.4)
CHD(n,%)	120(12.0)
Stroke(n,%)	162(16.1)
TC(mmol/l)	5.28 ± 1.02
TG(mmol/l)	2.30 ± 1.92
Fast Glucose(mmol/l)	6.36 ± 2.24
<i>BMI</i> body mass index, <i>WC</i> waist circumference, <i>SBP</i> systolic blood pressure, <i>DBP</i> diastolic blood pressure, <i>HR</i> heart rate, <i>CHD</i> coronary heart disease, <i>TC</i> total cholesterol, <i>TG</i> total triglyceride	

Table 2
Echocardiography parameters of the participants

Variables	Values
IVS(cm)	0.94 ± 0.13
LVPW(cm)	0.92 ± 0.07
LVEDD(cm)	4.30 ± 0.50
LVESD(cm)	2.65 ± 0.90
LVMI(g/m ²)	76.91 ± 16.20
LAVI(ml/m ²)	34.14 ± 9.83
E peak velocity (cm/s)	78.41 ± 17.91
A peak velocity (cm/s)	89.12 ± 18.39
DT(ms)	245.09 ± 48.22
E/A	0.91 ± 0.27
e'lat(cm/s)	9.37 ± 2.73
e'sep(cm/s)	6.54 ± 2.04
e'mean(cm/s)	7.96 ± 9.83
E/e'lat	8.99 ± 3.18
E/e'sep	12.83 ± 4.02
E/e'mean	10.91 ± 3.26
LVEF(%)	69.11 ± 8.25
EFT1(cm)	0.39 ± 1.26
EFT2(cm)	0.68 ± 1.77
<i>IVS</i> interventricular septum thickness, <i>LVPW</i> left ventricular posterior wall thickness, <i>LVEDD</i> left ventricular end diastolic diameter, <i>LVESD</i> left ventricular end systolic diameter, <i>LVMI</i> left ventricular mass index, <i>LAVI</i> left atrial volume index, <i>DT</i> E wave deceleration time, <i>e' sep</i> septal e', <i>e' lat</i> lateral e', <i>E/e' lat</i> lateral E/e', <i>E/e' sep</i> septal E/e', <i>LVEF</i> left ventricular ejection fraction, <i>EFT1</i> epicardial fat thickness 1 (measured perpendicularly on the right ventricular free wall at end-diastole in the extension line of the aortic root), <i>EFT2</i> epicardial fat thickness 2 (the maximum thickness measured perpendicularly on the right ventricular free wall at end-diastole)	

EFT1 and EFT2 were positively correlated with interventricular septum thickness and LV posterior wall thickness ($p < 0.05$), but not with LVMI and LAVI ($p > 0.05$). EFT1 and EFT2 were negatively correlated with E and E/A ($p < 0.05$), and positively correlated with A and DT ($p > 0.05$). EFT1 and EFT2 were negatively correlated with e' lat, e' sep, and e' mean ($p < 0.05$), and positively correlated with E/e' lat, E/e' sep, and E/e' mean ($p < 0.05$) (Table 3).

Table 3
Correlation analysis of epicardial fat thickness (EFT) with echocardiography parameters

	Epi1	Epi2
IVS	0.108 ^a	0.124 ^a
LVPW	0.115 ^a	0.136 ^a
LVEDD	-0.039	-0.060
LVESD	-0.012	-0.019
LVMl	0.030	0.016
LAVI	-0.029	-0.049
E peak velocity	-0.070 ^a	-0.113 ^a
A peak velocity	0.121 ^a	0.107 ^a
DT	0.073 ^a	0.089 ^a
E/A	-0.154 ^a	-0.206 ^a
e'lat	-0.136 ^a	-0.162 ^a
e'sep	-0.129 ^a	-0.146 ^a
e'mean	-0.138 ^a	-0.180 ^a
E/e'lat	0.093 ^a	0.090 ^a
E/e'sep	0.097 ^a	0.081 ^a
E/e'mean	0.100 ^a	0.090 ^a
LVEF	0.002	-0.009
^a <i>p</i> < 0.05		
<i>IVS</i> interventricular septum thickness, <i>LVPW</i> left ventricular posterior wall thickness, <i>LVEDD</i> left ventricular end diastolic diameter, <i>LVESD</i> left ventricular end systolic diameter, <i>LVMl</i> left ventricular mass index, <i>LAVI</i> left atrial volume index, <i>DT</i> E wave deceleration time, <i>e' sep</i> septal e', <i>e' lat</i> lateral e', <i>E/e' lat</i> lateral E/e', <i>E/e' sep</i> septal E/e', <i>LVEF</i> , left ventricular ejection fraction		

Univariate analysis showed that EFT1 and EFT2 were negatively correlated with e' mean, while multivariate regression analysis showed that the two EFTs were negatively correlated with e'mean after adjusting for age, sex, heart rate, WC, hypertension, diabetes, and CHD. EFT1 and EFT2 were not independently related with E/e' mean on univariate analysis or multivariate regression analysis (Table 4). Results of subgroup analysis of the relationship between e' mean and EFT2 are shown in Fig. 1. There were no interactions between EFT2 and any covariates, including age and heart groups, sex, BMI, or presence of hypertension, diabetes, or CHD. These findings were consistent with EFT1 (data were not shown).

Table 4

Univariate and multivariate regression analysis of epicardial fat thickness (EFT) and e' mean and E/e' mean

	e'mean			E/e'mean		
	Crude β (95%CI)	Adjusted β (95%CI)	p	Crude β (95%CI)	Adjusted β (95%CI)	p
EFT1	-0.235 (-0.341,-0.130)	-0.089(-0.177,-0.000)	0.050	0.212(0.052,0.372)	0.035(-0.116,0.186)	0.652
EFT2	-0.222 (-0.297,-0.146)	-0.078 (-0.143,-0.012)	0.020	0.181(0.066,0.296)	0.012(-0.099,0.123)	0.829
Adjusted for age, sex, heart rate, WC, hypertension, dia						

Discussion

There are many studies about the association of adipose tissue and LV diastolic dysfunction, including global adiposity [13], central adiposity [14], and visceral adiposity [15]. Epicardial fat plays an important role in lipid and energy metabolism, which can also have harmful effects because it can secrete many proatherogenic and proinflammatory cytokines [16]. In our study, EFT measured by echocardiography was independently correlated with e' mean, but not with E/e' mean, in a high-risk community population with an LVEF \geq 50%. Subgroup analysis showed no interaction with age, sex, BMI, heart rate, or preexisting conditions such as hypertension, diabetes, or CHD, and no correlation of EFT2 with e' mean.

Epicardial fat can be assessed by multiple imaging methods, including CT, MRI, and echocardiography. The thickness, area, and volume of epicardial fat can be measured by CT with manual or semi-automated methods. However, the high cost and radiation exposure associated with CT are disadvantageous, especially in large population studies. MRI is considered the golden standard for evaluating heart fat, although its use is also limited because of high cost and high requirement. By contrast, echocardiography is the most convenient method to evaluate epicardial fat, and is particularly suitable for epidemiological studies. The thickness and area of the epicardial fat can be measured with echocardiography [3], although the image quality has a marked influence on accuracy. Further, there can be differences between different readers. We used echocardiography to evaluate EFT in the present study. Echocardiography is widely used to evaluate epicardial fat [3]. Using ultrasound measurements, a Korean study reported a correlation between EFT and CT measurements of epicardial fat volume [17]. Echocardiographic epicardial fat measurements were also shown to have a strong correlation with MRI measurements [18]. It is important to note that paracardial fat consists of epicardial fat and pericardial fat, which should be distinguished during echocardiography examination. Echocardiographic EFT can be measured from the parasternal long and short axis views. However, EFT measured from the long axis view (but not the short axis view) was reported to be the independent predictor of e' septal and e' lateral [19].

In the present study, EFT was associated with most echocardiographic parameters. For example, EFT was positively correlated with ventricular septum and LV posterior wall thickness, but not with LVMI. One study showed that an increasing epicardial fat was significantly related to an increase in LVM because of high free fatty acids levels, insulin resistance and adrenergic activity, and that increased visceral fat directly affected LV output to perfuse the increased body mass [20]. In that study, the participants were younger (mean age, 46.9 years) and more obese (BMI, 30–30.5 kg/m²) than in our cohort. A further study reported that EFT was correlated with atria enlargement in morbidly obese

subjects [8]. By contrast, we found no association of EFT with LAVI, although our participants were older, with a higher prevalence of hypertension. Thus, LVMI and LAVI may be more significantly correlated with hypertension and age.

There are a few echocardiographic parameters used to evaluate LV diastolic function. Mitral inflow pattern, including E and A wave velocity, E/A, and E wave DT, are affected by many factors. A significant association of e' with LV relaxation was reported in human subjects [21]. The E/ e' ratio can also be used to evaluate LV filling pressures [22]. Further, e' is a powerful predictor of cardiac mortality in patients, independent of normal or abnormal LV systolic function [23, 24], while mitral E/ e' is a strong predictor of cardiac death or rehospitalization for CHF as well [25]. In our correlation analysis, EFT was negatively associated with E, E/A, and e' , and positively associated with A, DT, and E/ e' . In multivariate regression analysis, EFT was independently associated with e' mean, but not with E/ e' . Interestingly, in Japanese patients with known or suspected CAD, EFT was negatively associated with e' mean and positively correlated with E/ e' mean [10]. Konishi et al. also reported that epicardial fat volumes measured by CT were significantly and independently associated with E/ e' >10 in suspected CAD patients [15]. Further EFT was significantly associated with LV diastolic dysfunction in subjects with normal coronary artery [26]. Finally, Dabbah et al. reported that E/ e' was not associated with EFT [19], similar to that in the present study.

In our subgroup analysis, there was no effect of CHD on the relationship between EFT2 and e' mean. Cavalcante and Konishi reported an independent correlation of epicardial fat and E/ e' , although as that study included patients suspected of CAD, some patients may have advanced diastolic dysfunction since ischemia [15]. Hypertension is a risk factor for the occurrence of LV diastolic dysfunction. In patients with newly diagnosed and untreated hypertension, increased EFT was significantly and independently related to the degree of LV diastolic function [9]. In our subgroup analysis, we found no effect of hypertension on the relationship between EFT2 and e' mean. It was also reported that EFT was more common in women than men > 60 years old, and that EFT was significantly related to LV function in women, but not men [27]. By contrast, we found no interactions of different ages or sex on the correlation of EFT with e' mean. All of these contrasting findings may be related to the different imaging methods for evaluating epicardial fat, or to different populations of patients.

There are several limitations to our study. First, epicardial fat was measured by echocardiography rather than MRI or CT. However, EFT measured by echocardiography was shown to correlate with volumetric measurements. Second, because of the cross-sectional nature of our study, a causal relationship between EFT and e' cannot be determined. Prospective studies examining whether increased EFT is predictive of LV diastolic dysfunction are required. Finally, the majority of patients were > 40 years of age, and thus our findings may not reflect the characteristics of epicardial fat in a younger population.

In summary, EFT measured by echocardiography was independently correlated with the e' mean, but not with the E/ e' mean, in a high-risk community population with an LVEF \geq 50%. Subgroup analysis showed no interaction with age, sex, BMI, heart rate, or preexisting conditions such as hypertension, diabetes, or CHD, or of the correlation between EFT and e' mean.

Declarations

Ethical Approval and Consent to participate

The study was approved by the institutional review board of Peking University First Hospital, and informed consent was obtained from all participants.

Consent for publication

For all manuscripts that include details, images, or videos relating to an individual person, written informed consent for the publication of these details have been obtained from that person (or their parent or legal guardian in the case of children under 18).

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available due to the government policy but are available from the corresponding author on reasonable request.

Code availability

Not applicable

Competing interests

On behalf of all authors, the corresponding author declare that the authors have no conflicts of interest to disclose.

Funding

1. Peking University Medicine Seed Fund for Interdisciplinary Research (BMU2018MX002)
2. The National Key Research and Development Program of China (No.2017YFC1307704)

Authors' contributions

Wei Ma: Research design, echo data measurement, data analysis, paper writing

Baowei Zhang, Ying Yang, Litong Qi: Research design, performing echo , data analysis

Jin Zhou: Research design

Min Li, Jia Jia,: data analysis, paper writing

Yan Zhang, Huo Yong: Research design, data analysis

References

1. Hamdy O, Porramatikul S, Al-Ozairi E (2006) Metabolic obesity: the paradox between visceral and subcutaneous fat. *Current diabetes reviews* 2:367-373.
2. Sacks HS, Fain JN (2007) Human epicardial adipose tissue: a review. *American heart journal* 153:907-917.
3. Tekin I, Edem E (2018) Association of Epicardial Fat Tissue with Coronary Artery Disease and Left Ventricle Diastolic Function Indicators. *Medical science monitor : international medical journal of experimental and clinical research* 24:6367-6374.
4. Christensen RH, von Scholten BJ, Hansen CS, Jensen MT, Vilsbøll T, Rossing P, et al. (2019) Epicardial adipose tissue predicts incident cardiovascular disease and mortality in patients with type 2 diabetes. *Cardiovascular diabetology* 18:114.
5. Halley CM, Houghtaling PL, Khalil MK, Thomas JD, Jaber WA (2011) Mortality rate in patients with diastolic dysfunction and normal systolic function. *Archives of internal medicine* 171:1082-1087.
6. Haass M, Kitzman DW, Anand IS, Miller A, Zile MR, Massie BM, et al. (2011) Body mass index and adverse cardiovascular outcomes in heart failure patients with preserved ejection fraction: results from the Irbesartan in

- Heart Failure with Preserved Ejection Fraction (I-PRESERVE) trial. *Circulation Heart failure* 4:324-331.
7. Wu CK, Lee JK, Hsu JC, Su MYM, Wu YF, Lin TT, et al. (2020) Myocardial adipose deposition and the development of heart failure with preserved ejection fraction. *European journal of heart failure* 22:445-454.
 8. Iacobellis G, Leonetti F, Singh N, M Sharma A (2007) Relationship of epicardial adipose tissue with atrial dimensions and diastolic function in morbidly obese subjects. *International journal of cardiology* 115:272-273.
 9. Turak O, Özcan F, Canpolat U, İşleyen A, Cebeci M, Öksüz F, et al. (2013) Increased echocardiographic epicardial fat thickness and high-sensitivity CRP level indicate diastolic dysfunction in patients with newly diagnosed essential hypertension. *Blood pressure monitoring* 18:259-264.
 10. Hachiya K, Fukuta H, Wakami K, Goto T, Tani T, Ohte N (2014) Relation of epicardial fat to central aortic pressure and left ventricular diastolic function in patients with known or suspected coronary artery disease. *The international journal of cardiovascular imaging* 30:1393-1398.
 11. Liu L, Zhao F, Yang Y, Qi LT, Zhang BW, Chen F, et al. (2008) The clinical significance of carotid intima-media thickness in cardiovascular diseases: a survey in Beijing. *Journal of Human Hypertension* 22:259-265.
 12. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, et al. (2005) Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *Journal of the American Society of Echocardiography : official publication of the American Society of Echocardiography* 18:1440-1463.
 13. Russo C, Jin Z, Homma S, Rundek T, Elkind MSV, Sacco RL, et al. (2011) Effect of obesity and overweight on left ventricular diastolic function: a community-based study in an elderly cohort. *Journal of the American College of Cardiology* 57:1368-1374.
 14. Ammar KA, Redfield MM, Mahoney DW, Johnson M, Jacobsen SJ, Rodeheffer RJ (2008) Central obesity: association with left ventricular dysfunction and mortality in the community. *American heart journal* 156:975-981.
 15. Konishi M, Sugiyama S, Sugamura K, Nozaki T, Matsubara J, Akiyama E, et al. (2012) Accumulation of pericardial fat correlates with left ventricular diastolic dysfunction in patients with normal ejection fraction. *Journal of cardiology* 59:344-351.
 16. Iacobellis G, Corradi D, Sharma AM (2005) Epicardial adipose tissue: anatomic, biomolecular and clinical relationships with the heart. *Nature clinical practice Cardiovascular medicine* 2:536-543.
 17. Kim BJ, Kang JG, Lee SH, Lee JY, Sung KC, Kim BS, et al. (2017) Relationship of Echocardiographic Epicardial Fat Thickness and Epicardial Fat Volume by Computed Tomography with Coronary Artery Calcification: Data from the CAESAR Study. *Archives of Medical Research* 48:352-359.
 18. Iacobellis G, Assael F, Ribaudo MC, Zappaterreno A, Alessi G, Di Mario U, et al. (2003) Epicardial fat from echocardiography: a new method for visceral adipose tissue prediction. *Obesity research* 11:304-310.
 19. Dabbah S, Komarov H, Marmor A, Assy N (2014) Epicardial fat, rather than pericardial fat, is independently associated with diastolic filling in subjects without apparent heart disease. *Nutrition, metabolism, and cardiovascular diseases : NMCD* 24:877-882.
 20. Iacobellis G, Ribaudo MC, Zappaterreno A, Iannucci CV, Leonetti F (2004) Relation between epicardial adipose tissue and left ventricular mass. *The American journal of cardiology* 94:1084-1087.
 21. Sohn DW, Chai IH, Lee DJ, Kim HC, Kim HS, Oh BH, et al. (1997) Assessment of mitral annulus velocity by Doppler tissue imaging in the evaluation of left ventricular diastolic function. *Journal of the American College of Cardiology* 30:474-480.

22. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, et al. (2009) Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *Journal of the American Society of Echocardiography : official publication of the American Society of Echocardiography* 22:107-133.
23. Wang M, Yip GWK, Wang AYM, Zhang Y, Ho PY, Tse MK, et al. (2003) Peak early diastolic mitral annulus velocity by tissue Doppler imaging adds independent and incremental prognostic value. *Journal of the American College of Cardiology* 41:820-826.
24. Wang M, Yip G, Yu C-M, Zhang Q, Zhang Y, Tse D, et al. (2005) Independent and incremental prognostic value of early mitral annulus velocity in patients with impaired left ventricular systolic function. *Journal of the American College of Cardiology* 45:272-277.
25. Dokainish H, Zoghbi WA, Lakkis NM, Ambriz E, Patel R, Quinones MA, et al. (2005) Incremental predictive power of B-type natriuretic peptide and tissue Doppler echocardiography in the prognosis of patients with congestive heart failure. *Journal of the American College of Cardiology* 45:1223-1226.
26. Topuz M, Dogan A (2017) The effect of epicardial adipose tissue thickness on left ventricular diastolic functions in patients with normal coronary arteries. *Kardiologia polska* 75:196-203.
27. Kim SA, Kim MN, Shim WJ, Park SM (2017) Epicardial adipose tissue is related to cardiac function in elderly women, but not in men. *Nutrition, metabolism, and cardiovascular diseases : NMCD* 27:41-47.

Tables

Table 1 General characteristics of the study participants

Variables	Values
Sex[male]n[%]	516(51.4)
age[years]	63.91±9.02
BMI[kg/m ²]	25.88±3.40
WC[cm]	88.72±9.21
SBP[mmHg]	133.98±17.95
DBP[mmHg]	80.07±10.97
HR[beats/min]	71.09±10.04
Smoking[n(%)]	352(35.0)
Hypertension[n(%)]	803(80.0)
Diabetes[n(%)]	295(29.4)
CHD[n(%)]	120(12.0)
Stroke[n(%)]	162(16.1)
TC[mmol/l]	5.28±1.02
TG[mmol/l]	2.30±1.92
Fast Glucose[mmol/l]	6.36±2.24

BMI body mass index, *WC* waist circumference, *SBP* systolic blood pressure, *DBP* diastolic blood pressure, *HR* heart rate, *CHD* coronary heart disease, *TC* total cholesterol, *TG* total triglyceride

Table 2 Echocardiography parameters of the participants

Variables	Values
IVS(cm)	0.94±0.13
LVPW(cm)	0.92±0.07
LVEDD(cm)	4.30±0.50
LVESD(cm)	2.65±0.90
LVMI(g/m ²)	76.91±16.20
LAVI(ml/m ²)	34.14±9.83
E peak velocity (cm/s)	78.41±17.91
A peak velocity (cm/s)	89.12±18.39
DT(ms)	245.09±48.22
E/A	0.91±0.27
e'lat(cm/s)	9.37±2.73
e'sep(cm/s)	6.54±2.04
e'mean(cm/s)	7.96±9.83
E/e'lat	8.99±3.18
E/e'sep	12.83±4.02
E/e'mean	10.91±3.26
LVEF(%)	69.11±8.25
EFT1[cm]	0.39±1.26
EFT2[cm]	0.68±1.77

IVS interventricular septum thickness, *LVPW* left ventricular posterior wall thickness, *LVEDD* left ventricular end diastolic diameter, *LVESD* left ventricular end systolic diameter, *LVMI* left ventricular mass index, *LAVI* left atrial volume index, *DT* E wave deceleration time, *e' sep* septal e', *e' lat* lateral e', *E/e' lat* lateral E/e', *E/e' sep* septal E/e', *LVEF* left ventricular ejection fraction, *EFT1* epicardial fat thickness 1 (measured perpendicularly on the right ventricular free wall at end-diastole in the extension line of the aortic root), *EFT2* epicardial fat thickness 2 (the maximum thickness measured perpendicularly on the right ventricular free wall at end-diastole)

Table 3 Correlation analysis of epicardial fat thickness (EFT) with echocardiography parameters

	Epi1	Epi2
IVS	0.108 ^a	0.124 ^a
LVPW	0.115 ^a	0.136 ^a
LVEDD	-0.039	-0.060
LVESD	-0.012	-0.019
LVMi	0.030	0.016
LAVI	-0.029	-0.049
E peak velocity	-0.070 ^a	-0.113 ^a
A peak velocity	0.121 ^a	0.107 ^a
DT	0.073 ^a	0.089 ^a
E/A	-0.154 ^a	-0.206 ^a
e'lat	-0.136 ^a	-0.162 ^a
e'sep	-0.129 ^a	-0.146 ^a
e'mean	-0.138 ^a	-0.180 ^a
E/e'lat	0.093 ^a	0.090 ^a
E/e'sep	0.097 ^a	0.081 ^a
E/e'mean	0.100 ^a	0.090 ^a
LVEF	0.002	-0.009

^a $p < 0.05$

IVS interventricular septum thickness, LVPW left ventricular posterior wall thickness, LVEDD left ventricular end diastolic diameter, LVESD left ventricular end systolic diameter, LVMi left ventricular mass index, LAVI left atrial volume index, DTE wave deceleration time, e' sep septal e', e' lat lateral e', E/e' lat lateral E/e', E/e' sep septal E/e', LVEF, left ventricular ejection fraction

Table 4 Univariate and multivariate regression analysis of epicardial fat thickness (EFT) and e' mean and E/e' mean

	e'mean		E/e'mean			
	Crude β (95%CI)	Adjusted β (95%CI)	p	Crude β (95%CI)	Adjusted β (95%CI)	p
EFT1	-0.235 [-0.341;-0.130]	-0.089[-0.177;-0.000]	0.050	0.212[0.052;0.372]	0.035[-0.116;0.186]	0.652
EFT2	-0.222 [-0.297;-0.146]	-0.078 [-0.143;-0.012]	0.020	0.181[0.066;0.296]	0.012[-0.099;0.123]	0.829

Adjusted for age, sex, heart rate, WC, hypertension, dia

Figures




Figure 1

Subgroup analysis of the relationship between EFT2 and e' mean adjusted for age, sex, heart rate, WC, hypertension, diabetes, and coronary heart disease

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [conflictofinterest.pdf](#)