

Experiential learning for children's dental anxiety: a cluster randomized trial

Minmin Zhu (✉ zhuminmin1982@163.com)

Shenzhen Nanshan Center for Chronic Disease Control <https://orcid.org/0000-0002-1899-802X>

Hongbing Yu

Shenzhen Nanshan Center for Chronic Disease Control

Bo Xie

Shenzhen Nanshan Center for Chronic Disease Control

Hongwen Li

Shenzhen Nanshan Maternal and Child Health care Hospital

Qian He

Shenzhen Nanshan Center for Chronic Disease Control

Huimin Li

Shenzhen Nanshan Center for Chronic Disease Control

Jing Su

Shenzhen Nanshan Center for Chronic Disease Control

Xueqi Li

Shenzhen Nanshan Maternal and Child Health Care Hospital

Research article

Keywords: Dental anxiety, Child dentistry, Experiential learning, Efficacy, Tell-Show-Do, Cluster randomized trial

Posted Date: March 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-17207/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on July 31st, 2020. See the published version at <https://doi.org/10.1186/s12903-020-01204-5>.

Abstract

Background Dental anxiety (DA) has an impact on the quality of dental treatment and may have long-lasting implications for children. A recent study introducing experiential learning (EL) into children's oral health promotion resulted in better oral hygiene. The purpose of the study is to evaluate whether EL can reduce children's DA. **Methods** In September 2018, we recruited 988 children aged 7-8 years from 24 classes to participate in a cluster-randomized trial. Classes were randomly assigned to EL (in which children received a lively presentation on oral health and participated in a role play in a simulated dental clinic in the classroom) or the Tell-Show-Do (TSD) group (in which children received a conventional TSD behavior management). The primary outcome was the prevalence of DA after the procedure of pit and fissure sealant (PFS), assessed by a modified children's fear survey schedule-dental subscale. Secondary outcomes were changes in blood pressures (BP) and pulse rates (PR) before and after the PFS procedure. The intervention effects were estimated by means of mixed effect models, which included covariates of gender and school (and baseline value for BP and PR only), and a random cluster effect. **Results** In 396 children of the EL group who received the PFS treatment, the prevalence of DA (score > 38) was 18.5%, compared with 24.3% in 391 children of the TSD group (OR=0.65; 95% confidence interval, 0.46-0.93; P =0.019). The increases in BP and PR after the PFS were also significantly less in the EL group. **Conclusions** School-based experiential learning intervention before a dental visit is feasible and effective in lowering children's dental anxiety.

Background

Dental anxiety (DA), abnormal fear or dread of visiting the dentist and unwarranted anxiety over dental procedures, has an impact on the quality of dental treatment and may have long-lasting implications for children [1]. Cross-sectional and cohort studies published from 2000 to 2014 have reported prevalence rates of children's DA that ranged from 10–20% [2]. DA sometimes leads to a series of uncooperative or disruptive behaviors before or during dental procedures, termed dental behavior management problems (DBMP), that result in stressful and unpleasant experiences for both the child and the dentist [3]. In addition, studies have shown that DA has a consistent impact on pain throughout the entire dental treatment [4], and DA/DBMP are associated with children's dental caries [5–8].

Although it is possible to employ pharmacological interventions, such as anesthesia or sedation [9], dentists generally use communicational, behavioral and psychological techniques to manage children's DA/DBMP and achieve a high quality of dental care. These include methods such as "Tell-Show-Do" (TSD), voice control, distraction, modelling, restraint, and cognitive reconstructing [10–12]. Of these methods, however, some need specialist training, some cause psychological traumas to children, and most are only initiated just before or during dental treatment. In recent years, the most used non-pharmacological method has been TSD, which has generally been acceptable for the doctors, children and parents [13–16]. Nevertheless, the TSD is less useful for children with previous dental experience [17] and less effective than modelling [18, 19].

Experiential learning (EL) is an innovative learning technique by which knowledge or skill is gained through the experience of participating in real or simulated practical activities [20]. EL has been used effectively in enhancing knowledge, improving attitude and modifying behaviors in health education[21–24], and was recently introduced into children’s oral health promotion, resulting in better oral hygiene[25].

In this paper, we developed a school-based EL intervention and evaluated whether the EL was effective in reducing the prevalence of DA in primary school children.

Methods

Study design

This was a 2-arm cluster-randomized control trial comparing EL and TSD for their effectiveness in reducing DA in children. This study had the approved by the Ethics Committee of the Shenzhen Nanshan Maternal and Child Health Care Hospital. All parents of the children participated provided written informed consent. The trial was registered in Chinese Clinical Trial Registry (ChiCTR2000028878).

Participants

The participants were children aged 7–8 years, selected in September 2018 from the second grade of primary schools in the Nanshan District of the city of Shenzhen, China. At this age, children in Shenzhen routinely receive pit and fissure sealant (PFS) as a dental caries prophylaxis, which is a moderately invasive procedure.

Recruitment

The eligibility criteria for second grade classes were: 1) located in Nanshan District, Shenzhen City; 2) had 40 to 50 children aged 7–8 years; 3) agreed to participate. In September 2018, we used a multi-stage random sampling method to recruit the participants. We first used a table of random digits to select six primary schools in Nanshan that had four or more classes met eligibility criteria, and then random selected four eligible classes in each of the six schools. All children in the 24 selected classes were invited to take part in the study, and the informed consent were sent to their parents (Fig. 1). Children who already had a dental visit or refused to participate were excluded.

Randomization and Interventions

In this trial, the class was the unit of randomization. The sampled classes were randomly assigned into the EL or TSD groups in 1:1 ratio in school blocks with the use of a table of random digits (Fig. 1). Cluster randomization was necessary to avoid between-group contamination and it was feasibly conduct school-based intervention. After receiving the signed informed consent from the parents, the intervention was implemented on a class by class basis. Neither the dentists, nurses nor field interviewers who assessed the outcomes were aware of the randomization assignments of the classes, as the measurements were

the same for both groups. Only the dental nurses who performed the “Tell and Show” procedures were aware the assignments.

At the beginning of intervention, children in the EL group received a lively 40-minute presentation on oral health given by dentists. The standardized oral health education conveyed information about dental caries and their causes, proper toothbrushing and flossing, and the pit and fissure sealant, by using slides, cartoon videos, and dental models. One month later, the children participated in an EL activity in a simulated dental clinic in the classroom. In the activity, the dentists demonstrated the common tools of dentistry (mouth mirror, dental probe, dental handpiece and dental ejector, etc.). Then the children were organized in 4-person-groups for a five-minute role play around a dental bed, with which the children playing a patient, a dentist, a nurse and a parent. While one group of children performed their role play, the other children were in another room watching cartoon videos that delivered oral health education. And another month after that, they visited the dental clinic to receive the PFS treatment.

Children in the TSD group received a conventional 5-minute TSD behavior management session at the time of their visit to the dental clinic for the PFS procedure. The TSD behavior management process used in this study contained three procedures: Tell, the dental nurse explained to the child what the dentist would do during the PFS treatment; Show, the dentist nurse showed the equipment involved; Do, the dentist performed the PFS procedure.

Trial measurement

The dental nurse performed the measurements of the children’s blood pressures and pulse rates with an upper-arm electronic sphygmomanometer in the waiting room after a 15-minute quiet sitting period before the PFS treatment, and then remeasured them again after the treatment.

The field interviewers administered the modified children’s fear survey schedule-dental subscale (Modified CFSS-DS) [26] to the children after the PFS. The modified CFSS-DS was translated into Chinese from the original English version and was combined with a facial image scale (FIS) [27]. It had a Cronbach's alpha of 0.85 and a test-retest reliability ICC of 0.73 [26].

Outcomes

The primary outcome was the prevalence of DA, defined as a sum score of Modified CFSS-DS ≥ 38 . The secondary outcomes were mean differences of change in systolic blood pressures (SBP), diastolic blood pressures (DBP) and pulse rates (PR) in the measurements taken before and after PFS.

Statistical analysis

With an enrollment of 12 class per trial group, 40 participants per class, a rate of cohort retention of 80%, a rate of PFS of 90%, and an estimated intraclass correlation coefficient (0 – 0.01), the initial design yielded 80% power to detect a 10% difference in the prevalence of DA between TSD and EL groups, with a two-sided alpha level of 0.05.

The intervention effects were estimated by means of mixed effect logistic regression for binomial indicator (DA) and linear mixed-effects model for continuous variations (SBP, DBP and PR), which included a random cluster effect of class. The primary predictor was group (EL versus TSD) and included covariates of gender (male or female), school, and baseline value (for analysis of change of SBP, DBP or PR only), which were correlated with the dependent variable. All analyses were performed with the R project (Version 3.6.1 Patched for x64 Window system), and a P value < 0.05 was considered statistically significant.

Results

Characters of participants

A total of 24 classes from 6 schools were enrolled in the trial between Sept. 1st 2018 and Sept. 30th 2018, with a total initial enrollment of 988 children. The primary statistical analysis was based on the 396 participants in 12 EL classes and 391 participants in 11 TSD classes who completed the EL or TSD and the PFS treatment (Fig. 1). The two groups were similar in gender distribution and register residence (Table 1).

Table 1
Character of trial participants completing EL or TSD and PFS

Characters	EL group (n = 396)	TSD group (n = 391)	P value [§]
Gender			0.623
Male	221	226	
Female	175	165	
Register Residence*			0.654
Shenzhen	255	259	
Others	124	116	
*The register residence was unknown for 17 and 16 children in the EL and TSD groups, respectively.			
§ P value was calculated with the chi-square test.			

Primary Outcome

The prevalence of DA (Modified CFSS-DS scores ≥ 38) was 18.5% in the EL group and 24.3% in the TSD group. The mixed effect logistic regression showed that the OR was 0.65 (95% confidence interval [CI]: 0.46–0.93; P = 0.019), after adjusting gender and school (Table 2).

Table 2
Primary and secondary outcomes*

Outcomes	EL group	TSD group	Intervention effect [#]	P value [§]
DA (%)	18.5	24.3	0.65(0.46 – 0.93)	0.019
SBP (mmHg)				
At the waiting room	108.7 ± 12.1	107.8 ± 12.7		
After the PFS	106.1 ± 13.0	108.5 ± 14.2		
Change	-2.7 ± 12.8	0.7 ± 14.2	-2.6(-4.4 – -0.9)	0.036
DBP (mmHg)				
At the waiting room	66.2 ± 9.5	65.5 ± 10.0		
After the PFS	68.0 ± 12.1	70.0 ± 11.6		
Change	1.7 ± 12.3	4.5 ± 12.9	-2.0(-4.4–0.3)	0.086
PR (Beat per Minute)				
At the waiting room	93.1 ± 14.0	92.0 ± 13.5		
After the PFS	93.3 ± 13.9	94.4 ± 13.4		
Change	0.2 ± 14.0	2.5 ± 12.2	-1.7 (-3.4 – -0.1)	0.038
*DA: Dental anxiety, SBP: Systolic blood pressures, DBP: Diastolic blood pressures, PR: Pulse rates, and PFS: Pit and fissure sealant; Plus–minus value is Mean ± SD.				
[#] For DA, the result is the odds ratio and 95% confidence interval; For SBP, DBP and PR, the results are the mean changes and 95% confidence interval.				
[§] For DA, the P value was calculated with a mixed-effect logistic regression, which included a random cluster effect of class, and covariates of gender and school; while for changes of SBP, DBP and PR, P values were calculated by linear mixed-effect models, with a random cluster effect of class, and covariates of gender, school and baseline value.				
Figs				

Secondary Outcomes

The mean SBP measured in the waiting room before the PFS was 108.7 mmHg in the EL group and 107.8 in the TSD group. The mean change of SBP from before to after the PFS was – 2.7 in the EL and 0.7 in the TSD group, with a -2.6 mmHg adjusted mean difference (95% CI: -4.4 to -0.9 mmHg; P = 0.036). The mean DBP after the procedure increased in both groups, but by 1.7 mmHg in the EL group and 4.5 mm Hg in the TSD group, an adjusted mean difference of -2.0 mmHg (95% CI: -4.4 to 0.3 mmHg; P = 0.086) (Table 2).

The mean PR in the EL group increased by 0.2 Beats Per Minute (BPM), while in the TSD group it increased by 2.5 BPM, for an adjusted mean difference of -1.7 BPM (95% CI: -3.4 to -0.1 BPM; P = 0.038) (Table 2).

Detailed data of SBP, PR and DBP before and after the PFS according to the class cluster were illustrated (Fig. 2).

Discussion

To our knowledge, this is the first trial to assess the use of a school-based experiential learning intervention to reduce children's dental anxiety. On average, there were fewer children with dental anxiety, as defined by a score of ≥ 38 on the Modified CFSS-DS scale, in the EL group compared to the TSD group, with significantly lower increases in the SBP and PR after the PFS procedure.

There is evidence to suggest that classical conditioning plays a major role in the development of DA [1]. Children who have had a negative dental experience will have more DA [28], and can lead to DA in adulthood [29]. When enter an unfamiliar circumstance, the level of children's anxiety will increase. In the conventional TSD management, the children only listened and observed the process of treatment immediately before they were subjected to and might not have been emotionally ready for the treatment.

In contrast, in this study, the EL intervention educated the children about the dental procedures and invited them to role play in a simulated dental clinic a month before their dental visit. It allowed them to become familiar with the common dental tools and experience a simulated dental procedure, making them feel in control and will improve their self-efficacy. Similar to the EL intervention, Radhakrishna et al. modified the TSD technique by adding a component of learning through playing [30], and achieved lower DA scores. Meanwhile, in contrast to the clinic-based intervention, school-based EL intervention can be conveniently performed at any time before the dental visit, which can be accessed by massive scale of children.

In our trial we chose a cut-off value of 38 on the CFSS-DS to define DA, which was also used in the systemic review by Cianettic et al. [2]. The reduction of prevalence of DA obtained in our study, although not achieving the designed 10 percentage-point difference, was similar to that obtained during moderately invasive treatment with other non-pharmacologic techniques such as using modelling [19] or audiovisual distraction [31].

The blood pressure and pulse rate, objective indicators of anxiety during the dental procedure, would be expected to increase during the treatment and slowly descend after it concluded [32]. In this trial, we found that the SBP and PR increased less after the PFS in the EL group than in TSD group, and the mean differences of changes between the two groups were statistically significant. It is possible that the reductions in DA, BP and PR varied depending upon which role the children played in the EL (i.e. dentist, nurse, parent or patient), but we were not able to evaluate this.

Due to several considerations, we chose children in the second year of primary school as the target group for the EL intervention. These children are with high incidence of dental caries, some of them will probably visit the dentists soon. Also, they have better understanding than kids in kindergarten. And it is right before the planned dental prophylaxis of pit and fissure sealing for the first permanent molar, the intervention can be conveniently executed and evaluated.

Conclusions

School-based experiential learning intervention before a dental visit is feasible and effective in lowering children's dental anxiety. In Shenzhen, or other cities in China, the children receive health education presentations regularly at school. We believe that simply adding a special oral health education and short session of role playing in a simulated dental clinic in the class can have a significant effect in reducing DA in children. Not only will this make dental procedures easier and less traumatic for the dentists, nurses and children, but it could also reduce the potential for DA in adults.

List Of Abbreviations

BPM: Beats per minute;

BP: Blood pressures;

DA: Dental anxiety;

DBMP: Dental behavior management problems;

DBP: Diastolic blood pressures;

EL: Experiential learning;

Modified CFSS-DS: Modified children's fear survey schedule-dental subscale;

PFS: Pit and fissure sealant;

PR: Pulse rates;

SBP: Systolic blood pressures;

TSD: Tell-Show-Do.

Declarations

Ethics approval and consent to participate

This study had the approved by the Ethics Committee of the Shenzhen Nanshan Maternal and Child Health Care Hospital. All parents of the children participated provided written informed consent.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was granted by Scientific and Technology Research and Development fund (No. JCYJ20170306155909623) from the Science and Technology Innovation Committee of Shenzhen City.

Authors' contributions

MMZ contributed to conception, design, project coordination, data acquisition, analysis and interpretation, drafted and critically revised the manuscript; HBY, HWL contributed to conception, design, arranged and supervised the process of PFS, data interpretation, and critically revised the manuscript; BX contributed to design, guided measurement of the CFSS-DS, and critically revised the manuscript; QH, HML contributed to study design, data acquisition and analysis, and drafted the manuscript. JS contributed to design, intervention arrangement, data acquisition, and drafted the manuscript; XQL contributed to funding obtain, conception, design, project coordination, data acquisition and interpretation, and critically revised the manuscript. All authors gave their final approval and agree to be accountable for all aspects of the work.

Acknowledgements

Many thanks to analysis suggestions and language editing form Howard E. Takiff in Shenzhen Nanshan Center for Chronic Disease Control and anyone who helped the implementation of the project.

References

1. Seligman LD, Hovey JD, Chacon K, Ollendick TH. Dental anxiety: An understudied problem in youth. *Clin Psychol Rev.* 2017;55:25-40.
2. Cianetti S, Lombardo G, Lupatelli E, Pagano S, Abraha I, Montedori A, et al. Dental fear/anxiety among children and adolescents. A systematic review. *Eur J Paediatr Dent.* 2017;18(2):121-30.

3. Klingberg G, Broberg AG. Dental fear/anxiety and dental behaviour management problems in children and adolescents: a review of prevalence and concomitant psychological factors. *Int J Paediatr Dent*. 2007;17(6):391-406.
4. Lin CS, Wu SY, Yi CA. Association between Anxiety and Pain in Dental Treatment: A Systematic Review and Meta-analysis. *J Dent Res*. 2017;96(2):153-62.
5. Stenebrand A, Hakeberg M, Helkimo AN, Koch G, Boman UW. Dental anxiety and oral health in 15-year-olds: a repeated cross-sectional study over 30 years. *Community Dent Health*. 2015;32(4):221-5.
6. Torriani DD, Ferro RL, Bonow ML, Santos IS, Matijasevich A, Barros AJ, et al. Dental caries is associated with dental fear in childhood: findings from a birth cohort study. *Caries Res*. 2014;48(4):263-70.
7. Zinke A, Hannig C, Berth H. Comparing oral health in patients with different levels of dental anxiety. *Head Face Med*. 2018;14(1):25.
8. Alsadat FA, El-Housseiny AA, Alamoudi NM, Elderwi DA, Ainoso AM, Dardeer FM. Dental fear in primary school children and its relation to dental caries. *Niger J Clin Pract*. 2018;21(11):1454-60.
9. Ashley PF, Chaudhary M, Lourenco-Matharu L. Sedation of children undergoing dental treatment. *Cochrane Database Syst Rev*. 2018;2018(12):CD003877.
10. Armfield JM, Heaton LJ. Management of fear and anxiety in the dental clinic: a review. *Aust Dent J*. 2013;58(4):390-407.
11. Goettems ML, Zborowski EJ, Costa FD, Costa VP, Torriani DD. Nonpharmacologic Intervention on the Prevention of Pain and Anxiety During Pediatric Dental Care: A Systematic Review. *Acad Pediatr*. 2017;17(2):110-9.
12. Anthonappa RP, Ashley PF, Bonetti DL, Lombardo G, Riley P. Non-pharmacological interventions for managing dental anxiety in children. *Cochrane Database Syst Rev*. 2017;2017(6):CD012676.
13. Strom K, Ronneberg A, Skaare AB, Espelid I, Willumsen T. Dentists' use of behavioural management techniques and their attitudes towards treating paediatric patients with dental anxiety. *Eur Arch Paediatr Dent*. 2015;16(4):349-55.
14. Kantaputra PN, Chiewcharnvalijkit K, Wairatpanich K, Malikaew P, Aramrattana A. Children's attitudes toward behavior management techniques used by dentists. *J Dent Child (Chic)*. 2007;74(1):4-9.
15. Peretz B, Kharouba J, Blumer S. Pattern of parental acceptance of management techniques used in pediatric dentistry. *J Clin Pediatr Dent*. 2013;38(1):27-30.
16. Boka V, Arapostathis K, Vretos N, Kotsanos N. Parental acceptance of behaviour-management techniques used in paediatric dentistry and its relation to parental dental anxiety and experience. *Eur Arch Paediatr Dent*. 2014;15(5):333-9.
17. Samra-Quintero PA, Bernardoni-Socorro C, Borjas AM, Fuenmayor NR, Estevez J, Arteaga-Vizcaino M. Changes in blood pressure in children undergoing psychological treatment before dental procedures. *Acta Odontol Latinoam*. 2006;19(1):9-12.

18. Farhat-McHayleh N, Harfouche A, Souaid P. Techniques for managing behaviour in pediatric dentistry: comparative study of live modelling and tell-show-do based on children's heart rates during treatment. *J Can Dent Assoc.* 2009;75(4):283.
19. Karekar P, Bijle MN, Walimbe H. Effect of Three Behavior Guidance Techniques on Anxiety Indicators of Children Undergoing Diagnosis and Preventive Dental Care. *J Clin Pediatr Dent.* 2019;43(3):167-72.
20. Kolb DA. *Experiential Learning: Experience as the Source of Learning and Development.* Second ed. USA: Pearson FT Press; 2014 July 2015.
21. Dyck RA, Kleinman NJ, Funk DR, Yeung RO, Senior P, Yardley JE. We Can Work (It) Out Together: Type 1 Diabetes Boot Camp for Adult Patients and Providers Improves Exercise Self-Efficacy. *Can J Diabetes.* 2018;42(6):619-25.
22. Reagan L, Pereira K, Jefferson V, Evans Kreider K, Totten S, D'Eramo Melkus G, et al. Diabetes Self-management Training in a Virtual Environment. *Diabetes Educ.* 2017;43(4):413-21.
23. Battjes-Fries MCE, Haveman-Nies A, Zeinstra GG, van Dongen EJI, Meester HJ, van den Top-Pullen R, et al. Effectiveness of Taste Lessons with and without additional experiential learning activities on children's willingness to taste vegetables. *Appetite.* 2017;109:201-8.
24. Matthews JI, Zok AV, Quenneville EP, Dworatzek PD. Development and implementation of FRESH—a post-secondary nutrition education program incorporating population strategies, experiential learning and intersectoral partnerships. *Can J Public Health.* 2014;105(4):e306-11.
25. Angelopoulou MV, Kavvadia K, Taoufik K, Oulis CJ. Comparative clinical study testing the effectiveness of school based oral health education using experiential learning or traditional lecturing in 10 year-old children. *BMC Oral Health.* 2015;15:51.
26. Lu JX, Yu DS, Luo W, Xiao XF, Zhao W. [Development of Chinese version of children's fear survey schedule-dental subscale]. *Zhonghua Kou Qiang Yi Xue Za Zhi.* 2011;46(4):218-21.
27. Buchanan H, Niven N. Validation of a Facial Image Scale to assess child dental anxiety. *Int J Paediatr Dent.* 2002;12(1):47-52.
28. Townend E, Dimigen G, Fung D. A clinical study of child dental anxiety. *Behav Res Ther.* 2000;38(1):31-46.
29. Oliveira MA, Vale MP, Bendo CB, Paiva SM, Serra-Negra JM. Influence of negative dental experiences in childhood on the development of dental fear in adulthood: a case-control study. *J Oral Rehabil.* 2017;44(6):434-41.
30. Radhakrishna S, Srinivasan I, Setty JV, D RM, Melwani A, Hegde KM. Comparison of three behavior modification techniques for management of anxious children aged 4-8 years. *J Dent Anesth Pain Med.* 2019;19(1):29-36.
31. Liu Y, Gu Z, Wang Y, Wu Q, Chen V, Xu X, et al. Effect of audiovisual distraction on the management of dental anxiety in children: A systematic review. *Int J Paediatr Dent.* 2019;29(1):14-21.
32. Brand HS. Cardiovascular responses in patients and dentists during dental treatment. *Int Dent J.* 1999;49(1):60-6.

Figures




Figure 1

Flow diagram of the trial of EL vs TSD for reducing children's dental anxiety. EL: experiential learning, TSD: Tell-show-do.




Figure 2

SBP, DBP and PR before and after PFS according to class clusters. SBP: Systolic blood pressures, DBP: Diastolic blood pressures, PR: Pulse rates, and PFS: Pit and fissure sealant. The dots indicate the mean values, and I bars indicate the upper confidence value (mean plus 1.96 times the standard error) and the lower confidence value (mean minus 1.96 times the standard error). Those with green color represent values before PFS, and red indicate values after PFS.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CONSORTExtensionforClusterTrials2012Checklist.pdf](#)