Silencing of The *Slzf-31* Gene Decreases The Salt Stress Tolerance And Drought Tolerance Of Tomato

Tong Pei

Northeast Agricultural University

Yufang Bao

Northeast Agricultural University

Tairu Wu

Northeast Agricultural University

Ziyu Wang

Northeast Agricultural University

Yue Wang

Northeast Agricultural University

Qifeng Liu

Northeast Agricultural University

Huanhuan Yang

Northeast Agricultural University

Jingbin jiang

Northeast Agricultural University

He Zhang

Northeast Agricultural University

Jingfu Li

Northeast Agricultural University

Tingting Zhao

Northeast Agricultural University

Xiangyang Xu (✉ xy709@126.com)

Northeast Agricultural University

Research Note

Keywords: SIZF-31 gene, VIGS, salt stress, drought stress

Posted Date: February 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-172139/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Plant Cell, Tissue and Organ Culture (PCTOC) on March 19th, 2021. See the published version at <https://doi.org/10.1007/s11240-021-02048-9>.

Abstract

The *SIZF-31* gene is a member of the tomato C2H2 transcription factor family. Previous studies have shown that *SIZF-31* gene expression is upregulated under drought stress and salt stress, but the specific function of this gene in tomato plants in response to these two kinds of stress is still unclear. To further explore the function of the *SIZF-31* gene in tomato under drought stress and salt stress, we employed the virus-induced gene silencing (VIGS) method to reduce the expression of the *SIZF-31* gene in tomato. The results showed that TRV2-*SIZF-31* plants had higher levels of wilt and stem bending than CK and CK-TRV2 plants under drought and salt stress. The ABA content of TRV2-*SIZF-31* plants were lower than those of CK and CK-TRV2 plants. The analysis of physiological indexes showed that the SOD and POD activity and the PRO content of TRV2-*SIZF-31* plants were lower than those of CK and CK-TRV2 plants, while the MDA content of TRV2-*SIZF-31* plants was higher than those of CK and CK-TRV2 plants. The accumulation of H_2O_2 and O_2^{2-} in TRV2-*SIZF-31* plants was greater than those in CK and CK-TRV2 plants. The values of the chlorophyll fluorescence parameters (ΦII and qL) of TRV2-*SIZF-31* plants were significantly lower than those of CK and CK-TRV2 plants. These results showed that the silencing of the *SIZF-31* gene reduces the drought resistance and salt tolerance of tomato.

Introduction

Tomatoes are one of the important vegetables in the world, and they inevitably encounter adverse environmental factors during their growth and development, including both biological factors and nonbiological factors. Among these factors, drought causes the greatest economic losses in plants among all abiotic stresses (Placide et al. 2014) and can have a significant effect on water metabolism in tomato plants. For example, drought stress can reduce the stomatal opening rate to cause morphological, physiological and biochemical changes in tomato plants, thus hindering the normal growth and development of tomatoes (Torres-Ruiz et al. 2015). Tomatoes are a plant that is sensitive to salt. Salt treatment at an appropriate concentration can modulate the flavor, color and soluble substance content of tomato, which is beneficial for improving the sugar-acid ratio of tomato fruit and, thus, the quality of tomato. However, high salt concentrations often lead to slow growth and decrease the yield and quality of tomato (Xu et al. 2015). Therefore, salt stress disrupts the ion balance and cell structure in tomato, limits the absorption and utilization of water and nutrients in tomato roots, and thereby affects the growth and development of tomato. When plants suffer from drought and salt stress, their content of abscisic acid (ABA) will increase, and the ABA hormone signaling pathway will be activated, in turn activating corresponding transcription factors (TFs). TFs activate gene expression through cis-acting elements to improve the tolerance of plants to the stress response pathway (Wang et al. 2011). There are many TFs related to abiotic stress in the whole tomato genetic system, among which the C2H2 zinc finger protein family is composed of TFs associated with abiotic stress.

The C2H2 zinc finger protein forms a zinc finger structure composed of approximately 25–30 amino acids. The conserved amino acid sequence is C-X2 ~ 4-C-X3-P-X5-L-X2-H-X3 ~ 5-H (C: cysteine, H: histidine, X: amino acid, P: phenylalanine, L: leucine) (Shimeid 2008). Two cysteines and two histidines

combine with zinc ions to form a coordination bond. These amino acid sequences can form a compact finger-like tetrahedral structure in the presence of coordination bonds and remain stable (Pabo al. 2001). Within the zinc finger structure, there is a highly conserved sequence (QALGGH) located at the junction with DNA, and this conserved amino acid sequence is unique to plant zinc finger proteins, which indicates that C2H2 zinc finger proteins may play a regulatory role in the unique growth and development of plants (Huang et al. 2005).

In 1992, the first EPF1 gene of the C2H2 zinc finger was cloned from the petunia, and in recent years, the study of plant-type C2H2 zinc finger proteins in tobacco, Arabidopsis and tomato has been reported in plants; for example, C2H2-type zinc finger proteins have been found to regulate plant growth and development and to play an important role in abiotic stress (Amarjeet et al. 2013; Englbrecht et al. 2004). The overexpression of the TaZFP1B gene of C2H2 type zinc indicating protein in wheat, can stimulate oxidative reactions and hormone signal transduction pathways in plants and increase the drought tolerance of wheat (Cheuk et al. 2020). We previously performed a genome-wide analysis of the C2H2-ZFP TF family and found that 7 and 5 genes were specifically expressed during drought stress and salt stress, respectively (Zhao et al. 2020).

Virus-induced gene silencing (VIGS) is a genetic technique that inhibits the expression of endogenous genes in plants by inserting recombinant viruses into target gene fragments and is used mainly for the functional analysis of genes (Benedito et al. 2004). VIGS technology has been widely used in the study of plant gene function in fields such as adversity stress, disease defense and plant growth research during the 21st century (Benedito et al. 2004; Anand et al. 2007; Senthil-Kumar et al. 2007). According to a survey of the literature, this technology has been applied in more than 30 kinds of plants, such as Arabidopsis, rice, tobacco, tomato, cucumber, cabbage and other crops (Becker and Lange 2009). In recent years, the use of VIGS technology in tomato has also been increasing. In the present study, VIGS technology was used to reduce the expression level of the SlZF-31 gene to analyze and study the function of this gene under abiotic stress (drought stress and salt stress), which provides a theoretical basis and is of practical significance for future research on the function of this gene and C2H2 zinc finger proteins in general.

Materials And Methods

Plant materials

Seeds of the tomato variety "Moneymaker" were provided by the Tomato Research Institute of Northeast Agricultural University (Harbin, China) and were seeded in a multihole tray and incubated in the dark at 25°C with 75% humidity for 5 d until seedling emergence. The seedlings were cultured in incubators at 30°C with 75% humidity under a 16 h light, 8 h dark cycle until the first real leaf emerged, and a fraction of the seedlings were used for the VIGS experiment. The remaining tomato seedlings were treated with the ABA hormone when they had been cultured to the four-leaf stage.

VIGS vector construction for the SLZF-31 gene and preparation of the Agrobacterium suspension

Total RNA was extracted from tomato leaves via the TRIzol method. The specific steps were modified according to the instructions provided for the reagent by Invitrogen. cDNA extraction was performed in accordance with the cDNA Synthesis Kit of Vazyme Biotech Co., Ltd. The primers for the gene *SLZF-31* were designed according to the mRNA sequence of the *SLZF-31* gene (Soly08g063040.2.1) in the SGN tomato database. All primers employed in this study were designed using primer 5.0 software (Table 1). The target fragment of the *SLZF-31* gene was subjected to homologous recombination with the tobacco rattlesnake virus (pTRV2) vector, in which the specific steps were performed according to the ClonExpressII One Step Cloning Kit of Vazyme Biotech Co., Ltd. The ligated vector was transferred to an *Escherichia coli* DH5a culture in accordance with the instructions of the *Escherichia coli* transformation kit, and single colonies were selected for expansion in liquid LB medium (50µg/mL kanamycin) and then sequenced. The sequenced vector pTRV2-*SLZF-31* was transferred to *Agrobacterium* for the preparation of the immersion solution according to the method of Huang et al. (Huang et al. 2014). Single colonies were picked transferred to liquid LB medium (50µg/mL kanamycin, 50µg/mL rifampicin). Then, the bacteria were transferred to IM medium (50µg/mL kanamycin, 50µg/mL rifampicin, 200µM acetyleugenone) for 24 h, after which they were collected and transferred to IM medium (200µM acetyleugenone) containing MgCl2 to obtain an infection solution with an OD600 of 0.2–0.3.

Table 1
Primers used for target fragment amplification and qRT-PCR analysis

Primer name	(Restriction site sequence) Primer sequence (5'–3')
<i>SLZF-31</i> primer-F	(CGGAATTCCG)AGTGTCACCCCTGGGTTCAT
<i>SLZF-31</i> primer-R	(CGGGATCCCG)AGAGCAGCAGGTGGTGTAGT
<i>SLZF-31</i> primer-F(qRT-PCR)	GATGAAGGATTTGGTCGCGG
<i>SLZF-31</i> primer-R(qRT-PCR)	AAGAGCAGCAGGTGGTGTAG
<i>Actin</i> primer-F	GTCCTCTTCCAGCCATCCAT
<i>actin</i> primer-R	ACCACTGAGCACAATGTTACCG

Infection of tomato plants to reduce the SLZF-31 gene expression level

Agrobacterium infection was carried out on tomato seedlings via vacuum infiltration with a syringe; 50 seedlings were infected at a time, among which 10 were used as the CK control plant, 10 were used at the CK-TRV2 control, 15 were used for the drought stress treatment, and 15 were used for the salt stress treatment. The experiment was repeated 3 times. The experimental steps were performed in accordance with the method described by Velasquez et al. (Velasquez et al. 2009).

Drought And Salt Stress Treatment And Sample Collection

SIZF-31-silenced tomato plants at 25 d after infection were subjected to drought stress and salt stress (Zhang et al. 2014). The tomato plants (CK, CK-TRV₂ and *SIZF-31*-TRV₂) were treated with 15% PEG 6000 to simulate drought, and tomato leaves were collected at 0 h, 1.5 h, 3 h, 6 h and 12 h after treatment. Then, the tomato seedlings were treated with 100 mM NaCl as the salt treatment, and tomato leaves were collected at the same 5 time points after treatment. The collected tomato leaves were used to detect various indicators in the VIGS experiment.

***SIZF-31* gene expression pattern analysis**

The silencing efficiency of the *SIZF-31* gene was detected in tomato leaves collected at 25 d after infection. In this study, we analyzed *SIZF-31* gene expression patterns in tomato leaves at different time points (0 h, 1.5 h, 3 h, 6 h and 12 h) after the silencing efficiency and expression levels of the *SIZF-31* gene in TRV2-*SIZF-31* tomato leaves were also analyzed by qRT-PCR at different time points (0 h, 1.5 h, 3 h, 6 h and 12 h) after the initiation of drought stress and salt stress. The same methods described above were employed for RNA extraction and cDNA synthesis from tomato leaves. The qRT-PCR method specifically followed that described by Zhao et al. (Zhao et al. 2015; Zhao et al. 2016). The primers used in the qRT-PCR analysis were designed by using the primer design software Primer 5.0 (Table 1). According to the CT values obtained with the qTOWER3G system (Analytik Jena, Germany), the relative expression of genes was calculated according to the $2^{-\Delta\Delta CT}$ method with *actin* as the internal reference gene (Livak 2001; Rotenberg et al. 2006). Bar charts were generated with Office 2016 software.

Determination Of Aba Content

The content of ABA hormone in CK, CK-TRV₂ and TRV2-*SIZF-31* tomato plants under drought and salt stress (0h, 3h and 12h) was determined by high performance liquid chromatography (HPLC). For specific ABA hormone detection methods, refer to the 2018 article by Li et al (Li et al. 2018).

Determination Of Physiological Indexes (sod, Pod, Pro And Mda)

The physiological indexes of superoxide dismutase (SOD) and peroxidase (POD) activity and proline (PRO) and malondialdehyde MDA content were determined in tomato leaves (CK, CK-TRV₂, TRV2-*SIZF-31*) after drought stress and salt stress. The SOD activity in tomato leaves was determined according to Giannopolitis and Ries (1977). The POD activity in tomato leaves was determined according to Chance and Maehly (1955). The PRO content in tomato leaves was determined according to Cakmak and Horst (1991). The MDA content was determined with the thiobarbital acid method (Ali et al. 2017). These kits were purchased from Suzhou Keming Biotechnology Co., Ltd. A 0.2 g tomato leaf sample was weighed, and the extract was obtained by centrifugation, followed by incubation in a water bath and static treatment. The colorimetric results were quantified at OD560 nm for SOD, 470 nm for POD, 520 nm for PRO and 532 nm and 600 nm for MDA.

Nbt And Dab Staining

Tomato leaves (CK, CK-TRV2, TRV2-*SIZF-31*) under drought stress and salt stress were assessed by NBT and DAB staining. NBT (0.1 g) was dissolved in 50 ml of phosphate buffer, and 50 mg DAB was dissolved in 50 ml of distilled water, after which the pH was brought to 7.5. The leaves were placed in the NBT and DAB solutions overnight, and an appropriate amount of ethanol was added, followed by boiling in a water bath at 100°C for 15 min (Rao and Davis 1999; Freyeret et al. 2002). Finally, the results were observed on glass slides. A blue precipitate was observed in the tomato leaves after NBT staining, where a greater intensity of the blue precipitate indicated a higher $O_2^{\cdot -}$ content. A brown precipitate was observed in the tomato leaves after DAB staining, where a greater intensity of the brown precipitate indicated a higher H_2O_2 content.

Chlorophyll Fluorescence Detection

This evaluation was conducted under drought stress and salt stress at different time points (0 h, 1.5 h, 3 h, 6 h and 12 h) in CK, CK-TRV2 and TRV2-*SIZF-31* plants with a PhotosynQ MultispeQ multifunction plant measurement instrument made in the United States to measure the chlorophyll fluorescence parameters (PSII quantum yield (ΦII), non-photochemical quenching ($\Phi(NPQ)$), photochemical quenching (qL) and electron transfer efficiency (ETR)).

Results

Results of *SIZF-31* gene silencing efficiency analysis

The expression level of the *SIZF-31* gene in TRV2-*SIZF-31* plants was detected by qRT-PCR. Figure 1 shows the gene expression levels of partially silenced plants, and the results showed that the expression level of the *SIZF-31* gene in 10 TRV2-*SIZF-31* plants was lower than that in CK and CK-TRV₂ plants. *SIZF-31* gene expression in TRV2-*SIZF-31* plants was decreased by 42.6%-85%, where the greatest decrease was 85%, and the average decrease was 79.4%. Tomato plants in which gene expression was decreased by more than 50% were selected for subsequent drought stress and salt stress treatment.

Expression pattern analysis of the *SIZF-31* gene after drought stress and salt stress

The expression pattern of the *SIZF-31* gene at different time points (0 h, 1.5 h, 3 h, 6 h, 12 h and 24 h) after drought stress and salt stress was detected by qRT-PCR. The results showed that the expression of the *SIZF-31* gene in CK, CK-TRV2 and TRV2-*SIZF-31* plants first increased and then decreased under drought stress and salt stress treatment, and that its expression was always lower in TRV2-*SIZF-31* plants than in CK and CK-TRV2 plants. These results indicated that VIGS exerted a significant silencing effect on tomato *SIZF-31* (Fig. 2).

Phenotypic observations of tomato plants under drought stress and salt stress

The phenotypes of CK, CK-TRV2 and TRV2-*SIZF-31* plants were observed after PEG 6000 drought stress treatment, as shown in Fig. 3. CK and CK-TRV2 plants showed leaf wilting after 6 h of drought stress, which was aggravated after 12 h and was more serious after 24 h. The leaves of TRV2-*SIZF-31* plants were wilted after 1.5 h of drought treatment, and the leaves were wilted and dried from 3 h to 12 h. The stem became increasingly bent, the degree of leaf wilting reached its highest level, a portion of the leaves dried out, and the stem exhibited serious bending after 24 h of drought stress. CK, CK-TRV2 and TRV2-*SIZF-31* plants were observed after NaCl stress treatment. The CK plants showed leaf wilting after 12 h of treatment, and leaf wilting worsened after 24 h of treatment. CK-TRV2 plants showed leaf wilting after 3 h of treatment, and leaf wilting increased from 6 h to 24 h, with no obvious change in the stem. The leaves of TRV-*SIZF-31* plants began to wilt, and the stems were slightly bent after 3 h of salt stress. The degree of leaf wilting gradually increased from 6 h to 12 h, and after 24 h of treatment, the leaves had dried out, and the stems were severely bent. Thus, the *SIZF-31* gene plays a role in both drought and salt tolerance in tomato, and reducing the expression of the *SIZF-31* gene decreases the drought and salt tolerance of tomato.

Determination of ABA content in tomatoes

As shown in Fig. 4, The ABA content of TRV2-*SIZF-31* tomato silent plants before drought stress and salt stress is slightly lower than that of CK and CK-TRV2 plants, which indicates that TRV2-*SIZF-31* gene may be involved in the synthesis and accumulation of plant ABA hormone. After drought stress and salt stress, the ABA content in tomato plants (TRV2-*SIZF-31*, CK and CK-TRV2) showed a trend of first increasing and then decreasing, and the ABA content of TRV2-*SIZF-31* tomato plants was compared with CK and CK -TRV2 plants are significantly reduced.

Results of SOD, POD, PRO and MDA quantification

As shown in Fig. 5, the SOD and POD activities and the contents of PRO and MDA in CK, CK-TRV2 and TRV2-*SIZF-31* plants increased after drought and salt treatment. At all time points (0 h, 1.5 h, 3 h, 6 h and 12 h) during drought treatment, the SOD and POD activities and the content of PRO in TRV2-*SIZF-31* plants were lower than those in CK and CK-TRV2 plants, while the MDA content in TRV2-*SIZF-31* plants was higher than those in CK and CK-TRV2 plants. The results for the SOD activity, POD activity, PRO content and MDA content in plants (CK, CK-TRV2 and TRV2-*SIZF-31*) subjected to salt stress were similar to those obtained under drought stress. Therefore, the degree of cell damage was more severe in TRV2-*SLZF-31* plants than in CK and CK-TRV2 plants.

Results of NBT and DAB staining

As shown in Fig. 6, DAB and NBT staining was performed on CK, CK-TRV2 and TRV2-*SIZF-31* at different time points (0 h, 1.5 h, 3 h, 6 h, 12 h and 24 h) during drought stress and salt stress, respectively. The results showed that leaf color gradually deepened with the extension of the treatment time. The brown

precipitate and the blue precipitate appeared in the leaves of CK, CK-TRV2 and TRV2-*SIZF-31* plants stained with DAB and NBT, respectively, after drought stress. The area and color of the precipitates in the leaves of TRV2-*SIZF-31* plants were larger and darker than those in CK and CK-TRV2 plants, indicating that the accumulation of H_2O_2 and O^{2-} under drought stress was greater in TRV2-*SIZF-31* plants than in CK and CK-TRV2 plants. Under salt stress, the CK, CK-TRV2 and TRV2-*SIZF-31* plants stained with DAB and NBT showed similar characteristics, which proved that the accumulation of H_2O_2 and O^{2-} in *SIZF-31* plants was higher than that in CK and CK-TRV2 plants after salt stress. Therefore, reducing the expression level of the *SIZF-31* gene causes more damage to tomato under drought stress and salt stress.

Chlorophyll fluorescence parameter detection under drought stress and salt stress

As shown in Fig. 7, under drought stress, the chlorophyll fluorescence parameters (ΦII , qL and ETR) of CK, CK-TRV2 and TRV2-*SIZF-31* plants were reduced, among which the ΦII and qL values of TRV2-*SIZF-31* plants were significantly lower than those of CK and CK-TRV2 plants, and the ETR of TRV2-*SIZF-31* plants decreased the most. The Φ (NPQ) values of CK, CK-TRV2 and TRV2-*SIZF-31* plants were increased. The changes in the chlorophyll fluorescence parameters of CK, CK-TRV2 and TRV2-*SIZF-31* plants under salt stress were similar to those under drought stress.

Discussion

In the abiotic stress response, the regulatory network generated by the signal transduction pathways of different hormones is complex and diverse (Verma et al. 2016). ABA is one of the key hormones regulating plant growth, development and physiological activities, it not only regulates stomatal opening and controls organ shedding but also plays an important role in the response to adverse conditions and various types of stress (Yamaguchi-Shinozaki and Shinozaki 2006). When plants encounter drought, salt damage and other stresses, ABA levels increase, activating the ABA signal transduction pathway and inducing the expression of many genes related to stress resistance to reduce the damage to plants caused by stress and protect their normal growth (Wang et al. 2011). In this study, the ABA content of TRV2-*SIZF-31* plants that reduced the expression of the *SIZF-31* gene was lower than that of CK and CK-TRV2 plants before and after stress. Therefore, we speculate that the *SIZF-31* gene may be involved in the ABA hormone synthesis or metabolism pathway, so that this gene affects the drought and salt tolerance of tomato plants by affecting the signal transduction pathway of ABA hormone.

TRV2-*SIZF-31* plants showed no significant change in phenotype before stress treatment, indicating that the *SIZF-31* gene had no significant effect on the plant phenotype after silencing. We observed the phenotypes of CK, CK-TRV2 and TRV2-*SIZF-31* plants under drought and salt stress. It was found that the degrees of leaf wilting, drying and stem bending in TRV2-*SIZF-31* plants were more serious than those in CK and CK-TRV2 plants under drought stress and salt stress. Therefore, we speculate that the *SIZF-31* gene may play a role in both drought and salt tolerance in tomato, where reducing the expression of the gene *SIZF-31* will decrease the drought and salt tolerance of tomato.

ROS production is one of the earliest responses of plants to drought stress and salt stress. ROS damage the cell membrane, thus affecting cell metabolism. To remove ROS from plant cells and protect macromolecules in plant cells, a series of antioxidant systems involving reactive oxygen scavenging enzymes, such as SOD and POD, are activated in plants (Apel and Hirt 2004). SOD can convert superoxide in plants into H_2O_2 and O^{2-} to resist oxidative damage to plant cells (Ajithkumar et al. 2014). POD can protect the cell structure and reduce damage to the membrane system and plays a highly significant role in the maintenance of normal physiological activities under adverse conditions (Apel et al. 2004). PRO is an osmotic regulator that is accumulated by plants in the process of drought resistance and salt resistance and reflects the resistance abilities of plants to some extent. MDA is the final product of membrane peroxidation in plants, and the accumulation of MDA damages the structure and function of the cell membrane. The MDA content represents the degree of membrane lipid peroxidation and the degree of damage to the membrane system and is therefore an important indicator of stress resistance in the plant (Kishor et al. 2014). The results of this experiment showed that under drought stress and salt stress, the SOD and POD activities and the PRO content of TRV2-*SIZF-31* plants were higher than those of CK and CK-TRV2 plants. The MDA content of TRV2-*SIZF-31* plants was also higher than that of CK and CK-TRV2 plants, indicating that tomato plant cells suffered more damage after the expression of the *SIZF-31* gene was reduced. Similar results were found in a study by Zhao et al., who reported that the SOD and POD activity and the PRO content were lower in TRV2-*SL-ZH13* plants than in CK plants after *SL-ZH13* gene expression was reduced, while the content of MDA was higher than that of CK plants (Zhao et al. 2018; Zhao et al. 2019).

The accumulation of superoxide radicals (O^{2-}) in tomato leaves was examined by NBT staining. The higher the O^{2-} accumulation is, the darker the blue precipitate will be in leaves stained with NBT (Ivan et al. 2009). DAB staining was used to detect the accumulation of H_2O_2 . The higher the accumulation of H_2O_2 is, the more intense the brown precipitate will be in leaves stained with DAB (Fryer et al. 2002). The experimental results showed that after drought stress and salt stress, the blue precipitate in the leaves of TRV2-*SIZF-31* plants stained with NBT was darker than that in the leaves of CK and CK-TRV2 plants, and the brown precipitate in the leaves of TRV2-*SIZF-31* plants stained with DAB was darker than that in the leaves of CK and CK-TRV2 plants. Therefore, the accumulation of O^{2-} and H_2O_2 in TRV2-*SIZF-31* plants was higher than those in CK and CK-TRV2 plants. This phenomenon indicated that *SIZF-31* gene silencing did not directly affect the accumulation of reactive oxygen species in leaves, but after drought and salt stress, *SIZF-31* gene silencing led to more intense accumulation of reactive oxygen species in leaves.

Drought, salt and other abiotic stresses can degrade chlorophyll in plants, thus reducing the absorption and transmission of light energy by chloroplasts (Kuang et al. 1980). Chlorophyll fluorescence parameters can reflect the PSII plant photosynthetic system of light absorption and utilization (Demmig-Adams and Adams 1996). Φ represents the actual photosynthetic efficiency of PSII (the PSII quantum efficiency of energy conversion). QL reflects the proportion of open PSII reaction centers (i.e., the degree of openness of the PSII reaction center). Φ (NPQ) represents the PSII regulatory energy dissipation

quantum yield. ETR is the relative linear electron flow rate through the PSII optical system. In this study, the values of the chlorophyll fluorescence parameters (Φ_{II} and q_L) of TRV2-*SIZF-31* plants were found to be significantly lower than those of CK and CK-TRV2 plants under drought stress and salt stress, and the ETR of TRV2-*SIZF-31* plants showed the greatest decrease. We speculated that this might have occurred because of the decreased drought resistance and salt resistance of TRV2-*SIZF-31* plants after the *SIZF-31* gene was downregulated. Therefore, the chloroplast photosynthetic mechanism of TRV2-*SIZF-31* plants under abiotic stress was more severely damaged than those of CK and CK-TRV2 plants, and the photosynthetic capacity of TRV2-*SIZF-31* plants was significantly decreased. Lu C et al. showed that under abiotic stress, the plant chloroplast photosynthetic mechanism is damaged, the PSII original light energy conversion efficiency (Φ_{II}) is reduced, the photosynthetic ETR is reduced, the q_L light energy available for photosynthesis is reduced, and the excess excitation energy ($\Phi(NPQ)$) is increased, leading to a decrease in PSII function and photosynthetic capacity (Lu and Zhang 1999).

Prior to the abiotic stress treatment (drought stress and salt stress), there were no significant differences in the phenotype, physiological indexes (SOD, POD, PRO and MDA), NBT staining or DAB staining between control plants and TRV2-*SIZF-31* plants. After drought stress and salt stress, the phenotypes and related indexes of the control plants and TRV2-*SIZF-31*-silenced plants were significantly different, indicating that *SIZF-31* gene silencing had no significant influence on the growth and development of tomato plants, while this gene did play a role in plant resilience to stress.

Conclusion

In conclusion, the degrees of withering and stem bending in TRV2-*SIZF-31* plants were greater than those in CK and CK-TRV2 plants under drought stress and salt stress. The ABA content of TRV2-*SIZF-31* plants were lower than those of CK and CK-TRV2 plants. The analysis of physiological indexes showed that the SOD and POD activity and the PRO content of TRV2-*SIZF-31* plants were lower than those of CK and CK-TRV2 plants, while the MDA content of TRV2-*SIZF-31* plants was higher than those in CK and CK-TRV2 plants. The accumulation of H_2O_2 and O_2^- in TRV2-*SIZF-31* plants was higher than those in CK and CK-TRV2 plants. The values of the chlorophyll fluorescence parameters (Φ_{II} and q_L) of TRV2-*SIZF-31* plants were significantly lower than those of CK and CK-TRV2 plants, and the ETR of TRV2-*SIZF-31* plants showed the greatest decrease. These results indicated that the downregulation of the *SIZF-31* gene affected the response of tomato under drought and salt stress and reduced the drought and salt resistance of tomato plants.

Declarations

Funding:

This work was supported by the Hundreds of millions project science and technology major special project [grant numbers 2019ZX16B02]; the National Natural Science Foundation of China [grant numbers 32072589]; the China Agriculture Research System [grant numbers CARS-23-A-16]; the Postdoctoral

science foundation of heilongjiang province [grant numbers LBH-Z19046]; and the China postdoctoral science foundation grant [grant numbers 2018 M630333].

Author contributions:

Tong Pei: Conceived and wrote the manuscript;

Acknowledgments

This work was funded by Hundreds of millions project science and technology major special project (Grant No. 2019ZX16B02); National Natural Science Foundation of China (Grant No. 32072589); China Agriculture Research System (Grant No. CARS-23-A-16); Postdoctoral science foundation of heilongjiang province (Grant LBH-Z19046) and China postdoctoral science foundation grant (Grant No. 2018 M630333).

References

Ajithkumar IP, Panneerselvam R (2014) ROS scavenging system, osmotic maintenance, pigment and growth status of *Panicum sumatrense* Roth. under drought stress. *Cell Biochemistry & Biophysics* 68: 587–595

Ali Q, Javed MT, Noman A, Haider MZ, Waseem M, Iqbal N, Waseem M, Shah MS, Shahzad F, Perveen R (2017) . Assessment of drought tolerance in mung bean cultivars/lines as depicted by the activities of germination enzymes, seedling's antioxidative potential and nutrient acquisition. *Arch. Agron. Soil Sci* 64: 84–102

Amarjeet S, Poonam K, Amita P, Tyagi AK, Sopory SK, Sanjay K (2013) Comprehensive genomic analysis and expression profiling of phospholipase c gene family during abiotic stresses and development in rice. *Plos One* 8: 62494

Anand A, Vaghchhipawala ZR, Ryu CM, Kang L, Wang K, Del-PozoO, Martin GB, Mysore KS (2007) Identification and characterization of plant genes involved in *Agrobacterium*-mediated plant transformation by virus-induced gene silencing. *MPMI* 187: 41–54

Apel K, Hirt H (2004) Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annual Review of Plant Biology* 55: 373–399

Becker A, Lange M (2009) VIGS genomics goes functional. *Trends Plant Sci* 15: 1–4

Benedito VA, Visser PB, Angenent GC, Krens FA (2004) The potential of virusinduced gene silencing for speeding up functional characterization of plant genes. *Genet Mol Res* 3: 323–413

- Cakmak I, Horst WJ (1991) Effect of aluminium on lipid peroxidation, superoxide dismutase, catalase, and peroxidase activities in root tips of soybean (*Glycine max*). *Physiol Plant* 83: 463–468
- Chance B, Maehly A (1955) Assay of catalases and peroxidases. *Methods Enzymol* 2: 764–775
- Cheuk A, Ouellet F, Houde M (2020) The barley stripe mosaic virus expression system reveals the wheat c2h2 zinc finger protein tazfp1b as a key regulator of drought tolerance. *BMC Plant Biology* 20(1)
- Demmig-Adams B, Adams WW (1996) Xanthophyll cycle and light stress in nature: Uniform response to excess direct sunlight among higher plant species. *Planta* 198: 460-470
- Englbrecht CC, Schoof H, Bohm S (2004) Conservation, diversification and expansion of C2H2 zinc finger proteins in the *Arabidopsis thaliana* genome. *BMC Genomics* 5: 39
- Fryer MJ, Oxborough K, Mullineaux PM, Baker NR (2002) Imaging of photo-oxidative stress responses in leaves. *Journal of Experimental Botany* 53: 1249–1254
- Giannopolitis CN, Ries SK (1977) Superoxide dismutase occurrence in higher plants. *Plant Physiol* 59: 309–314
- Huang J, Wang J, Zhang H (2005) Rice ZFP15 gene encoding for a novel C2H2-type Zinc Finger Pprotein lacking DLN box, is regulated by spike development but not by abiotic stresses. *Molecular Biology Reports* 32: 177-183
- Huang Y, Mei M, Mao Z, Lv S, Zhou J, Chen S (2014) Molecular cloning and virus induced gene silencing of MiASB, in the southern root-knot nematode, *Meloidogyne incognita*. *European Journal of Plant Pathology* 138: 181-193
- Ivan C, Matthieu B, Cécile SC, Fanny R, Gwenola G (2009) Differential patterns of reactive oxygen species and antioxidative mechanisms during atrazine injury and sucrose-induced tolerance in *Arabidopsis thaliana* plantlets. *BMC Plant Biol* 9: 28
- Kishor PBK, Sreenivasulu N (2014). Is proline accumulation per se correlated with stress tolerance or is proline homeostasis a more critical issue. *Plant Cell Environ* 37: 300-311
- Kuang TY, Zuo PY, Chang CT, Lon SJ, Li TS, Hao HP, Lin SC (1980) Structure and function of chloroplast membranes. Effects of potassium and magnesium ions on the structure and the absorption spectrum of two kinds of chloroplast membranes. *Photobiochemistry Photobiophys* 1: 73-82
- Livak KJ (2001) Schmittgen Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25:402-408
- Li XY, Zhao QQ, Liu Y, Liu ZL, Yang JM, Zhang, GJ (2018) Determination of abscisic acid in begonia fruit by HPLC. *Northern horticulture* 4: 138-144

- Lu CM, Zhang JH (1999) Effects of water stress on photosystem II photochemistry and its thermostability in wheat plants. *Journal of Experimental Botany* 50: 1199-1206
- Pabo CO, Peisach E, Grant RA (2001) Design and selection of novel Cys2His2 zinc finger proteins. *Annual Review of Biochemistry* 70: 313-340
- Placide R., Hirut GB, Stephan N, Fekadu B (2014) Assessment of drought stress tolerance in root and tuber crops. *African Journal of Plant Science* 8: 214–224
- Rao MV, Davis KR (1999) Ozone-induced cell death occurs via two distinct mechanisms in Arabidopsis: The role of salicylic acid. *Plant Journal* 17: 603–614
- Rotenberg D, Thompson TS, German TL, Willis DK (2006) Methods for effective real-time RT-PCR analysis of virus-induced gene silencing. *Journal of Virological Methods* 138: 49-59
- Senthil-Kumar M, Hema R, Anand A, Kang L, Udayakumar M, Mysore KS (2007) A systematic study to determine the extent of gene silencing in *Nicotiana benthamiana* and other Solanaceae species when heterologous gene sequences are used for virus-induced gene silencing. *New Phytologist* 176: 782–91
- Shimeid SM (2008) C2H2 zinc finger genes of the Gli, Zic, KLF, SP, Wilms' tumour, Hucklebein, Snail, Ovo, Spalt, Odd, Blimp-1, Fez and related gene families from *Branchiostoma floridae*. *Development Genes and Evolution* 218: 639-649
- Torres-Ruiz JM, Antonio DE, Alfonso PM, Virginia HS (2015) Role of hydraulic and chemical signals in leaves, stems and roots in the stomatal behaviour of olive trees under water stress and recovery conditions. *Tree Physiology* 4: 415-424
- Velasquez AC, Chakravarthy S, Martin GB (2009) Virus-induced gene silencing (VIGS) in *Nicotiana benthamiana* and tomato. *Journal of Visualized Experiments* 28
- Verma V, Ravinran P, Kumar PP (2016) Plant hormone-mediated regulation of stress responses. *BMC Plant Biology* 16: 86
- Wang ZY, Xiong LM, Li WB, Zhu JK, Zhu JH (2011) The Plant Cuticle Is Required for Osmotic Stress Regulation of Abscissic Acid Biosynthesis and Osmotic Stress Tolerance in Arabidopsis. *Plant Cell* 23: 1971-1984
- Xu JN, Xu YQ, Yan Z, Zhang LY, Gao JW (2015) Research progress on response mechanism of tomato to abiotic stress. *Shandong agricultural science* 47: 120-124
- Yamaguchi-Shinozaki K, Shinozaki K (2006) Transcriptional Regulatory Networks in Cellular Responses and Tolerance to Dehydration and Cold Stresses. *Annual Review of Plant Biology* 57: 781-803

- Zhang ZZ, Chen XL, Guan X, Liu Y, Chen HY, Wang TT, Mouekouba LD, Li JF, Wang AX (2014) A genome-wide survey of homeodomain-leucine zipper genes and analysis of coldresponsive HD-Zip1 members' expression in tomato. *Bioscience Biotechnology and Biochemistry* 78: 1337–1349
- Zhao TT, Liu G, Li S, Li JF, Jang JB, Zhang H, Chen X, Xu XY (2015) Differentially expressed gene transcripts related to the *Cf-19*-mediated resistance response to *Cladosporium fulvum*, infection in tomato. *Physiological & Molecular Plant Pathology* 89: 8-15
- Zhao TT, Jiang JB, Liu G, He S, Zhang H, Chen X, Li JF, Xu XY (2016) Mapping and candidate gene screening of tomato *Cladosporium fulvum*-resistant gene *Cf-19*, based on high-throughput sequencing technology. *Bmc Plant Biology* 16: 51
- Zhao TT, Hu JK, Gao YM, Wang ZY, Xu XY (2018) Silencing of the *SL-ZH13* Transcription Factor Gene Decreases the Salt Stress Tolerance of Tomato. *Journal of the American Society for Horticultural Science* 143: 1–6
- Zhao TT, Wang ZY, Bao YF, Zhang XC, Yang HH, Zhang DY, Jiang JB, Zhang H, Li J F, Chen QS, Xu XY (2019) Downregulation of SL-ZH13 transcription factor gene expression decreases drought tolerance of tomato. *Journal of Integrative Agriculture* 18: 1579–1586
- Zhao TT, Wu TR, Wang ZY, Pei T, Yang HH, Li JF, Xu XY (2020) Genome-Wide Analyses of the Genetic Screening of C2H2-Type Zinc Finger Transcription Factors and Abiotic and Biotic Stress Responses in Tomato (*Solanum lycopersicum*) Based on RNA-Seq Data. *Frontiers in Genetics* 11: 540

Figures

Figure 1

Detection of the silencing efficiency of the SIZF-31 gene. Ten TRV2-SIZF-31 tomato plants were analyzed (VIGS-1 to VIGS-10); CK: control plants; CK-TRV2: empty vector control plants

Figure 2

Expression levels of SlZF-31 under drought stress and salt stress. a: Relative expression of the SlZF-31 gene under drought stress; b: Relative expression of the SlZF-31 gene under salt stress

Figure 3

Phenotypic observation of tomato plants under drought stress and salt stress. a: Phenotypic observation of tomato plants (CK, CK-TRV2 and TRV2-SlZF-31) under drought stress; b: Phenotypic observation of tomato plants (CK, CK-TRV2 and TRV2-SlZF-31) under salt stress

Figure 4

Determination of ABA content under drought stress and salt stress. a: ABA content under drought stress; b: ABA content under salt stress

Figure 5

Quantification of SOD, POD, PRO and MDA under drought stress and salt stress. a: SOD activity under drought stress; b: POD activity under drought stress; c: PRO content under drought stress; d: MDA content

under drought stress; e: SOD activity under salt stress; f: POD activity under salt stress; g: PRO content under salt stress; h: MDA content under salt stress

Figure 6

Images of the NBT and DAB staining of CK, CK-TRV2 and TRV2-SIZF-31 plants after drought stress and salt stress. a: DAB staining images of CK, CK-TRV2 and TRV2-SIZF-31 plants after drought treatment; b: NBT staining images of CK, CK-TRV2 and TRV2-SIZF-31 plants after drought treatment; c: DAB staining images of CK, CK-TRV2 and TRV2-SIZF-31 plants after salt treatment; d: NBT staining images of CK, CK-TRV2 and TRV2-SIZF-31 plants after salt treatment

Figure 7

Chlorophyll fluorescence parameters of the control group and gene-silenced group under drought stress and salt stress. a: Φ_{II} -drought stress; b: $\Phi(NPQ)$ -drought stress; c: q_L -drought stress; d: ETR-drought stress; e: Φ_{II} -salt stress; f: $\Phi(NPQ)$ -salt stress; g: q_L -salt stress; b: ETR-salt stress