

Geographical distribution of family physicians in Japan: A nationwide cross-sectional study

Shuhei Yoshida (✉ yoshida.shuhei.0810@gmail.com)

"Hiroshima Daigaku" <https://orcid.org/0000-0002-6747-1857>

Masatoshi Matsumoto

"Hiroshima Daigaku"

Saori Kashima

"Hiroshima Daigaku"

Soichi Soichi Koike

Jichi Ika Daigaku

Susumu Tazuma

"Hiroshima Daigaku"

Takahiro Maeda

Nagasaki Daigaku

Research article

Keywords: geography; health policy; Japan; family physician

Posted Date: July 1st, 2019

DOI: <https://doi.org/10.21203/rs.2.10782/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on October 29th, 2019. See the published version at <https://doi.org/10.1186/s12875-019-1040-6>.

Abstract

Background Geographical maldistribution of physicians, and their subsequent shortage in rural areas, has been a serious problem in Japan and in other countries. Family Medicine, a new board-certified specialty started 10 years ago in Japan by Japan Primary Care Association (JPCA), may be a solution to this problem. **Methods** We obtained the workplace information of 527 (78.4%) of the 672 JPCA-certified family physicians from an online database. From the national census data, we also obtained the workplace information of board-certified general internists, surgeons, obstetricians/gynaecologists and paediatricians and of all physicians as the same-generation comparison group (ages 30 to 49). Chi-squared test and residual analysis were conducted to compare the distribution between family physicians and other specialists. **Results** 519 JPCA-certified family physicians and 137,587 same-generation physicians were analysed. The distribution of family physicians was skewed to municipalities with a lower population density, which shows a sharp contrast to the urban-biased distribution of other specialists. The proportion of family physicians in non-metropolitan municipalities was significantly higher than that expected based on the distribution of all same-generation physicians ($p < 0.001$). **Conclusions** Family physicians distributed in favour of rural areas much more than any other specialists in Japan. The better balance of family physician distribution reported from countries with a strong primary care orientation seems to hold even in a country where primary care orientation is weak, physician distribution is not regulated, and patients have free access to healthcare. Family physicians comprise only 0.2% of all Japanese physicians. However, if their population grows, they can potentially rectify the imbalance of physician distribution. Government support is mandatory to promote family medicine in Japan.

Background

The geographical maldistribution of physicians, and their subsequent shortage in rural areas, is a serious social problem in Japan and internationally.¹ The dearth of physicians in rural areas has persisted in Japan despite various large-scale political interventions by the government such as the 'one medical school in each prefecture policy' in 1970s and 1980s and foundation of a special medical school, Jichi Medical University, for producing rural physicians in 1972²⁻⁴. This maldistribution has even worsened since 2004 when the new residency training programme for postgraduate year 1 and year 2 physicians was implemented nationwide^{5,6}.

Primary care physicians reportedly distribute better than other physicians in some countries with a strong primary care orientation, but it is unknown whether this is the case in a country like Japan with a weak primary care orientation, no regulation of physician distribution, and patients' free access to physicians^{7,8}. Primary care has not traditionally been recognised as an independent academic discipline in Japan, but in 2009 the Japan Primary Care Association (JPCA) introduced board certification as the only qualification for primary care physicians in the country⁹. However, family medicine remains

unpopular in Japan. The total number of JPCA-certified family physicians was only 672 (0.2%) of 311,205 physicians in 2018¹⁰.

Then a new board-certification system by the Japanese Medical Specialty Board, which is independent of existing specialist bodies, started from 2018 and 'general practice' was added as the 19th major clinical discipline. The certification of family physicians/general practitioners thus will undergo transition from JPCA to the Japanese Medical Specialty Board in 2021. With the reform of board certification, the number of trained general practitioners is expected to increase. In actuality, however, of the 8,217 applicants to specialist training programs in 2019, only 158 (1.9%) applied to 'general practice'¹¹. This means that the minority status of primary care physicians in Japan will remain largely unchanged.

Japan's national and local governments have very limited amount of authority to regulate the distribution of physicians or cap the number of physicians in cities. Consequently, physicians, regardless of their specialties, tend to concentrate in urban areas. Moreover, Japanese patients can choose any physician that they like, which encourages the movement of patients from rural to urban areas. This makes Japan an ideal place to examine the distribution of family physicians. If Japanese family physicians turn out to distribute better than other physicians, the fairness of the distribution would be more in the nature of family medicine than in that of the healthcare or economic system that regulates the distribution of primary care physicians.

In this study we evaluated the geographic distribution of Japanese family physicians and compared it with same-generation physicians of other specialties. We then discussed the potential effectiveness of promotion of family medicine in Japan for the more equal access of its population to healthcare.

Methods

Study setting

Nationwide

Data of JPCA-certified family physicians

We used an online database, current as of 31 July 2018, of 527 (78.4%) of the 672 JPCA-certified family physicians who gave the permission to the JPCA office¹⁰. The website discloses each physician's name, workplace (municipality and medical institution) and area of interest.

Data on other physicians in Japan

To estimate the age of the certified family physicians, we searched the year of physician license registration of each JPCA-certified family physician, using the open online database of the Ministry of Health, Labour and Welfare¹². We found the registration year of 467 (88.6%) physicians. As a result, their median (95th percentile) age in 2018 was estimated to be 37 (31-48). We extracted, as the comparison group of the family physicians, physicians between the ages of 30 and 49 from all physicians in the dataset of the 'Survey of Physicians, Dentists and Pharmacists' (Physician Census) conducted by the Ministry of Health, Labour and Welfare in December 2016. The Census contained information on the major board-certification for all registered physicians in Japan, but the information on JPCA certification was not included because family medicine was not recognised as a formal clinical discipline until 2018. Within the comparison group we identified all board-certified general internists (Fellow of the Japanese Society of Internal Medicine: *sougou-naika-senmoni*), surgeons (Japan Surgical Society), obstetricians/gynaecologists (Japan Society of Obstetrics and Gynecology), and paediatricians (Japan Pediatric Society) in the Census. Individual data in the Census was used with special permission of the Ministry (permission no. 0411-3).

Data on municipalities

Japan has three levels of government: national, prefectural and municipal. Municipalities comprise cities, towns and villages. We compared the rurality of workplace municipality of family physicians with that of the comparison group physicians. Using the data on population and land area in each municipality published by the Statistics Bureau, Ministry of Internal Affairs and Communications,¹³ we divided municipalities into quintiles sorted by population density (Quintile 1 \leq 433.14, Quintile 2 \leq 1195.59, Quintile 3 \leq 3155.03, Quintile 4 \leq 10845.34, Quintile 5 10845.34+ people per square kilometer) so that each quintile has 20% of all municipalities. To measure rurality in another way, we divided municipalities into 'metropolis,' 'city' and 'town/village.' 'Metropolis' was all of the ordinance-designated cities (*seirei-shitei-toshi*) and 23 special wards of Tokyo ($n = 171$). 'City' was the other cities (*shi*) ($n = 756$). 'Town/village' was towns (*cho*) and villages (*son*) ($n = 884$).

Statistical analysis

A chi-square test was conducted to compare the distribution of categorical data between two groups. Residual analysis was used to compare the real value and the expected value derived from the distribution of all physicians. All statistical analyses were performed using Microsoft Excel and STATA/SE version 15 (StataCorp, 2017).

Results

Eight family physicians were excluded due to lack of precise workplace information, leaving a total of 519 family physicians and 137,587 same-generation physicians (including 11,947 general internists, 11,570 surgeons, 5268 obstetricians/gynaecologists, and 6919 paediatricians).

Table 1 shows the distribution of each group of physicians among the quintiles of municipalities sorted by population density. The proportion of family physicians in municipalities with a low population density was significantly higher than the proportion expected based on the distribution of all same-generation physicians. General internists and surgeons distributed almost similarly to all the physicians, and obstetricians/gynaecologists and paediatricians distributed slightly more to urban areas than all physicians.

Figure 1 shows the number of board-certified physicians per 100,000 population classified by population density. The population of family physicians was tiny, but their distribution was skewed more to rural areas than to urban areas, which is quite contrary to the urban-biased distribution of physicians in other specialties.

Table 2 shows the distribution of each group of physicians among the three municipality types. The proportion of family physicians in rural municipalities was significantly higher than expected based on the distribution of all same-generation physicians. The distributions of physicians in other specialties were not largely different from the distribution of all physicians.

Figure 2 shows the number of board-certified physicians per 100,000 population classified by municipality types. As with the population density, the distribution of family physicians was much more rural-biased than the distributions of other specialties although the population of family physicians was tiny.

Discussion

Although the total number of family physicians was much lower than those of other physicians, family physicians in Japan distributed more to rural than to urban areas, in stark contrast to the urban-biased distribution of other physicians. This study suggests primary care physicians have a very favourable distribution even in a country with a poor primary care orientation, no regulation on the distribution of physicians, and free access of people to physicians.

The better distribution of family physicians and urban-biased distribution of other specialists were also observed in the United States and Europe¹⁴⁻¹⁷. In countries with a strong primary care orientation and a public sector-based healthcare system such as the United Kingdom and the Netherlands, everyone is

required to register with his/her general practitioner. It is quite reasonable, in such countries, for general practitioners to distribute according to population. In countries with a private sector-based healthcare system such as the United States, a substantial proportion of population is gate-kept by a family physician designated by each person's health insurance company. In such a country, it is not surprising that family physicians distribute more equally than other specialists. However, the Japanese healthcare system is unique. The Japanese do not need to register with a certain primary care physician. Their access to any physician is not limited by insurance companies or the government. All the Japanese are covered by public health insurance and they can choose their physicians of any specialty. In addition, physicians can choose where to practice without any governmental regulation. This study showed, even in Japan, primary care physicians distributed quite equally, or even disproportionately to rural areas.

Accessibility, both financial and geographical, is a principle of primary care¹⁸. To ensure the quality of primary care, physicians need to ensure equal and good accessibility to their patients¹⁹. The egalitarian principle of primary care might help family physicians to voluntarily distribute equally or more to places where they are in greatest demand.

Another potential reason for a rural-biased distribution of Japanese family physicians is the cost of practicing in urban areas, especially for young family physicians. In Japan, most clinics are managed by an individual or a family. A relatively few physicians are hired to work in clinics. The initial cost to establish a clinic is estimated, on average, at 94 million yen (758,000 euro), 20 million of which come from the physician's own savings²⁰. In Japan, early-career physicians are usually hired by a hospital and then open their own clinics in the middle or towards the end of their career. The median age of our study participants was 37. Because early-career physicians do not have 20 million yen in savings, if they become family physicians, they may be limited to working in public clinics established by municipalities in areas with a physician shortage. This might tilt the distribution of Japanese family physicians towards rural areas.

The results of this study suggest that the increase of family physicians might rectify the present urban-skewed distribution of physicians, which has long been a serious social problem in Japan³. However, the number of Japanese family physicians was negligible, because of the short history of JPCA certification and the lack of governmental support for increasing the population of family physicians. A small but important first step to the expansion of this group of physicians was the introduction of 'general practice' as one of 19 major clinical disciplines under the new training system for board certification starting in 2018. However, against expectations, the number of applicants to 'general practice' was only 1774 (2.1%) of the 8604 applicants in 2019. The low popularity is probably caused by low awareness among population and lack of governmental political support for general practice in Japan²¹.

Family physicians are pivotal in providing care in Japan where the population is rapidly ageing and more patients than ever are presenting with numerous chronic conditions. The government and professional bodies need to counteract the limited popularity of family medicine. By increasing the number of family

physicians, Japan will be in a better position to handle the multi- and complex-morbidity of patients and the geographic maldistribution of physicians in a quickly ageing society.

This study has the following limitations. First, the age of JPCA-certified family physicians was estimated. Therefore, we could not assert that physicians between the ages of 30 and 49 were the most appropriate comparison group. In addition, we could not adjust the factors related to physicians' choice of practice location such as gender and birthplace because these data were not available.

Article 25 of the Japanese Constitution states that everybody has a right to be healthy regardless of where they live or how much they earn. However, the geographic barrier to healthcare for rural residents has persisted despite half a century of financial and political investment by the government³. To improve the quality of care, streamline the provision of care, and especially to equalise the distribution of care, the Japanese government should increase the number of family physicians through national policies including offering financial incentives to medical students and physicians in training who hope to enter family practice.

Abbreviations

JPCA: The Japan Primary Care Association

SD: standard deviation

Declarations

Ethics approval and consent to participate: Ethical approval was granted by the Ethics Committee for epidemiological research at Hiroshima University (ref. no. 778) and the Research Ethics Committee of Nagasaki University Graduate School of Biomedical Sciences (ref. no. 13091342).

Consent for publication: Not applicable

Availability of data and material: Conditions of the ethical approvals permit the cohort office (Department of Community-Based Medical Systems, Graduate School of Biomedical and Health Sciences, Hiroshima University) and the suboffice (Department of Community Medicine, Nagasaki University Graduate School of Biomedical Science) to share the aggregated data with stakeholders or researchers.

Competing interests: The authors declare that they have no competing interests.

Funding: This study is funded by the Ministry of Education, Culture, Sports, Science and Technology KAKENHI Grant-in-Aid for Scientific Research (C), grant number (18K10084) and by the Satake Fund. This study was also funded by the Pfizer Health Research Foundation.

Authors' contributions: SY assisted with analysis and interpretation of data and writing of the draft. MM contributed to the study design, tools, study administration, data collection, analysis and interpretation of

data and writing of the draft. TM assisted with the study design, tools, study administration and writing of the draft. SaK conducted data analysis, data interpretation and writing of the draft. SoK and ST contributed to the study design, data collection and writing of the draft.

Acknowledgements: Not applicable

References

1. Makaroff LA, Green LA, Petterson SM, Bazemore AW. Trends in physician supply and population growth. *American family physician*. 2013;87(7):Online.
2. Kobayashi Y, Takaki H. Geographic distribution of physicians in Japan. *Lancet (London, England)*. 1992;340(8832):1391-1393.
3. Matsumoto M, Inoue K, Kajii E, Takeuchi K. Retention of physicians in rural Japan: concerted efforts of the government, prefectures, municipalities and medical schools. *Rural and remote health*. 2010;10(2):1432.
4. Matsumoto M, Inoue K, Bowman R, Noguchi S, Toyokawa S, Kajii E. Geographical distributions of physicians in Japan and US: Impact of healthcare system on physician dispersal pattern. *Health policy (Amsterdam, Netherlands)*. 2010;96(3):255-261.
5. Hara K, Otsubo T, Kunisawa S, Imanaka Y. Examining sufficiency and equity in the geographic distribution of physicians in Japan: a longitudinal study. *BMJ open*. 2017;7(3):e013922.
6. Toyabe S. Trend in geographic distribution of physicians in Japan. *International journal for equity in health*. 2009;8:5.
7. Agency for Healthcare Research and Quality [Internet]. The Distribution of the U.S. Primary Care Workforce. Content last reviewed July 2018. Agency for Healthcare Research and Quality, Rockville, MD. [cited 2019 Jan 25]. Available from: <https://www.ahrq.gov/research/findings/factsheets/primary/pcwork3/index.html>.
8. Matsumoto M, Inoue K, Bowman R, Noguchi S, Kajii E. Physician scarcity is a predictor of further scarcity in US, and a predictor of concentration in Japan. *Health policy (Amsterdam, Netherlands)*. 2010;95(2-3):129-136.
9. Onishi H, Park YS, Takayanagi R, Fujinuma Y. Combining Scores Based on Compensatory and Noncompensatory Scoring Rules to Assess Resident Readiness for Unsupervised Practice: Implications From a National Primary Care Certification Examination in Japan. *Academic medicine : journal of the Association of American Medical Colleges*. 2018;93(11S Association of American Medical Colleges Learn Serve Lead: Proceedings of the 57th Annual Research in Medical Education Sessions):S45-s51.
10. Japan Primary Care Association [Internet]. Certified system [cited 2018 Jan 25]. Available from: http://www.primary-care.or.jp/nintei_fp/fp_list.html.
11. Nihon Senmoni Kikou (Japanese Board of Medical Specialties)[Internet]. Application information about the training for each specialized area [cited 2019 May 7]. Available from:

<https://www.mhlw.go.jp/content/10803000/000491858.pdf>.

12. Ministry of Health, Labour and Welfare [Internet]. Check and search system of physician qualification [cited 2019 February 28]. Available from: https://licenseif.mhlw.go.jp/search_isei/
13. Statistics Bureau, Ministry of Internal Affairs and Communications [Internet]. Statistical Observations of Shi,Ku,Machi,Mura 2018. [cited 2019 Jan 28]. Available from: <https://www.stat.go.jp/data/s-sugata/index.html>.
14. Rosenthal MB, Zaslavsky A, Newhouse JP. The geographic distribution of physicians revisited. *Health services research*. 2005;40(6 Pt 1):1931-1952.
15. Schwartz WB, Newhouse JP, Bennett BW, Williams AP. The changing geographic distribution of board-certified physicians. *The New England journal of medicine*. 1980;303(18):1032-1038.
16. Isabel C, Paula V. Geographic distribution of physicians in Portugal. *The European Journal of Health Economics*. 2010;11(4):383-393.
17. Kroezen M, Dussault G, Craveiro I, et al. Recruitment and retention of health professionals across Europe: A literature review and multiple case study research. *Health policy (Amsterdam, Netherlands)*. 2015;119(12):1517-1528.
18. Medicine Io. *A Manpower Policy for Primary Health Care: Report of a Study*. Washington, DC: The National Academies Press; 1978.
19. Institute of Medicine Committee on Quality of Health Care in A. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington (DC): National Academies Press (US) Copyright 2001 by the National Academy of Sciences. All rights reserved.; 2001:360.
20. *Clinic Keiei Kenkyukai (Clinic Management Research Group)*. *linkaigyō handbook Handbook of opening private practice*. *Nihon Iryō Kikaku: Tokyo; 2005 (in Japanese)*.
21. Maeno T. Sougoushinryō ga tiikiiryō ni okeru senmoni ya tashokusyūrenkeinado ni ataeru kouka ni tuiteno kenkyū (Research of the effect that general practice gives to other specialists and interprofessional working in community medicine). 2017; <https://mhlw-grants.niph.go.jp/niph/search/NIDD00.do?resrchNum=201706032A>. Accessed May 28, 2019.

Tables

Table 1: Workplace municipalities of board-certified physicians ages 30 to 49, classified by population density.

		Quintile of municipalities sorted by population density ^a					Total
		1	2	3	4	5	
Family physicians	N	27 ^{**+}	49 ^{**+}	75 ^{**+}	138 ^{**+}	230 ^{**-}	519
	%	5.2%	9.4%	14.5%	26.6%	44.3%	100.0%
General internists	N	62	271	873	2599	8142	11947
	%	0.5%	2.3%	7.3%	21.8%	68.1%	100.0%
Surgeons	N	51	302	894	26009 ^{*+}	7723 ^{**-}	11570
	%	0.4%	2.6%	7.7%	22.5%	66.8%	100.0%
Obstetricians and gynaecologists	N	7 ^{**-}	121	318 ^{**-}	1085	1435 ^{**+}	5268
	%	0.1%	2.3%	6.0%	20.6%	70.9%	100.0%
Paediatricians	N	17 ^{**-}	148	545	1435 ^{*-}	4774 ^{*+}	6919
	%	0.2	2.1	7.9	20.7	69.0	100.0%
All physicians ^b	N	747	3846	10830	29489	92675	137587
	%	0.5	2.8	7.9	21.4	67.4	100.0%

^aQuintile 1 <= 433.14, Quintile 2 <= 1195.59, Quintile 3 <= 3155.03, Quintile 4 <= 10845.34, Quintile 5 10845.34+ people per square kilometer

^bAll physicians were not included in the residual analysis.

Residual analysis based on chi square test which examines the difference between the real and expected value at each cell.

*+: greater than expected value. P<0.05

**+: greater than expected value. P<0.001

*-: less than expected value. P<0.05

** -: less than expected value. P<0.001

Table 2: Workplace municipalities of board-certified physicians ages 30 to 49, classified by municipality type.

		Municipality type ^a			Total
		Metropolis ^a	City ^b	Town/village ^c	
Family physicians	N	194**-	260***+	65***+	519
	%	37.4%	50.1%	12.5%	100.0%
General internists	N	7299	4194	454	11947
	%	61.1%	35.1%	3.8%	100.0%
Surgeons	N	6928	4167	475	11570
	%	59.9%	36.0%	4.1%	100.0%
Obstetricians and gynaecologists	N	3299**-	1811*-	158**-	5268
	%	62.6%	34.4%	3.0%	100.0%
Paediatricians	N	4164	2493	262	6919
	%	60.2	36.0	3.8	100.0%
All physicians ^d	N	81794	49858	5935	137587
	%	59.4	36.2	4.3	100.0%

^a Metropolis: all the ordinance-designated cities (seirei-shitei-toshi and chukaku-shi) and 23 special wards of Tokyo (ku) (n = 97)

^bCity: the other cities (shi) (n = 717)

^cTown/village: towns (cho) and villages (son) (n = 923)

^dAll physicians were not included in the residual analysis.

Residual analysis based on chi square test which examines the difference between the real and expected value at each cell.

*+: greater than expected value. P<0.05

***+: greater than expected value. P<0.001

*-: less than expected value. P<0.05

** -: less than expected value. P<0.001

Figures




Figure 1

Number of board-certified physicians ages 30 to 49 per 100,000 population, classified by population density.




Figure 2

Number of board-certified physicians ages 30 to 49 per 100,000 population, classified by municipality type.