

High-gradient Magnetic Fields and Starch Metabolism - Results from a Space Experiment

Karl H. Hasenstein (✉ hasenstein@louisiana.edu)

University of Louisiana at Lafayette

MR Park

National Institute of Crop Science

John SP

University of Louisiana at Lafayette

Chitra Ajala

University of Louisiana at Lafayette

Article

Keywords:

Posted Date: July 6th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1731472/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Directing plant growth in weightlessness requires understanding the processes that establish plant orientation and how to manipulate them. Both gravi- and phototropism determine directional growth and previous experiments showed that high gradient magnetic fields (HGMF) can induce curvature in roots and shoots. Experiments with *Brassica rapa* verified that that gravitropism-like induction of curvature is possible in space and that the HGMF-responsive organelles are amyloplasts. We assessed the effect of space and HGMF based on 16 genes and compared their transcription with static growth and clinorotation. Amyloplasts size in root tips increased under weightlessness but decreased under clinorotation but not in response to magnetic fields. Amyloplast size changes were correlated with reduced amylase transcription in space samples and enhanced transcription after clinorotation. Mechanostimulation and weightlessness have opposite effects on the size of amyloplasts. The data show that plants perceive weightlessness, and that their metabolism adjusts to microgravity and mechanostimulation. Thus, clinorotation as surrogate for space research may lead to incorrect interpretations.

Introduction

The perception of the gravity stimulus and the involvement of amyloplasts have long been the focus of plant biology because of the cryptic nature of essential steps in the response of a biological system to physical stimuli such as gravity^{1–4}. While the first step of the gravitropic response primarily depends on the interaction of intracellular particles with Earth's gravitational field, the ability to respond to the gravity signal depends on biological conditions such as auxin sensitivity and transport⁵, time^{6,7}, the cytoskeleton⁸ and physical parameters such as temperature⁹, orientation^{10,11}, osmotic conditions¹², mechanical noise¹³, cytoplasmic viscosity¹⁴, and high-gradient magnetic fields^{15–17}. The redundancy and complexity contribute to the stability of the graviresponse system and can integrate additional signals such as hydrotropism^{18,19} and touch^{20,21}. Because any physiological response is a function of the accumulated signal (auxin, altered gene expression, pH, ion, or charge shifts), control of the response shifts from the original signal to downstream events such as signal processing (translation) and response variables that are related to signal strength and persistence²².

The complexity of the sensing process is amplified by physiological and genetic conditions that affect the (gravi)response; therefore, the analysis of mechano-sensing processes depends on the elimination of the gravity effects by the weightlessness conditions in space. The study of amyloplasts movement in the absence of gravity by HGMF was the focus of a shuttle experiment in 2003 (STS-107). However, the disintegration of the Shuttle during reentry made the intended examination impossible. Nonetheless, this experiment provided strong evidence that mechano-sensitivity increased in microgravity²³.

Here we report data from a space experiment on the Space-X Crew Resupply Service 3 mission. The aim of this investigation was to repeat the ill-fated experiment on STS-107. However, instead of flax (*Linum usitatissimum*) we used *Brassica rapa* seeds to study the effect of magnetic gradients and to take

advantage of its *Arabidopsis*-related genome to assess transcriptional responses of genes related to growth, metabolism, auxin, and stress.

HGMF

Induction of curvature by magneto-mechanical forces depends on the magnetic susceptibility (χ), which is characteristic of a given substance and describes its ability to acquire magnetization I as a result of the (inducing) magnetic field H such that $I = \chi H$. Diamagnetic substances have negative susceptibility and include the vast majority of biological substances. However, some metal-containing proteins such as hemoglobin, cytochrome, ferritin etc., can have positive susceptibility and are paramagnetic^{24–26}. The magnetic susceptibility of dia- and paramagnetic compounds is proportional to their density and their behavior in a magnetic gradient is analogous to that in the gravity field^{16,24,27}. Magnetic susceptibility and density of the cytoplasm are equal to that of water, $\chi_w = 7.2 \times 10^{-7}$ emu (electromagnetic units, table value), and that of starch $\chi_{st} = (8 \pm 0.2) \times 10^{-7}$, thus the differential susceptibility between starch and cytoplasm $\Delta\chi \approx 8 \times 10^{-8}$ emu. The density of starch $\rho_{st} = 1.5 \text{ g/cm}^3$, and cytoplasm $\sim 1 \text{ g/cm}^3$; therefore $\Delta\rho = 0.5 \text{ g/cm}^3$ and $\Delta\rho/\Delta\chi \approx -6 \times 10^6 \text{ (g/cm}^3\text{)/emu}^{24}$.

Although other methods of generating HGMF are available^{24,27}, the generation of a magnetic gradient in the experiments described here, was realized by inserting into a uniform, strong magnetic field ferromagnetic wedges. The wedges become magnetized and create a HGMF (Fig. 1) such that the gradient ∇H is directed toward the tip of the wedge. Therefore, the force acting on a diamagnetic body ($\chi < 0$) repels starch-filled amyloplasts from the inserted object. In contrast, a para- or ferromagnetic body ($\chi > 0$) would experience an attractive force. The force generated by the HGMF utilized in the employed setup has been determined previously to be about 0.6 g²⁸ and exceeds the acceleration that plants respond to (about 10^{-3} g²⁹).

In contrast to cultivation on earth where orienting factors are provided by gravity as constant and light as intermittent influence, plant growth in space cannot rely on gravitational clues; it depends only on the plants' phototropic response. If light is eliminated, elongation is variable and plant growth responds mostly to water gradients (hydrotropism)^{19,30}. Removing the two most consequential vectors for plant orientation, i.e., growing plants under weightlessness conditions in the dark, provides an opportunity to examine the role of amyloplasts for directional growth analogous to their role in gravity sensing. Probing magnetophoretic effects and the space conditions on root growth was the objective for the experiment described here.

The investigation included transcription analyses of 16 genes that represent auxin effects (*IAA5*, *PIN 1, 3, 7*) as the main growth regulator of differential elongation^{31–33}, starch catabolism (*Amy1*)³⁴, sucrose synthase (*SUS*, cell wall formation)³⁵, *UBQ1* (ubiquitin1; a putative reference³⁶ and indicator of protein metabolism)³⁷, *TUB1* (tubulin1; reference³⁶, cell growth, mitosis), *ACT7* (actin7; putative reference³⁸, root growth and germination³⁹), *ADH1* (alcohol dehydrogenase; stress response⁴⁰), *COX* (cytochrome oxidase

subunit1; metabolism, stress⁴¹), *G6PDH5* (glucose-6-phosphate dehydrogenase5; oxidative pentose phosphate pathway, basic metabolism⁴²), *GLK* (glucokinase; carbohydrate metabolism⁴³) HXK (hexokinase 1; glucose metabolism and developmental regulator⁴⁴), *PFK* (phosphofructokinase; carbohydrate metabolism, hypoxia stress⁴⁵), and *TAGL* (triacylglycerol lipase; lipid metabolism^{46,47}). The transcription pattern of these genes under conditions such as space (weightlessness), clinorotation (enhanced mechanostimulation), and static controls provided information on how seedlings respond to the respective conditions.

This report describes an experiment that built on prior experience (i.e., Biotube1 on STS-107) but used genetically better-characterized plant material, *Brassica rapa*. Our data support HGMFs as suitable method to establish plant orientation in space but also show that mechanostimulation affects gravisensing-related metabolism. The report demonstrates that plants perceive weightlessness ("microgravity") and respond by reduced starch reduction in gravity perceiving cells.

Material And Methods

Plant material

Brassica rapa, var. rapa seeds were commercially sourced. No approvals were required for the study, which complied with all relevant regulations. Seeds were attached to germination paper with polyvinyl acetate (clear Elmer's) glue such that the micropyle was oriented toward the opening of seed cassettes (Sup. Figure 1). The seeds cassettes were designed to fit into specially designed magnetic field chambers (MFCs, Fig. 1).

Experimental Setup

The experiment was a follow-up of the STS-107 experiment that was lost as a result of the shuttle accident in 2003. A computer (PC104 stack) controlled all aspects and was remotely operated such that no crew involvement was necessary. The experimental principle of the hardware (Fig. 1) and the arrangement of the components is shown in Sup. Figure 2. Experiment initialization, water dispensation, image acquisition, temperature and pressure recording, and termination of the experiment by fixation were remotely initiated and controlled by the hardware's own computer. The entire hardware was custom built at the Kennedy Space Center using several iterations and named Biotube-MICRo (Magnetic Induction of Curvature in Roots).

Hardware

The MFCs consisted of machined aluminum cases each containing 10 Neodymium Iron Boron (NdFeB) magnets, 12.7 mm thick with a magnetization of about 30000 Oe (~ 3 Tesla, Magnet Sales and Manufacturing, Culver City CA). The magnetic circuit was closed by two yokes. To the surface of each

magnet, two wedges of ferromagnetic steel (equilateral cross section, 6 mm high, 50 mm long) were attached that generated the magnetic gradients (Fig. 1). Between each pair of magnets seed cassettes were inserted that contained 10 *Brassica rapa* seeds. Five seeds each were glued (Elmer's clear glue) along the long edges of germination paper. Roots that emerged from the seed cassettes were expected to curve away from the wedges.

A camera system consisting of eight cameras on either side was installed between two of the three MFCs such that the cameras were positioned opposite a window that permitted viewing the seed cassettes. Illumination was achieved by a single IR LED (750–800 nm) to avoid phototropic stimulation. The LEDs were activated only during imaging. The cameras acquired images in two-minute intervals and the video signal together with temperature, and atmospheric pressure data and position number were transmitted to the Kennedy Space Center (Florida) as well as recorded internally on a hard drive.

Space Flight Parameters

The experiment was launched on Space-X3 (April 18, 2014) and initiated after 19 days by dispensing 400 µL deionized water to each seed cassette from a Micro Effusion Device for Space Applications (MEDUSA). The seeds germinated after about 22 hours. The experiment was terminated after 48 hours by the injection of RNA-Later® into MFC-C (first in the fixation sequence, not imaged), injection of 4% formaldehyde in PHEMD buffer⁴⁸ in MFC-A (image sequences designated 'Y'). MFC-B received RNALater, was designated 'Z' and third in the fixation sequence. MFC-C did not contain magnets but Al blocks of identical dimensions as the magnets and served as control. The dispensation of the respective fixative was initiated by pressurizing an aluminum chamber that contained sealed bags with fixative, one for each MFC. A solenoid valve controlled fixative flow to one MFC at a time. Excess liquid was collected in a second bag housed in a plastic cage. The entire hardware and fixed plant material deorbited on May 20, 2014. After two days the hardware was

received at the Kennedy Space Center, and sample processing was completed during the next three days. A ground control using the space hardware was performed at the Kennedy Space Center on a large clinostat (8/12–8/15/14, 1.5 rpm) and at the University of Louisiana using single clinorotated MFCs without imaging but under otherwise identical conditions.

Data analysis

Image sequences were obtained from HD-stored files and compiled as video sequences (Sup. Video 1 and Sup. Video 2). Root curvature was recorded from seed cassettes that were submersed in the same buffer as during fixation in an upright position and photographed from four sides. After disassembly of the seed cassettes, root appearance was categorized as germinated, straight, or curved. Curvature was evaluated as affected by HGMF only if the root length was sufficient to reach the HGMF area and corresponding curvature was visible.

Microscopy was performed on seedlings fixed in 4% (v/v) formaldehyde in PHEMD buffer⁴⁸. Samples were dehydrated in a graded ethanol series and 100% acetone and embedded in Spurr's resin. Longitudinal median sections (2 µm thick) were cut on an ultra-microtome (Sorvall MT2-B) and stained with toluidine blue (0.1% [w/v] in 0.1% [w/v] boric acid). The serial sections were photographed with a digital camera (Sony DKC-ST5) and measured using ImageJ (v.1.53).

RNA extraction was performed in four separate sections of seedlings (root tip, root proper, hypocotyl, and cotyledons) using a kit (Spectrum™ Plant Total RNA Kit, Sigma STRN250) following the manufacturer's protocol. Transcription analysis was performed after reverse transcription (High-Capacity cDNA Reverse Transcription Kit, Applied Biosystems, USA) and qPCR reactions for 16 different genes using specific primers, with stable and suitable efficiencies (Sup. Table 1). Conditions for the amplification on a Step-one real-time PCR system (ThermoFisher Scientific) included a 2 min incubation at 95°C, followed by 40 cycles (95°C for 10 s; 60°C for 10 s) with fluorescent readings taken at the end of the annealing cycle. Quality control included melt curve analyses and capillary electrophoresis (QIAxcel Advanced).

The assessment of the space environment on seedling biology was examined using qPCR data of 16 genes. Typically, qPCR data rely on so-called reference or housekeeping genes, but this approach does not consider that the environmental conditions (space flight, clinorotation, and static growth conditions) are likely to affect reference genes themselves. Therefore, we compared the entire data sets against each other based on all studied conditions. If no difference exists between the two data sets, then such plots result in a diagonal line with a slope of one. The evaluation based on slope and variance (R^2 value) as indicator of overall stability of gene activity also provides information on significantly affected genes. Such data points will show up as 'outliers' from the bulk distribution⁴⁹. This approach was evaluated based on four individual tissues of the seedlings, resulting in 64 data points (16 genes × 4 tissue types), for each treatment.

The most relevant comparison of the explained type focuses on the effect of weightlessness and HGMFs in addition to the effect of clinorotation and static, 1g controls. The complete set of comparisons is provided as Sup. Table 2. Transcription data were evaluated based on correspondence plots of pairwise comparisons. Significant changes in transcriptions were evaluated by the Mean Method⁴⁹, and statistical significance was based on Z-scores.

Results And Discussion

Growth and seedling development

Each MFC contained eight seed cassettes with ten seeds each. Seed germination in the two MFCs with HGMFs was higher than in the non-magnetic chamber (Table 1), indicating that the presence of magnets and magnetic fields did not negatively affect germination.

Table 1
Germination and root length of *Brassica rapa* seeds in magnetic field chambers with (A&B) and without magnets (C). There was no difference between chambers or seed cassettes.

		Cassette:								
		1	2	3	4	5	6	7	8	
MFC-A (HGMF)	Average:	14.3	14.6	15.4	11.4	14.4	15.1	16.2	12.6	14.7
	SE:	4.2	3.2	3.5	3.7	4.4	2.8	4.8	4.8	5.0
	N=	77	9	10	10	10	10	9	9	
MFC-B (HGMF)	Average:	16.1	16.6	15.0	17.3	16.8	15.5	16.3	16.4	14.6
	SE:	3.5	2.5	3.0	3.6	4.9	3.5	3.7	4.0	2.8
	N=	80	10	10	10	10	10	10	10	
MFC-C: (no magnets)	Average:	14.7	19.1	14.2	13.4	15.2	15.3	14.5	13.0	13.4
	SE:	4.4	3.9	4.7	4.0	2.3	4.3	6.4	3.3	3.2
	N=	70	8	10	9	8	9	10	7	9

Curvature Induction By Hgmf:

Exposure to HGMF induced curvature (Sup. Video 1, Table 2). However, substantial curvature occurred in both presence and absence of magnetic gradients. The non-magnetic chamber showed 81% curvature in roots within the seed cassette around the germination paper (Fig. 2).

Table 2
Germination and curvature in the presence and absence of HGMFs of a total of 80 seeds per MFC

	not germinated	Straight*	curved	curved (HGMF)
MFC-A (HGMF)	3	8	40	28
MFC-B (HGMF)	0	0	50	30
MFC-C (No magnets)	10	5	65	0

*refers to roots that never emerged from the seed cassette or were too short to reach the HGMF area (as shown in Fig. 2).

MFCs with magnets showed curvature that was independent of the HGMF because of growth patterns or curvature occurring inside the seed cassettes where the magnetic gradient was too weak to affect curvature.

The comparison of space-grown or static ground control seedlings exposed to HGMF with seedlings grown in the non-magnetic field chamber (#1 and #23 in Sup. Table 2) shows strong correspondence between the two conditions (Fig. 3), indicating that magnetic fields do not affect general metabolism or transcription activity. This notion is in line with reports that failed to detect effects of magnetic fields on growth^{50,51} but contrasts with effects of weak magnetic fields on root curvature⁵² and effects of the geomagnetic field on stress response and hormesis^{53,54}. The geomagnetic field (typically about 0.5 G) is orders of magnitude weaker than the employed magnetic fields in this research (ca. 30 kG).

These data indicate that factors other than HGMF induce curvature and the most likely factor is hydrotropism^{18,30,55}. The lack of a gravity stimulus and the distance between the germination paper and the HGMF-generating wedges resulted in a large number of roots confined to the germination paper. The same factor also applied to seedlings in the magnetic chambers and resulted in reduced numbers of roots reaching the HGMF. However, the unambiguous evidence for curvature in space (see Sup. Video 1) and clinorotated seedlings (Sup. Video 2) indicates that HGMFs do have the ability to induce curvature as had been demonstrated earlier in roots, coleoptiles, inflorescences, and hypocotyls^{16,17,24,56,57}.

The presumably hydrotropic growth reduced the number of roots that reached the influence of the HGMFs. Therefore, the overall number of roots curving in response to the magnetic gradient was low (Table 2).

Examining the gravisensing mechanism includes measuring the size of the presumptive sensors as previous work has shown that gravisensitivity depends on the amount of starch in amyloplasts⁵⁸. Assuming equal density, the diameter of amyloplasts determines the relative mass. Measurements of

amyloplasts in columella cells differed between the space-grown, clinorotated, and statically grown ground controls (Fig. 4) and indicates that the amyloplast size and therefore the gravisensing mechanism is responsive to the growth condition, notably weightlessness and clinorotation that respectively reduces and enhances mechanostimulation.

Earlier analyses of starch from space and ground controls found that ethylene reduces starch particle size and that in cotyledons starch particles sizes in space and ground controls were of equal size⁵⁹. Since we used activated charcoal cloth to absorb volatile organic compounds (Fig. 1), we assume that ethylene was not a factor in the different experiments. In addition, we examined the size distribution of amyloplasts in the root columella. Earlier work showed that amyloplasts in gravisensing tissues (root cap and endodermis) are about twice the size of other tissues⁷. Our data (Fig. 4) indicate that the amyloplast size is responsive to the gravitational and mechanical stimulation. The reduced size after clinorotation, the average size in static 1-g controls, and the enlarged size in roots grown under 'micro-gravity' conditions support the notion that the extend of gravitational stimulation is inversely proportional to the mass of the amyloplasts. Thus, plants not only perceive the direction of an accelerative force but adjust their (starch) metabolism according to the amount of stimulation. If this notion is correct, then gene transcription data should support this notion and the following section confirms that amylase activity indeed is a function of mechanostimulation.

Transcript Analyses

Since four different tissue types were analyzed, tissue variability and response to spaceflight and clinorotation can be assessed for all examined tissues and genes. Based on distributions of transcription patterns (Figs. 3 and 5), a comprehensive analysis of all treatment and tissue combinations was performed such that the scatter for each comparison and gene was determined based on the formula

$$\sum_{i=1}^n \sum_{j=1}^4 \sqrt{\left(X_{ij} - \bar{X}_j \right)^2 + \left(Y_{ij} - \bar{Y}_j \right)^2}$$

where j represents the tissue types (root tip, root proper, hypocotyl, cotyledons) and i the individual experimental comparisons (28, Sup. Table 2) or examined genes (16). This value was calculated for each analyzed gene and the smallest value (scatter in Figs. 3 & 5) was identified as the most stably transcribed gene (Table 3).

The distribution of the least variable gene transcription varied greatly (Table 3). Although common reference genes (e.g., *TUB1*, *ACT7*) are represented (Table 3), the observation that individual tissues differed in the genes of greatest stability and that the average of all tissues resulted in different assortments suggested that referring transcription data to the average of all measurements (i.e., the regression lines in Figs. 3 and 5) is superior to relying on a single gene. Because *PFK* showed the greatest

stability for all tissues, we placed *PFK* at the origin of the coordinate system (0/0). However, the results explained below are independent of this selection.

Table 3
Assessment of the consistency of transcription data based on the average of examined tissues. The top four choices of all combinations (as in Sup. Table 2) are based on the percentage that resulted in the least scatter of all 448 combinations (16 genes by 28 comparisons).

	Gene	n	%
All tissues	<i>PFK</i>	60	13.4
	<i>PIN1</i>	47	10.5
	<i>TUB1</i>	45	10.0
	<i>GLK</i>	33	7.4
Root tip	<i>PIN1</i>	43	9.6
	<i>PIN3</i>	41	9.2
	<i>SUS</i>	40	8.9
	<i>GLK</i>	35	7.8
Root proper	<i>COX</i>	51	11.4
	<i>ACT7</i>	42	9.4
	<i>SUS</i>	42	9.4
	<i>HXK</i>	39	8.7
Hypocotyl	<i>HXK</i>	42	9.4
	<i>ACT7</i>	37	8.3
	<i>SUS</i>	37	8.3
	<i>ADH1</i>	27	6.0
Cotyledons	<i>PFK</i>	49	10.9
	<i>ACT7</i>	45	10.0
	<i>ADH1</i>	40	8.9
	<i>PIN7</i>	35	7.8

The comparisons (Sup. Table 2) show the least and most significant effect on gene expression and can be used to identify the conditions that induce physiological responses in brassica seedlings. The greatest stability in gene transcription was seen for treatments with similar mechanical load, for example, static growth with and without HGMF, or clinorotated samples (KSC and Lab). The largest scatter or least consistent transcription pattern was associated with different mechanical loads such as flight (i.e., no mechanical load) and clinorotation (enhanced mechanical load), or static and space flight conditions. The main conclusion of these evaluations is that HGMFs or more generally, strong magnetic fields, do not affect transcription; HGMF data are equally present in the most and least affected conditions (Table 4). However, a comparison between clinorotated experiments with the original flight hardware at KSC and experiments in our lab (comparison # 17 and # 22) show statistically significant differences (Sup. Figure 4).

Sensitivity of *AMY1*

Data sets comparing space-grown with clinorotated seedlings show large scatter and offset (Fig. 5A and Sup. Figure 3A), suggesting that the shift in the transcription pattern is not dependent on HGMF but differences in mechanostimulation (i.e., clinorotation). This effect is especially noticeable for *AMY1*. The effect of clinorotation on amylase transcription is independent of the HGMF because transcription of *AMY1* in the absence of HGMF was higher in static seedlings than clinorotated seedlings (Fig. 5B). Space flight and clinorotation resulted in reduced and elevated transcription, respectively. A similar pattern was observed between clinorotated and static samples; however, HGMF had no effect (Sup. Figure 3). The altered *AMY1* transcription between clinorotation at KSC and our lab is related to different stabilities of the hardware. The flight hardware contained a webbing-like base mount (implemented because of weight concerns), which provided stable support during space flight but flexed readily during clinorotation compared to a rigid assembly on the laboratory clinostat.

The transcription data correspond with the observed size distribution of amyloplasts (Fig. 4) and indicate that amyloplast size is regulated by starch degradation in clinorotated seedlings and amylase repression (starch accumulation) in space samples. Together these observations indicate that plant adapt to the weightlessness of space by increasing their amyloplast size which likely enhances their gravisensitivity. In contrast, clinorotation represents excessive mechanostimulation and leads to a reduction of amyloplast size through enhanced degradation (amylase transcription).

The observed *AMY1* levels between space-grown and clinorotated seedlings were independent of reference genes; the effect persisted regardless of whatever gene was used as reference. This observation supports using transcription of all available genes as a reliable approach to identifying transcriptional changes of individual genes.

The *AMY1* data correspond to earlier observations of enhanced gravisensitivity of space-grown lentil seedlings^{60,61} and reduced starch after clinorotation⁶². Although the STS-107 flight experiment could not be retrieved to measure amyloplast size, the image analysis of flax seedlings indicated that the magnetic

gradient had stronger effects than during previous ground experiments because the root curvature started at a greater distance from the HGMF-inducing wedge²³ than during ground controls. This observation is in line with the present data and strongly supports greater (gravi)sensitivity of plants growing in a microgravity environment.

The current report is the first to associate starch metabolism with amyloplasts size and gravisensitivity. The larger amyloplasts in space-grown plants suggest that the application of HGMFs in space is more effective than in earth-grown and especially in clinorotated plants. However, the unreliable growth direction of roots makes HGMF difficult to implement.

Gravisensitivity has also been linked to changes in calcium in statocytes^{63–65} and calcium has been shown to stabilize α -amylase^{66–68}. Therefore, Ca^{2+} and amylase are controlling element for the starch content in statocytes. However, starch content is the result of homeostasis for catabolic and anabolic metabolism. Starch biosynthesis depends on a complex set of enzymes that include phosphoglucose isomerase (PGI)⁶⁹, phosphoglucomutase (PGM)^{70,71}, and starch synthases (SSs)^{69,72} among others. Data on the balance between starch degradation and starch biosynthesis undoubtedly would provide a more comprehensive assessment of the sensitivity of starch metabolism to reduced and enhanced mechanostimulation. However, the limited set of transcriptionally analyzed genes prevents a more thorough assessment of starch biosynthesis in response to altered mechanostimulation. Future space experiments might remedy this shortcoming.

Table 4

The similarity of gene transcription between pairs of treatments based on high gradient magnetic fields (HGMF) or no magnetic fields (no-MF) that were applied during the space flight (Flight), on ground controls (static), or during clinorotation (Clino) at the Kennedy Space Center (KSC) or in the laboratory (Lab). The values were obtained by averaging the R^2 values that resulted from using each of the 16 genes as reference (i.e., origin in graphical comparisons such as Figs. 3&5 and Sup. Figure 3&4).

	#*	R^2 , %	STDEV, %	Comparisons (all genes)
least stable ←————→ most stable	23	94.8	1.9	KSC static no-MF vs. KSC static HGMF
	17	90.8	2.7	Lab Clino HGMF vs. KSC Clino HGMF
	28	82.6	7.5	Lab Clino no-MF vs. Lab Clino HGMF
	1	80.2	7.2	Flight-no MF vs. Flight-HGMF
	15	80.0	3.3	KSC static HGMF vs. KSC Clino HGMF
	18	78.8	3.7	Lab Clino no-MF vs. KSC Clino HGMF
	16	76.6	3.3	KSC static no-MF vs. KSC Clino HGMF
	25	76.3	2.6	Lab Clino no-MF vs. KSC static HGMF
	27	75.6	4.7	Lab Clino no-MF vs. KSC static no-MF
	24	73.1	5.2	Lab Clino HGMF vs. KSC static HGMF
	19	72.8	7.0	KSC static HGMF vs. KSC Clino no-MF
	26	70.7	5.5	Lab Clino HGMF vs. KSC static no-MF
	20	70.4	6.7	KSC static no-MF vs. KSC Clino no-MF

* Numbers refer to Sup. Table 2.

	#*	R ² , %	STDEV, %	Comparisons (all genes)
	14	60.6	7.7	KSC Clino no-MF vs. KSC Clino HGMF
	13	59.2	11.1	Lab Clino no-MF vs. Flight-no MF
	22	56.9	6.2	Lab Clino no-MF vs. KSC Clino no-MF
	21	56.2	7.3	Lab Clino HGMF vs. KSC Clino no-MF
	7	56.1	11.2	Lab Clino no-MF vs. Flight-HGMF
	12	53.6	7.0	Lab Clino HGMF vs. Flight-no MF
	9	52.3	10.0	KSC Clino no-MF vs. Flight-no MF
	10	51.0	8.2	KSC static HGMF vs. Flight-no MF
	11	50.7	8.4	KSC static no-MF vs. Flight-no MF
	6	49.9	7.5	Lab Clino HGMF vs. Flight-HGMF
	8	46.0	7.1	KSC Clino HGMF vs. Flight-no MF
	2	40.6	6.7	KSC Clino HGMF vs. Flight-HGMF
	5	38.0	7.4	KSC static no-MF vs. Flight-HGMF
	3	37.4	10.6	KSC Clino no-MF vs. Flight-HGMF
	4	37.4	10.6	KSC static HGMF vs. Flight-HGMF

* Numbers refer to Sup. Table 2.

Genes other than AMY1

Although weightlessness is the dominant difference between ground and space flights, the lack of density-driven gas exchange and water distribution are equally significant alterations for plant growth in

space. Because the flight hardware was enclosed in a hermetically sealed (triple-contained) chamber, atmospheric effects can be excluded. Therefore, the following considerations only apply to gravity effects. Comparing the number of significantly ($|p| < 0.05$) affected combinations (genes by reference), shows the largest effect on *AMY1* transcription (Fig. 6).

Genes other than *AMY1* responded to specific conditions but at reduced power. *ACT7* showed modifications only when comparing clinorotation at KSC with static growth (Comparisons #19 & #20); *ACT7* was not affected by HGMF but sensitive to vibrations (higher frequency movements in excess to the 1.5 rpm of the clinostat). This relationship is similar to the enhanced *AMY1* transcription addressed in Sup. Figure 4. The enhanced oscillation can be verified by the jitter in Sup. Video 2. To improve visibility, the individual frames were aligned, but the inconsistent position of imprinted data confirms the added vibrational stimulation that affected *ACT7* activation (and enhanced *AMY1* transcription) compared to regular (smooth) clinorotation.

COX showed the strongest upregulation in space flight material compared with static growth conditions (#4, #5, #10 #11) but was not affected by HGMF. This observation further supports the claim that magnetic fields do not affect transcription. Instead, *COX* responds to space flight associated stress as has been shown previously for fish brain⁷³ and skeletal muscle⁷⁴. Changes in *G6PDH5* were limited to treatment differences between space flight and clinorotation (#8, #12, #15, #16) and correspond to earlier reports of altered enzyme activities of pine seedlings after exposure to clinorotation and hyper-g⁷⁵ and enzyme activities in artemia cysts after space flight⁷⁶. *IAA5* and *PIN7* responded to HGMF both under flight (*PIN7*) and clinorotation (*IAA5*) and underscores their relevance for auxin modification of growth. Although the number of roots that curved in response to HGMF was low (Table 2), it is possible that differential growth (curvature in response to hydrotropism?) affected the transcription of these genes. *GLK* did not respond significantly; changes were not limited to any particular condition and therefore cannot be associated with specific experimental conditions but likely represent natural variations. *UBq1*, *Tub1*, *ADH1*, *HXK*, *PFK*, *SUS*, *PIN1* showed no significant changes in transcription and therefore could all serve as references. The lack of response of *ADH1* is surprising as this gene was previously identified as space stress indicator^{77,78}. *TAGL* is the only gene that shows reduced transcription in the presence of magnetic fields but only in clinorotated samples. While there is precedence that the lipid metabolism is affected by hypergravity⁷⁹ and clinorotation⁸⁰, this is the first observation that magnetic fields might contribute to such changes.

In summary, our data support plant proprioception of weightlessness and metabolic control of amyloplast size. The adjustment of size and mass of amyloplasts indicates that plants perceive gravity and ponderomotive forces, which provide not only enhance gravisensitivity but also explain some metabolic responses to space conditions. The advantage of a sealed environment suggests that this effect is not an artefact but related to gravitational and mechanosensory response. The results also demonstrate that static magnetic fields do not influence gene transcription. Future work needs to

investigate the role of other starch-related genes to understand the entire dynamic of metabolic plasticity that relates to weightlessness.

Declarations

Data availability

All data are available upon request from the corresponding author

Acknowledgements

This work was supported by the National Aeronautics and Space Administration grants NNX10AP91G, NNH18ZHA004C, and 80NSSC17K0344 to KHH.

The authors are grateful for the help of KSC personnel, especially Allison Caron, Ralph Fritsche, Howard Levine, Carole Miller, Bradley Burns ,and Dave Reed. We also acknowledge the assistance of Allison Mjoen, Carlos Gill, George Guerra, James Smodell.

Author Contributions

KHH received funding, designed the experiment, analyzed the data, and wrote the paper MRP performed qPCR work, SPJ and CA performed microscopy and measured amyloplast size. All authors reviewed the text.

Competing Interests:

The authors declare no competing interests.

Additional Information

Supplementary information for this paper is provided as separate files (2 video files and one Word file).

References

1. Hasenstein, K. H. Plant responses to gravity - Insights and extrapolations from ground studies. *Gravitational and Space Biology* **22**, 21–32 (2009).
2. Blancaflor, E. B. & Masson, P. H. Plant gravitropism. Unraveling the ups and downs of a complex process. *Plant Physiol.* **133**, 1677–1690, doi:10.1104/pp.103.032169 (2003).
3. Su, S. H., Gibbs, N. M., Jancewicz, A. L. & Masson, P. H. Molecular Mechanisms of Root Gravitropism. *Curr. Biol.* **27**, R964-R972, doi:10.1016/j.cub.2017.07.015 (2017).
4. Toyota, M. & Gilroy, S. Gravitropism and mechanical signaling in plants. *Am. J. Bot.* **100**, 111–125, doi:10.3732/ajb.1200408 (2013).

5. Levernier, N., Pouliquen, O. & Forterre, Y. An Integrative Model of Plant Gravitropism Linking Statoliths Position and Auxin Transport. *Front. Plant Sci.* **12**, 651928, doi:10.3389/fpls.2021.651928 (2021).
6. Ma, Z. & Hasenstein, K. H. The onset of gravisensitivity in the embryonic root of flax. *Plant Physiol.* **140**, 159–166, doi:10.1104/pp.105.073296 (2006).
7. Ajala, C. & Hasenstein, K. H. Augmentation of root gravitropism by hypocotyl curvature in *Brassica rapa* seedlings. *Plant Sci.* **285**, 214–223, doi:10.1016/j.plantsci.2019.05.017 (2019).
8. Blancaflor, E. B. The Cytoskeleton and Gravitropism in Higher Plants. *J. Plant Growth Regul.* **21**, 120–136 (2002).
9. Withers, J. C. *et al.* Gravity persistent signal 1 (gps1) reveals novel cytochrome p450s involved in gravitropism. *Am. J. Bot.* **100**, 183–193, doi:10.3732/ajb.1200436 (2013).
10. Toal, T. W. *et al.* Regulation of Root Angle and Gravitropism. *Genes Genomes Genetics* **8**, 3841, doi:10.1534/g3.118.200540 (2018).
11. Dummer, M., Forreiter, C. & Galland, P. Gravitropism in *Arabidopsis thaliana*: Root-specific action of the EHB gene and violation of the resultant law. *J. Plant Physiol.* **189**, 24–33, doi:10.1016/j.jplph.2015.09.008 (2015).
12. Blancaflor, E. B. & Hasenstein, K. H. Growth and Microtubule Orientation of Zea-Mays Roots Subjected to Osmotic-Stress. *Int. J. Plant Sci.* **156**, 774–783 (1995).
13. Ma, Z. & Hasenstein, K. H. Noise amplification of plant gravisensing. *Adv. Space Res.* **39**, 1119–1126 (2007).
14. Scherp, P. & Hasenstein, K. H. Microinjection - A tool to study gravitropism. *Adv. Space Res.* **31**, 2221–2227 (2003).
15. Hasenstein, K. H. & Kuznetsov, O. A. The response of lazy-2 tomato seedlings to curvature-inducing magnetic gradients is modulated by light. *Planta* **208**, 59–65, doi:10.1007/s004250050534 (1999).
16. Kuznetsov, O. A. & Hasenstein, K. H. Magnetophoretic induction of curvature in coleoptiles and hypocotyls. *J. Exp. Bot.* **48**, 1951–1957, doi:10.1093/jexbot/48.316.1951 (1997).
17. Weise, S. E., Kuznetsov, O. A., Hasenstein, K. H. & Kiss, J. Z. Curvature in *Arabidopsis* inflorescence stems is limited to the region of amyloplast displacement. *Plant & cell physiology* **41**, 702–709 (2000).
18. Kobayashi, A. *et al.* A gene essential for hydrotropism in roots. *Proc. Natl. Acad. Sci. USA* **104**, 4724–4729 (2007).
19. Dietrich, D. *et al.* Root hydrotropism is controlled via a cortex-specific growth mechanism. *Nature Plants* **3**, 17057 (2017).
20. Massa, G. D. & Gilroy, S. Touch modulates gravity sensing to regulate the growth of primary roots of *Arabidopsis thaliana*. *Plant J.* **33**, 435–445 (2003).
21. Braam, J. In Touch: Plant Responses to Mechanical Stimuli. *New Phytol.* **165**, 373–389 (2005).
22. John, S. P. & Hasenstein, K. H. Effects of mechanostimulation on gravitropism and signal persistence in flax roots. *Plant Signal Behav* **6**, 1–6 (2011).

23. Hasenstein, K. H., Scherp, P. & Ma, Z. Gravisensing in flax roots - results from STS-107. *Adv. Space Res.* **36**, 1189–1195 (2005).
24. Kuznetsov, O. A. & Hasenstein, K. H. Intracellular magnetophoresis of amyloplasts and induction of root curvature. *Planta* **198**, 87–94 (1996).
25. Senftle, F. & Hambright, W. in *Biological Effects of Magnetic Fields (2. vol)* (ed Madeleine F Barnothy) 261–306 (Plenum Press, 1969).
26. Theil, E. Ferritin: structure, gene regulation, and cellular function in animals, plants, and microorganisms. *Annu. Rev. Biochem.* **56**, 289–315 Annual Reviews, INC (1987).
27. Kuznetsov, O. A. & H., H. K. in *Scientific and Clinical Applications of magnetic carriers* (eds Urs Häfeli, Wolfgang Schütt, Joachim Teller, & Maceij Zborowski) Ch. 33, 429–444 (Plenum Press, 1997).
28. Hasenstein, K. H., John, S., Scherp, P., Povinelli, D. & Mopper, S. Analysis of magnetic gradients to study gravitropism. *Am. J. Bot.* **100**, 249–255, doi:10.3732/ajb.1200304 (2013).
29. Shen-Miller, J., Hinchman, R. & Gordon, S. A. Thresholds for Georesponse to Acceleration in Gravity-Compensated *Avena* Seedlings. *Plant Physiol.* **43**, 338–344 (1968).
30. Takahashi, H. & Scott, T. K. Hydrotropism and Its Interaction With Gravitropism in Maize Roots. *Plant Physiol.* **96**, 558–564 (1991).
31. Blilou, I. *et al.* The PIN auxin efflux facilitator network controls growth and patterning in *Arabidopsis* roots. *Nature* **433**, 39–44 (2005).
32. Hasenstein, K. H. & Evans, M. L. Effects of cations on hormone transport in primary roots of *Zea mays*. *Plant Physiol.* **86**, 890–894, doi:10.1104/pp.86.3.890 (1988).
33. Adamowski, M. & Friml, J. PIN-Dependent Auxin Transport: Action, Regulation, and Evolution. *Plant Cell* **27**, 20–32, doi:10.1105/tpc.114.134874 (2015).
34. Deikman, J. & Jones, R. Control of alpha amylase messenger RNA accumulation by gibberellic acid and calcium in barley aleurone layers. *Plant Physiol.* **78**, 192–198 (1985).
35. Coleman, H. D., Yan, J. & Mansfield, S. D. Sucrose synthase affects carbon partitioning to increase cellulose production and altered cell wall ultrastructure. *Proc Natl Acad Sci U S A.* **106**, 13118–13123 (2009).
36. da Silva Santos, P. H. *et al.* Selection and validation of reference genes by RT-qPCR under photoperiodic induction of flowering in sugarcane (*Saccharum* spp.). *Sci Rep* **11**, 4589, doi:10.1038/s41598-021-83918-2 (2021).
37. Xu, F. Q. & Xue, H. W. The ubiquitin-proteasome system in plant responses to environments. *Plant Cell Environ* **42**, 2931–2944, doi:10.1111/pce.13633 (2019).
38. Yang, H. L. *et al.* Selection and evaluation of novel reference genes for quantitative reverse transcription PCR (qRT-PCR) based on genome and transcriptome data in *Brassica napus* L. *Gene* **538**, 113–122, doi:10.1016/j.gene.2013.12.057 (2014).
39. Gilliland, L. U., Pawloski, L. C., Kandasamy, M. K. & Meagher, R. B. *Arabidopsis* actin gene ACT7 plays an essential role in germination and root growth. *Plant J.* **33**, 319–328 (2003).

40. Papdi, C. *et al.* Functional identification of Arabidopsis stress regulatory genes using the controlled cDNA overexpression system. *Plant Physiol.* **147**, 528–542, doi:10.1104/pp.108.116897 (2008).
41. Garcia, L. *et al.* The cytochrome c oxidase biogenesis factor AtCOX17 modulates stress responses in Arabidopsis. *Plant Cell & Environment* **39**, 628–644, doi:10.1111/pce.12647 (2016).
42. Wakao, S. & Benning, C. Genome-wide analysis of glucose-6-phosphate dehydrogenases in Arabidopsis. *Plant J.* **41**, 243–256, doi:10.1111/j.1365-313X.2004.02293.x (2005).
43. Klepikova, A. V., Kasianov, A. S., Gerasimov, E. S., Logacheva, M. D. & Penin, A. A. A high resolution map of the Arabidopsis thaliana developmental transcriptome based on RNA-seq profiling. *Plant J.* **88**, 1058–1070, doi:10.1111/tpj.13312 (2016).
44. Aguilera-Alvarado, G. P. & Sanchez-Nieto, S. Plant Hexokinases are Multifaceted Proteins. *Plant & cell physiology* **58**, 1151–1160, doi:10.1093/pcp/pcx062 (2017).
45. Hwang, J. H. *et al.* Expression Profile Analysis of Hypoxia Responses in Arabidopsis Roots and Shoots. *Journal of Plant Biology* **54**, 373–383, doi:10.1007/s12374-011-9172-9 (2011).
46. McGlew, K. *et al.* An annotated database of Arabidopsis mutants of acyl lipid metabolism. *Plant Cell Rep.* **34**, 519–532, doi:10.1007/s00299-014-1710-8 (2015).
47. Kelly, A. A. & Feussner, I. Oil is on the agenda: Lipid turnover in higher plants. *Biochim. Biophys. Acta* **1861**, 1253–1268, doi:10.1016/j.bbalip.2016.04.021 (2016).
48. Blancaflor, E. & Hasenstein, K. Organization of cortical microtubules in graviresponding maize roots. *Planta* **191**, 231–237 (1993).
49. Houston, L. M., Hasenstein, K. H. & Deoli, N. T. The Mean Method: A Specific Outlier Boundary for Arbitrary Distributions. *Theoretical Mathematics & Applications* **5**, 1–12 (2015).
50. Chaplin, A. The effects of a one Tesla magnet on human fibroblast growth. *BIOS* **76**, 193–203, doi:[https://doi.org/10.1893/0005-3155\(2005\)076\[0193:TEOAOT\]2.0.CO;2](https://doi.org/10.1893/0005-3155(2005)076[0193:TEOAOT]2.0.CO;2) (2005).
51. Gellrich, D., Becker, S. & Strieth, S. Static magnetic fields increase tumor microvessel leakiness and improve antitumoral efficacy in combination with paclitaxel. *Cancer Lett.* **343**, 107–114, doi:10.1016/j.canlet.2013.09.021 (2014).
52. Kordyum, E. L., Bogatina, N. I., Kalinina, Y. M. & Sheykina, N. V. A weak combined magnetic field changes root gravitropism. *Adv. Space Res.* **36**, 1229–1236, doi:10.1016/j.asr.2005.05.103 (2005).
53. Maffei, M. E. Magnetic field effects on plant growth, development, and evolution. *Front. Plant Sci.* **5**, doi:10.3389/fpls.2014.00445 (2014).
54. Paponov, I. A., Fliegmann, J., Narayana, R. & Maffei, M. E. Differential root and shoot magnetoresponses in Arabidopsis thaliana. *Sci Rep* **11**, 9195, doi:10.1038/s41598-021-88695-6 (2021).
55. Miao, R. *et al.* Comparative Analysis of Arabidopsis Ecotypes Reveals a Role for Brassinosteroids in Root Hydrotropism. *Plant Physiol.* **176**, 2720–2736, doi:10.1104/pp.17.01563 (2018).
56. Kuznetsov, O. A. & Hasenstein, K. H. The response of the lazy-2 tomato mutant to HGMF and red light. *Plant Physiol.* **114**, 1480–1480 (1997).

57. Weise, S. E. & Kiss, J. Z. Gravitropism of Inflorescence Stems in Starch-Deficient Mutants of *Arabidopsis*. *Int. J. Plant Sci.* **160**, 521 (1999).
58. Kiss, J. Z., Hertel, R. & Sack, F. D. Amyloplasts are necessary for full gravitropic sensitivity in roots of *Arabidopsis thaliana*. *Planta* **177**, 198–206 (1989).
59. Kuznetsov, O. A. *et al.* Composition and physical properties of starch in microgravity-grown plants. *Adv. Space Res.* **28**, 651–658 (2001).
60. Driss-Ecole, D., Legue, V., Carnero-Diaz, E. & Perbal, G. Gravisensitivity and automorphogenesis of lentil seedling roots grown on board the International Space Station. *Physiol. Plant.* **134**, 191–201, doi:10.1111/j.1399-3054.2008.01121.x (2008).
61. Volkmann, D. & Tewinkel, M. Gravisensitivity of Cress Roots. *21*, 1209–1217.
62. Brown, C. S. & Piastuch, W. C. Starch metabolism in germinating soybean cotyledons is sensitive to clinorotation and centrifugation. *Plant Cell & Environment* **17**, 341–344 (1994).
63. Bizet, F. *et al.* Both gravistimulation onset and removal trigger an increase of cytoplasmic free calcium in statocytes of roots grown in microgravity. *Sci Rep* **8**, doi:10.1038/s41598-018-29788-7 (2018).
64. Wendt, M. & Sievers, A. The polarity of statocytes and the gravisensitivity of roots are dependent on the concentration of calcium in statocytes. *Plant & cell physiology* **30**, 929–932 (1989).
65. Kordyum, E. L. A role for the cytoskeleton in plant cell gravisensitivity and Ca^{2+} -signaling in microgravity. *Cell Biol. Int.* **27**, 219–221, doi:10.1016/s1065-6995(02)00315-3 (2003).
66. Chen, X. F., Chang, M. C., Wang, B. Y. & Wu, R. Cloning of a Ca^{2+} -ATPase gene and the role of cytosolic Ca^{2+} in the gibberellin-dependent signaling pathway in aleurone cells. *Plant J.* **11**, 363–371, doi:10.1046/j.1365-313X.1997.11030363.x (1997).
67. Saboury, A. A. Stability, activity and binding properties study of alpha-amylase upon interaction with Ca^{2+} and Co^{2+} . *Biologia* **57**, 221–228 (2002).
68. Kashem, M. A., Itoh, K., Iwabuchi, S., Hori, H. & Mitsui, T. Possible involvement of phosphoinositide- Ca^{2+} signaling in the regulation of alpha-amylase expression and germination of rice seed (*Oryza sativa* L.). *Plant Cell Physiol.* **41**, 399–407 (2000).
69. Bahaji, A. *et al.* Plastidial Phosphoglucose Isomerase Is an Important Determinant of Seed Yield through Its Involvement in Gibberellin-Mediated Reproductive Development and Storage Reserve Biosynthesis in *Arabidopsis*. *Plant Cell* **30**, 2082–2098, doi:10.1105/tpc.18.00312 (2018).
70. Caspar, T., Huber, S. & Somerville, C. Alterations in growth, photosynthesis, and respiration in a starchless mutant of *Arabidopsis thaliana* (L.) deficient in chloroplast phosphoglucomutase activity. *Plant Physiol.* **79**, 11–17 (1985).
71. Caspar, T. & Pickard, B. G. Gravitropism in a starchless mutant of *Arabidopsis* - Implications for the starch-statolith theory of gravity sensing. *Planta* **177**, 185–197 (1989).
72. Tsai, H. L. *et al.* Starch synthesis in *Arabidopsis* is achieved by spatial cotranscription of core starch metabolism genes. *Plant Physiol.* **151**, 1582–1595, doi:10.1104/pp.109.144196 (2009).

73. Paulus, U., Kortje, K. H., Slenzka, K. & Rahmann, H. Correlation of altered gravity and cytochrome oxidase activity in the developing fish brain. *Journal of Brain Research-Journal Fur Hirnforschung* **37**, 103–107 (1996).
74. Fitts, R. H. *et al.* Effects of prolonged space flight on human skeletal muscle enzyme and substrate profiles. *J. Appl. Physiol.* **115**, 667–679, doi:10.1152/japplphysiol.00489.2013 (2013).
75. Faraoni, P. *et al.* Glyoxylate cycle activity in *Pinus pinea* seeds during germination in altered gravity conditions. *Plant Physiol. Biochem.* **139**, 389–394, doi:10.1016/j.plaphy.2019.03.042 (2019).
76. Gaubin, Y. *et al.* Enzyme activities and membrane lipids in *Artemia* cysts after a long duration space flight. *Advances in Space Research* **18**, 221–227 (1996).
77. Porterfield, D. M., Matthews, S. W. & Musgrave, M. E. Spaceflight Exposure Effects on Transcription, Activity, and Localization of Alcohol Dehydrogenase in the Roots of *Arabidopsis thaliana*. *Plant Physiol.* **113**, 685–693 (1997).
78. Paul, A. L. *et al.* Transgene expression patterns indicate that spaceflight affects stress signal perception and transduction in *Arabidopsis*. *Plant Physiol.* **126**, 613–621, doi:10.1104/pp.126.2.613 (2001).
79. Kolzumi, T. *et al.* Changes in membrane lipid composition in azuki bean epicotyls under hypergravity conditions: Possible role of membrane sterols in gravity resistance. *Adv. Space Res.* **39**, 1198–1203, doi:10.1016/j.asr.2007.02.040 (2007).
80. Nedukha, O., Kordyum, E. & Vorobyova, T. Sensitivity of Plant Plasma Membrane to Simulated Microgravity. *Microgravity Science and Technology* **33**, doi:10.1007/s12217-020-09865-1 (2021).

Figures




Figure 1

The design of the magnetic field chamber (MFC) with 10 magnets arranged in two stories with a total of eight compartments. The magnetic field was contained with two ferromagnetic yokes. Each compartment contained two ferromagnetic wedges and two pieces of activated charcoal cloth (ACC), and one seed cassette each. The diagram on the upper right shows the distortion of the magnetic field by the wedges (applicable to all compartments) and the projected growth of the seedling roots. The flow path of the fixative from a single fill port (F) to the drain port (D) is shown as white canals for front-visible paths; connections on the rear side between the individual compartments are shown as dark lines.




Figure 2

Example of root orientation inside seed cassettes. The predominant curvature and elongation occurred onto and along the surface of the germination paper.




Figure 3

Transcription values between space grown *Brassica rapa* seedlings (A, #1) and static ground controls (B, #23) in the presence and absence of HGMFs show strong correlation (unity slope) and low scatter ($R^2 = 0.85$ and 0.95), indicating no or low effect of magnetic fields on transcription. Panels C & D identify the examined tissues in A and B, respectively. The data sets use *PFK* as reference and show efficiency-corrected ΔC_q and log(2) transformed values.




Figure 4

Size distribution of amyloplasts in the root cap of *Brassica rapa* seedlings grown in space, on the ground or in a clinostat. The data were compiled from 2 μm thick longitudinal sections of root cap tissue after fixation in paraformaldehyde. The difference between space and clinostat amyloplasts was highly significant ($P < 10^{-4}$); the difference between static ground control and clinorotated amyloplasts was significant at the 5% level. Data were normalized based on 290, 243, and 210 amyloplast measurements for space, clinorotated, and static ground control plants, respectively.




Figure 5

Transcription values for space grown and clinorotated KSC ground controls. *Brassica rapa* seedlings (A, #2) show strong reduction of *AMY1* transcription in clinorotated seedlings (circle). A comparison between static and clinorotated seedlings (B, #27) shows higher *AMY1* transcription in non-clinorotated seedlings. Panels C & D identify the tissues in A and B, respectively. The circled data indicate *AMY1* transcription relative to *PFK*. In both cases average *AMY1* transcription was significantly different ($p = 0.002$, left and $p < 0.001$, right; $n = 64$). The data sets use *PFK* as reference and show efficiency-corrected ΔC_q and $\log(2)$ transformed values.




Figure 6

The percentage of significant ($P < 0.05$) changes in gene transcription in response to space, HGMF and clinorotation of *Brassica rapa* seedlings. Compared to *AMY1*, all other genes showed fewer significant alterations (% of 16 genes \times 16 references).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupVideo1Y5.mp4](#)
- [SupVideo2Y2.mp4](#)
- [SupplementaryInformation.docx](#)