Affine connection representation of gauge fields

Zhao-Hui Man (✉ shetcslion@163.com)

Yinchuan Hua Yang Nian Hua Bei Qu <https://orcid.org/0000-0003-4202-8741>

Research Article

Keywords: affine connection representation of gauge fields, geometric meaning of coupling constant, time metric, reference-system, distribution of gradient directions

Posted Date: April 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-173799/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Affine connection representation of gauge fields

Zhao-Hui Man

the date of receipt and acceptance should be inserted later

Abstract There are two ways to unify gravitational field and gauge field. One is to represent gravitational field as principal bundle connection, and the other is to represent gauge field as affine connection. Poincaré gauge theory and metric-affine gauge theory adopt the first approach. This paper adopts the second.

An affine connection is used to establish a unified coordinate description of gauge field and gravitational field. This theory has the following advantages.

(i) Gauge field and gravitational field can both be represented by affine connection; they can be described by a unified spatial frame.

(ii) Time can be regarded as the total metric with respect to all dimensions of internal coordinate space and external coordinate space. On-shell can be regarded as gradient direction. Quantum theory can be regarded as a geometric theory of distribution of gradient directions. Hence, gravitational theory and quantum theory obtain the same view of time and space and a unified description of evolution in affine connection representation of gauge fields.

(iii) Chiral asymmetry, coupling constants, MNS mixing and CKM mixing can appear spontaneously as geometric properties in affine connection representation, whereas in $U(1) \times SU(2) \times SU(3)$ principal bundle connection representation they can just only be artificially set up. Some postulates of the Standard Model can be turned into theorems in affine connection representation, so they are not necessary to be regarded as postulates anymore.

(iv) The unification theory of gauge fields that are represented by affine connection can avoid the problem that a proton decays into a lepton.

(v) Since the concept of point particle is thoroughly abandoned, this theory is not required to be renormalized.

(iv) There exists a possible geometric interpretation to the color confinement of quarks.

The Standard Model is not possessed of the above advantages. In the affine connection representation, we can get better interpretations of these physical properties. This is probably a necessary step towards the ultimate theory of physics.

Keywords affine connection representation of gauge fields · geometric meaning of coupling constant · time metric · reference-system · distribution of gradient directions

Mathematics Subject Classification (2010) Primary 51P05, 70A05, 53Z05 · Secondary 53C05, 58A05

Contents

1	Introduction	2
1.1	Background and purpose	2
1.2	Ideas and methods	3
1.3	Content and organization	4
2	Mathematical preparations	5
2.1	Geometric manifold	5
2.2	Metric and semi-metric	5
2.3	Gauge transformation in affine connection representation	5
2.4	Coordinate transformation of holonomic connection and frame transformation of gauge connection	6
3	The evolution in affine connection representation of gauge fields	7

3.1	The relation between time and space	7
3.2	Evolution path as a submanifold	8
3.3	Evolution lemma	9
3.4	On-shell evolution as a gradient	10
3.5	On-shell evolution of potential field and affine connection representation of Yang-Mills equation	10
3.6	On-shell evolution of general charge and affine connection representation of mass, energy, momentum and action	11
3.7	Inversion transformation in affine connection representation	13
3.8	Two dual descriptions of gradient direction field	14
3.9	Quantum evolution as a distribution of gradient directions	15
4	Affine connection representation of gauge fields in classical spacetime	18
4.1	Classical spacetime submanifold	18
4.2	Classical spacetime reference-system	19
4.3	Affine connection representation of classical spacetime evolution	20
4.4	Affine connection representation of Dirac equation	21
4.5	From classical spacetime back to full-dimensional space	23
5	Affine connection representation of the gauge field of weak-electromagnetic interaction	24
6	Affine connection representation of the gauge field of strong interaction	28
7	Affine connection representation of the unified gauge field	29
8	Conclusions	34

1 Introduction

1.1 Background and purpose

We know that in gauge theory, the field strength and the gauge-covariant derivative

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc}A_\mu^b A_\nu^c, \quad D_\mu = \partial_\mu - igT^a A_\mu^a$$

both contain a coupling constant g , which measures the strength of interaction. A problem is that why is there a coupling constant g ?

If representing gauge fields by affine connection, we can obtain a nice interpretation. For example, if we use Γ_{MNP} to represent gauge potentials, it is not hard to find some specific conditions to turn the curvature tensor R_{NPQ}^M to

$$\begin{aligned} R_{MNPQ} &= \partial_P \Gamma_{MNQ} - \partial_Q \Gamma_{MNP} + \Gamma_{MHP} \Gamma_{NQ}^H - \Gamma_{NP}^H \Gamma_{MHQ} \\ &= \partial_P \Gamma_{MNQ} - \partial_Q \Gamma_{MNP} + G^{RH} (\Gamma_{MHP} \Gamma_{RNQ} - \Gamma_{RNP} \Gamma_{MHQ}). \end{aligned} \quad (1)$$

Thus, R_{MNPQ} can be used to represent field strength. In addition, for any ρ_M , we see that

$$\rho_{M;P} = \partial_P \rho_M - \Gamma_{MP}^H \rho_H = \partial_P \rho_M - G^{RH} \Gamma_{RMP} \rho_H. \quad (2)$$

Eq.(1) and Eq.(2) mean that the coupling constant g may have a geometric meaning, which originates from G^{RH} .

This implies that only when affine connection is adopted to represent gauge field can some physical properties be better interpreted. On the other hand, in the general relativity theory, gravitational field is also described by affine connection, so it is convenient to describe gravitational field and gauge field uniformly by affine connection. So it is necessary to study the affine connection representation of gauge fields. This is the basic motivation of this paper. To represent gauge fields by affine connection is probably a necessary step towards the ultimate theory of physics.

We know that there are the following two ways to unify gravitational field and gauge field.

One way is to represent gravitational field as principal bundle connection. We can take the transformation group $Gravi(3, 1)$ of gravitational field as the structure group of principal bundle to establish a gauge theory of gravitational field, the local transformation group of which is in the form of $Gravi(3, 1) \otimes Gauge(n)$, e.g. Poincaré gauge theory [1–11] and metric-affine gauge theory [12–23]. Gravitational field and gauge field can be uniformly described by connections on principal bundle.

The other way is to represent gauge field as affine connection. This is the approach adopted by this paper. Gravitational field and gauge field can both be described by affine connection. Besides, we also establish an affine connection representation of elementary particles. In this way, Elementary particle fields obtain explicit descriptions of geometric entity. The MNS mixing of three generations of leptons obtains a new geometric interpretation. Some postulates of the Standard Model that are based on principal bundle connection representation can be turned into theorems in affine connection representation, so they are not necessary to be regarded as postulates anymore.

1.2 Ideas and methods

We divide the problem of establishing affine connection representation of gauge fields into three parts as follows.

(I) Which affine connection is suitable for describing not only gravitational field, but also gauge field and elementary particle field?

(II) How to describe the evolution of these fields in affine connection representation?

(III) What are the concrete forms of electromagnetic, weak and strong interaction fields in affine connection representation?

For the problem (I). On a Riemannian manifold (M, G) , the metric tensor can be expressed as $G_{MN} = \delta_{AB} B_M^A B_N^B$ and $G^{MN} = \delta^{AB} C_A^M C_B^N$, where B_M^A and C_A^M are semi-metrics, or to say frame fields. It is evident that semi-metric is more fundamental than metric, so we hope B_M^A or C_A^M is regarded as a unified frame field of gravitational field and gauge field, and the frame transformation of B_M^A or C_A^M is regarded as gauge transformation. Hence, we need a more general manifold (M, B_M^A) rather than the Riemannian manifold (M, G) .

Next, we put metric and semi-metric together to construct a new affine connection, in this way, gravitational field and various gauge fields can be unified on a manifold (M, B_M^A) that is defined by semi-metric.

In addition, we notice that in the theories based on principal bundle connection representation:

(1) Several complex-valued functions which satisfy the Dirac equation, are sometimes used to refer to a charged lepton field l , and sometimes a neutrino field ν . It is not clear how to distinguish these field functions l and ν by inherent geometric constructions.

(2) In the Standard Model, gauge potentials are abstract; they have no inherent geometric constructions. In other words, the Levi-Civita connection $\Gamma_{\nu\rho}^\mu$ of gravity is constructed by the metric $g_{\mu\nu}$, however it is not explicit what geometric quantity the connection A_μ^a of gauge field is constructed by.

By contrast, in the affine connection representation of this paper, we are able to use the semi-metrics B_M^A and C_A^M of internal coordinate space to endow particle fields l and ν and gauge fields A_μ^a with geometric constructions. Thus, they are not only irreducible representations of group, but also possessed of concrete geometric entities.

For the problem (II). There is a fundamental difficulty that time is a component of gravitational field, but not a component of gauge field. This leads to an essential difference between the description of evolution of gravitational field and that of gauge field. In this case, it seems difficult to obtain a unified theory of evolution in affine connection representation. Nevertheless, we find that, we can define time as the total metric with respect to all dimensions of internal coordinate space and external coordinate space, and define evolution as one-parameter group of diffeomorphism, to overcome the above difficulty.

Now that gauge field and gravitational field are both represented as affine connection, then the properties that are related to gauge field, such as charge, current, mass, energy, momentum and action, must have corresponding affine connection representations. Thus, Yang-Mills equation, energy-momentum equation and Dirac equation are turned into geometric properties in gradient direction, in other words, on-shell evolution is characterized by gradient direction. Correspondingly, quantum theory can be interpreted as a geometric theory of distribution of gradient directions.

For the problem (III). The basic idea is that on a $\mathfrak{D}$ -dimensional manifold, the components B_m^a and C_a^m of semi-metrics B_M^A and C_A^M with $m, a \in \{4, 5, \dots, \mathfrak{D}\}$ are regarded as the frame field of electromagnetic, weak and strong interactions. The other components of B_M^A and C_A^M are regarded as the frame field of gravity.

We take the affine connection as

$$\begin{aligned} \Gamma_{NP}^M &\triangleq \frac{1}{2} ([{}^M_{NP}] + \{{}^M_{NP}\}) = \frac{1}{2} [C_A^M (D_P B_N^A) + \{{}^M_{NP}\}] = \frac{1}{2} [C_A^M (D_C B_N^A) b_P^C + \{{}^M_{NP}\}] \\ &= \frac{1}{2} \left[C_A^M \left(\frac{\partial B_N^A}{\partial \zeta^C} + ({}^A_{BC}) B_N^B \right) b_P^C + \frac{1}{2} G^{MQ} \left(\frac{\partial G_{NQ}}{\partial x^P} + \frac{\partial G_{PQ}}{\partial x^N} - \frac{\partial G_{NP}}{\partial x^Q} \right) \right] \\ &= \frac{1}{2} \left[\left(C_A^M \frac{\partial B_N^A}{\partial x^P} + C_A^M ({}^A_{BP}) B_N^B \right) + \frac{1}{2} G^{MQ} \left(\frac{\partial G_{NQ}}{\partial x^P} + \frac{\partial G_{PQ}}{\partial x^N} - \frac{\partial G_{NP}}{\partial x^Q} \right) \right], \end{aligned} \quad (3)$$

where $b_P^C \triangleq \frac{\partial \zeta^C}{\partial x^P}$ is a local coordinate transformation, $\{{}^M_{NP}\}$ is Christoffel symbol, $G_{MN} = \delta_{AB} B_M^A B_N^B$,

$$[{}^M_{NP}] \triangleq C_A^M (D_P B_N^A) = C_A^M \frac{\partial B_N^A}{\partial x^P} + C_A^M ({}^A_{BP}) B_N^B$$

is said to be a gauge connection, and Γ_{NP}^M is said to be a holonomic connection. $({}^A_{BP}) \triangleq ({}^A_{BC}) b_P^C$.

$$({}^A_{BC}) \triangleq \frac{1}{2} C_{A'}^A \left(\frac{\partial B_B^{A'}}{\partial \zeta^C} + \frac{\partial B_C^{A'}}{\partial \zeta^B} \right)$$

is said to be a torsion-free simple connection. Thus,

$$\Gamma_{MNP} = \frac{1}{2} ([MNP] + \{MNP\}) = \frac{1}{2} \left[\delta_{AD} B_M^D \left(\frac{\partial B_N^A}{\partial x^P} + \binom{A}{BP} B_N^B \right) + \frac{1}{2} \left(\frac{\partial G_{NM}}{\partial x^P} + \frac{\partial G_{PM}}{\partial x^N} - \frac{\partial G_{NP}}{\partial x^M} \right) \right].$$

For the sake of simplicity, we firstly consider the affine connection representation of gauge fields without gravitation. That is to say, let

$$s, i, j = 1, 2, 3; \quad a, m, n, l, q = 4, 5, \dots, \mathfrak{D}; \quad A, B, M, N, P = 1, 2, \dots, \mathfrak{D};$$

and consider a $\mathfrak{D}$ -dimensional manifold $(M, (B_f)_M^A)$ that satisfies the following conditions:

- (i) $B_i^s = \delta_i^s$, $B_i^a = 0$, $B_m^s = 0$;
- (ii) $G_{ij} = \delta_{ij}$, $G_{mn} = \text{const}$, $G_{mi} = 0$;
- (iii) When $m \neq n$, $G_{mn} = 0$.

Thus, $\{MNP\} = 0$, $[MNP] \neq 0$ in general. The components Γ_{mnP} of $\Gamma_{MNP} = \frac{1}{2} [MNP]$ with $m, n \in \{4, 5, \dots, \mathfrak{D}\}$ describe gauge potentials of electromagnetic, weak and strong interactions. We also use the affine connection Γ_{NP}^M to construct elementary particle fields ρ_{MN} . The components ρ_{mn} of ρ_{MN} with $m, n \in \{4, 5, \dots, \mathfrak{D}\}$ describe field functions of leptons and quarks.

The components G^{mn} of G^{MN} with $m, n \in \{4, 5, \dots, \mathfrak{D}\}$ describe coupling constants of particle fields ρ_{mn} and gauge potentials Γ_{mnP} . The other components of G^{MN} are the metrics of gravitational field. The other components of ρ_{MN} and Γ_{MNP} provide possible candidates for dark matters and their interactions.

1.3 Content and organization

In this paper, we are going to show how to construct the affine connection representation of gauge fields. Sections are organized as follows.

Corresponding to the problem (I), in section 2 we make some necessary mathematical preparations, and discuss the coordinate transformation and frame transformation of the above connection. Meanwhile, in order to make the languages that are used to describe gauge field and gravitational field unified and harmonized, we generalize the notion of reference system, and give it a strict mathematical definition. The reference system in conventional sense is just only defined on a local coordinate neighborhood, and it has only $(1 + 3)$ dimensions. But in this paper we define the concept of reference-system over the entire manifold. It is possessed of more dimensions but different from Kaluza-Klein theory [24–26] and string theories [27–39]. Thus, both of gravitational field and gauge field are regarded as special cases of such a concept of reference-system.

Corresponding to the problem (II), in section 3 we establish the general theory of evolution in affine connection representation of gauge fields, and in section 4 we discuss the application of this general theory of evolution to $(1 + 3)$ -dimensional classical spacetime.

Corresponding to the problem (III), in sections 5 to 7 we show concrete forms of affine connection representations of electromagnetic, weak and strong interaction fields.

Some important topics are organized as follows.

(1) Time is regarded as the total metric with respect to all spatial dimensions including external coordinate space and internal coordinate space, see Definition 3.1.1 and Remark 4.2.1 for detail. The *CPT* inversion is interpreted as the composition of the total inversion of coordinates and the total inversion of metrics, see section 3.7 for detail. The conventional $(1 + 3)$ -dimensional Minkowski coordinate x^μ originates from the general $\mathfrak{D}$ -dimensional coordinate x^M . The construction method of extra dimensions is different from those of Kaluza-Klein theory and string theory, see section 4.2 for detail.

(2) On-shell evolution is characterized by gradient direction field. See sections 3.4, 3.5, 3.6 and 4.3 for detail. Quantum theory is regarded as a geometric theory of distribution of gradient directions. We show two dual descriptions of gradient direction. They just exactly correspond to the Schrödinger picture and the Heisenberg picture. In these points of view, the gravitational theory and quantum theory become coordinated. They have a unified description of evolution, and the definition of Feynman propagator is simplified to a stricter form. See sections 3.8 and 3.9 for detail.

(3) Yang-Mills equation originates from a geometric property of gradient direction. We show the affine connection representation of Yang-Mills equation. See section 3.5 and section 4.5 for detail.

(4) Energy-momentum equation originates from a geometric property of gradient direction. We show the affine connection representation of mass, energy, momentum and action, see section 3.6, Definition 4.3.1 and Discussion 4.3.1 for detail. Furthermore, we also show the affine connection representation of Dirac equation, see section 4.4 for detail.

(5) Why do not neutrinos participate in the electromagnetic interactions? And why do not right-handed neutrinos participate in the weak interactions with W bosons? In the Standard Model, they are both postulates. However, in the new framework, they are natural and geometric results of affine connection representation of gauge fields, therefore not necessary to be regarded as postulates anymore. See Proposition 5.2 and Proposition 7.1 for detail.

(6) Why do leptons and quarks both have three generations? We give an idea in section 7. It is a natural geometric result of affine connection representation of gauge fields.

2 Mathematical preparations

2.1 Geometric manifold

In order to make the languages that are used to describe gauge field and gravitational field unified and harmonized, we adopt the following definition.

Definition 2.1.1. Let M be a $\mathfrak{D}$ -dimensional connected smooth real manifold. $\forall p \in M$, take a coordinate chart (U_p, φ_{U_p}) on a neighborhood U_p of p . They constitute a coordinate covering

$$\varphi \triangleq \{(U_p, \varphi_{U_p})\}_{p \in M},$$

which is said to be a **point-by-point covering**. For the sake of simplicity, U_p can be denoted by U , and φ_{U_p} by φ_U .

Let φ and ψ be two point-by-point coverings. For the two coordinate frames φ_U and ψ_U on the neighborhood U of point p , if

$$f_p \triangleq \varphi_U \circ \psi_U^{-1} : \psi_U(U) \rightarrow \varphi_U(U), \quad \xi^A \mapsto x^M$$

is a smooth homeomorphism, f_p is called a **local reference-system**, where ψ_U is said to be the **basis coordinate frame** of f_p , and φ_U is said to be the **performance coordinate frame** of f_p .

If every $p \in M$ is endowed with a local reference-system $f(p)$, and we require the semi-metrics B_M^A and C_A^M in Eq.(6) to be smooth real functions on M , then

$$f : M \rightarrow REF, \quad p \mapsto f(p) \quad (4)$$

is said to be a **reference-system** on M , and (M, f) is said to be a **geometric manifold**.

2.2 Metric and semi-metric

In the absence of a special declaration, the indices take values as $A, B, C, D, E = 1, 2, \dots, \mathfrak{D}$ and $M, N, P, Q, R = 1, 2, \dots, \mathfrak{D}$. The derivative functions

$$b_M^A \triangleq \frac{\partial \xi^A}{\partial x^M}, \quad c_A^M \triangleq \frac{\partial x^M}{\partial \xi^A} \quad (5)$$

of $f(p)$ on U_p define the semi-metrics (or to say frame field) B_M^A and C_A^M of f on the manifold M , that are

$$B_M^A : M \rightarrow \mathbb{R}, \quad p \mapsto B_M^A(p) \triangleq (b_{f(p)})_M^A(p), \quad C_A^M : M \rightarrow \mathbb{R}, \quad p \mapsto C_A^M(p) \triangleq (c_{f(p)})_A^M(p). \quad (6)$$

Let $\delta_{AB} = \delta^{AB} = \delta_B^A = \text{Kronecker}(A, B)$ and $\varepsilon_{MN} = \varepsilon^{MN} = \varepsilon_N^M = \text{Kronecker}(M, N)$. The metric tensors of f are

$$G_{MN} = \delta_{AB} B_M^A B_N^B, \quad H_{AB} = \varepsilon_{MN} C_A^M C_B^N. \quad (7)$$

Similarly, it can also be defined that $\bar{b}_A^M \triangleq \frac{\partial \xi_A}{\partial x^M}$, $\bar{c}_M^A \triangleq \frac{\partial x_M}{\partial \xi_A}$ and corresponding $\bar{B}_A^M$, $\bar{C}_M^A$.

2.3 Gauge transformation in affine connection representation

$\forall p \in M$, $f(p) \triangleq \rho_U \circ \psi_U^{-1}$ induces **local reference-system transformations**

$$\begin{aligned} L_{f(p)} : k(p) &\triangleq \psi_U \circ \varphi_U^{-1} \mapsto \rho_U \circ \varphi_U^{-1} = f(p) \circ k(p), \\ R_{f(p)} : h(p) &\triangleq \varphi_U \circ \rho_U^{-1} \mapsto \varphi_U \circ \psi_U^{-1} = h(p) \circ f(p), \end{aligned}$$

and **reference-system transformations** on the manifold M

$$L_f : p \mapsto L_{f(p)}, \quad R_f : p \mapsto R_{f(p)}. \quad (8)$$

We also speak of L_f and R_f as **(affine) gauge transformations**.

- (i) L_f and R_f are identical transformations if and only if $[B_M^A]$ of f is an identity matrix.
- (ii) L_f and R_f are flat transformations if and only if $\forall p_1, p_2 \in M, B_M^A(p_1) = B_M^A(p_2)$.
- (iii) L_f and R_f are orthogonal transformations if and only if $\delta_{AB} B_M^A B_N^B = \varepsilon_{MN}$.

The totality of all reference-system transformations on M is denoted by $GL(M)$, which is a subgroup of $\bigotimes_{p \in M} GL(\mathfrak{D}, \mathbb{R})_p$, where $\bigotimes$ represents external direct product.

2.4 Coordinate transformation of holonomic connection and frame transformation of gauge connection

Suppose there are reference-systems g and $\mathfrak{g}$ on the manifold M , denote $\mathcal{G} \triangleq \mathfrak{g} \circ g$, and $\forall p \in M$, on the neighborhood U of p , $g(p)$ and $\mathfrak{g}(p)$ satisfy

$$(U, x^M) \xleftarrow{g(p)} (U, \zeta^A) \xleftarrow{\mathfrak{g}(p)} (U, \beta^{A'}).$$

On the geometric manifold $(M, \mathfrak{g})$ we define **torsion-free simple connection** D and its coefficients $(\frac{A}{BC})_{\mathfrak{g}}$ by

$$D \frac{\partial}{\partial \zeta^B} \triangleq (\omega_{\mathfrak{g}})_B^A \otimes \frac{\partial}{\partial \zeta^A} = (\frac{A}{BC})_{\mathfrak{g}} d\zeta^C \otimes \frac{\partial}{\partial \zeta^A} = \frac{1}{2} (C_{\mathfrak{g}})_{A'}^A \left(\frac{\partial (B_{\mathfrak{g}})_{B'}^A}{\partial \zeta^C} + \frac{\partial (B_{\mathfrak{g}})_{C'}^A}{\partial \zeta^B} \right) d\zeta^C \otimes \frac{\partial}{\partial \zeta^A}. \quad (9)$$

Then, we can compute the absolute derivative of the frame field $\frac{\partial}{\partial x^N}$

$$\begin{aligned} D \frac{\partial}{\partial x^N} &= D \left((B_g)_N^B \frac{\partial}{\partial \zeta^B} \right) = d(B_g)_N^B \otimes \frac{\partial}{\partial \zeta^B} + (B_g)_N^B D \frac{\partial}{\partial \zeta^B} \\ &= \frac{\partial (B_g)_N^B}{\partial \zeta^C} d\zeta^C \otimes \frac{\partial}{\partial \zeta^B} + (B_g)_N^B (\frac{A}{BC})_{\mathfrak{g}} d\zeta^C \otimes \frac{\partial}{\partial \zeta^A} = \left(\frac{\partial (B_g)_N^A}{\partial \zeta^C} + (B_g)_N^B (\frac{A}{BC})_{\mathfrak{g}} \right) d\zeta^C \otimes \frac{\partial}{\partial \zeta^A}, \end{aligned}$$

Thus, it is obtained that

$$D_C (B_g)_N^A = \frac{\partial (B_g)_N^A}{\partial \zeta^C} + (B_g)_N^B (\frac{A}{BC})_{\mathfrak{g}}.$$

Denote $D_P \triangleq (b_{g(p)})_P^C D_C$, thus we can define on $(M, \mathcal{G})$ the required **gauge connection**, which is

$$[{}^M_{NP}]_{\mathcal{G}} \triangleq (C_g)_A^M D_P (B_g)_N^A = (C_g)_A^M \frac{\partial (B_g)_N^A}{\partial x^P} + (C_g)_A^M (\frac{A}{BP})_{\mathfrak{g}} (B_g)_N^B. \quad (10)$$

It is important that the gauge connection $[{}^M_{NP}]_{\mathcal{G}}$ that is defined in this way is not only an affine connection on $(M, \mathcal{G})$, but also a connection on frame bundle.

(I). As an affine connection, under the coordinate transformation $L_{k(p)} : (U, x^M) \rightarrow (U, x^{M'})$, $b_{M'}^M \triangleq \frac{\partial x^M}{\partial x^{M'}}$, $c_M^{M'} \triangleq \frac{\partial x^{M'}}{\partial x^M}$, $(B_g)_M^A \mapsto (B_g)_{M'}^A = b_{M'}^M (B_g)_M^A$, $(C_g)_A^M \mapsto (C_g)_{A'}^{M'} = c_M^{M'} (C_g)_A^M$. Consequently, the gauge connection $[{}^M_{NP}]_{\mathcal{G}}$ is transformed according to

$$L_{k(p)} : [{}^M_{NP}]_{\mathcal{G}} \mapsto [{}^{M'}_{N'P'}]_{\mathcal{G}} = c_M^{M'} [{}^M_{NP}]_{\mathcal{G}} b_{N'}^N b_{P'}^P + c_M^{M'} \frac{\partial b_{N'}^M}{\partial x^{P'}}. \quad (11)$$

Due to Eq.(11), under the coordinate transformation, the **holonomic connection**

$$\begin{aligned} (\Gamma_g)_{NP}^M &\triangleq \frac{1}{2} \left([{}^M_{NP}]_{\mathcal{G}} + \{{}^M_{NP}\}_{\mathcal{G}} \right) \\ &= \frac{1}{2} \left[\left((C_g)_A^M \frac{\partial (B_g)_N^A}{\partial x^P} + (C_g)_A^M (\frac{A}{BP})_{\mathfrak{g}} (B_g)_N^B \right) + \frac{1}{2} (G_g)^{MQ} \left(\frac{\partial (G_g)_{NQ}}{\partial x^P} + \frac{\partial (G_g)_{PQ}}{\partial x^N} - \frac{\partial (G_g)_{NP}}{\partial x^Q} \right) \right] \end{aligned} \quad (12)$$

is transformed according to

$$L_{k(p)} : (\Gamma_g)_{NP}^M \mapsto (\Gamma_g)_{N'P'}^{M'} = c_M^{M'} (\Gamma_g)_{NP}^M b_{N'}^N b_{P'}^P + c_M^{M'} \frac{\partial b_{N'}^M}{\partial x^{P'}}. \quad (13)$$

(II). As a connection on frame bundle, under the frame transformation $L_k : (M, \mathcal{G}) \mapsto (M, \mathcal{G}')$, $\frac{\partial}{\partial x^M} \mapsto \frac{\partial}{\partial x^{M'}} = (B_k)_{M'}^M \frac{\partial}{\partial x^M}$, $(B_g)_M^A \mapsto (B_{g'})_{M'}^A = (B_k)_{M'}^M (B_g)_M^A$, $(C_g)_A^M \mapsto (C_{g'})_{A'}^{M'} = (C_k)_M^{M'} (C_g)_A^M$. Consequently, the gauge connection $[{}^M_{NP}]_{\mathcal{G}}$ is transformed according to

$$L_k : [{}^M_{NP}]_{\mathcal{G}} \mapsto [{}^{M'}_{N'P'}]_{\mathcal{G}'} = [{}^{M'}_{N'P}]_{\mathcal{G}'} b_{P'}^P = \left((C_k)_M^{M'} [{}^M_{NP}]_{\mathcal{G}} (B_k)_{N'}^N + (C_k)_M^{M'} \frac{\partial (B_k)_{N'}^M}{\partial x^P} \right) b_{P'}^P, \quad (14)$$

Eq.(11) and Eq.(14) show that $[{}^M_{NP}]_{\mathcal{G}}$ is not only an affine connection, but also a connection on frame bundle. Apply Eq.(11)~(14) to the curvature tensors

$$\begin{aligned} [{}^M_{NPQ}] &\triangleq \frac{\partial [{}^M_{NQ}]}{\partial x^P} - \frac{\partial [{}^M_{NP}]}{\partial x^Q} + [{}^M_{HP}] [{}^H_{NQ}] - [{}^H_{NP}] [{}^M_{HQ}], \\ \{ {}^M_{NPQ} \} &\triangleq \frac{\partial \{ {}^M_{NQ} \}}{\partial x^P} - \frac{\partial \{ {}^M_{NP} \}}{\partial x^Q} + \{ {}^M_{HP} \} \{ {}^H_{NQ} \} - \{ {}^H_{NP} \} \{ {}^M_{HQ} \}, \\ R_{NPQ}^M &\triangleq \frac{\partial \Gamma_{NQ}^M}{\partial x^P} - \frac{\partial \Gamma_{NP}^M}{\partial x^Q} + \Gamma_{HP}^M \Gamma_{NQ}^H - \Gamma_{NP}^H \Gamma_{HQ}^M, \end{aligned}$$

then it is obtained that

$$\begin{aligned} L_k : [{}^M_{NPQ}]_{\mathcal{G}} &\mapsto [{}^{M'}_{N'P'Q'}]_{\mathcal{G}'} = [{}^{M'}_{N'PQ}]_{\mathcal{G}'} b_{P'}^P b_{Q'}^Q = \left((C_k)_M^{M'} [{}^M_{NPQ}]_{\mathcal{G}} (B_k)_{N'}^N \right) b_{P'}^P b_{Q'}^Q, \\ L_{k(p)} : [{}^M_{NPQ}]_{\mathcal{G}} &\mapsto [{}^{M'}_{N'P'Q'}]_{\mathcal{G}'} = c_M^{M'} [{}^M_{NPQ}]_{\mathcal{G}} b_{N'}^N b_{P'}^P b_{Q'}^Q, \\ L_{k(p)} : \{ {}^M_{NPQ} \}_{\mathcal{G}} &\mapsto \{ {}^{M'}_{N'P'Q'} \}_{\mathcal{G}'} = c_M^{M'} \{ {}^M_{NPQ} \}_{\mathcal{G}} b_{N'}^N b_{P'}^P b_{Q'}^Q, \\ L_{k(p)} : (R_{\mathcal{G}})_{NPQ}^M &\mapsto (R_{\mathcal{G}})_{N'P'Q'}^{M'} = c_M^{M'} (R_{\mathcal{G}})_{NPQ}^M b_{N'}^N b_{P'}^P b_{Q'}^Q. \end{aligned} \quad (15)$$

We see from Eq.(15) that the $[{}^M_{NPQ}]_{\mathcal{G}}$ without gravitation is both a curvature tensor of affine connection, and a curvature tensor on frame bundle, and that the $(R_{\mathcal{G}})_{NPQ}^M$ with gravitation is a curvature tensor of affine connection, but not a curvature tensor on frame bundle. In other words, under the gauge transformation L_k , $[{}^M_{NPQ}]_{\mathcal{G}}$ and $[{}^{M'}_{N'P'Q'}]_{\mathcal{G}'}$ represent the same physical state, while $(R_{\mathcal{G}})_{NPQ}^M$ and $(R_{\mathcal{G}'})_{N'P'Q'}^{M'}$ represent different physical states. This shows that the gravitational field in $(R_{\mathcal{G}})_{NPQ}^M$ makes the gauge frames B_M^A and C_A^M have physical effects.

3 The evolution in affine connection representation of gauge fields

Now that we have the required affine connection, next we have to solve the problem that how to describe the evolution in affine connection representation.

In the existing theories, time is a component of gravitational field, but not a component of gauge field. This leads to an essential difference between the description of evolution of gravitational field and that of gauge field. In this case, it is difficult to obtain a unified theory of evolution in affine connection representation. We adopt the following way to overcome this difficulty.

3.1 The relation between time and space

Definition 3.1.1. Suppose $M = P \times N$ and $r \triangleq \dim P = 3$. Let

$$A, B, M, N = 1, \dots, \mathfrak{D}; \quad s, i = 1, \dots, r; \quad a, m = r + 1, \dots, \mathfrak{D}.$$

On a geometric manifold (M, f) , the $d\xi^0$ and dx^0 which are defined by

$$\begin{aligned} (d\xi^0)^2 &\triangleq \sum_{A=1}^{\mathfrak{D}} (d\xi^A)^2 = \delta_{AB} d\xi^A d\xi^B = G_{MN} dx^M dx^N, \\ (dx^0)^2 &\triangleq \sum_{M=1}^{\mathfrak{D}} (dx^M)^2 = \varepsilon_{MN} dx^M dx^N = H_{AB} d\xi^A d\xi^B \end{aligned} \quad (16)$$

are said to be **total space metrics** or **time metrics**.

We also suppose

$$\begin{aligned} (d\xi^{(P)})^2 &\triangleq \sum_{s=1}^r (d\xi^s)^2, & (d\xi^{(N)})^2 &\triangleq \sum_{a=r+1}^{\mathfrak{D}} (d\xi^a)^2, \\ (dx^{(P)})^2 &\triangleq \sum_{i=1}^r (dx^i)^2, & (dx^{(N)})^2 &\triangleq \sum_{m=r+1}^{\mathfrak{D}} (dx^m)^2. \end{aligned}$$

$d\xi^{(N)}$ and $dx^{(N)}$ are regarded as proper-time metrics. For convenience, P is said to be **external space** and N is said to be **internal space**.

Remark 3.1.1. The above definition implies a new viewpoint about time and space. The relation between time and space in this way is different from the Minkowski coordinates x^μ ($\mu = 0, 1, 2, 3$). Time and space are not the components on an equal footing anymore, but have a relation of total to component. It can be seen later that time reflects the total evolution in the full-dimensional space, while a specific spatial dimension reflects just a partial evolution in a specific direction.

3.2 Evolution path as a submanifold

Definition 3.2.1. Let there be reference-systems $f, g, \mathfrak{f}, \mathfrak{g}$ on a manifold M , such that $\forall p \in M$, on the neighborhood U of p ,

$$(U, \alpha^{A'}) \xrightarrow{\mathfrak{f}(p)} (U, \xi^A) \xrightarrow{f(p)} (U, x^M) \xleftarrow{g(p)} (U, \zeta^A) \xleftarrow{\mathfrak{g}(p)} (U, \beta^{A'}). \quad (17)$$

Denote $\mathcal{F} \triangleq \mathfrak{f} \circ f$ and $\mathcal{G} \triangleq \mathfrak{g} \circ g$, then we say $\mathcal{F}$ and $\mathcal{G}$ **motion relatively and interact mutually**, and also say that $\mathcal{F}$ evolves in $\mathcal{G}$, or $\mathcal{F}$ evolves on the geometric manifold $(M, \mathcal{G})$. Meanwhile, $\mathcal{G}$ evolves in $\mathcal{F}$, or to say $\mathcal{G}$ evolves on $(M, \mathcal{F})$.

From Eq.(10) we know that in $\mathcal{F}$ and $\mathcal{G}$, gauge fields originate from $\mathfrak{f}$ and $\mathfrak{g}$, and gravitational fields $(G_{\mathcal{F}})_{MN}$ and $(G_{\mathcal{G}})_{MN}$ are effected by $\mathfrak{f}$ and $\mathfrak{g}$, respectively. We are going to describe their evolutions step by step in the following sections.

Let there be a one-parameter group of diffeomorphisms

$$\varphi_X : M \times \mathbb{R} \rightarrow M$$

acting on M , such that $\varphi_X(p, 0) = p$. Thus, φ_X determines a smooth tangent vector field X on M . If X is nonzero everywhere, we say φ_X is a **set of evolution paths**, and X is an **evolution direction field**. Let $T \subseteq \mathbb{R}$ be an interval, then the regular imbedding

$$L_p \triangleq \varphi_{X,p} : T \rightarrow M, \quad t \mapsto \varphi_X(p, t) \quad (18)$$

is said to be an **evolution path** through p . The tangent vector $\frac{d}{dt} \triangleq [L_p] = X(p)$ is called an **evolution direction** at p . For the sake of simplicity, we also denote $L_p \triangleq L_p(T) \subset M$, then

$$\pi : L_p \rightarrow M, \quad q \mapsto q \quad (19)$$

is also a regular imbedding. If it is not necessary to emphasize the point p , L_p is denoted by L concisely.

In order to describe physical evolution, next we are going to strictly describe the mathematical properties of the reference-systems f and g sending onto the evolution path L .

Definition 3.2.2. Let the time metrics of (U, ξ^A) , (U, x^M) , (U, ζ^A) be $d\xi^0$, dx^0 , $d\zeta^0$, respectively. On $U_L \triangleq U \cap L_p$ we have parameter equations

$$\begin{aligned} \xi^A &= \xi^A(x^0), & x^M &= x^M(\xi^0), & \zeta^A &= \zeta^A(x^0), \\ \xi^0 &= \xi^0(x^0), & x^0 &= x^0(\xi^0), & \zeta^0 &= \zeta^0(x^0). \end{aligned} \quad (20)$$

Take f for example, according to Eq.(20), on U_L we define

$$\begin{aligned} b_0^A &\triangleq \frac{d\xi^A}{dx^0}, & b_0^0 &\triangleq \frac{d\xi^0}{dx^0}, & \varepsilon_0^M &\triangleq \frac{dx^M}{dx^0} = b_0^0 c_0^M = b_0^A c_A^M, \\ c_0^M &\triangleq \frac{dx^M}{d\xi^0}, & c_0^0 &\triangleq \frac{dx^0}{d\xi^0}, & \delta_0^A &\triangleq \frac{d\xi^A}{d\xi^0} = c_0^0 b_0^A = c_0^M b_M^A. \end{aligned}$$

Define $d\xi_0 \triangleq \frac{dx^0}{d\xi^0} dx^0$ and $dx_0 \triangleq \frac{d\xi^0}{dx^0} d\xi^0$, which induce $\frac{d}{d\xi_0}$ and $\frac{d}{dx_0}$, such that $\left\langle \frac{d}{d\xi_0}, d\xi_0 \right\rangle = 1$, $\left\langle \frac{d}{dx_0}, dx_0 \right\rangle = 1$. So we can also define

$$\begin{aligned} \bar{b}_A^0 &\triangleq \frac{d\xi_A}{dx_0}, & \bar{b}_0^0 &\triangleq \frac{d\xi_0}{dx_0}, & \varepsilon_M^0 &\triangleq \frac{dx_M}{dx_0} = \bar{b}_0^0 \bar{c}_M^0 = \bar{b}_A^0 \bar{c}_M^A, \\ \bar{c}_M^0 &\triangleq \frac{dx_M}{d\xi_0}, & \bar{c}_0^0 &\triangleq \frac{dx_0}{d\xi_0}, & \bar{\delta}_A^0 &\triangleq \frac{d\xi_A}{d\xi_0} = \bar{c}_0^0 \bar{b}_A^0 = \bar{c}_M^0 \bar{b}_A^M. \end{aligned}$$

They determine the following smooth functions on the entire L , similar to section 2.2, that

$$\begin{aligned} B_0^A : L &\rightarrow \mathbb{R}, \quad p \mapsto B_0^A(p) \triangleq (b_{f(p)})_0^A(p), & C_0^M : L &\rightarrow \mathbb{R}, \quad p \mapsto C_0^M(p) \triangleq (c_{f(p)})_0^M(p), \\ \bar{B}_A^0 : L &\rightarrow \mathbb{R}, \quad p \mapsto \bar{B}_A^0(p) \triangleq (\bar{b}_{f(p)})_A^0(p), & \bar{C}_M^0 : L &\rightarrow \mathbb{R}, \quad p \mapsto \bar{C}_M^0(p) \triangleq (\bar{c}_{f(p)})_M^0(p), \\ B_0^0 : L &\rightarrow \mathbb{R}, \quad p \mapsto B_0^0(p) \triangleq (b_{f(p)})_0^0(p), & C_0^0 : L &\rightarrow \mathbb{R}, \quad p \mapsto C_0^0(p) \triangleq (c_{f(p)})_0^0(p), \\ \bar{B}_0^0 : L &\rightarrow \mathbb{R}, \quad p \mapsto \bar{B}_0^0(p) \triangleq (\bar{b}_{f(p)})_0^0(p), & \bar{C}_0^0 : L &\rightarrow \mathbb{R}, \quad p \mapsto \bar{C}_0^0(p) \triangleq (\bar{c}_{f(p)})_0^0(p). \end{aligned}$$

For convenience, we still use the notations ε and δ , and have the following smooth functions.

$$\begin{aligned} \varepsilon_0^M &\triangleq B_0^0 C_0^M = B_0^A C_A^M, & \delta_0^A &\triangleq C_0^0 B_0^A = C_0^M B_M^A, & G_{00} &\triangleq B_0^0 B_0^0 = G_{MN} \varepsilon_0^M \varepsilon_0^N, \\ \varepsilon_M^0 &\triangleq \bar{B}_0^0 \bar{C}_M^0 = \bar{B}_A^0 \bar{C}_M^A, & \bar{\delta}_A^0 &\triangleq \bar{C}_0^0 \bar{B}_A^0 = \bar{C}_M^0 \bar{B}_A^M, & G^{00} &\triangleq C_0^0 C_0^0 = G^{MN} \varepsilon_M^0 \varepsilon_N^0. \end{aligned}$$

It is easy to verify that $dx_0 = G_{00} dx^0$ and $\frac{d}{dx_0} = G^{00} \frac{d}{dx^0}$ are both true on L by a simple calculation.

3.3 Evolution lemma

We have the following two evolution lemmas. The affine connection representations of Yang-Mills equation, energy-momentum equation and Dirac equation are dependent on them.

Definition 3.3.1. $\forall p \in L$, suppose $T_p(M)$ and $T_p(L)$ are tangent spaces, $T_p^*(M)$ and $T_p^*(L)$ are cotangent spaces. The regular imbedding $\pi : L \rightarrow M$, $q \mapsto q$ induces the tangent map and the cotangent map

$$\begin{aligned} \pi_* : T_p(L) &\rightarrow T_p(M), \quad [\gamma_L] \mapsto [\pi \circ \gamma_L], \\ \pi^* : T_p^*(M) &\rightarrow T_p^*(L), \quad df \mapsto d(f \circ \pi). \end{aligned} \tag{21}$$

Evidently, π_* is an injection, and π^* is a surjection. $\forall \frac{d}{dt_L} \in T_p(L)$, $\frac{d}{dt} \in T_p(M)$, $df \in T_p^*(M)$, $df_L \in T_p^*(L)$, if and only if

$$\frac{d}{dt} = \pi_* \left(\frac{d}{dt_L} \right), \quad df_L = \pi^*(df) \tag{22}$$

are true, we denote

$$\frac{d}{dt} \cong \frac{d}{dt_L}, \quad df \cong df_L. \tag{23}$$

Then, we have the following two propositions that are evidently true.

Proposition 3.3.1. If $\frac{d}{dt} \cong \frac{d}{dt_L}$ and $df \cong df_L$, then

$$\left\langle \frac{d}{dt}, df \right\rangle = \left\langle \frac{d}{dt_L}, df_L \right\rangle. \tag{24}$$

Proposition 3.3.2. The following conclusions are true.

$$\begin{cases} w^M \frac{\partial}{\partial x^M} \cong w^0 \frac{d}{dx^0} & \Leftrightarrow w^M = w^0 \varepsilon_0^M, \\ w_M dx^M \cong w_0 dx^0 & \Leftrightarrow w_M \varepsilon_0^M = w_0, \end{cases} \quad \begin{cases} \bar{w}_M \frac{\partial}{\partial x_M} \cong \bar{w}_0 \frac{d}{dx_0} & \Leftrightarrow \bar{w}_M = \bar{w}_0 \varepsilon_M^0, \\ \bar{w}^M dx_M \cong \bar{w}^0 dx_0 & \Leftrightarrow \bar{w}^M \varepsilon_M^0 = \bar{w}^0. \end{cases} \tag{25}$$

3.4 On-shell evolution as a gradient

Let $\mathbf{T}$ be an smooth n -order tensor field. The restriction on (U, x^M) is $\mathbf{T} \triangleq t \left\{ \frac{\partial}{\partial x} \otimes dx \right\}$, where $\left\{ \frac{\partial}{\partial x} \otimes dx \right\}$ represents the tensor basis generated by several $\frac{\partial}{\partial x^M}$ and dx^M , and the tensor coefficients of $\mathbf{T}$ are concisely denoted by $t : U \rightarrow \mathbb{R}$.

Let D be a holonomic connection. Consider $D\mathbf{T} \triangleq t_{;Q} dx^Q \otimes \left\{ \frac{\partial}{\partial x} \otimes dx \right\}$. Denote

$$Dt \triangleq t_{;Q} dx^Q, \quad \nabla t \triangleq t_{;Q} \frac{\partial}{\partial x^Q}.$$

$\forall p \in M$, the integral curve of ∇t , that is $L_p \triangleq \varphi_{\nabla t, p}$, is a **gradient line** of $\mathbf{T}$. It can be seen later that the above gradient operator ∇ characterizes the **on-shell evolution**.

For any evolution path L , let $U_L \triangleq U \cap L$. Denote $t_L \triangleq t|_{U_L}$ and $t_{L;0} \triangleq t_{;Q} \varepsilon_0^Q$, as well as

$$D_L t_L \triangleq t_{L;0} dx^0, \quad \nabla_L t_L \triangleq t_{L;0} \frac{d}{dx^0}.$$

Proposition 3.4.1. The following conclusions are evidently true.

- (i) $Dt \simeq D_L t_L$ if and only if L is an arbitrary evolution path.
- (ii) $\nabla t \cong \nabla_L t_L$ if and only if L is a gradient line of $\mathbf{T}$.

Remark 3.4.1. More generally, suppose there is a tensor $\mathbf{U} \triangleq u_Q dx^Q \otimes \left\{ \frac{\partial}{\partial x} \otimes dx \right\}$. In such a notation, all the indices are concisely ignored except Q . $u_Q dx^Q$ uniquely determines a characteristic direction $u_Q \frac{\partial}{\partial x^Q}$.

If the system of 1-order linear partial differential equations $t_{;Q} = u_Q$ has a solution t , then it is true that $Dt = u_Q dx^Q$ and $\nabla t = u_Q \frac{\partial}{\partial x^Q}$. Thus, in the evolution direction $[L] = u_Q \frac{\partial}{\partial x^Q}$, the following conclusions are true.

$$Dt \simeq D_L t_L, \quad \nabla t \cong \nabla_L t_L, \quad (26)$$

where $D_L t_L \triangleq u_0 dx^0$, $\nabla_L t_L \triangleq u_0 \frac{d}{dx^0}$ and $u_0 \triangleq u_Q \varepsilon_0^Q$.

Now for any geometric property in the form of tensor $\mathbf{U}$, we are able to express its on-shell evolution in the form of ∇t .

Next, two important on-shell evolutions are discussed in the following two sections. One is the on-shell evolution of the potential field of a reference-system. The other is the one that a general charge of a reference-system evolves in the potential field of another reference-system.

3.5 On-shell evolution of potential field and affine connection representation of Yang-Mills equation

The table I of the article[40] proposes a famous correspondence between gauge field terminologies and fibre bundle terminologies. However, it does not find out the corresponding mathematical object to the source J_μ^K . In this section, we give an answer to this problem, and show the affine connection representation of Yang-Mills equation.

In order to obtain the general Yang-Mills equation with gravitation, we have to adopt holonomic connection to construct it. Suppose $\mathcal{F}$ evolves in $\mathcal{G}$ according to Definition 3.2.1, that is, $\forall p \in M$,

$$(U, \alpha^{A'}) \xrightarrow{f(p)} (U, \xi^A) \xrightarrow{f(p)} (U, x^M) \xleftarrow{g(p)} (U, \zeta^A) \xleftarrow{g(p)} (U, \beta^{A'}).$$

We always take the following notations in the coordinate frame (U, x^M) .

(i) Let the holonomic connections, which are defined by Eq.(12), of geometric manifolds $(M, \mathcal{F})$ and $(M, \mathcal{G})$ be $(\Gamma_{\mathcal{F}})_{NP}^M$ and $(\Gamma_{\mathcal{G}})_{NP}^M$, respectively. The colon ":" and semicolon ";" are used to express the covariant derivatives on $(M, \mathcal{F})$ and $(M, \mathcal{G})$, respectively, e.g.:

$$u^Q_{;P} = \frac{\partial u^Q}{\partial x^P} + (\Gamma_{\mathcal{F}})_{HP}^Q u^H, \quad u^Q_{;P} = \frac{\partial u^Q}{\partial x^P} + (\Gamma_{\mathcal{G}})_{HP}^Q u^H.$$

(ii) Let the coefficients of curvature tensor of $(M, \mathcal{F})$ and $(M, \mathcal{G})$ be K_{NPQ}^M and R_{NPQ}^M , respectively, i.e.

$$\begin{aligned} K_{NPQ}^M &\triangleq \frac{\partial(\Gamma_{\mathcal{F}})_{NQ}^M}{\partial x^P} - \frac{\partial(\Gamma_{\mathcal{F}})_{NP}^M}{\partial x^Q} + (\Gamma_{\mathcal{F}})_{NQ}^H (\Gamma_{\mathcal{F}})_{HP}^M - (\Gamma_{\mathcal{F}})_{NP}^H (\Gamma_{\mathcal{F}})_{HQ}^M, \\ R_{NPQ}^M &\triangleq \frac{\partial(\Gamma_{\mathcal{G}})_{NQ}^M}{\partial x^P} - \frac{\partial(\Gamma_{\mathcal{G}})_{NP}^M}{\partial x^Q} + (\Gamma_{\mathcal{G}})_{NQ}^H (\Gamma_{\mathcal{G}})_{HP}^M - (\Gamma_{\mathcal{G}})_{NP}^H (\Gamma_{\mathcal{G}})_{HQ}^M. \end{aligned} \quad (27)$$

Denote $K_{NPQ}^M \cdot^P \triangleq (G_{\mathcal{F}})^{PP'} K_{NPQ}^M \cdot^{P'}$. On an arbitrary evolution path L , we define

$$\rho_{N0}^M dx^0 \triangleq \pi^* \left(K_{NPQ}^M \cdot^P dx^Q \right) \in T^*(L).$$

Then, according to Definition 3.3.1 and the evolution lemma of Proposition 3.3.2, we obtain $\rho_{N0}^M = K_{NPQ}^M \cdot^P \varepsilon_0^Q$ and

$$K_{NPQ}^M \cdot^P dx^Q \simeq \rho_{N0}^M dx^0.$$

Let $\nabla t = K_{NPQ}^M \cdot^P \frac{\partial}{\partial x_Q}$. Then, according to Proposition 3.4.1, if and only if $\forall p \in M, [L_p] = \nabla t|_p$, we have

$$K_{NPQ}^M \cdot^P \frac{\partial}{\partial x_Q} \cong \rho_{N0}^M \frac{d}{dx_0}.$$

Applying the evolution lemma of Proposition 3.3.2 again, we obtain

$$K_{NPQ}^M \cdot^P = \rho_{N0}^M \varepsilon_Q^0.$$

Denote $j_{NQ}^M \triangleq \rho_{N0}^M \varepsilon_Q^0$, then if and only if $[L_p] = \nabla t|_p$ we have

$$K_{NPQ}^M \cdot^P = j_{NQ}^M, \quad (28)$$

which is said to be **(affine) Yang-Mills equation** of $\mathcal{F}$. It contains effects of gravitation, which makes the gauge frames $(B_f)_M^A$ and $(C_f)_A^M$ have physical effects. According to Eq.(15), we know Eq.(28) is coordinate covariant, and if gravitation is removed, it is also gauge covariant.

Thus, we have the following two results.

(i) The Yang-Mills equation originates from a geometric property in the direction ∇t . In other words, the on-shell evolution of gauge field is described by the direction field ∇t .

(ii) We obtain the mathematical origination of charge and current. We know that the evolution path L is an imbedding submanifold of M . Thus, the charge ρ_{N0}^M originates from the pull-back π^* from M to L , and the current j_{NQ}^M originates from ∇t that is associated to ρ_{N0}^M .

If we let (M, f) be completely flat, i.e. $(B_f)_M^A = \delta_M^A$, $(C_f)_A^M = \delta_A^M$, then by calculation we find ρ_{N0}^M can still be non-vanished. This shows that ρ_{N0}^M originates from (M, f) ultimately.

Definition 3.5.1. We speak of the real-valued

$$\rho_{MN0} \triangleq G_{MH} \rho_{N0}^H \quad (29)$$

as the **field function of a general charge**, or speak of it as a **charge** of $\mathcal{F}$ for short.

3.6 On-shell evolution of general charge and affine connection representation of mass, energy, momentum and action

In order to be compatible with the affine connection representation of gauge fields, we also have to define mass, energy, momentum and action in the form associated to affine connection. We are going to show them in this section and section 4.3.

Let $F_0 \triangleq \rho_{MN0} dx^M \otimes dx^N$. For the sake of simplicity, denote the charge ρ_{MN0} of $\mathcal{F}$ by ρ_{MN} concisely. Let D be the holonomic connection of $(M, \mathcal{G})$, then

$$D\mathbf{F}_0 \triangleq D\rho_{MN} \otimes dx^M \otimes dx^N, \quad \nabla\mathbf{F}_0 \triangleq \nabla\rho_{MN} \otimes dx^M \otimes dx^N,$$

where $D\rho_{MN} \triangleq \rho_{MN;Q} dx^Q$ and $\nabla\rho_{MN} \triangleq \rho_{MN;Q} \frac{\partial}{\partial x_Q}$. According to Proposition 3.4.1, if and only if $\forall p \in L$, the evolution direction is taken as $[L_p] = \nabla\rho_{MN}|_p$, we have

$$D\rho_{MN} \simeq D_L \rho_{MN}, \quad \nabla\rho_{MN} \cong \nabla_L \rho_{MN},$$

that is

$$\rho_{MN;Q} dx^Q \simeq \rho_{MN;0} dx^0, \quad \rho_{MN;Q} \frac{\partial}{\partial x_Q} \cong \rho_{MN;0} \frac{d}{dx_0}. \quad (30)$$

Definition 3.6.1. For more convenience, the notation ρ_{MN} is further abbreviated as ρ . In affine connection representation, energy and momentum of ρ are defined as

$$\begin{aligned} E_0 &\triangleq \rho_{;0} \triangleq \rho_{;Q} \varepsilon_0^Q, & p_Q &\triangleq \rho_{;Q}, & H_0 &\triangleq \frac{d\rho}{dx^0}, & P_Q &\triangleq \frac{\partial \rho}{\partial x^Q}, \\ E^0 &\triangleq \rho^{;0} \triangleq \rho^{;Q} \varepsilon_Q^0, & p^Q &\triangleq \rho^{;Q}, & H^0 &\triangleq \frac{d\rho}{dx_0}, & P^Q &\triangleq \frac{\partial \rho}{\partial x_Q}. \end{aligned} \quad (31)$$

Proposition 3.6.1. At any point p on M , the equation

$$E_0 E^0 = p_Q p^Q \quad (32)$$

holds if and only if the evolution direction $[L_p] = \nabla \rho|_p$. Eq.(32) is the **(affine) energy-momentum equation** of ρ .

Proof. According to the above discussion, $\forall p \in M$, $[L_p] = \nabla \rho|_p$ is equivalent to

$$p_Q dx^Q \simeq E_0 dx^0, \quad p_Q \frac{\partial}{\partial x_Q} \simeq E_0 \frac{d}{dx_0}. \quad (33)$$

Then due to Proposition 3.3.1 we obtain the directional derivative in the gradient direction $\nabla \rho$:

$$\left\langle p_Q \frac{\partial}{\partial x_Q}, p_M dx^M \right\rangle = \left\langle E_0 \frac{d}{dx_0}, E_0 dx^0 \right\rangle,$$

i.e. $G^{QM} p_Q p_M = G^{00} E_0 E_0$, or $p_Q p^Q = E_0 E^0$. $\square$

Proposition 3.6.2. At any point p on M , the equations

$$p^Q = E^0 \frac{dx^Q}{dx^0}, \quad p_Q = E_0 \frac{dx_Q}{dx_0} \quad (34)$$

hold if and only if the evolution direction $[L_p] = \nabla \rho|_p$.

Proof. Due to the evolution lemma of Proposition 3.3.2, we immediately obtain Eq.(34) from Eq.(33). $\square$

Remark. In the gradient direction $\nabla \rho$, Eq.(34) is consistent with the conventional

$$p = mv.$$

Thus, in affine connection representation, the energy-momentum equation and the conventional definition of momentum both originate from a geometric property in gradient direction. In other words, the on-shell evolution of the particle field ρ is described by the gradient direction field $\nabla \rho$.

Definition 3.6.2. Let $\mathcal{P}(b, a)$ be the totality of paths from a to b . And suppose $L \in \mathcal{P}(b, a)$, and the evolution parameter x^0 satisfies $t_a \triangleq x^0(a) < x^0(b) \triangleq t_b$. The **elementary affine action** of ρ is defined as

$$\mathfrak{s}(L) \triangleq \int_L D\rho = \int_L p_Q dx^Q = \int_{t_a}^{t_b} E_0 dx^0. \quad (35)$$

Thus, $\delta \mathfrak{s}(L) = 0$ if and only if L is a gradient line of ρ .

In particular, in the case where $\mathcal{G}$ is orthogonal, we can also define action in the following way.

On $(M, \mathcal{G})$ let there be Dirac algebras γ^M and γ_N such that

$$\gamma^M \gamma^N + \gamma^N \gamma^M = 2G^{MN}, \quad \gamma_M \gamma_N + \gamma_N \gamma_M = 2G_{MN}, \quad \gamma_M \gamma^M = 1.$$

In a gradient direction of ρ , from Eq.(32) we obtain that

$$\begin{aligned} p_Q p^Q = E_0 E^0 &\Leftrightarrow \rho_{;Q} \rho^{;Q} = \rho_{;0} \rho^{;0} \\ &\Leftrightarrow G^{PQ} \rho_{;P} \rho_{;Q} = G^{00} \rho_{;0} \rho_{;0} \\ &\Leftrightarrow (\gamma^P \gamma^Q + \gamma^Q \gamma^P) \rho_{;P} \rho_{;Q} = 2 \rho_{;0} \rho_{;0} \\ &\Leftrightarrow (\gamma^P \rho_{;P})(\gamma^Q \rho_{;Q}) + (\gamma^Q \rho_{;Q})(\gamma^P \rho_{;P}) = 2 \rho_{;0} \rho_{;0} \\ &\Leftrightarrow (\gamma^P \rho_{;P})^2 = (\rho_{;0})^2. \end{aligned}$$

Take $\gamma^P \rho_{;P} = \rho_{;0}$ without loss of generality, then, in the gradient direction of ρ we have

$$\gamma^P \rho_{;P} dx^0 = \rho_{;0} dx^0 = \varepsilon_0^P \rho_{;P} dx^0 = D\rho. \quad (36)$$

So we can define the following **(orthogonal affine) action** of ρ .

$$s(L) \triangleq \int_L (\gamma^P \rho_{;P} dx^0 + D\rho) = \int_{t_a}^{t_b} (\gamma^P \rho_{;P} + \varepsilon_0^P \rho_{;P}) dx^0 = \int_{t_a}^{t_b} (\gamma^P \rho_{;P} + E_0) dx^0. \quad (37)$$

Remark 3.6.1 and Remark 4.4.1 explain the rationality of this definition. We have $s(L) = 2\mathfrak{s}(L)$ in the gradient direction of ρ , so $\mathfrak{s}(L)$ and $s(L)$ are consistent.

Remark 3.6.1. In the Minkowski coordinate frame of section 4.2, the evolution parameter x^0 is replaced by $\tilde{x}^\tau$, then there still exists a concept of gradient direction $\tilde{\nabla}\tilde{\rho}$. Correspondingly,

$$\mathfrak{s}(L) \triangleq \int_L D\rho = \int_L p_Q dx^Q = \int_{t_a}^{t_b} E_0 dx^0, \quad s(L) = \int_{t_a}^{t_b} (\gamma^P \rho_{;P} + E_0) dx^0$$

present as

$$\tilde{\mathfrak{s}}(L) \triangleq \int_L \tilde{D}\tilde{\rho} = \int_L \tilde{p}_\mu d\tilde{x}^\mu = \int_{\tau_a}^{\tau_b} \tilde{m}_\tau d\tilde{x}^\tau, \quad \tilde{s}(L) = \int_{\tau_a}^{\tau_b} (\gamma^\mu \tilde{\rho}_{;\mu} + \tilde{m}_\tau) d\tilde{x}^\tau,$$

where $\tilde{m}_\tau$ is the rest-mass and $\tilde{x}^\tau$ is the proper-time.

Remark 3.6.2. Define the following notations.

$$[\rho\Gamma_G] \triangleq \frac{\partial\rho_{MN}}{\partial x^G} - \rho_{MN;G} = \rho_{MH}\Gamma_{NG}^H + \rho_{HN}\Gamma_{MG}^H, \quad [\rho R_{PQ}] \triangleq \rho_{MH}R_{NPQ}^H + \rho_{HN}R_{MPQ}^H.$$

Then, through some calculations, we can obtain that

$$f_P \triangleq p_{P;0} = E_{0;P} - p_Q \varepsilon_{0;P}^Q + [\rho R_{PQ}] \varepsilon_0^Q,$$

which is the affine connection representations of general Lorentz force equation. See Discussion 4.3.1 for further illustrations.

3.7 Inversion transformation in affine connection representation

In affine connection representation, *CPT* inversion is interpreted as a total inversion of coordinates and metrics. Let $i, j = 1, 2, 3$ and $m, n = 4, 5, \dots, \mathfrak{D}$.

Let the local coordinate representation of reference-system k be $x'^j = -\delta_i^j x^i$, $x'^m = \delta_m^n x^n$, then parity inversion can be represented as

$$P \triangleq L_k : x^i \rightarrow -x^i, x^m \rightarrow x^m.$$

Let the local coordinate representation of reference-system h be $x'^j = \delta_i^j x^i$, $x'^m = -\delta_m^n x^n$, then charge conjugate inversion can be represented as

$$C \triangleq L_h : x^i \rightarrow x^i, x^m \rightarrow -x^m.$$

Time coordinate inversion can be represented as

$$T_0 : x^0 \rightarrow -x^0.$$

Total inversion of coordinates can be represented as

$$CPT_0 : x^Q \rightarrow -x^Q, x^0 \rightarrow -x^0. \quad (38)$$

The positive or negative sign of metric marks two opposite directions of evolution. Let N be a closed submanifold of M , and let its metric be $dx^{(N)}$. Denote the totality of closed submanifolds of M by $\mathfrak{B}(M)$, then **total inversion of metrics** can be expressed as

$$T^{(M)} \triangleq \prod_{N \in \mathfrak{B}(M)} (dx^{(N)} \rightarrow -dx^{(N)}). \quad (39)$$

Denote time inversion by

$$T \triangleq T^{(M)} T_0,$$

then the joint transformation of the total inversion of coordinates CPT_0 and the total inversion of metrics $T^{(M)}$ is

$$(CPT_0)(T^{(M)}) = CPT. \quad (40)$$

Summarize the above discussions, then we have

$$\begin{aligned} CPT_0 : x^Q &\rightarrow -x^Q, \quad x^0 \rightarrow -x^0, \quad dx^Q \rightarrow dx^Q, \quad dx^0 \rightarrow dx^0, \\ T^{(M)} : x^Q &\rightarrow x^Q, \quad x^0 \rightarrow x^0, \quad dx^Q \rightarrow -dx^Q, \quad dx^0 \rightarrow -dx^0, \\ CPT : x^Q &\rightarrow -x^Q, \quad x^0 \rightarrow -x^0, \quad dx^Q \rightarrow -dx^Q, \quad dx^0 \rightarrow -dx^0. \end{aligned}$$

The CPT invariance in affine connection representation is very clear. Concretely, on $(M, \mathcal{G})$ we consider the CPT transformation acting on $\mathcal{G}$. Denote $s \triangleq \int_L D\rho$ and $D_P e^{is} \triangleq (\frac{\partial}{\partial x^P} - i[\rho \Gamma_P]) e^{is}$, then through simple calculations we obtain that

$$CPT : D\rho \rightarrow D\rho, \quad D_P e^{is} \rightarrow -D_P e^{is}.$$

Remark 3.7.1. In quantum mechanics there is a complex conjugation in the time inversion of wave function $T : \psi(x, t) \rightarrow \psi^*(x, -t)$. In affine connection representation, we know the complex conjugation can be interpreted as a straightforward mathematical result of the total inversion of metrics $T^{(M)}$.

3.8 Two dual descriptions of gradient direction field

Discussion 3.8.1. Let X and Y be non-vanishing smooth tangent vector fields on the manifold M . And let L_Y be the Lie derivative operator induced by the one-parameter group of diffeomorphism φ_Y . Then, according to a well-known theorem[41], we obtain the Lie derivative equation

$$[X, Y] = L_Y X. \quad (41)$$

Suppose $\forall p \in M$, $Y(p)$ is a unit-length vector, i.e. $\|Y(p)\| = 1$. Let the parameter of φ_Y be x^0 . Then, on the evolution path $L \triangleq \varphi_{Y,p}$, we have

$$Y \cong \frac{d}{dx^0}. \quad (42)$$

Thus, Eq.(41) can also be represented as

$$[X, Y] = \frac{d}{dx^0} X. \quad (43)$$

On the other hand, $\forall df \in T(M)$ and $df_L \triangleq \pi^*(df)$, due to (42) and Proposition 3.3.1 we have $\langle Y, df \rangle = \langle \frac{d}{dx^0}, df_L \rangle$, that is

$$Y f = \frac{d}{dx^0} f_L. \quad (44)$$

Definition 3.8.1. Let $H \triangleq \|\nabla \rho\|^{-1} \nabla \rho = \varepsilon_0^M \frac{\partial}{\partial x^M} \cong \frac{d}{dx^0}$. It is evident that $\forall p \in M$, $\|H(p)\| = 1$. If and only if taking $Y = H$, we speak of (43) and (44) as real-valued **(affine) Heisenberg equation** and **(affine) Schrödinger equation**, respectively, that is

$$[X, H] = \frac{d}{dx^0} X, \quad H f = \frac{d}{dx^0} f_L. \quad (45)$$

Discussion 3.8.2. The above two equations both describe the gradient direction field, and thereby reflect on-shell evolution. Such two dual descriptions of gradient direction show the real-valued affine connection representation of Heisenberg picture and Schrödinger picture.

It is not hard to find out several different kinds of complex-valued representations of gradient direction. For examples, one is the affine Dirac equation in section 4.4, another is as follows.

Let $\psi \triangleq f e^{is_L}$, where it is fine to take either $s_L \triangleq s(L)$ or $s_L \triangleq \mathfrak{s}(L)$ from Definition 3.6.2. According to Eq.(45), it is easy to obtain on L , that

$$[X, H] = \frac{d}{dx^0} X, \quad H \psi = \frac{d\psi}{dx^0}. \quad (46)$$

This is consistent with the conventional Heisenberg equation and Schrödinger equation (taking the natural units that $\hbar = 1$, $c = 1$)

$$[X, -iH] = \frac{\partial}{\partial t} X, \quad -iH \psi = \frac{\partial \psi}{\partial t} \quad (47)$$

and they have a coordinate correspondence

$$\frac{\partial}{\partial(ix^k)} \leftrightarrow \frac{\partial}{\partial x^k}, \quad \frac{\partial}{\partial t} \leftrightarrow \frac{d}{dx^0}.$$

We know that $\frac{\partial}{\partial t} \leftrightarrow \frac{d}{dx^0}$ originates from the difference that the evolution parameter is x^τ or x^0 . The imaginary unit i originates from the difference between the regular coordinates x^1, x^2, x^3, x^τ and the Minkowski coordinates x^1, x^2, x^3, x^0 . That is to say, the regular coordinates satisfy

$$(dx^0)^2 = (dx^1)^2 + (dx^2)^2 + (dx^3)^2 + (dx^\tau)^2,$$

and the Minkowski coordinates satisfy

$$(dx^\tau)^2 = (dx^0)^2 - (dx^1)^2 - (dx^2)^2 - (dx^3)^2 = (dx^0)^2 + (d(ix^1))^2 + (d(ix^2))^2 + (d(ix^3))^2.$$

This causes the appearance of the imaginary unit i in the correspondence

$$ix^k \leftrightarrow x^k.$$

So Eq.(46) and Eq.(47) have exactly the same essence, and their differences only come from different coordinate representations.

The differences between coordinate representations have nothing to do with the geometric essence and the physical essence. We notice that the value of a gradient direction is dependent on geometry, but independent of that the equations are real-valued or complex-valued. Therefore, it is unnecessary for us to confine to such algebraic forms as real-valued or complex-valued forms, but we should focus on such geometric essence as gradient direction.

The advantage of complex-valued form is that it is applicable for describing the coherent superposition of propagator. However, this is independent of the above discussions, and we are going to discuss it in section 3.9.

3.9 Quantum evolution as a distribution of gradient directions

From Proposition 3.6.1 we see that, in affine connection representation, the classical on-shell evolution is described by gradient direction. Then, naturally, quantum evolution should be described by the distribution of gradient directions.

The distribution of gradient directions on a geometric manifold $(M, \mathcal{G})$ is effected by the bending shape of $(M, \mathcal{G})$, in other words, the distribution of gradient directions can be used to reflect the shape of $(M, \mathcal{G})$. This is the way that the quantum theory in affine connection representation describes physical reality.

In order to know the full picture of physical reality, it is necessary to fully describe the shape of the geometric manifold. For a single observation:

(1) It is the reference-system, not a point, that is used to describe the physical reality, so the coordinate of an individual point is not enough to fully describe the location information about the physical reality.

(2) Through a single observation of momentum, we can only obtain information about an individual gradient direction, this cannot reflect the full picture of the shape of the geometric manifold.

Quantum evolution provides us with a guarantee that we can obtain the distribution of gradient directions through multiple observations, so that we can describe the full picture of the shape of the geometric manifold.

Next, we are going to carry out strict mathematical descriptions for the quantum evolution in affine connection representation.

Definition 3.9.1. Let ρ be a geometric property on M , such as a charge of f . Then $H \triangleq \nabla \rho$ is a gradient direction field of ρ on $(M, \mathcal{G})$.

Let $\mathfrak{T}$ be the totality of all flat transformations L_k defined in section 2.3. $\forall T \in \mathfrak{T}$, the flat transformation $T : f \mapsto Tf$ induces a transformation $T^* : \rho \mapsto T^* \rho$. Denote

$$|\rho| \triangleq \{\rho_T \triangleq T^* \rho | T \in \mathfrak{T}\}, \quad |H| \triangleq \{H_T \triangleq \nabla \rho_T | T \in \mathfrak{T}\}.$$

$\forall a \in M$, the restriction of $|H|$ at a are denoted by $|H(a)| \triangleq \{H_T(a) | T \in \mathfrak{T}\}$.

We say $|H|$ is the **total distribution** of the gradient direction field H .

$\forall T \in \mathfrak{T}$, we say H_T is a **position distribution of gradient directions in momentum representation**.

$\forall a \in M$, we say $|H(a)|$ is a **momentum distribution of gradient directions in position representation**.

Remark 3.9.1. When T is fixed, H_T can reflect the shape of $(M, \mathcal{G})$. When a is fixed, $|H(a)|$ can reflect the shape of $(M, \mathcal{G})$.

However, when T and a are both fixed, $H_T(a)$ is a fixed individual gradient direction, which cannot reflect the shape of $(M, \mathcal{G})$. In other words, if the momentum p_T and the position x_a of ρ are both definitely observed, the physical reality $\mathcal{G}$ would be unknowable, therefore this is unacceptable. This is the embodiment of quantum uncertainty in affine connection representation.

Definition 3.9.2. Let φ_H be the one-parameter group of diffeomorphisms corresponding to H . The parameter of φ_H is x^0 . $\forall a \in M$, according to Definition 3.2.1, let $\varphi_{H,a}$ be the evolution path through a , such that $\varphi_{H,a}(0) = a$. $\forall t \in \mathbb{R}$, denote

$$\varphi_{|H|,a} \triangleq \{\varphi_{X,a} | X \in |H|\}, \quad \varphi_{|H|,a}(t) \triangleq \{\varphi_{X,a}(t) | X \in |H|\}.$$

$\forall \Omega \subseteq \mathfrak{T}$, we also denote $|H_\Omega| \triangleq \{H_T | T \in \Omega\} \subseteq |H|$ and

$$\varphi_{|H_\Omega|,a} \triangleq \{\varphi_{X,a} | X \in |H_\Omega|\} \subseteq \varphi_{|H|,a}, \quad \varphi_{|H_\Omega|,a}(t) \triangleq \{\varphi_{H,a}(t) | X \in |X_\Omega|\} \subseteq \varphi_{|H|,a}(t).$$

$\forall a \in M$, the restriction of $|H_\Omega|$ at a are denoted by $|H_\Omega(a)| \triangleq \{H_T(a) | T \in \Omega\}$.

Remark 3.9.2. At the beginning $t = 0$, intuitively, the gradient directions $|H(a)|$ of $|\rho|$ start from a and point to all directions around a uniformly. If $(M, \mathcal{G})$ is not flat, when evolving to a certain time $t > 0$, the distribution of gradient directions on $\varphi_{|H|,a}(t)$ is no longer as uniform as beginning. The following definition precisely characterizes this kind of ununiformity.

Definition 3.9.3. Let the transformation $L_{\mathcal{G}^{-1}}$ act on $\mathcal{G}$, then we obtain the trivial $e \triangleq L_{\mathcal{G}^{-1}}(\mathcal{G})$. Now $(M, \mathcal{G})$ is sent to a flat (M, e) , and the gradient direction field $|H|$ of $|\rho|$ on $(M, \mathcal{G})$ is sent to a gradient direction field $|O|$ of $|\rho|$ on (M, e) . Correspondingly, $\forall t \in \mathbb{R}$, $\varphi_{|H|,a}(t)$ is sent to $\varphi_{|O|,a}(t)$. In a word, $L_{\mathcal{G}^{-1}}$ induces the following two maps:

$$\mathcal{G}_*^{-1} : |H| \rightarrow |O|, \quad \mathcal{G}_{**}^{-1} : \varphi_{|H|,a} \rightarrow \varphi_{|O|,a}.$$

$\forall T \in \mathfrak{T}$, deontic $\mathfrak{N} \triangleq \{N \in \mathfrak{T} \mid \det N = \det T\}$. Due to $\mathfrak{T} \cong GL(\mathfrak{D}, \mathbb{R})$, let $\mathfrak{U}$ be a neighborhood of T , with respect to the topology of $GL(\mathfrak{D}, \mathbb{R})$.

Take $\Omega = \mathfrak{N} \cap \mathfrak{U}$, then

$$|O_\Omega| = \mathcal{G}_*^{-1}(|H_\Omega|), \quad \varphi_{|O_\Omega|,a} = \mathcal{G}_{**}^{-1}(\varphi_{|H_\Omega|,a}).$$

Let μ be a Borel measure on the manifold M . $\forall t \in \mathbb{R}$, we know

$$\varphi_{|H_\Omega|,a}(t) \simeq \varphi_{|O_\Omega|,a}(t) \simeq \mathbb{S}^{\mathfrak{D}-1}.$$

Thus, $\varphi_{|H_\Omega|,a}(t) \subseteq \varphi_{|H_\Omega|,a}(t)$ and $\varphi_{|O_\Omega|,a}(t) \subseteq \varphi_{|O_\Omega|,a}(t)$ are Borel sets, so they are measurable. Denote

$$\mu_a(\varphi_{|H_\Omega|,a}(t)) \triangleq \mu(\mathcal{G}_{**}^{-1}(\varphi_{|H_\Omega|,a}(t))) = \mu(\varphi_{|O_\Omega|,a}(t)).$$

When $\mathfrak{U} \rightarrow T$, we have $\Omega \rightarrow T$, $|H_\Omega| \rightarrow H_T$, $|H_\Omega(a)| \rightarrow H_T(a)$, and $\varphi_{|H_\Omega|,a}(t) \rightarrow b \triangleq \varphi_{H_T,a}(t)$.

For the sake of simplicity, denote $L \triangleq \varphi_{H_T,a}$. Thus, we have $a = L(0)$, $b = L(t)$, and denote $p_a \triangleq [L_a] = H_T(a)$, $p_b \triangleq [L_b] = H_T(b)$.

Because μ_a is absolutely continuous with respect to μ , Radon-Nikodym theorem[42] ensures the existence of the following limit. The Radon-Nikodym derivative

$$W_L(b, a) \triangleq \frac{d\mu_a}{d\mu_b} \triangleq \lim_{\mathfrak{U} \rightarrow T} \frac{\mu_a(\varphi_{|H_\Omega|,a}(t))}{\mu(\varphi_{|H_\Omega|,a}(t))} = \lim_{\mathfrak{U} \rightarrow T} \frac{\mu(\mathcal{G}_{**}^{-1}(\varphi_{|H_\Omega|,a}(t)))}{\mu(\varphi_{|H_\Omega|,a}(t))} = \lim_{\mathfrak{U} \rightarrow T} \frac{\mu(\varphi_{|O_\Omega|,a}(t))}{\mu(\varphi_{|H_\Omega|,a}(t))} \quad (48)$$

is called the **distribution density of $|H|$ along L in position representation**.

On a neighborhood U of a , $\forall T \in \mathfrak{T}$, denote the normal section of $H_T(a)$ by $N_{H_T,a}$, that is

$$N_{H_T,a} \triangleq \{n \in U \mid H_T(a) \cdot (n - a) = 0\}, \quad N_{H_T,a}(t) \triangleq \{\varphi_{H_T,x}(t) \mid x \in N_{H_T,a}\}.$$

Thus, $N_{H_T,a} = N_{H_T,a}(0)$ and $N_{H_T,b} \triangleq N_{H_T,a}(t)$. If $U \rightarrow a$, we have $N_{H_T,a} \rightarrow a$ and $N_{H_T,a}(t) \rightarrow b \triangleq \varphi_{H_T,a}(t)$. The Radon-Nikodym derivative

$$Z_L(b, a) \triangleq \frac{d\mu(a)}{d\mu(b)} \triangleq \lim_{U \rightarrow a} \frac{\mu(N_{H_T,a})}{\mu(N_{H_T,b})} = \lim_{U \rightarrow a} \frac{\mu(N_{H_T,a})}{\mu(N_{H_T,a}(t))} \quad (49)$$

is called the **distribution density of $|H|$ along L in momentum representation**.

In a word, $W_L(b, a)$ and $Z_L(p_b, p_a)$ describe the density of the gradient lines that are adjacent to b in two different ways. They have the following property that is evidently true.

Proposition 3.9.1. Let L be a gradient line. $\forall a, b, c \in L$ such that $L(x_a^0) = a$, $L(x_b^0) = b$, $L(x_c^0) = c$ and $x_b^0 > x_c^0 > x_a^0$, then

$$W_L(b, a) = W_L(b, c)W_L(c, a), \quad Z_L(b, a) = Z_L(b, c)Z_L(c, a).$$

Definition 3.9.4. If L is a gradient line of some $\rho' \in |\rho|$, we also say L is a gradient line of $|\rho|$.

Remark 3.9.3. For any a and b , it anyway makes sense to discuss the gradient line of $|\rho|$ from a to b . It is because even if the gradient line of ρ starting from a does not pass through b , it just only needs to carry out a certain flat transformation T defined in section 2.3 to obtain a $\rho' \triangleq T_*\rho$, thus the gradient line of ρ' starting from a can just exactly pass through b . Due to $\rho, \rho' \in |\rho|$, we do not distinguish them, it is just fine to uniformly use $|\rho|$. Intuitively speaking, when $|\rho|$ takes two different initial momentums, $|\rho|$ presents as ρ and ρ' , respectively.

Discussion 3.9.1. With the above preparations, we obtain a new way to describe the construction of the propagator strictly.

For any path L that starts at a and ends at b , we denote $||L|| \triangleq \int_L dx^0$ concisely. Let $\mathcal{P}(b, a)$ be the totality of all the paths from a to b . Denote

$$\mathcal{P}(b, x_b^0; a, x_a^0) \triangleq \{L \mid L \in \mathcal{P}(b, a), ||L|| = x_b^0 - x_a^0\}.$$

$\forall L \in \mathcal{P}(b, x_b^0; a, x_a^0)$, we can let $L(x_a^0) = a$ and $L(x_b^0) = b$ without loss of generality. Thus, $\mathcal{P}(b, x_b^0; a, x_a^0)$ is the totality of all the paths from $L(x_a^0) = a$ to $L(x_b^0) = b$.

Abstractly, the propagator is defined as the Green function of the evolution equation. Concretely, the propagator still needs a constructive definition. One method is the Feynman path integral

$$K(b, x_b^0; a, x_a^0) \triangleq \int_{\mathcal{P}(b, x_b^0; a, x_a^0)} e^{is} dL. \quad (50)$$

However, there are so many redundant paths in $\mathcal{P}(b, x_b^0; a, x_a^0)$ that: (i) it is difficult to generally define a measure dL on $\mathcal{P}(b, x_b^0; a, x_a^0)$, (ii) it may cause unnecessary infinities when carrying out some calculations.

In order to solve this problem, we try to reduce the scope of summation from $\mathcal{P}(b, x_b^0; a, x_a^0)$ to $H(b, x_b^0; a, x_a^0)$, where $H(b, x_b^0; a, x_a^0)$ is the totality of all the gradient lines of $|\rho|$ from $L(x_a^0) = a$ to $L(x_b^0) = b$. Thus, the (50) is turned into

$$K(b, x_b^0; a, x_a^0) = \int_{H(b, x_b^0; a, x_a^0)} \Psi(L) e^{is} dL.$$

We notice that as long as we take the probability amplitude $\Psi(L)$ of the gradient line L such that $[\Psi(L)]^2 = W_L(b, a)$ in position representation, or take $[\Psi(L)]^2 = Z_L(b, a)$ in momentum representation, it can exactly be consistent with the Copenhagen interpretation. This provides the following new constructive definition for the propagator.

Definition 3.9.5. Suppose $|\rho|$ is defined as Definition 3.9.1, and denote $H \triangleq \nabla\rho$.

Let $\mathcal{L}(b, a)$ be the totality of all the gradient lines of $|\rho|$ from a to b . Denote

$$H(b, x_b^0; a, x_a^0) \triangleq \{L \mid L \in \mathcal{L}(b, a), ||L|| = x_b^0 - x_a^0\}.$$

Let $\mathcal{L}(p_b, p_a)$ be the totality of all the gradient lines of $|\rho|$, whose starting-direction is p_a and ending-direction is p_b . Denote

$$H(p_b, x_b^0; p_a, x_a^0) \triangleq \{L \mid L \in \mathcal{L}(p_b, p_a), ||L|| = x_b^0 - x_a^0\}.$$

Let dL be a Borel measure on $H(b, x_b^0; a, x_a^0)$. In consideration of Remark 4.4.1, we let s be the affine action $s(L)$ in Definition 3.6.2. We say the geometric property

$$K(b, x_b^0; a, x_a^0) \triangleq \int_{H(b, x_b^0; a, x_a^0)} \sqrt{W_L(b, a)} e^{is} dL \quad (51)$$

is the **propagator** of $|\rho|$ from (a, x_a^0) to (b, x_b^0) **in position representation**. If we let dL be a Borel measure on $H(p_b, x_b^0; p_a, x_a^0)$, then we say

$$\mathcal{K}(p_b, x_b^0; p_a, x_a^0) \triangleq \int_{H(p_b, x_b^0; p_a, x_a^0)} \sqrt{Z_L(b, a)} e^{is} dL \quad (52)$$

is the **propagator** of $|\rho|$ from (p_a, x_a^0) to (p_b, x_b^0) **in momentum representation**.

Discussion 3.9.2. Now (51) and (52) are strictly defined, but the Feynman path integral (50) has not been possessed of a strict mathematical definition until now. This makes it impossible at present to obtain (e.g. in position representation) a strict mathematical proof of

$$\int_{H(b, x_b^0; a, x_a^0)} \sqrt{W_L(b, a)} e^{is} dL = \int_{\mathcal{P}(b, x_b^0; a, x_a^0)} e^{is} dL.$$

We notice that the distribution densities $W_L(b, a)$ and $Z_L(b, a)$ of gradient directions establish an association between probability interpretation and geometric interpretation of quantum evolution. Therefore, we can base on probability interpretation to intuitively consider both sides of "=" as the same thing.

Discussion 3.9.3. The quantization methods of QFT are successful, and they are also applicable in affine connection representation for orthogonal $\mathcal{F}$ and $\mathcal{G}$, but in this paper we do not discuss them. We try to propose some more ideas for general $\mathcal{F}$ and $\mathcal{G}$ to understand the quantization of field in affine connection representation.

(1) If we take

$$\mathfrak{s} = \int_L D\rho = \int_L p_Q dx^Q = \int_L E_0 dx^0$$

according to Definition 3.6.2, where D is the holonomic connection of $(M, \mathcal{G})$, then consider the distribution of $H \triangleq \nabla\rho$, we know that

$$K(b, x_b^0; a, x_a^0) \triangleq \int_{\nabla\rho(b, x_b^0; a, x_a^0)} \sqrt{W_L(b, a)} e^{is} dL, \quad \mathcal{K}(p_b, x_b^0; p_a, x_a^0) \triangleq \int_{\nabla\rho(p_b, x_b^0; p_a, x_a^0)} \sqrt{Z_L(b, a)} e^{is} dL$$

describe the quantization of energy-momentum. Every gradient line in $\nabla\rho(b, x_b^0; a, x_a^0)$ corresponds to a set of eigenvalues of energy and momentum.

(2) If we take

$$\mathfrak{s} = \int_L Dt = \int_L K_{NPQ}^M :^P dx^Q = \int_L \rho_{N0}^M dx^0,$$

according to section 3.5, where D is the holonomic connection of $(M, \mathcal{F})$, then consider the distribution of $H \triangleq \nabla t$, we know that

$$K(b, x_b^0; a, x_a^0) \triangleq \int_{\nabla t(b, x_b^0; a, x_a^0)} \sqrt{W_L(b, a)} e^{is} dL, \quad \mathcal{K}(p_b, x_b^0; p_a, x_a^0) \triangleq \int_{\nabla t(p_b, x_b^0; p_a, x_a^0)} \sqrt{Z_L(b, a)} e^{is} dL$$

describe the quantizations of charge and current. It should be emphasized that this is not the quantization of the energy-momentum of the field, but the quantization of the field itself, which presents as quantized (e.g. discrete) charges and currents.

4 Affine connection representation of gauge fields in classical spacetime

The new framework established in section 3 is discussed in the $\mathfrak{D}$ -dimensional general coordinate x^M , which is more general than the $(1+3)$ -dimensional conventional Minkowski coordinate x^μ .

$(dx^0)^2 = \sum_{M=1}^{\mathfrak{D}} (dx^M)^2$ is the total metric of internal space and external space, $(dx^\tau)^2 = \sum_{m=4}^{\mathfrak{D}} (dx^m)^2$ is the metric of internal space.

(i) The evolution parameter of the $\mathfrak{D}$ -dimensional general coordinate x^M ($M = 1, 2, \dots, \mathfrak{D}$) is x^0 . The parameter equation of an evolution path L is represented as $x^M = x^M(x^0)$.

(ii) The evolution parameter of the $(1+3)$ -dimensional Minkowski coordinate x^μ ($\mu = 0, 1, 2, 3$) is x^τ . The parameter equation of L is represented as $x^\mu = x^\mu(x^\tau)$.

The coordinate x^μ works on the $(1+3)$ -dimensional classical spacetime submanifold defined as follows.

4.1 Classical spacetime submanifold

Let there be a smooth tangent vector field X on (M, f) . If $\forall p \in M$, $X(p) = b^A \frac{\partial}{\partial \xi^A} \Big|_p = c^M \frac{\partial}{\partial x^M} \Big|_p$ satisfies that b^a are not all zero and c^m are not all zero, then we say X is **internal-directed**. For any evolution path $L \triangleq \varphi_{X,p}$, we also say L is **internal-directed**.

Suppose $M = P \times N$, $\mathfrak{D} \triangleq \dim M$ and $r \triangleq \dim P = 3$. X is a smooth tangent vector field on M . Fix a point $o \in M$. If X is internal-directed, then there exist a unique $(1 + 3)$ -dimensional imbedding submanifold $\gamma : \tilde{M} \rightarrow M$, $p \mapsto p$ and a unique smooth tangent vector field $\tilde{X}$ on $\tilde{M}$ such that:

(i) $P \times \{o\}$ is a closed submanifold of $\tilde{M}$.

(ii) The tangent map $\gamma_* : T(\tilde{M}) \rightarrow T(M)$ satisfies that $\forall q \in \tilde{M}$, $\gamma_* : \tilde{X}(q) \mapsto X(q)$.

Such an $\tilde{M}$ is said to be a **classical spacetime submanifold**.

Let $\varphi_X : M \times \mathbb{R} \rightarrow M$ and $\varphi_{\tilde{X}} : \tilde{M} \times \mathbb{R} \rightarrow \tilde{M}$ be the one-parameter groups of diffeomorphisms corresponding to X and $\tilde{X}$, respectively. Thus, we have

$$\varphi_{\tilde{X}} = \varphi_X|_{\tilde{M} \times \mathbb{R}}.$$

So the evolution in classical spacetime can be described by $\varphi_{\tilde{X}}$. It should be noticed that:

(i) $\tilde{M}$ inherits a part of geometric properties of M , but not all. The physical properties reflected by $\tilde{M}$ are incomplete.

(ii) The correspondence between $\tilde{X}$ and the restriction of X to $\tilde{M}$ is one-to-one. For convenience, next we are not going to distinguish the notations X and $\tilde{X}$ on $\tilde{M}$, but uniformly denote them by X .

(iii) An arbitrary path $\tilde{L} : T \rightarrow \tilde{M}$, $t \mapsto p$ on $\tilde{M}$ uniquely corresponds to a path $L \triangleq \gamma \circ \tilde{L} : T \rightarrow M$, $t \mapsto p$ on M . Evidently the image sets of L and $\tilde{L}$ are the same, that is, $L(T) = \tilde{L}(T)$. For convenience, later we are not going to distinguish the notations L and $\tilde{L}$ on $\tilde{M}$, but uniformly denote them by L .

4.2 Classical spacetime reference-system

Let there be a geometric manifold (M, f) and its classical spacetime submanifold $\tilde{M}$. And let $L \triangleq \varphi_{\tilde{X}, a}$ be an evolution path on $\tilde{M}$. Suppose $p \in L$ and U is a coordinate neighborhood of p . According to Definition 3.2.2, suppose the $f(p)$ on U and the $f_L(p)$ on $U_L \triangleq U \cap L$ satisfy that

$$f(p) : \xi^A = \xi^A(x^M) = \xi^A(x^0), \quad \xi^0 = \xi^0(x^0), \quad A, M = 1, 2, \dots, \mathfrak{D}. \quad (53)$$

Thus, it is true that:

(1) There exists a unique local reference-system $\tilde{f}(p)$ on $\tilde{U} \triangleq U \cap \tilde{M}$ such that

$$\tilde{f}(p) : \xi^U = \xi^U(x^K) = \xi^U(x^0), \quad \xi^0 = \xi^0(x^0), \quad U, K = 1, 2, 3, \tau. \quad (54)$$

(2) If L is internal-directed, then the above coordinate frames $(\tilde{U}, \xi^U)$ and $(\tilde{U}, x^K)$ of $\tilde{f}(p)$ uniquely determine the coordinate frames $(\tilde{U}, \tilde{\xi}^\alpha)$ and $(\tilde{U}, \tilde{x}^\mu)$ such that

$$\tilde{f}(p) : \tilde{\xi}^\alpha = \tilde{\xi}^\alpha(\tilde{x}^\mu) = \tilde{\xi}^\alpha(\tilde{x}^\tau), \quad \tilde{\xi}^\tau = \tilde{\xi}^\tau(\tilde{x}^\tau), \quad \alpha, \mu = 0, 1, 2, 3 \quad (55)$$

and the coordinates satisfy

$$\tilde{\xi}^s = \xi^s, \quad \tilde{\xi}^\tau = \xi^\tau, \quad \tilde{\xi}^0 = \xi^0, \quad \tilde{x}^i = x^i, \quad \tilde{x}^\tau = x^\tau, \quad \tilde{x}^0 = x^0.$$

That is to say, $\tilde{f}(p)$ is just exactly the reference system in conventional sense, which has two different coordinate representations (54) and (55).

We speak of

$$\tilde{f} : \tilde{M} \rightarrow REF_{\tilde{M}}, \quad p \mapsto \tilde{f}(p) \in REF_p$$

as a **classical spacetime reference-system**. Thus, inertial system can be strictly interpreted as follows. Suppose we have a geometric manifold $(\tilde{M}, \tilde{g})$. $F_{\tilde{g}}$ is a transformation induced by $\tilde{g}$.

(1) If $\tilde{\delta}_{\alpha\beta} \tilde{B}_\mu^\alpha \tilde{B}_\nu^\beta = \tilde{\varepsilon}_{\mu\nu}$, then $\tilde{g}$ is said to be **(Lorentz) orthogonal**. In this case, $F_{\tilde{g}}$ is just exactly a local Lorentz transformation.

(2) If $\tilde{B}_\mu^\alpha$ and $\tilde{C}_\alpha^\mu$ are constants on $\tilde{M}$, then $\tilde{g}$ is said to be **flat**.

(3) If $\tilde{g}$ is both orthogonal and flat, then $\tilde{g}$ is said to be an **inertial-system**. In this case, $F_{\tilde{g}}$ is just exactly a Lorentz transformation.

Remark 4.2.1. Due to

$$\begin{aligned} (d\tilde{\xi}^\tau)^2 &= (d\xi^0)^2 - \sum_{s=1}^3 (d\xi^s)^2 = \tilde{\delta}_{\alpha\beta} d\tilde{\xi}^\alpha d\tilde{\xi}^\beta = \tilde{G}_{\mu\nu} d\tilde{x}^\mu d\tilde{x}^\nu, \quad \tilde{G}_{\mu\nu} \triangleq \tilde{\delta}_{\alpha\beta} \tilde{B}_\mu^\alpha \tilde{B}_\nu^\beta, \\ (d\tilde{x}^\tau)^2 &= (dx^0)^2 - \sum_{i=1}^3 (dx^i)^2 = \tilde{\varepsilon}_{\mu\nu} d\tilde{x}^\mu d\tilde{x}^\nu = \tilde{H}_{\alpha\beta} d\tilde{\xi}^\alpha d\tilde{\xi}^\beta, \quad \tilde{H}_{\alpha\beta} \triangleq \tilde{\varepsilon}_{\mu\nu} \tilde{C}_\alpha^\mu \tilde{C}_\beta^\nu, \end{aligned}$$

it is easy to know that $\tilde{g}$ is orthogonal if and only if $d\tilde{x}^\tau = d\tilde{x}^\tau$, i.e. $\tilde{G}_{\tau\tau} \triangleq \tilde{B}_\tau^\tau \tilde{B}_\tau^\tau = 1$. It is only in this case that we can denote $d\tilde{\xi}^\tau$ and $d\tilde{x}^\tau$ uniformly by $d\tau$, otherwise we should be aware of the difference between $d\tilde{\xi}^\tau$ and $d\tilde{x}^\tau$ in non-trivial gravitational field. No matter whether $\tilde{g}$ is an inertial-system or not, and whether there is a non-trivial gravitation field or not, $(d\tilde{\xi}^\tau)^2 = (d\xi^0)^2 - \sum_{s=1}^3 (d\xi^s)^2$ and $(d\tilde{x}^\tau)^2 = (dx^0)^2 - \sum_{i=1}^3 (dx^i)^2$ are always both true in their respective coordinate frames.

Remark 4.2.2. The evolution lemmas in section 3.3 can be expressed in Minkowski coordinate as:

- (i) If $\frac{d}{dt} \cong \frac{d}{d\tilde{t}_L}$ and $d\tilde{f} \simeq d\tilde{f}_L$, then $\left\langle \frac{d}{dt}, d\tilde{f} \right\rangle = \left\langle \frac{d}{d\tilde{t}_L}, d\tilde{f}_L \right\rangle$.
- (ii) The following conclusions are true.

$$\begin{aligned} w^\mu \frac{\partial}{\partial \tilde{x}^\mu} &\cong w^\tau \frac{d}{d\tilde{x}^\tau} \Leftrightarrow w^\mu = w^\tau \tilde{\varepsilon}_\tau^\mu, & \bar{w}_\mu \frac{\partial}{\partial \tilde{x}_\mu} &\cong \bar{w}_\tau \frac{d}{d\tilde{x}_\tau} \Leftrightarrow \bar{w}_\mu = \bar{w}_\tau \tilde{\varepsilon}_\mu^\tau, \\ w_\mu d\tilde{x}^\mu &\simeq w_\tau d\tilde{x}^\tau \Leftrightarrow \tilde{\varepsilon}_\tau^\mu w_\mu = w_\tau, & \bar{w}^\mu d\tilde{x}_\mu &\simeq \bar{w}^\tau d\tilde{x}_\tau \Leftrightarrow \tilde{\varepsilon}_\mu^\tau \bar{w}^\mu = \bar{w}^\tau. \end{aligned}$$

4.3 Affine connection representation of classical spacetime evolution

Let $\tilde{D}$ be the holonomic connection on $(\tilde{M}, \tilde{G})$, and denote $\tilde{t}_{L;\tau} \triangleq \tilde{t}_{;\sigma} \tilde{\varepsilon}_\tau^\sigma$, then the absolute differential and gradient of section 3.4 can be expressed on $\tilde{M}$ in Minkowski coordinate as

$$\begin{aligned} \tilde{D}\tilde{t} &\triangleq \tilde{t}_{;\sigma} d\tilde{x}^\sigma, & \tilde{D}_L \tilde{t}_L &\triangleq \tilde{t}_{L;\tau} d\tilde{x}^\tau, \\ \tilde{\nabla} \tilde{t} &\triangleq \tilde{t}_{;\sigma} \frac{\partial}{\partial \tilde{x}^\sigma}, & \tilde{\nabla}_L \tilde{t}_L &\triangleq \tilde{t}_{L;\tau} \frac{d}{d\tilde{x}_\tau}. \end{aligned}$$

Evidently, $\tilde{D}\tilde{t} \simeq \tilde{D}_L \tilde{t}_L$ if and only if L is an arbitrary path. $\tilde{\nabla} \tilde{t} \cong \tilde{\nabla}_L \tilde{t}_L$ if and only if L is the gradient line.

Definition 4.3.1. Similar to section 3.6, suppose a charge $\tilde{\rho}$ of $\tilde{\mathcal{F}}$ evolves on $(\tilde{M}, \tilde{G})$. We have the following definitions.

- (1) $\tilde{m}^\tau \triangleq \tilde{\rho}^{;\tau}$ and $\tilde{m}_\tau \triangleq \tilde{\rho}_{;\tau}$ are said to be the **rest mass** of $\tilde{\rho}$.
- (2) $\tilde{p}^\mu \triangleq -\tilde{\rho}^{;\mu}$ and $\tilde{p}_\mu \triangleq -\tilde{\rho}_{;\mu}$ are said to be the **energy-momentum** of $\tilde{\rho}$, and $\tilde{E}^0 \triangleq \tilde{\rho}^{;0}$, $\tilde{E}_0 \triangleq \tilde{\rho}_{;0}$ are said to be the **energy** of $\tilde{\rho}$.
- (3) $\tilde{M}^\tau \triangleq \frac{d\tilde{\rho}}{d\tilde{x}^\tau}$ and $\tilde{M}_\tau \triangleq \frac{d\tilde{\rho}}{d\tilde{x}_\tau}$ are said to be the **canonical rest mass** of $\tilde{\rho}$.
- (4) $\tilde{P}^\mu \triangleq -\frac{\partial \tilde{\rho}}{\partial \tilde{x}_\mu}$ and $\tilde{P}_\mu \triangleq -\frac{\partial \tilde{\rho}}{\partial \tilde{x}^\mu}$ are said to be the **canonical energy-momentum** of $\tilde{\rho}$, and $\tilde{H}^0 \triangleq \frac{\partial \tilde{\rho}}{\partial \tilde{x}_0}$, $\tilde{H}_0 \triangleq \frac{\partial \tilde{\rho}}{\partial \tilde{x}^0}$ are said to be the **canonical energy** of $\tilde{\rho}$.

Discussion 4.3.1. Similar to Proposition 3.6.1, $\forall p \in \tilde{M}$, if and only if the evolution direction $[L_p] = \tilde{\nabla} \tilde{\rho}|_p$, the directional derivative is

$$\left\langle \tilde{m}_\tau \frac{d}{d\tilde{x}_\tau}, \tilde{m}_\tau d\tilde{x}^\tau \right\rangle = \left\langle \tilde{p}_\mu \frac{\partial}{\partial \tilde{x}_\mu}, \tilde{p}_\mu d\tilde{x}^\mu \right\rangle,$$

that is $\tilde{G}^{\tau\tau} \tilde{m}_\tau \tilde{m}_\tau = \tilde{G}^{\mu\nu} \tilde{p}_\mu \tilde{p}_\nu$, or

$$\tilde{m}_\tau \tilde{m}^\tau = \tilde{p}_\mu \tilde{p}^\mu,$$

which is the affine connection representation of energy-momentum equation.

Similar to Proposition 3.6.2, according to the evolution lemma, $\forall p \in \tilde{M}$, if and only if the evolution direction $[L_p] = \tilde{\nabla} \tilde{\rho}|_p$, we have $\tilde{p}_\mu = -\tilde{m}_\tau \frac{d\tilde{x}_\mu}{d\tilde{x}_\tau}$, that is $\tilde{E}_0 = \tilde{m}_\tau \frac{d\tilde{x}_0}{d\tilde{x}_\tau} = \tilde{m}_\tau \frac{dx_0}{dx_\tau}$ and $\tilde{p}_i = -\tilde{m}_\tau \frac{d\tilde{x}_i}{d\tilde{x}_\tau} = \tilde{m}_\tau \frac{-d\tilde{x}_i}{d\tilde{x}_\tau} = \tilde{m}_\tau \frac{dx_i}{dx_\tau} = \tilde{E}_0 \frac{dx_i}{dx_0}$. This can also be regarded as the origin of $p = mv$.

Similar to Remark 3.6.2, denote

$$[\tilde{\rho} \tilde{\Gamma}_\omega] \triangleq \frac{\partial \tilde{\rho}_{\mu\nu}}{\partial \tilde{x}^\omega} - \tilde{\rho}_{\mu\nu;\omega} = \tilde{\rho}_{\mu\chi} \tilde{\Gamma}_{\nu\omega}^\chi + \tilde{\rho}_{\chi\nu} \tilde{\Gamma}_{\mu\omega}^\chi, \quad [\tilde{\rho} \tilde{R}_{\rho\sigma}] \triangleq \tilde{\rho}_{\mu\chi} \tilde{R}_{\nu\rho\sigma}^\chi + \tilde{\rho}_{\chi\nu} \tilde{R}_{\mu\rho\sigma}^\chi.$$

Then for the same reason as Remark 3.6.2, based on Definition 4.3.1, we can strictly obtain

$$\tilde{f}_\rho \triangleq \tilde{p}_{\rho;\tau} = \tilde{m}_{\tau;\rho} - \tilde{p}_\sigma \tilde{\varepsilon}_{\tau;\rho}^\sigma + [\tilde{\rho} \tilde{R}_{\rho\sigma}] \tilde{\varepsilon}_\tau^\sigma. \quad (56)$$

In the mass-point model, $\tilde{m}_{\tau;\rho}$ and $\tilde{\varepsilon}_{\tau;\rho}^\sigma$ do not make sense, so Eq.(56) turns into

$$\tilde{f}_\rho = [\tilde{\rho} \tilde{R}_{\rho\sigma}] \tilde{\varepsilon}_\tau^\sigma.$$

This is the affine connection representation of the force of interaction (e.g. the Lorentz force $\mathbf{f} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$ or $f_\rho = j^\sigma F_{\rho\sigma}$ of the electrodynamics).

Similar to Definition 3.6.2, let $\tilde{\mathcal{P}}(b, a)$ be the totality of paths on $\tilde{M}$ from point a to point b . And let $L \in \tilde{\mathcal{P}}(b, a)$, and parameter $\tilde{x}^\tau$ satisfy $\tau_a \triangleq \tilde{x}^\tau(a) < \tilde{x}^\tau(b) \triangleq \tau_b$. The affine connection representation of action in Minkowski coordinates can be defined as

$$\tilde{s}(L) \triangleq \int_L \tilde{D}\tilde{\rho} = \int_L \tilde{\rho}_\mu d\tilde{x}^\mu = \int_{\tau_a}^{\tau_b} \tilde{m}_\tau d\tilde{x}^\tau, \quad \tilde{s}(L) \triangleq \int_{\tau_a}^{\tau_b} (\gamma^\mu \tilde{\rho}_{;\mu} + \tilde{m}_\tau) d\tilde{x}^\tau. \quad (57)$$

There are more illustrations in Remark 4.4.1.

4.4 Affine connection representation of Dirac equation

Discussion 4.4.1. Define Dirac algebras γ^μ and γ^α such that

$$\gamma^\mu = \tilde{C}_\alpha^\mu \gamma^\alpha, \quad \gamma^\alpha \gamma^\beta + \gamma^\beta \gamma^\alpha = 2\tilde{\delta}^{\alpha\beta}, \quad \gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2\tilde{G}^{\mu\nu}.$$

Suppose $(\tilde{M}, \tilde{G})$ is orthogonal. According to Remark 4.2.1, $\tilde{G}_{\tau\tau} = 1$. Due to Discussion 4.3.1, in a gradient direction of $\tilde{\rho} \triangleq \tilde{\rho}_{\omega\nu}$, we have

$$\begin{aligned} \tilde{\rho}_{;\mu} \tilde{\rho}^{;\mu} = \tilde{\rho}_{;\tau} \tilde{\rho}^{;\tau} &\Leftrightarrow \tilde{G}^{\mu\nu} \tilde{\rho}_{;\mu} \tilde{\rho}_{;\nu} = \tilde{m}_\tau^2 \\ &\Leftrightarrow (\gamma^\mu \tilde{\rho}_{;\mu})(\gamma^\nu \tilde{\rho}_{;\nu}) + (\gamma^\nu \tilde{\rho}_{;\nu})(\gamma^\mu \tilde{\rho}_{;\mu}) = 2\tilde{m}_\tau^2 \\ &\Leftrightarrow (\gamma^\mu \tilde{\rho}_{;\mu})(\gamma^\nu \tilde{\rho}_{;\nu}) = \tilde{m}_\tau^2 \\ &\Leftrightarrow (\gamma^\mu \tilde{\rho}_{;\mu})^2 = \tilde{m}_\tau^2. \end{aligned}$$

Without loss of generality, take $\gamma^\mu \tilde{\rho}_{;\mu} = \tilde{m}_\tau$, that is

$$\gamma^\mu \tilde{\rho}_{\omega\nu;\mu} = \tilde{m}_{\omega\nu\tau}. \quad (58)$$

Next, denote

$$[g\tilde{I}_\mu]^{\omega\nu} \triangleq \sum_\sigma \tilde{G}^{\nu\nu'} \tilde{I}_{\nu'\sigma\mu} + \sum_\kappa \tilde{G}^{\omega\omega'} \tilde{I}_{\omega'\kappa\mu}, \quad \tilde{D}_\mu^{\omega\nu} \triangleq \partial_\mu - [g\tilde{I}_\mu]^{\omega\nu}.$$

From Eq.(58), it is obtained that

$$\begin{aligned} \sum_{\omega,\nu} \gamma^\mu \tilde{\rho}_{\omega\nu;\mu} &= \sum_{\omega,\nu} \tilde{m}_{\omega\nu\tau} \Leftrightarrow \sum_{\omega,\nu} \gamma^\mu \left(\partial_\mu \rho_{\omega\nu} - \tilde{\rho}_{\omega\chi} \tilde{I}_{\nu\mu}^\chi - \tilde{\rho}_{\chi\nu} \tilde{I}_{\omega\mu}^\chi \right) = \sum_{\omega,\nu} \tilde{m}_{\omega\nu\tau} \\ &\Leftrightarrow \sum_{\omega,\nu} \gamma^\mu \left(\partial_\mu \rho_{\omega\nu} - \tilde{\rho}_{\omega\nu} \sum_\sigma \tilde{I}_{\sigma\mu}^\nu - \tilde{\rho}_{\omega\nu} \sum_\kappa \tilde{I}_{\kappa\mu}^\omega \right) = \sum_{\omega,\nu} \tilde{m}_{\omega\nu\tau} \\ &\Leftrightarrow \sum_{\omega,\nu} \gamma^\mu \left(\partial_\mu \rho_{\omega\nu} - \tilde{\rho}_{\omega\nu} [g\tilde{I}_\mu]^{\omega\nu} \right) = \sum_{\omega,\nu} \tilde{m}_{\omega\nu\tau} \\ &\Leftrightarrow \sum_{\omega,\nu} \gamma^\mu \left(\partial_\mu - [g\tilde{I}_\mu]^{\omega\nu} \right) \tilde{\rho}_{\omega\nu} = \sum_{\omega,\nu} \tilde{m}_{\omega\nu\tau}, \end{aligned} \quad (59)$$

that is

$$\sum_{\omega,\nu} \gamma^\mu \tilde{D}_\mu^{\omega\nu} \tilde{\rho}_{\omega\nu} = \sum_{\omega,\nu} \tilde{m}_{\omega\nu\tau}, \quad \tilde{D}_\mu^{\omega\nu} \triangleq \partial_\mu - [g\tilde{I}_\mu]^{\omega\nu}. \quad (60)$$

We speak of the real-valued Eq.(58) and (60) as **affine Dirac equations**.

Discussion 4.4.2. Next, we construct a kind of complex-valued representation of affine Dirac equation. The restriction of the charge $\tilde{\rho}_{\omega\nu}$ to $(\tilde{U}, \tilde{x}^\mu)$ is a function $\tilde{\rho}_{\omega\nu}(\tilde{x}^\mu)$ with respect to the coordinates $(\tilde{x}^\mu) \triangleq (\tilde{x}^0, \tilde{x}^1, \tilde{x}^2, \tilde{x}^3)$. Let

$$\tilde{\mathcal{P}}_{\omega\nu}(\tilde{x}^0) \triangleq \int_{(\tilde{x}^1, \tilde{x}^2, \tilde{x}^3)} \tilde{\rho}_{\omega\nu}(\tilde{x}^\mu) d^3\tilde{x}.$$

Suppose a function $f_{\omega\nu} = f_{\omega\nu}(\tilde{x}^\mu)$ on $(\tilde{U}, \tilde{x}^\mu)$ satisfies that

$$\tilde{\rho}_{\omega\nu} = (f_{\omega\nu})^2 \tilde{\mathcal{P}}_{\omega\nu}, \quad \int_{(\tilde{x}^1, \tilde{x}^2, \tilde{x}^3)} (f_{\omega\nu})^2 d^3\tilde{x} = 1, \quad \varepsilon_\tau^\mu \frac{\partial f_{\omega\nu}}{\partial \tilde{x}^\mu} = 0, \quad \gamma^\mu \frac{\partial f_{\omega\nu}}{\partial \tilde{x}^\mu} = 0.$$

We define $\psi_{\omega\nu}$ and $\tilde{\mathbb{M}}_{\omega\nu\tau}$ in the following way.

$$\begin{aligned}\tilde{y}_{\omega\nu} &\triangleq \int_L d\tilde{\rho}_{\omega\nu} = \int_L \frac{d\tilde{\rho}_{\omega\nu}}{d\tilde{x}^\tau} d\tilde{x}^\tau = \int_L \left(\frac{d(f_{\omega\nu}^2)}{d\tilde{x}^\tau} \tilde{\mathbb{P}}_{\omega\nu} + f_{\omega\nu}^2 \frac{d\tilde{\mathbb{P}}_{\omega\nu}}{d\tilde{x}^\tau} \right) d\tilde{x}^\tau = f_{\omega\nu}^2 \int_L \frac{d\tilde{\mathbb{P}}_{\omega\nu}}{d\tilde{x}^\tau} d\tilde{x}^\tau \triangleq f_{\omega\nu}^2 \tilde{Y}_{\omega\nu}, \\ \psi_{\omega\nu} &\triangleq f_{\omega\nu} e^{i\tilde{Y}_{\omega\nu}}, \quad \tilde{m}_{\omega\nu\tau} \triangleq \tilde{\rho}_{\omega\nu;\tau} = (f_{\omega\nu}^2)_{,\tau} \tilde{\mathbb{P}}_{\omega\nu} + f_{\omega\nu}^2 \tilde{\mathbb{P}}_{\omega\nu;\tau} = f_{\omega\nu}^2 \tilde{\mathbb{P}}_{\omega\nu;\tau} \triangleq f_{\omega\nu}^2 \tilde{\mathbb{M}}_{\omega\nu\tau}.\end{aligned}$$

In the QFT propagator, we usually take S in the path integral $\int e^{iS} \mathcal{D}\psi$ of a fermion in the form of

$$- \int \left(i\bar{\psi} \gamma^\mu D_\mu \psi - \bar{\psi} \tilde{\mathbb{M}}_\tau \psi \right) d^4 \tilde{x},$$

where S and $d^4 \tilde{x}$ are both covariant. We believe that the external spatial integral $\int_{(\tilde{x}^1, \tilde{x}^2, \tilde{x}^3)} d^3 \tilde{x}$ is not an essential part for evolution, so for the sake of simplicity, we do not take into account the external spatial part $\int_{(\tilde{x}^1, \tilde{x}^2, \tilde{x}^3)} d^3 \tilde{x}$, but only consider the evolution part $\int_L d\tilde{x}^0$. Meanwhile, in order to remain the covariance, $\int_L d\tilde{x}^0$ has to be replaced by $\int_L d\tilde{x}^\tau$. Thus, in affine connection representation of gauge fields, we shall consider an action in the form of

$$- \int_L \left(i\bar{\psi} \gamma^\mu D_\mu \psi - \bar{\psi} \tilde{\mathbb{M}}_\tau \psi \right) d\tilde{x}^\tau.$$

Concretely speaking, denote

$$\tilde{D}_{\omega\nu\mu} \triangleq \frac{\partial}{\partial \tilde{x}^\mu} - i[\tilde{\mathbb{P}} \tilde{F}_\mu]_{\omega\nu}, \quad [\tilde{\mathbb{P}} \tilde{F}_\mu]_{\omega\nu} \triangleq \sum_\sigma \tilde{\mathbb{P}}_{\omega\nu} \tilde{F}_{\sigma\mu}^\nu + \sum_\kappa \tilde{\mathbb{P}}_{\omega\nu} \tilde{F}_{\kappa\mu}^\omega.$$

From Eq.(57), we have

$$\tilde{s}_{\omega\nu}(L) \triangleq \int_L (\gamma^\mu \tilde{\rho}_{\omega\nu;\mu} + \tilde{m}_{\omega\nu\tau}) d\tilde{x}^\tau.$$

And from Eq.(59) we know $\sum_{\omega,\nu} \gamma^\mu \tilde{\rho}_{\omega\nu;\mu} = \sum_{\omega,\nu} \gamma^\mu \tilde{D}_\mu^{\omega\nu} \tilde{\rho}_{\omega\nu}$. Then it is obtained that

$$\begin{aligned}\tilde{s}_{\tilde{\rho}}(L) &\triangleq \sum_{\omega,\nu} \tilde{s}_{\omega\nu}(L) = \int_L \sum_{\omega,\nu} (\gamma^\mu \tilde{\rho}_{\omega\nu;\mu} + \tilde{m}_{\omega\nu\tau}) d\tilde{x}^\tau = \int_L \sum_{\omega,\nu} (\gamma^\mu \tilde{D}_\mu^{\omega\nu} \tilde{\rho}_{\omega\nu} + \tilde{m}_{\omega\nu\tau}) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(\gamma^\mu (\partial_\mu \tilde{\rho}_{\omega\nu} - [g \tilde{F}_\mu]^{\omega\nu} \tilde{\rho}_{\omega\nu}) + \tilde{m}_{\omega\nu\tau} \right) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(\gamma^\mu (\partial_\mu \tilde{y}_{\omega\nu} - [g \tilde{F}_\mu]^{\omega\nu} \tilde{\rho}_{\omega\nu}) + \tilde{m}_{\omega\nu\tau} \right) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(\gamma^\mu (\partial_\mu (f_{\omega\nu}^2 \tilde{Y}_{\omega\nu}) - [g \tilde{F}_\mu]^{\omega\nu} f_{\omega\nu}^2 \tilde{\mathbb{P}}_{\omega\nu}) + f_{\omega\nu}^2 \tilde{\mathbb{M}}_{\omega\nu\tau} \right) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(\gamma^\mu (\partial_\mu \tilde{Y}_{\omega\nu} - [\tilde{\mathbb{P}} \tilde{F}_\mu]_{\omega\nu}) f_{\omega\nu}^2 + f_{\omega\nu}^2 \tilde{\mathbb{M}}_{\omega\nu\tau} \right) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(f_{\omega\nu} e^{-i\tilde{Y}_{\omega\nu}} \gamma^\mu \left(f_{\omega\nu} e^{i\tilde{Y}_{\omega\nu}} \partial_\mu \tilde{Y}_{\omega\nu} - [\tilde{\mathbb{P}} \tilde{F}_\mu]_{\omega\nu} f_{\omega\nu} e^{i\tilde{Y}_{\omega\nu}} \right) + f_{\omega\nu} e^{-i\tilde{Y}_{\omega\nu}} \tilde{\mathbb{M}}_{\omega\nu\tau} f_{\omega\nu} e^{i\tilde{Y}_{\omega\nu}} \right) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(-\bar{\psi}_{\omega\nu} i \gamma^\mu \left(e^{i\tilde{Y}_{\omega\nu}} \partial_\mu f_{\omega\nu} + f_{\omega\nu} e^{i\tilde{Y}_{\omega\nu}} i \partial_\mu \tilde{Y}_{\omega\nu} - i [\tilde{\mathbb{P}} \tilde{F}_\mu]_{\omega\nu} \psi_{\omega\nu} \right) + \bar{\psi}_{\omega\nu} \tilde{\mathbb{M}}_{\omega\nu\tau} \psi_{\omega\nu} \right) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(-\bar{\psi}_{\omega\nu} i \gamma^\mu \left(\partial_\mu (f_{\omega\nu} e^{i\tilde{Y}_{\omega\nu}}) - i [\tilde{\mathbb{P}} \tilde{F}_\mu]_{\omega\nu} \psi_{\omega\nu} \right) + \bar{\psi}_{\omega\nu} \tilde{\mathbb{M}}_{\omega\nu\tau} \psi_{\omega\nu} \right) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(-\bar{\psi}_{\omega\nu} i \gamma^\mu \left(\partial_\mu - i [\tilde{\mathbb{P}} \tilde{F}_\mu]_{\omega\nu} \right) \psi_{\omega\nu} + \bar{\psi}_{\omega\nu} \tilde{\mathbb{M}}_{\omega\nu\tau} \psi_{\omega\nu} \right) d\tilde{x}^\tau \\ &= \int_L \sum_{\omega,\nu} \left(-\bar{\psi}_{\omega\nu} i \gamma^\mu \tilde{D}_{\omega\nu\mu} \psi_{\omega\nu} + \bar{\psi}_{\omega\nu} \tilde{\mathbb{M}}_{\omega\nu\tau} \psi_{\omega\nu} \right) d\tilde{x}^\tau = - \int_L \sum_{\omega,\nu} \bar{\psi}_{\omega\nu} \left(i \gamma^\mu \tilde{D}_{\omega\nu\mu} - \tilde{\mathbb{M}}_{\omega\nu\tau} \right) \psi_{\omega\nu} d\tilde{x}^\tau.\end{aligned}\tag{61}$$

Thus, we have obtained a complex-valued representation of gradient direction of $\tilde{\rho}_{\omega\nu}$.

Remark 4.4.1. From the above discussion, we know in the gradient direction of $\rho_{\omega\nu}$, that

$$-\sum_{\omega,\nu} \bar{\psi}_{\omega\nu} i\gamma^\mu \tilde{D}_{\omega\nu\mu} \psi_{\omega\nu} d\tilde{x}^\tau = \sum_{\omega,\nu} \tilde{D} \tilde{\rho}_{\omega\nu}.$$

This shows that $s(L)$ and $\tilde{s}(L)$ in Definition 3.6.2 and Remark 3.6.1 are indeed applicable for constructing propagator by $e^{is(L)}$ and $e^{i\tilde{s}(L)}$ in affine connection representation of gauge fields. Therefore, the idea in Discussion 3.9.3 are reasonable.

4.5 From classical spacetime back to full-dimensional space

Discussion 4.5.1. Now there is a problem. $(\tilde{M}, \tilde{\mathcal{F}})$ and $(\tilde{M}, \tilde{\mathcal{G}})$ cannot totally reflect the geometric properties of internal space of $(M, \mathcal{F})$ and $(M, \mathcal{G})$. Concretely speaking:

In the previous section, we discuss the affine Dirac equation $\gamma^\mu \tilde{\rho}_{\omega\nu;\mu} = \tilde{m}_{\omega\nu\tau}$ on $(\tilde{M}, \tilde{\mathcal{G}})$. Similar to section 3.5, we have the affine Yang-Mills equation $\tilde{K}_{\nu\rho\sigma}^{\mu\cdot\rho} = \tilde{\rho}_\nu^\mu \gamma_\sigma$ on $(\tilde{M}, \tilde{\mathcal{F}})$. Suppose there is no gravitational field, then the remaining non-vanish equations are just only

$$\gamma^\mu \tilde{\rho}_{00;\mu} = \tilde{m}_{00\tau}, \quad \tilde{K}_{0\rho\sigma}^0{}^{\cdot\rho} = \tilde{\rho}_0^0 \gamma_\sigma.$$

There are multiple internal charges

$$\rho_{mn} (m, n = 4, 5, \dots, \mathfrak{D})$$

on $(M, \mathcal{F})$. We intend to use these ρ_{mn} to describe leptons and hadrons. However, via encapsulation of classical spacetime, $(\tilde{M}, \tilde{\mathcal{F}})$ remains only one internal charge $\tilde{\rho}_{00}$, it falls short. It is impossible for the only one real-valued field function $\tilde{\rho}_{00}$ to describe so many leptons and hadrons.

On the premise of not abandoning the $(1+3)$ -dimensional spacetime, if we want to describe gauge fields, there is a method that to use some non-coordinate abstract degrees of freedom on the phase of $e^{iT_a\theta^a}$ of a complex-valued field function ψ , just like that in Standard Model. This way is effective, but not natural. It is not satisfying for a theory to adopt a coordinate representation for external space but a non-coordinate representation for internal space.

A logically more natural way is required to abandon the framework of $(1+3)$ -dimensional spacetime $(\tilde{M}, \tilde{\mathcal{F}})$ and $(\tilde{M}, \tilde{\mathcal{G}})$. We should put internal space and external space together to describe their unified geometry with the same spatial frame. On $(M, \mathcal{F})$ and $(M, \mathcal{G})$, there are enough real-valued field functions ρ_{mn} to describe leptons and hadrons, and enough internal components $[mnP]$ of affine connection to describe gauge potentials.

Therefore, only on the full-dimensional $(M, \mathcal{F})$ and $(M, \mathcal{G})$ can total advantages of affine connection representation of gauge fields be brought into full play, and thereby show complete details of geometric properties of gauge field. So we are going to stop the discussions about the classical spacetime $\tilde{M}$, but to focus on the full-dimensional manifold M .

Discussion 4.5.2. On M , due to $\Gamma_{MNP} = \frac{1}{2} ([MNP] + \{MNP\})$, $[MNP] = \delta_{AD} B_M^D \left(\frac{\partial B_N^A}{\partial x^P} + \left(\frac{A}{BP} \right) B_N^B \right)$ and $G_{MN} = \delta_{AB} B_M^A B_N^B$ we know that gauge field and gravitational field can both be described by spatial frames B_M^A and C_A^M in a reference-system. Reference-system is the common origination of gauge field and gravitational field. The invariance under reference-system transformation is the common origination of gauge covariance and general covariance.

We adopt the components $[mnP]$ of $[MNP]$ with $m, n \in \{4, 5, \dots, \mathfrak{D}\}$ to describe the gauge potentials of typical gauge fields such as electromagnetic, weak and strong interaction fields, and adopt the components ρ_{mn} of ρ_{MN} with $m, n \in \{4, 5, \dots, \mathfrak{D}\}$ to describe the charges of leptons and hadrons. The physical meanings of the other components of ρ_{MN} and $[MNP]$ are not clear at present, maybe they could be used to describe dark matters and their interactions.

On orthogonal $(M, \mathcal{G})$ and $(M, \mathcal{F})$, there are full-dimensional field equations, i.e. affine Dirac equation and affine Yang-Mills equation

$$\gamma^P \rho_{MN;P} = \rho_{MN;0}, \quad K_{NPQ}^M{}^{\cdot P} = \rho_N^M \gamma_Q, \quad (62)$$

which reflect the on-shell evolution directions $\nabla\rho$ and ∇t , respectively. Their quantum evolutions are described by the propagators in Definition 3.9.5 or Discussion 3.9.3.

Discussion 4.5.3. On an orthogonal $(M, \mathcal{G})$, Eq.(61) presents as a full-dimensional action

$$s_\rho(L) = \int_L \sum_{M,N} (\gamma^P \rho_{MN;P} + \varepsilon_0^P \rho_{MN;P}) dx^0 = -i \int_L \sum_{M,N} \bar{\psi}_{MN} (\gamma^P D_{MNP} + \varepsilon_0^P D_{MNP}) \psi_{MN} dx^0. \quad (63)$$

If and only if $L_k : g \rightarrow g'$ is an orthogonal transformation, L_k sends $s_\rho(L)$ to

$$s'_\rho(L) = \int_L \sum_{M,N} (\gamma^{P'} \rho_{MN;P'} + \varepsilon_0^{P'} \rho_{MN;P'}) dx^{0'} = -i \int_L \sum_{M,N} \bar{\psi}'_{MN} (\gamma^{P'} D'_{MNP'} + \varepsilon_0^{P'} D'_{MNP'}) \psi'_{MN} dx^{0'},$$

where ρ_{MN} is determined by the reference-system $f \circ f$ but not $g \circ g$, so ρ_{MN} does not vary with the transformation $L_k : g \rightarrow g'$. We see that in affine connection representation of gauge fields, the gauge transformations $\psi \mapsto \psi'$ and $D \mapsto D'$ essentially boil down to the reference-system transformation L_k .

Remark 1. For a general $(M, \mathcal{G})$, $\mathcal{G}$ is not necessarily orthogonal, so the corresponding action should be described by

$$s_{MN}(L) = \int_L (B_0^0 \gamma^P \rho_{MN;P} + \varepsilon_0^P \rho_{MN;P}) dx^0.$$

In this general case, Definition 3.6.2 and the method in Discussion 3.9.3 are also available and effective, where we take

$$s_{MN}(L) = \int_L D\rho_{MN}.$$

Remark 2. We see that the real-valued representation of action is more concise than the complex-valued representation of action. Hence, it is more convenient to adopt real-valued representations for field function, field equation and action.

In the following sections, we are going to use $[MNP]$ to show the affine connection representations of electromagnetic, weak and strong interaction fields, and to adopt the real-valued representation $\rho_{MN;P}$ to discuss the interactions between gauge fields and elementary particles. They are based on the following definition.

Definition 4.5.1. Let $M = P \times N$, $r \triangleq \dim P = 3$ and $\mathfrak{D} \triangleq \dim M = 5$ or 6 or 8 . Consider $\mathcal{F} = f \circ f$ and $\mathcal{G} = g \circ g$ that are defined by Eq.(17), that is, $\forall p \in M$,

$$(U, \alpha^{A'}) \xrightarrow{f(p)} (U, \xi^A) \xrightarrow{f(p)} (U, x^M) \xleftarrow{g(p)} (U, \zeta^A) \xleftarrow{g(p)} (U, \beta^{A'})$$

and furthermore let

$$\begin{aligned} f(p) : \xi^a &= \xi^a(x^m), \quad \xi^s = \delta_i^s x^i; & f(p) : \alpha^{a'} &= \alpha^{a'}(\xi^a), \quad \alpha^{s'} = \delta_s^{s'} \xi^s; \\ g(p) : \zeta^a &= \zeta^a(x^m), \quad \zeta^s = \delta_i^s x^i; & g(p) : \beta^{a'} &= \beta^{a'}(\zeta^a), \quad \beta^{s'} = \delta_s^{s'} \zeta^s; \end{aligned} \quad (64)$$

$(s', s, i = 1, 2, 3; a', a, m = 4, 5, \dots, \mathfrak{D})$ and both of $\mathcal{F}$ and $\mathcal{G}$ satisfy

$$(i) G_{mn} = \text{const}, \quad (ii) \text{ when } m \neq n, G_{mn} = 0. \quad (65)$$

In the above case, we use $\mathcal{F}$ and $\mathcal{G}$ to describe electromagnetic, weak and strong interactions without gravitation.

5 Affine connection representation of the gauge field of weak-electromagnetic interaction

Definition 5.1. Suppose $(M, \mathcal{F})$ and $(M, \mathcal{G})$ conform to Definition 4.5.1. Let $\mathfrak{D} = r + 2 = 5$ and both of $\mathcal{F}$ and $\mathcal{G}$ satisfy

$$G^{(\mathfrak{D}-1)(\mathfrak{D}-1)} = G^{\mathfrak{D}\mathfrak{D}}.$$

Thus, $\mathcal{F}$ and $\mathcal{G}$ can describe weak and electromagnetic interactions.

Proposition 5.1. Let the holonomic connection of $(M, \mathcal{F})$ be Γ_{NP}^M and Γ_{MNP} . And let the coefficients of curvature tensor of $(M, \mathcal{F})$ be K_{NPPQ}^M and K_{MNPQ} . Denote

$$\begin{cases} B_P \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}\mathfrak{D}P} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P}), & B_{PQ} \triangleq \frac{1}{\sqrt{2}} (K_{\mathfrak{D}\mathfrak{D}PQ} + K_{(\mathfrak{D}-1)(\mathfrak{D}-1)PQ}), \\ A_P^3 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}\mathfrak{D}P} - \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P}), & F_{PQ}^3 \triangleq \frac{1}{\sqrt{2}} (K_{\mathfrak{D}\mathfrak{D}PQ} - K_{(\mathfrak{D}-1)(\mathfrak{D}-1)PQ}), \\ A_P^1 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} + \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}), & F_{PQ}^1 \triangleq \frac{1}{\sqrt{2}} (K_{(\mathfrak{D}-1)\mathfrak{D}PQ} + K_{\mathfrak{D}(\mathfrak{D}-1)PQ}), \\ A_P^2 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} - \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}), & F_{PQ}^2 \triangleq \frac{1}{\sqrt{2}} (K_{(\mathfrak{D}-1)\mathfrak{D}PQ} - K_{\mathfrak{D}(\mathfrak{D}-1)PQ}). \end{cases}$$

And denote $g \triangleq \sqrt{(G^{(\mathfrak{D}-1)(\mathfrak{D}-1)})^2 + (G^{\mathfrak{D}\mathfrak{D}})^2}$. Thus, the following equations hold spontaneously.

$$\begin{aligned} B_{PQ} &= \frac{\partial B_Q}{\partial x^P} - \frac{\partial B_P}{\partial x^Q}, \\ F_{PQ}^3 &= \frac{\partial A_Q^3}{\partial x^P} - \frac{\partial A_P^3}{\partial x^Q} + g(A_P^1 A_Q^2 - A_P^2 A_Q^1), \\ F_{PQ}^1 &= \frac{\partial A_Q^1}{\partial x^P} - \frac{\partial A_P^1}{\partial x^Q} + g(A_P^2 A_Q^3 - A_P^3 A_Q^2), \\ F_{PQ}^2 &= \frac{\partial A_Q^2}{\partial x^P} - \frac{\partial A_P^2}{\partial x^Q} + g(A_P^1 A_Q^3 - A_P^3 A_Q^1). \end{aligned}$$

Proof. Due to Eq.(64) it is obtained that the semi-metric of $(M, \mathfrak{f})$ satisfies

$$(B_{\mathfrak{f}})_{a'}^{s'} = 0, \quad (C_{\mathfrak{f}})_{s'}^a = 0, \quad (B_{\mathfrak{f}})_{s'}^{a'} = 0, \quad (C_{\mathfrak{f}})_{a'}^s = 0, \quad (B_{\mathfrak{f}})_{s'}^{s'} = \delta_s^{s'}, \quad (C_{\mathfrak{f}})_{s'}^s = \delta_s^s.$$

Then, compute $(\frac{A}{BC})_{\mathfrak{f}} \triangleq \frac{1}{2}(C_{\mathfrak{f}})_{A'}^A \left(\frac{\partial(B_{\mathfrak{f}})_{B'}^{A'}}{\partial \xi^C} + \frac{\partial(B_{\mathfrak{f}})_{C'}^{A'}}{\partial \xi^B} \right)$ and we obtain

$$(\frac{s}{BC})_{\mathfrak{f}} = 0, \quad (\frac{a}{tu})_{\mathfrak{f}} = 0, \quad (\frac{a}{bC})_{\mathfrak{f}} \neq 0; \quad s, t, u = 1, 2, 3; \quad a, b = 4, 5, \dots, \mathfrak{D}; \quad A, B, C = 1, 2, \dots, \mathfrak{D}.$$

It is obtained from Eq.(64) again that the semi-metric of (M, f) satisfies

$$B_m^s = 0, \quad C_s^m = 0, \quad B_i^a = 0, \quad C_a^i = 0, \quad B_i^s = \delta_i^s, \quad C_s^i = \delta_s^i.$$

Let $s', t', i, j, k = 1, 2, 3$; $a', b', m, n, p = 4, 5, \dots, \mathfrak{D}$. Compute the metric of $(M, \mathcal{F})$ and we obtain

$$\begin{cases} G_{ij} = \delta_{s't'} B_i^{s'} B_j^{t'} + \delta_{a'b'} B_i^{a'} B_j^{b'} = \delta_{s't'} \delta_i^{s'} \delta_j^{t'} = \delta_{ij}, \\ G_{in} = \delta_{s't'} B_i^{s'} B_n^{t'} + \delta_{a'b'} B_i^{a'} B_n^{b'} = 0, \\ G_{mj} = \delta_{s't'} B_m^{s'} B_j^{t'} + \delta_{a'b'} B_m^{a'} B_j^{b'} = 0, \\ G_{mn} = B_m^{\mathfrak{D}-1} B_n^{\mathfrak{D}-1} + B_m^{\mathfrak{D}} B_n^{\mathfrak{D}} = \text{const}, \end{cases} \quad \begin{cases} G^{ij} = \delta^{s't'} C_{s'}^i C_{t'}^j = \delta^{s't'} \delta_s^i \delta_t^j = \delta^{ij}, \\ G^{in} = \delta^{s't'} C_{s'}^i C_{t'}^n = 0, \\ G^{mj} = \delta^{s't'} C_{s'}^m C_{t'}^j = 0, \\ G^{mn} = C_{\mathfrak{D}-1}^m C_{\mathfrak{D}-1}^n + C_{\mathfrak{D}}^m C_{\mathfrak{D}}^n = \text{const}. \end{cases}$$

Compute the holonomic connection of $\mathcal{F}$ according to $\Gamma_{NP}^M \triangleq \frac{1}{2} ([M_P] + \{M_P\}) = \frac{1}{2} \left(C_A^M \frac{\partial B_N^A}{\partial x^P} + C_A^M (\frac{A}{BP})_{\mathfrak{f}} B_N^B \right)$, and it is obtained that

$$\begin{cases} \Gamma_{NP}^i = 0, \\ \Gamma_{jk}^m = 0, \\ \Gamma_{nP}^m = \frac{1}{2} \left(C_a^m \frac{\partial B_n^a}{\partial x^P} + C_a^m (\frac{a}{bP})_{\mathfrak{f}} B_n^b \right), \\ \Gamma_{Np}^m = \frac{1}{2} \left(C_a^m \frac{\partial B_N^a}{\partial x^p} + C_a^m (\frac{a}{Bp})_{\mathfrak{f}} B_N^B \right), \end{cases} \quad \begin{cases} \Gamma_{iNP} = G_{iM'} \Gamma_{NP}^{M'} = G_{ii'} \Gamma_{NP}^{i'} = 0, \\ \Gamma_{mjk} = G_{mM'} \Gamma_{jk}^{M'} = G_{mm'} \Gamma_{jk}^{m'} = 0, \\ \Gamma_{mnp} = \frac{1}{2} \delta_{ab} B_m^b \left(\frac{\partial B_n^a}{\partial x^p} + (\frac{a}{bP})_{\mathfrak{f}} B_n^b \right), \\ \Gamma_{mNp} = \frac{1}{2} \delta_{ab} B_m^b \left(\frac{\partial B_N^a}{\partial x^p} + (\frac{a}{Bp})_{\mathfrak{f}} B_N^B \right). \end{cases} \quad (66)$$

Compute the coefficients of curvature of $\mathcal{F}$, that is

$$K_{nPQ}^m \triangleq \frac{\partial \Gamma_{nP}^m}{\partial x^Q} - \frac{\partial \Gamma_{nQ}^m}{\partial x^P} + \Gamma_{HP}^m \Gamma_{nQ}^H - \Gamma_{nQ}^H \Gamma_{HP}^m, \quad K_{mnpQ} \triangleq G_{mM'} K_{nPQ}^{M'} = G_{mm'} K_{nPQ}^{m'},$$

then we obtain

$$\begin{aligned} K_{(\mathfrak{D}-1)(\mathfrak{D}-1)PQ} &= \frac{\partial \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)Q}}{\partial x^P} - \frac{\partial \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P}}{\partial x^Q} + G^{\mathfrak{D}\mathfrak{D}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} \Gamma_{\mathfrak{D}(\mathfrak{D}-1)Q} - \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P} \Gamma_{(\mathfrak{D}-1)\mathfrak{D}Q}), \\ K_{\mathfrak{D}(\mathfrak{D}-1)PQ} &= \frac{\partial \Gamma_{\mathfrak{D}(\mathfrak{D}-1)Q}}{\partial x^P} - \frac{\partial \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}}{\partial x^Q} + G^{\mathfrak{D}\mathfrak{D}} (\Gamma_{\mathfrak{D}\mathfrak{D}P} \Gamma_{\mathfrak{D}(\mathfrak{D}-1)Q} - \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P} \Gamma_{\mathfrak{D}\mathfrak{D}Q}) \\ &\quad + G^{(\mathfrak{D}-1)(\mathfrak{D}-1)} (\Gamma_{\mathfrak{D}(\mathfrak{D}-1)P} \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)Q} - \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} \Gamma_{\mathfrak{D}(\mathfrak{D}-1)Q}), \\ K_{(\mathfrak{D}-1)\mathfrak{D}PQ} &= \frac{\partial \Gamma_{(\mathfrak{D}-1)\mathfrak{D}Q}}{\partial x^P} - \frac{\partial \Gamma_{(\mathfrak{D}-1)\mathfrak{D}P}}{\partial x^Q} + G^{\mathfrak{D}\mathfrak{D}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} \Gamma_{\mathfrak{D}\mathfrak{D}Q} - \Gamma_{\mathfrak{D}\mathfrak{D}P} \Gamma_{(\mathfrak{D}-1)\mathfrak{D}Q}) \\ &\quad + G^{(\mathfrak{D}-1)(\mathfrak{D}-1)} (\Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} \Gamma_{(\mathfrak{D}-1)\mathfrak{D}Q} - \Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)Q}), \\ K_{\mathfrak{D}\mathfrak{D}PQ} &= \frac{\partial \Gamma_{\mathfrak{D}\mathfrak{D}Q}}{\partial x^P} - \frac{\partial \Gamma_{\mathfrak{D}\mathfrak{D}P}}{\partial x^Q} + G^{(\mathfrak{D}-1)(\mathfrak{D}-1)} (\Gamma_{\mathfrak{D}(\mathfrak{D}-1)P} \Gamma_{(\mathfrak{D}-1)\mathfrak{D}Q} - \Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} \Gamma_{\mathfrak{D}(\mathfrak{D}-1)Q}). \end{aligned}$$

Hence,

$$\begin{aligned}
B_{PQ} &\triangleq \frac{1}{\sqrt{2}} (K_{\mathfrak{D}\mathfrak{D}PQ} + K_{(\mathfrak{D}-1)(\mathfrak{D}-1)PQ}) \\
&= \frac{1}{\sqrt{2}} \frac{\partial (\Gamma_{\mathfrak{D}\mathfrak{D}Q} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)Q})}{\partial x^P} - \frac{1}{\sqrt{2}} \frac{\partial (\Gamma_{\mathfrak{D}\mathfrak{D}P} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P})}{\partial x^Q} = \frac{\partial B_Q}{\partial x^P} - \frac{\partial B_P}{\partial x^Q}. \\
F_{PQ}^3 &\triangleq \frac{1}{\sqrt{2}} (K_{\mathfrak{D}\mathfrak{D}PQ} - K_{(\mathfrak{D}-1)(\mathfrak{D}-1)PQ}) \\
&= \frac{1}{\sqrt{2}} \left(\frac{\partial \Gamma_{\mathfrak{D}\mathfrak{D}Q}}{\partial x^P} - \frac{\partial \Gamma_{\mathfrak{D}\mathfrak{D}P}}{\partial x^Q} + G^{(\mathfrak{D}-1)(\mathfrak{D}-1)} (\Gamma_{\mathfrak{D}(\mathfrak{D}-1)P} \Gamma_{(\mathfrak{D}-1)\mathfrak{D}Q} - \Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} \Gamma_{\mathfrak{D}(\mathfrak{D}-1)Q}) \right) \\
&\quad - \frac{1}{\sqrt{2}} \left(\frac{\partial \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)Q}}{\partial x^P} - \frac{\partial \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P}}{\partial x^Q} + G^{\mathfrak{D}\mathfrak{D}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} \Gamma_{\mathfrak{D}(\mathfrak{D}-1)Q} - \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P} \Gamma_{(\mathfrak{D}-1)\mathfrak{D}Q}) \right) \\
&= \frac{\partial A_Q^3}{\partial x^P} - \frac{\partial A_P^3}{\partial x^Q} + g (\Gamma_{\mathfrak{D}(\mathfrak{D}-1)P} \Gamma_{(\mathfrak{D}-1)\mathfrak{D}Q} - \Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} \Gamma_{\mathfrak{D}(\mathfrak{D}-1)Q}) \\
&= \frac{\partial A_Q^3}{\partial x^P} - \frac{\partial A_P^3}{\partial x^Q} + g (A_P^1 A_Q^2 - A_P^2 A_Q^1).
\end{aligned}$$

Then, F_{PQ}^1 and F_{PQ}^2 can also be computed similarly. $\square$

Remark 5.1. Comparing the above conclusion and $U(1) \times SU(2)$ principal bundle theory, we know this proposition shows that the reference-system $\mathcal{F}$ indeed can describe weak and electromagnetic field.

The following proposition shows an advantage of affine connection representation, that is, affine connection representation spontaneously implies the chiral asymmetry of weak interaction, but $U(1) \times SU(2)$ principal bundle connection representation cannot imply it spontaneously.

Definition 5.2. According to Definition 3.5.1, let the charges of the above reference-system $\mathcal{F}$ be ρ_{mn} , where $m, n \in \{\mathfrak{D}-1, \mathfrak{D}\} = \{4, 5\}$. Then, $l \triangleq (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)}, \rho_{\mathfrak{D}\mathfrak{D}})^T$ is said to be an **electric charged lepton**, $\nu \triangleq (\rho_{\mathfrak{D}(\mathfrak{D}-1)}, \rho_{(\mathfrak{D}-1)\mathfrak{D}})^T$ is said to be a **neutrino**. l and ν are collectively denoted by L . Thus, $\frac{1}{\sqrt{2}}(1, 1)L$ is said to be a **left-handed lepton**, $\frac{1}{\sqrt{2}}(1, -1)L$ is said to be a **right-handed lepton**, denoted by

$$\begin{cases} l_L \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} + \rho_{\mathfrak{D}\mathfrak{D}}), & \begin{cases} \nu_L \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)\mathfrak{D}}), \\ l_R \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} - \rho_{\mathfrak{D}\mathfrak{D}}), & \begin{cases} \nu_R \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-1)} - \rho_{(\mathfrak{D}-1)\mathfrak{D}}). \end{cases} \end{cases} \end{cases} \quad (67)$$

Denote $(\Gamma_G)_{MNP}$ by Γ_{MNP} concisely. Then, we define on $(M, \mathcal{G})$ that

$$\begin{cases} Z_P \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} + \Gamma_{\mathfrak{D}\mathfrak{D}P}), & \begin{cases} W_P^1 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} + \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}), \\ A_P \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} - \Gamma_{\mathfrak{D}\mathfrak{D}P}), & \begin{cases} W_P^2 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} - \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}), \end{cases} \end{cases} \end{cases} \quad (68)$$

and say A_P is **(affine) electromagnetic potential**, while Z_P , W_P^1 and W_P^2 are **(affine) weak gauge potentials**. Then denote

$$\begin{cases} W_P^+ \triangleq \frac{1}{\sqrt{2}} (W_P^1 - iW_P^2), & \begin{cases} l_L^+ = \frac{1}{\sqrt{2}} (l_L - il_R), & \begin{cases} l_R^+ \triangleq \frac{1}{\sqrt{2}} (l_R - il_L), \\ W_P^- \triangleq \frac{1}{\sqrt{2}} (W_P^1 + iW_P^2), & \begin{cases} l_L^- = \frac{1}{\sqrt{2}} (l_L + il_R), & \begin{cases} l_R^- \triangleq \frac{1}{\sqrt{2}} (l_R + il_L). \end{cases} \end{cases} \end{cases} \end{cases}$$

Proposition 5.2. If $(M, \mathcal{G})$ satisfies the symmetry condition $\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} = \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}$, then the geometric properties l and ν of $\mathcal{F}$ satisfy the following conclusions on $(M, \mathcal{G})$,

$$\begin{cases} l_{L;P} = \partial_P l_L - g l_L Z_P - g l_R A_P - g \nu_L W_P^1, \\ l_{R;P} = \partial_P l_R - g l_R Z_P - g l_L A_P, \\ \nu_{L;P} = \partial_P \nu_L - g \nu_L Z_P - g l_L W_P^1, \\ \nu_{R;P} = \partial_P \nu_R - g \nu_R Z_P. \end{cases} \quad (69)$$

Proof. Due to $\rho_{mn;P} = \partial_P \rho_{mn} - \rho_{Hn} \Gamma_{mP}^H - \rho_{mH} \Gamma_{nP}^H = \partial_P \rho_{mn} - \rho_{hn} \Gamma_{mP}^h - \rho_{mh} \Gamma_{nP}^h$, we have

$$\begin{aligned} & \begin{cases} \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1);P} = \partial_P \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} - 2\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} \Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-1} - (\rho_{\mathfrak{D}(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)\mathfrak{D}}) \Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}}, \\ \rho_{\mathfrak{D}\mathfrak{D};P} = \partial_P \rho_{\mathfrak{D}\mathfrak{D}} - 2\rho_{\mathfrak{D}\mathfrak{D}} \Gamma_{\mathfrak{D}P}^{\mathfrak{D}} - (\rho_{\mathfrak{D}(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)\mathfrak{D}}) \Gamma_{\mathfrak{D}P}^{\mathfrak{D}-1}, \\ \rho_{\mathfrak{D}(\mathfrak{D}-1);P} = \partial_P \rho_{\mathfrak{D}(\mathfrak{D}-1)} - (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} \Gamma_{\mathfrak{D}P}^{\mathfrak{D}-1} + \rho_{\mathfrak{D}\mathfrak{D}} \Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}}) - \rho_{\mathfrak{D}(\mathfrak{D}-1)} (\Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-1} + \Gamma_{\mathfrak{D}P}^{\mathfrak{D}}), \\ \rho_{(\mathfrak{D}-1)\mathfrak{D};P} = \partial_P \rho_{(\mathfrak{D}-1)\mathfrak{D}} - (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} \Gamma_{\mathfrak{D}P}^{\mathfrak{D}-1} + \rho_{\mathfrak{D}\mathfrak{D}} \Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}}) - \rho_{(\mathfrak{D}-1)\mathfrak{D}} (\Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-1} + \Gamma_{\mathfrak{D}P}^{\mathfrak{D}}). \end{cases} \\ \Rightarrow & \begin{cases} l_{L;P} = \partial_P l_L - \sqrt{2} \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} \Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-1} - \sqrt{2} \rho_{\mathfrak{D}\mathfrak{D}} \Gamma_{\mathfrak{D}P}^{\mathfrak{D}} - g\nu_L W_P^1, \\ l_{R;P} = \partial_P l_R - \sqrt{2} \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} \Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-1} + \sqrt{2} \rho_{\mathfrak{D}\mathfrak{D}} \Gamma_{\mathfrak{D}P}^{\mathfrak{D}}, \\ \nu_{L;P} = \partial_P \nu_L - g l_L W_P^1 - \nu_L (\Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-1} + \Gamma_{\mathfrak{D}P}^{\mathfrak{D}}), \\ \nu_{R;P} = \partial_P \nu_R - \nu_R (\Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-1} + \Gamma_{\mathfrak{D}P}^{\mathfrak{D}}). \end{cases} \\ \Rightarrow & \begin{cases} l_{L;P} = \partial_P l_L - g l_L Z_P - g l_R A_P - g\nu_L W_P^1, \\ l_{R;P} = \partial_P l_R - g l_R Z_P - g l_L A_P, \\ \nu_{L;P} = \partial_P \nu_L - g\nu_L Z_P - g l_L W_P^1, \\ \nu_{R;P} = \partial_P \nu_R - g\nu_R Z_P. \end{cases} \end{aligned}$$

□

Remark 5.2. The above proposition shows that the chiral asymmetry of leptons is a direct mathematical result of affine connection representation, and that it is not necessary to be postulated artificially like that in $U_Y(1) \times SU_L(2)$ theory.

Remark 5.3. Let the totality of all geometric manifolds be $\mathcal{M}$. Suppose there is a geometric property P , and the value of P is dependent on the values of B_M^A and C_A^M . Thus, P is an invariant under the identical transformation $e : B_M^A \mapsto B_M^A$. The corresponding transformation group is the trivial group $\{e\}$, which is the smallest transformation group, with the fewest symmetry. Therefore, $\mathcal{M}/\{e\}$ is the largest geometry, with the richest geometric properties. From Proposition 5.2 we notice that:

(1) Principal bundle connection representation starts from a very large group, and reduces symmetries in way of "symmetry breaking" to approach the target geometry.

(2) Affine connection representation starts from the smallest group $\{e\}$, and adds symmetries in way of "symmetry condition" to approach the target geometry.

Such two ways must lead to the same destination. They both go towards the same specific geometry.

Remark 5.4. Proposition 5.2 shows that:

(1) In affine connection representation of gauge fields, the coupling constant g is possessed of a geometric meaning, that is in fact the metric of internal space. But it does not have such a clear geometric meaning in $U(1) \times SU(2)$ principal bundle connection representation.

(2) At the most fundamental level, the coupling constant of Z_P and that of A_P are equal, i.e.

$$g_Z = g_A = g.$$

Suppose there is a kind of medium. Z boson and photon move in it. Suppose Z field has interaction with the medium, but electromagnetic field A has no interaction with the medium. Then we have coupling constants

$$\tilde{g}_Z \neq g_A = g$$

in the medium, and the Weinberg angle appears.

It is quite reasonable to consider a Higgs boson as a zero-spin pair of neutrinos, because in Standard Model, Higgs boson only couples with Z field and W field, but does not couple with electromagnetic field and gluon field. If so, Higgs boson would lose its fundamentality and it would not have enough importance in a theory at the most fundamental level.

(3) The mixing of leptons of three generations does not appear in Proposition 5.2, but it can spontaneously appear in Proposition 7.1 due to the affine connection representation of the gauge field that is given by Definition 7.1.

6 Affine connection representation of the gauge field of strong interaction

Definition 6.1. Suppose $(M, \mathcal{F})$ and $(M, \mathcal{G})$ conform to Definition 4.5.1. Let $\mathfrak{D} = r + 3 = 6$ and both of $\mathcal{F}$ and $\mathcal{G}$ satisfy

$$G^{(\mathfrak{D}-2)(\mathfrak{D}-2)} = G^{(\mathfrak{D}-1)(\mathfrak{D}-1)} = G^{\mathfrak{D}\mathfrak{D}}.$$

Thus, $\mathcal{F}$ and $\mathcal{G}$ can describe strong interaction.

Definition 6.2. According to Definition 3.5.1, let the charges of $\mathcal{F}$ be ρ_{mn} , where $m, n = 4, 5, \dots, \mathfrak{D}$. Define

$$\begin{cases} d_1 \triangleq (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)}, \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)})^T, \\ d_2 \triangleq (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)}, \rho_{\mathfrak{D}\mathfrak{D}})^T, \\ d_3 \triangleq (\rho_{\mathfrak{D}\mathfrak{D}}, \rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)})^T, \end{cases} \quad \begin{cases} u_1 \triangleq (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-1)}, \rho_{(\mathfrak{D}-1)(\mathfrak{D}-2)})^T, \\ u_2 \triangleq (\rho_{(\mathfrak{D}-1)\mathfrak{D}}, \rho_{\mathfrak{D}(\mathfrak{D}-1)})^T, \\ u_3 \triangleq (\rho_{\mathfrak{D}(\mathfrak{D}-2)}, \rho_{(\mathfrak{D}-2)\mathfrak{D}})^T. \end{cases}$$

We say d_1 and u_1 are **red color charges**, d_2 and u_2 are **blue color charges**, d_3 and u_3 are **green color charges**. Then d_1, d_2, d_3 are said to be **down-type color charges**, and u_1, u_2, u_3 are said to be **up-type color charges**. Their left-handed and right-handed charges are

$$\begin{cases} d_{1L} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)}), \\ d_{2L} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} + \rho_{\mathfrak{D}\mathfrak{D}}), \\ d_{3L} \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}\mathfrak{D}} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)}), \end{cases} \quad \begin{cases} d_{1R} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)} - \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)}), \\ d_{2R} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} - \rho_{\mathfrak{D}\mathfrak{D}}), \\ d_{3R} \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}\mathfrak{D}} - \rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)}), \end{cases}$$

$$\begin{cases} u_{1L} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-2)}), \\ u_{2L} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-1)}), \\ u_{3L} \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)\mathfrak{D}}), \end{cases} \quad \begin{cases} u_{1R} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-1)} - \rho_{(\mathfrak{D}-1)(\mathfrak{D}-2)}), \\ u_{2R} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)\mathfrak{D}} - \rho_{\mathfrak{D}(\mathfrak{D}-1)}), \\ u_{3R} \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-2)} - \rho_{(\mathfrak{D}-2)\mathfrak{D}}). \end{cases}$$

On $(M, \mathcal{G})$ we denote

$$\begin{aligned} g_s &\triangleq \sqrt{(G^{(\mathfrak{D}-1)(\mathfrak{D}-1)})^2 + (G^{\mathfrak{D}\mathfrak{D}})^2} = \sqrt{(G^{(\mathfrak{D}-1)(\mathfrak{D}-1)})^2 + (G^{(\mathfrak{D}-2)(\mathfrak{D}-2)})^2} \\ &= \sqrt{(G^{(\mathfrak{D}-2)(\mathfrak{D}-2)})^2 + (G^{\mathfrak{D}\mathfrak{D}})^2}. \end{aligned}$$

$$\begin{cases} U_P^1 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P}), \\ V_P^1 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P} - \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P}), \end{cases} \quad \begin{cases} X_P^{23} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-1)P} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-2)P}), \\ Y_P^{23} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-1)P} - \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-2)P}), \end{cases}$$

$$\begin{cases} U_P^2 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} + \Gamma_{\mathfrak{D}\mathfrak{D}P}), \\ V_P^2 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} - \Gamma_{\mathfrak{D}\mathfrak{D}P}), \end{cases} \quad \begin{cases} X_P^{31} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} + \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}), \\ Y_P^{31} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} - \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}), \end{cases}$$

$$\begin{cases} U_P^3 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}\mathfrak{D}P} + \Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P}), \\ V_P^3 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}\mathfrak{D}P} - \Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P}), \end{cases} \quad \begin{cases} X_P^{12} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}(\mathfrak{D}-2)P} + \Gamma_{(\mathfrak{D}-2)\mathfrak{D}P}), \\ Y_P^{12} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}(\mathfrak{D}-2)P} - \Gamma_{(\mathfrak{D}-2)\mathfrak{D}P}). \end{cases}$$

We notice that there are just only three independent ones in $U_P^1, U_P^2, U_P^3, V_P^1, V_P^2$ and V_P^3 . Without loss of generality, let

$$\begin{cases} R_P \triangleq a_R U_P^1 + b_R U_P^2 + c_R U_P^3, \\ S_P \triangleq a_S U_P^1 + b_S U_P^2 + c_S U_P^3, \\ T_P \triangleq a_T U_P^1 + b_T U_P^2 + c_T U_P^3, \end{cases} \quad \begin{cases} U_P^1 \triangleq \alpha_R R_P + \alpha_S S_P + \alpha_T T_P, \\ U_P^2 \triangleq \beta_R R_P + \beta_S S_P + \beta_T T_P, \\ U_P^3 \triangleq \gamma_R R_P + \gamma_S S_P + \gamma_T T_P, \end{cases}$$

where the coefficients matrix is non-singular. Thus, it is not hard to find the following proposition true.

Proposition 6.1. Let λ_a ($a = 1, 2, \dots, 8$) be the Gell-Mann matrices, and $T_a \triangleq \frac{1}{2}\lambda_a$ the generators of $SU(3)$ group. When $(M, \mathcal{G})$ satisfies the symmetry condition $\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} + \Gamma_{\mathfrak{D}\mathfrak{D}P} = 0$, denote

$$\begin{aligned} A_P^{11} &\triangleq S_P + \frac{1}{\sqrt{6}}T_P, & A_P^{12} &\triangleq X_P^{12} - iY_P^{12}, & A_P^{13} &\triangleq X_P^{31} - iY_P^{31}, \\ A_P &\triangleq \frac{1}{2} \begin{pmatrix} A_P^{11} & A_P^{12} & A_P^{13} \\ A_P^{21} & A_P^{22} & A_P^{23} \\ A_P^{31} & A_P^{32} & A_P^{33} \end{pmatrix}, & A_P^{21} &\triangleq X_P^{12} + iY_P^{12}, & A_P^{22} &\triangleq -S_P + \frac{1}{\sqrt{6}}T_P, & A_P^{23} &\triangleq X_P^{23} - iY_P^{23}, \\ & & A_P^{31} &\triangleq X_P^{31} + iY_P^{31}, & A_P^{32} &\triangleq X_P^{23} + iY_P^{23}, & A_P^{33} &\triangleq -\frac{2}{\sqrt{6}}T_P. \end{aligned}$$

Thus, $A_P = T_a A_P^a$ if and only if $A_P^1 = X_P^{12}$, $A_P^2 = Y_P^{12}$, $A_P^3 = S_P$, $A_P^4 = X_P^{31}$, $A_P^5 = Y_P^{31}$, $A_P^6 = X_P^{23}$, $A_P^7 = Y_P^{23}$ and $A_P^8 = T_P$.

Proof. We just need to substitute the Gell-Mann matrices

$$\begin{aligned} \lambda_1 &\triangleq \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \lambda_2 &\triangleq \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \lambda_3 &\triangleq \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \lambda_4 &\triangleq \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\ \lambda_5 &\triangleq \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, & \lambda_6 &\triangleq \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, & \lambda_7 &\triangleq \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, & \lambda_8 &\triangleq \frac{1}{\sqrt{6}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}. \end{aligned}$$

into $A_P = T_a A_P^a$ and directly verify them. $\square$

Remark 6.1. On one hand, the above proposition shows that Definition 6.1 is an affine connection representation of strong interaction field. It does not define the gauge potentials abstractly like QCD based on $SU(3)$ theory, but gives concrete geometric constructions to the gauge potentials.

On the other hand, the above proposition implies that if we take appropriate symmetry conditions, the algebraic properties of $SU(3)$ group can be described by the transformation group $GL(3, \mathbb{R})$ of internal space of $\mathcal{G}$. In other words, the exponential map $\exp : GL(3, \mathbb{R}) \rightarrow U(3)$, $[B_m^a] \mapsto e^{iT_m^a B_m^a}$ defines a covering homomorphism, and $SU(3)$ is a subgroup of $U(3)$.

7 Affine connection representation of the unified gauge field

Definition 7.1. Suppose $(M, \mathcal{F})$ and $(M, \mathcal{G})$ conform to Definition 4.5.1. Let $\mathfrak{D} = r + 5 = 8$ and both of $\mathcal{F}$ and $\mathcal{G}$ satisfy

$$G^{(\mathfrak{D}-4)(\mathfrak{D}-4)} = G^{(\mathfrak{D}-3)(\mathfrak{D}-3)}, \quad G^{(\mathfrak{D}-2)(\mathfrak{D}-2)} = G^{(\mathfrak{D}-1)(\mathfrak{D}-1)} = G^{\mathfrak{D}\mathfrak{D}}.$$

Thus, $\mathcal{F}$ and $\mathcal{G}$ can describe the unified field of electromagnetic, weak and strong interactions.

Definition 7.2. According to Definition 3.5.1, let the charges of $\mathcal{F}$ be ρ_{mn} , where $m, n = 4, 5, \dots, \mathfrak{D}$. Define

$$\begin{cases} l \triangleq (\rho_{(\mathfrak{D}-4)(\mathfrak{D}-4)}, \rho_{(\mathfrak{D}-3)(\mathfrak{D}-3)})^T, \\ d_1 \triangleq (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)}, \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)})^T, \\ d_2 \triangleq (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)}, \rho_{\mathfrak{D}\mathfrak{D}})^T, \\ d_3 \triangleq (\rho_{\mathfrak{D}\mathfrak{D}}, \rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)})^T, \end{cases} \quad \begin{cases} \nu \triangleq (\rho_{(\mathfrak{D}-3)(\mathfrak{D}-4)}, \rho_{(\mathfrak{D}-4)(\mathfrak{D}-3)})^T, \\ u_1 \triangleq (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-1)}, \rho_{(\mathfrak{D}-1)(\mathfrak{D}-2)})^T, \\ u_2 \triangleq (\rho_{(\mathfrak{D}-1)\mathfrak{D}}, \rho_{\mathfrak{D}(\mathfrak{D}-1)})^T, \\ u_3 \triangleq (\rho_{\mathfrak{D}(\mathfrak{D}-2)}, \rho_{(\mathfrak{D}-2)\mathfrak{D}})^T. \end{cases}$$

And Denote

$$\begin{cases} l_L \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-4)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-3)}), \\ l_R \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-4)(\mathfrak{D}-4)} - \rho_{(\mathfrak{D}-3)(\mathfrak{D}-3)}), \end{cases} \quad \begin{cases} \nu_L \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-3)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-3)}), \\ \nu_R \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-3)(\mathfrak{D}-4)} - \rho_{(\mathfrak{D}-4)(\mathfrak{D}-3)}), \end{cases}$$

$$\begin{cases} d_{1L} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)}), \\ d_{2L} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} + \rho_{\mathfrak{D}\mathfrak{D}}), \\ d_{3L} \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}\mathfrak{D}} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)}), \end{cases} \quad \begin{cases} d_{1R} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)} - \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)}), \\ d_{2R} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} - \rho_{\mathfrak{D}\mathfrak{D}}), \\ d_{3R} \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}\mathfrak{D}} - \rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)}), \end{cases}$$

$$\begin{cases} u_{1L} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-2)}), \\ u_{2L} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-1)}), \\ u_{3L} \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)\mathfrak{D}}), \end{cases} \quad \begin{cases} u_{1R} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-1)} - \rho_{(\mathfrak{D}-1)(\mathfrak{D}-2)}), \\ u_{2R} \triangleq \frac{1}{\sqrt{2}} (\rho_{(\mathfrak{D}-1)\mathfrak{D}} - \rho_{\mathfrak{D}(\mathfrak{D}-1)}), \\ u_{3R} \triangleq \frac{1}{\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-2)} - \rho_{(\mathfrak{D}-2)\mathfrak{D}}). \end{cases}$$

On $(M, \mathcal{G})$ we denote

$$\begin{cases} g \triangleq \sqrt{(G^{(\mathfrak{D}-4)(\mathfrak{D}-4)})^2 + (G^{(\mathfrak{D}-3)(\mathfrak{D}-3)})^2}, \\ g_s \triangleq \sqrt{(G^{(\mathfrak{D}-1)(\mathfrak{D}-1)})^2 + (G^{\mathfrak{D}\mathfrak{D}})^2} = \sqrt{(G^{(\mathfrak{D}-1)(\mathfrak{D}-1)})^2 + (G^{(\mathfrak{D}-2)(\mathfrak{D}-2)})^2} \\ \quad = \sqrt{(G^{(\mathfrak{D}-2)(\mathfrak{D}-2)})^2 + (G^{\mathfrak{D}\mathfrak{D}})^2}, \end{cases}$$

$$\begin{cases} Z_P \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-4)(\mathfrak{D}-4)P} + \Gamma_{(\mathfrak{D}-3)(\mathfrak{D}-3)P}), \\ A_P \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-4)(\mathfrak{D}-4)P} - \Gamma_{(\mathfrak{D}-3)(\mathfrak{D}-3)P}), \end{cases} \quad \begin{cases} W_P^1 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-4)(\mathfrak{D}-3)P} + \Gamma_{(\mathfrak{D}-3)(\mathfrak{D}-4)P}), \\ W_P^2 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-4)(\mathfrak{D}-3)P} - \Gamma_{(\mathfrak{D}-3)(\mathfrak{D}-4)P}), \end{cases}$$

$$\begin{cases} U_P^1 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P}), \\ V_P^1 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P} - \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P}), \end{cases} \quad \begin{cases} X_P^{23} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-1)P} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-2)P}), \\ Y_P^{23} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-1)P} - \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-2)P}), \end{cases}$$

$$\begin{cases} U_P^2 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} + \Gamma_{\mathfrak{D}\mathfrak{D}P}), \\ V_P^2 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} - \Gamma_{\mathfrak{D}\mathfrak{D}P}), \end{cases} \quad \begin{cases} X_P^{31} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} + \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}), \\ Y_P^{31} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} - \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P}), \end{cases}$$

$$\begin{cases} U_P^3 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}\mathfrak{D}P} + \Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P}), \\ V_P^3 \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}\mathfrak{D}P} - \Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P}), \end{cases} \quad \begin{cases} X_P^{12} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}(\mathfrak{D}-2)P} + \Gamma_{(\mathfrak{D}-2)\mathfrak{D}P}), \\ Y_P^{12} \triangleq \frac{1}{\sqrt{2}} (\Gamma_{\mathfrak{D}(\mathfrak{D}-2)P} - \Gamma_{(\mathfrak{D}-2)\mathfrak{D}P}). \end{cases}$$

Definition 7.3. Define the symmetry condition of unification:

(1) Basic conditions, No.1:

$$\begin{cases} G^{(\mathfrak{D}-4)(\mathfrak{D}-4)} = G^{(\mathfrak{D}-3)(\mathfrak{D}-3)}, \\ G^{(\mathfrak{D}-2)(\mathfrak{D}-2)} = G^{(\mathfrak{D}-1)(\mathfrak{D}-1)} = G^{\mathfrak{D}\mathfrak{D}}, \end{cases}$$

(2) Basic conditions, No.2:

$$\begin{cases} \Gamma_{(\mathfrak{D}-3)(\mathfrak{D}-4)P} = \Gamma_{(\mathfrak{D}-4)(\mathfrak{D}-3)P}, \\ \Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P} + \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} + \Gamma_{\mathfrak{D}\mathfrak{D}P} = 0, \end{cases}$$

(3) MNS mixing conditions of weak interaction, No.1:

$$\begin{cases} \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}-2} = c_{\mathfrak{D}-3}^{\mathfrak{D}-2} \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}-3}, \\ \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}-1} = c_{\mathfrak{D}-3}^{\mathfrak{D}-1} \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}-3}, \\ \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}} = c_{\mathfrak{D}-3}^{\mathfrak{D}} \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}-3}, \end{cases} \quad \begin{cases} \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}-2} = c_{\mathfrak{D}-4}^{\mathfrak{D}-2} \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}-4}, \\ \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}-1} = c_{\mathfrak{D}-4}^{\mathfrak{D}-1} \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}-4}, \\ \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}} = c_{\mathfrak{D}-4}^{\mathfrak{D}} \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}-4}, \end{cases} \quad \begin{cases} c_{\mathfrak{D}-3}^{\mathfrak{D}-2} = c_{\mathfrak{D}-4}^{\mathfrak{D}-2}, \\ c_{\mathfrak{D}-3}^{\mathfrak{D}-1} = c_{\mathfrak{D}-4}^{\mathfrak{D}-1}, \\ c_{\mathfrak{D}-3}^{\mathfrak{D}} = c_{\mathfrak{D}-4}^{\mathfrak{D}}, \end{cases}$$

(4) MNS mixing conditions of weak interaction, No.2:

$$\begin{cases} \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} = \rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)}, \\ \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} = \rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)}, \\ \rho_{\mathfrak{D}(\mathfrak{D}-3)} = \rho_{\mathfrak{D}(\mathfrak{D}-4)}, \end{cases} \quad \begin{cases} \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} = \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)}, \\ \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} = \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)}, \\ \rho_{(\mathfrak{D}-3)\mathfrak{D}} = \rho_{(\mathfrak{D}-4)\mathfrak{D}}, \end{cases}$$

(5) CKM mixing conditions of strong interaction, No.1:

$$\begin{cases} \Gamma_{(\mathfrak{D}-2)P}^{\mathfrak{D}-3} = c_{\mathfrak{D}-2}^{\mathfrak{D}-4} \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}-3}, & \Gamma_{(\mathfrak{D}-2)P}^{\mathfrak{D}-4} = c_{\mathfrak{D}-2}^{\mathfrak{D}-3} \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}-4}, \\ \Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-3} = c_{\mathfrak{D}-1}^{\mathfrak{D}-4} \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}-3}, & \Gamma_{(\mathfrak{D}-1)P}^{\mathfrak{D}-4} = c_{\mathfrak{D}-1}^{\mathfrak{D}-3} \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}-4}, \\ \Gamma_{\mathfrak{D}P}^{\mathfrak{D}-3} = c_{\mathfrak{D}}^{\mathfrak{D}-4} \Gamma_{(\mathfrak{D}-4)P}^{\mathfrak{D}-3}, & \Gamma_{\mathfrak{D}P}^{\mathfrak{D}-4} = c_{\mathfrak{D}}^{\mathfrak{D}-3} \Gamma_{(\mathfrak{D}-3)P}^{\mathfrak{D}-4}, \end{cases} \quad \begin{cases} c_{\mathfrak{D}-2}^{\mathfrak{D}-4} = c_{\mathfrak{D}-1}^{\mathfrak{D}-4} = c_{\mathfrak{D}}^{\mathfrak{D}-4}, \\ c_{\mathfrak{D}-2}^{\mathfrak{D}-3} = c_{\mathfrak{D}-1}^{\mathfrak{D}-3} = c_{\mathfrak{D}}^{\mathfrak{D}-3}, \end{cases}$$

(6) CKM mixing conditions of strong interaction, No.2:

$$\begin{cases} \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} = \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} = \rho_{\mathfrak{D}(\mathfrak{D}-3)}, & \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} = \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} = \rho_{(\mathfrak{D}-3)\mathfrak{D}}, \\ \rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} = \rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} = \rho_{\mathfrak{D}(\mathfrak{D}-4)}, & \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)} = \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)} = \rho_{(\mathfrak{D}-4)\mathfrak{D}}, \end{cases}$$

where c_n^m are constants.

Proposition 7.1. Denote

$$\begin{aligned} l' &\triangleq \left(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-4)} + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}-2}}{2} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)}) \right. \\ &\quad + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}-1}}{2} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)}) + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}}}{2} (\rho_{\mathfrak{D}(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)\mathfrak{D}}), \\ &\quad \rho_{(\mathfrak{D}-3)(\mathfrak{D}-3)} + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}-2}}{2} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)}) \\ &\quad \left. + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}-1}}{2} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)}) + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}}}{2} (\rho_{\mathfrak{D}(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)\mathfrak{D}}) \right)^T, \\ \nu' &\triangleq \left(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-4)} + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}-2}}{2} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)}) \right. \\ &\quad + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}-1}}{2} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)}) + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}}}{2} (\rho_{\mathfrak{D}(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)\mathfrak{D}}), \\ &\quad \rho_{(\mathfrak{D}-4)(\mathfrak{D}-3)} + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}-2}}{2} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)}) \\ &\quad \left. + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}-1}}{2} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)}) + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}}}{2} (\rho_{\mathfrak{D}(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)\mathfrak{D}}) \right)^T. \end{aligned}$$

When $(M, \mathcal{G})$ satisfies the symmetry conditions of Definition 7.3, the geometric properties l and ν of $\mathcal{F}$ satisfy the following conclusions on $(M, \mathcal{G})$.

$$\begin{cases} l_{L;P} = \partial_P l_L - gl_L Z_P - gl_R A_P - g\nu'_L W_P^1, \\ l_{R;P} = \partial_P l_R - gl_R Z_P - gl_L A_P, \\ \nu_{L;P} = \partial_P \nu_L - g\nu_L Z_P - gl'_L W_P^1, \\ \nu_{R;P} = \partial_P \nu_R - g\nu_R Z_P. \end{cases} \quad (70)$$

Proof. First, we compute the covariant differential of ρ_{mn} of $\mathcal{F}$.

$$\begin{aligned} \rho_{mn;P} &= \partial_P \rho_{mn} - \rho_{Hn} \Gamma_{mP}^H - \rho_{mH} \Gamma_{nP}^H \\ &= \partial_P \rho_{mn} - \rho_{(\mathfrak{D}-4)n} \Gamma_{mP}^{\mathfrak{D}-4} - \rho_{(\mathfrak{D}-3)n} \Gamma_{mP}^{\mathfrak{D}-3} - \rho_{(\mathfrak{D}-2)n} \Gamma_{mP}^{\mathfrak{D}-2} - \rho_{(\mathfrak{D}-1)n} \Gamma_{mP}^{\mathfrak{D}-1} - \rho_{\mathfrak{D}n} \Gamma_{mP}^{\mathfrak{D}} \\ &\quad - \rho_{m(\mathfrak{D}-4)} \Gamma_{nP}^{\mathfrak{D}-4} - \rho_{m(\mathfrak{D}-3)} \Gamma_{nP}^{\mathfrak{D}-3} - \rho_{m(\mathfrak{D}-2)} \Gamma_{nP}^{\mathfrak{D}-2} - \rho_{m(\mathfrak{D}-1)} \Gamma_{nP}^{\mathfrak{D}-1} - \rho_{m\mathfrak{D}} \Gamma_{nP}^{\mathfrak{D}}. \end{aligned}$$

According to Definition 7.2 and Definition 7.3, by calculation we obtain that

$$\begin{aligned} l_{L;P} &= \partial_P l_L - gl_L Z_P - gl_R A_P - g\nu_L W_P^1 \\ &\quad - \frac{1}{2} \left[c_{\mathfrak{D}-4}^{\mathfrak{D}-2} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)}) + c_{\mathfrak{D}-3}^{\mathfrak{D}-2} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)}) \right] \frac{g}{\sqrt{2}} W_P^1 \\ &\quad - \frac{1}{2} \left[c_{\mathfrak{D}-4}^{\mathfrak{D}-1} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)}) + c_{\mathfrak{D}-3}^{\mathfrak{D}-1} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)}) \right] \frac{g}{\sqrt{2}} W_P^1 \\ &\quad - \frac{1}{2} \left[c_{\mathfrak{D}-4}^{\mathfrak{D}} (\rho_{\mathfrak{D}(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)\mathfrak{D}}) + c_{\mathfrak{D}-3}^{\mathfrak{D}} (\rho_{\mathfrak{D}(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)\mathfrak{D}}) \right] \frac{g}{\sqrt{2}} W_P^1, \\ l_{R;P} &= \partial_P l_R - gl_R Z_P - gl_L A_P, \end{aligned}$$

$$\begin{aligned}
\nu_{L;P} &= \partial_P \nu_L - g \nu_L Z_P - g l_L W_P^1 \\
&\quad - \frac{1}{2} \left[c_{\mathfrak{D}-4}^{\mathfrak{D}-2} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)}) + c_{\mathfrak{D}-3}^{\mathfrak{D}-2} (\rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)}) \right] \frac{g}{\sqrt{2}} W_P^1 \\
&\quad - \frac{1}{2} \left[c_{\mathfrak{D}-4}^{\mathfrak{D}-1} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)}) + c_{\mathfrak{D}-3}^{\mathfrak{D}-1} (\rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)}) \right] \frac{g}{\sqrt{2}} W_P^1 \\
&\quad - \frac{1}{2} \left[c_{\mathfrak{D}-4}^{\mathfrak{D}} (\rho_{\mathfrak{D}(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)\mathfrak{D}}) + c_{\mathfrak{D}-3}^{\mathfrak{D}} (\rho_{(\mathfrak{D}-3)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-3)}) \right] \frac{g}{\sqrt{2}} W_P^1, \\
\nu_{R;P} &= \partial_P \nu_R - g \nu_R Z_P.
\end{aligned}$$

Then, according to definitions of l' and ν' , we obtain that

$$\begin{aligned}
l'_L &= l_L \\
&\quad + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}-2}}{2\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)}) + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}-1}}{2\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)}) + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}}}{2\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)\mathfrak{D}}) \\
&\quad + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}-2}}{2\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)}) + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}-1}}{2\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)}) + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}}}{2\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)\mathfrak{D}}), \\
\nu'_L &= \nu_L \\
&\quad + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}-2}}{2\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)}) + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}-1}}{2\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)}) + \frac{c_{\mathfrak{D}-3}^{\mathfrak{D}}}{2\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-4)} + \rho_{(\mathfrak{D}-4)\mathfrak{D}}) \\
&\quad + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}-2}}{2\sqrt{2}} (\rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)}) + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}-1}}{2\sqrt{2}} (\rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)}) + \frac{c_{\mathfrak{D}-4}^{\mathfrak{D}}}{2\sqrt{2}} (\rho_{\mathfrak{D}(\mathfrak{D}-3)} + \rho_{(\mathfrak{D}-3)\mathfrak{D}}).
\end{aligned}$$

Substitute them into the previous equations, and we obtain that

$$\begin{cases} l_{L;P} = \partial_P l_L - g l_L Z_P - g l_R A_P - g \nu'_L W_P^1, \\ l_{R;P} = \partial_P l_R - g l_R Z_P - g l_L A_P, \end{cases}, \quad \begin{cases} \nu_{L;P} = \partial_P \nu_L - g \nu_L Z_P - g l'_L W_P^1, \\ \nu_{R;P} = \partial_P \nu_R - g \nu_R Z_P. \end{cases}$$

□

Remark 7.1. Reviewing Remark 5.3, we know the above proposition shows the geometric origin of MNS mixing of weak interaction. In affine connection representation of gauge fields, MNS mixing spontaneously appears as a geometric property, therefore it is not necessary to be postulated artificially like that in the Standard Model.

In conventional physics, e , μ and τ have just only ontological differences, but they have no difference in mathematical connotation. By contrast, Proposition 7.1 tells us that leptons of three generations should be constructed by different linear combinations of $\{\rho_{pq}, \rho_{qp}\}_{p=4,5; q=6,7,8}$. Thus, e , μ and τ may have concrete and distinguishable mathematical connotations. For example, suppose $a_\mu, b_\mu, a_\mu^m, b_\mu^m, a_\tau, b_\tau, a_\tau^m, b_\tau^m$ are constants, then we can imagine that

$$\begin{cases} e \triangleq l = (\rho_{(\mathfrak{D}-4)(\mathfrak{D}-4)}, \rho_{(\mathfrak{D}-3)(\mathfrak{D}-3)})^T, \\ \nu_e \triangleq \nu = (\rho_{(\mathfrak{D}-3)(\mathfrak{D}-4)}, \rho_{(\mathfrak{D}-4)(\mathfrak{D}-3)})^T. \\ \left\{ \begin{aligned} \mu &\triangleq a_\mu e + \frac{1}{2} \left(a_\mu^{\mathfrak{D}-2} \rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} + a_\mu^{\mathfrak{D}-1} \rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} + a_\mu^{\mathfrak{D}} \rho_{\mathfrak{D}(\mathfrak{D}-4)}, \right. \\ &\quad \left. a_\mu^{\mathfrak{D}-2} \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} + a_\mu^{\mathfrak{D}-1} \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} + a_\mu^{\mathfrak{D}} \rho_{\mathfrak{D}(\mathfrak{D}-3)} \right)^T. \\ \nu_\mu &\triangleq b_\mu \nu_e + \frac{1}{2} \left(b_\mu^{\mathfrak{D}-2} \rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)} + b_\mu^{\mathfrak{D}-1} \rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)} + b_\mu^{\mathfrak{D}} \rho_{\mathfrak{D}(\mathfrak{D}-4)}, \right. \\ &\quad \left. b_\mu^{\mathfrak{D}-2} \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)} + b_\mu^{\mathfrak{D}-1} \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)} + b_\mu^{\mathfrak{D}} \rho_{\mathfrak{D}(\mathfrak{D}-3)} \right)^T. \end{aligned} \right. \\ \left\{ \begin{aligned} \tau &\triangleq a_\tau \mu + \frac{1}{2} \left(a_\tau^{\mathfrak{D}-2} \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)} + a_\tau^{\mathfrak{D}-1} \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)} + a_\tau^{\mathfrak{D}} \rho_{(\mathfrak{D}-4)\mathfrak{D}}, \right. \\ &\quad \left. a_\tau^{\mathfrak{D}-2} \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} + a_\tau^{\mathfrak{D}-1} \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} + a_\tau^{\mathfrak{D}} \rho_{(\mathfrak{D}-3)\mathfrak{D}} \right)^T. \\ \nu_\tau &\triangleq b_\tau \nu_\mu + \frac{1}{2} \left(b_\tau^{\mathfrak{D}-2} \rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)} + b_\tau^{\mathfrak{D}-1} \rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)} + b_\tau^{\mathfrak{D}} \rho_{(\mathfrak{D}-4)\mathfrak{D}}, \right. \\ &\quad \left. b_\tau^{\mathfrak{D}-2} \rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} + b_\tau^{\mathfrak{D}-1} \rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} + b_\tau^{\mathfrak{D}} \rho_{(\mathfrak{D}-3)\mathfrak{D}} \right)^T. \end{aligned} \right. \end{cases}$$

Proposition 7.2. When $(M, \mathcal{G})$ satisfies the symmetry conditions of Definition 7.3, denote

$$\begin{aligned}
d'_{1L} &\triangleq \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-1}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-2}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)}) \\
&\quad + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-1}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-2}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)}), \\
d'_{2L} &\triangleq \frac{1}{2\sqrt{2}}c_{\mathfrak{D}}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-1}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-4)}) \\
&\quad + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-1}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-3)}), \\
d'_{3L} &\triangleq \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-2}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)}) \\
&\quad + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-2}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-3)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)}), \\
u'_{1L} &\triangleq \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-2}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-2}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)}) \\
&\quad + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-1}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-1}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)}), \\
u'_{2L} &\triangleq \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-1}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-1}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-1)} + \rho_{(\mathfrak{D}-1)(\mathfrak{D}-3)}) \\
&\quad + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-3)}), \\
u'_{3L} &\triangleq \frac{1}{2\sqrt{2}}c_{\mathfrak{D}}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)\mathfrak{D}} + \rho_{\mathfrak{D}(\mathfrak{D}-3)}) \\
&\quad + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-2}^{\mathfrak{D}-3}(\rho_{(\mathfrak{D}-4)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-4)}) + \frac{1}{2\sqrt{2}}c_{\mathfrak{D}-2}^{\mathfrak{D}-4}(\rho_{(\mathfrak{D}-3)(\mathfrak{D}-2)} + \rho_{(\mathfrak{D}-2)(\mathfrak{D}-3)}).
\end{aligned}$$

Then the geometric properties $d_1, d_2, d_3, u_1, u_2, u_3$ of $\mathcal{F}$ satisfy the following conclusions on $(M, \mathcal{G})$.

$$\begin{aligned}
d_{1L;P} &= \partial_P d_{1L} - g_s d_{1L} U_P^1 + g_s d_{2L} V_P^1 - g_s d_{3L} V_P^1 \\
&\quad - g_s u_{1L} X_P^{23} - \frac{g_s}{2} u_{2L} X_P^{31} + \frac{g_s}{2} u_{2L} Y_P^{31} - \frac{g_s}{2} u_{3L} X_P^{12} - \frac{g_s}{2} u_{3L} Y_P^{12} - g u'_{1L} W_P^1, \\
d_{2L;P} &= \partial_P d_{2L} - g_s d_{2L} U_P^2 + g_s d_{3L} V_P^2 - g_s d_{1L} V_P^2 \\
&\quad - g_s u_{2L} X_P^{31} - \frac{g_s}{2} u_{3L} X_P^{12} + \frac{g_s}{2} u_{3L} Y_P^{12} - \frac{g_s}{2} u_{1L} X_P^{23} - \frac{g_s}{2} u_{1L} Y_P^{23} - g u'_{2L} W_P^1, \\
d_{3L;P} &= \partial_P d_{3L} - g_s d_{3L} U_P^3 + g_s d_{1L} V_P^3 - g_s d_{2L} V_P^3 \\
&\quad - g_s u_{3L} X_P^{12} - \frac{g_s}{2} u_{1L} X_P^{23} + \frac{g_s}{2} u_{1L} Y_P^{23} - \frac{g_s}{2} u_{2L} X_P^{31} - \frac{g_s}{2} u_{2L} Y_P^{31} - g u'_{3L} W_P^1, \\
d_{1R;P} &= \partial_P d_{1R} - g_s d_{1L} V_P^1 + g_s d_{2L} U_P^1 - g_s d_{3L} U_P^1 \\
&\quad + g_s u_{1L} Y_P^{23} + \frac{g_s}{2} u_{2L} X_P^{31} - \frac{g_s}{2} u_{2L} Y_P^{31} - \frac{g_s}{2} u_{3L} X_P^{12} - \frac{g_s}{2} u_{3L} Y_P^{12}, \\
d_{2R;P} &= \partial_P d_{2R} - g_s d_{2L} V_P^2 + g_s d_{3L} U_P^2 - g_s d_{1L} U_P^2 \\
&\quad + g_s u_{2L} Y_P^{31} + \frac{g_s}{2} u_{3L} X_P^{12} - \frac{g_s}{2} u_{3L} Y_P^{12} - \frac{g_s}{2} u_{1L} X_P^{23} - \frac{g_s}{2} u_{1L} Y_P^{23}, \\
d_{3R;P} &= \partial_P d_{3R} - g_s d_{3L} V_P^3 + g_s d_{1L} U_P^3 - g_s d_{2L} U_P^3 \\
&\quad + g_s u_{3L} Y_P^{12} + \frac{g_s}{2} u_{1L} X_P^{23} - \frac{g_s}{2} u_{1L} Y_P^{23} - \frac{g_s}{2} u_{2L} X_P^{31} - \frac{g_s}{2} u_{2L} Y_P^{31}, \\
u_{1L;P} &= \partial_P u_{1L} - g_s u_{1L} U_P^1 - \frac{g_s}{2} u_{2L} X_P^{12} - \frac{g_s}{2} u_{2L} Y_P^{12} - \frac{g_s}{2} u_{3L} X_P^{31} + \frac{g_s}{2} u_{3L} Y_P^{31} \\
&\quad - g_s d_{1L} X_P^{23} + g_s d_{2L} Y_P^{23} - g_s d_{3L} Y_P^{23} - g d'_{1L} W_P^1, \\
u_{2L;P} &= \partial_P u_{2L} - g_s u_{2L} U_P^2 - \frac{g_s}{2} u_{3L} X_P^{23} - \frac{g_s}{2} u_{3L} Y_P^{23} - \frac{g_s}{2} u_{1L} X_P^{12} + \frac{g_s}{2} u_{1L} Y_P^{12} \\
&\quad - g_s d_{2L} X_P^{31} + g_s d_{3L} Y_P^{31} - g_s d_{1L} Y_P^{31} - g d'_{2L} W_P^1, \\
u_{3L;P} &= \partial_P u_{3L} - g_s u_{3L} U_P^3 - \frac{g_s}{2} u_{1L} X_P^{31} - \frac{g_s}{2} u_{1L} Y_P^{31} - \frac{g_s}{2} u_{2L} X_P^{23} + \frac{g_s}{2} u_{2L} Y_P^{23} \\
&\quad - g_s d_{3L} X_P^{12} + g_s d_{1L} Y_P^{12} - g_s d_{2L} Y_P^{12} - g d'_{3L} W_P^1,
\end{aligned}$$

$$\begin{aligned}
u_{1R;P} &= \partial_P u_{1R} - g_s u_{1R} U_P^1 + \frac{g_s}{2} u_{2R} X_P^{12} + \frac{g_s}{2} u_{2R} Y_P^{12} + \frac{g_s}{2} u_{3R} X_P^{31} - \frac{g_s}{2} u_{3R} Y_P^{31}, \\
u_{2R;P} &= \partial_P u_{2R} - g_s u_{2R} U_P^2 + \frac{g_s}{2} u_{3R} X_P^{23} + \frac{g_s}{2} u_{3R} Y_P^{23} + \frac{g_s}{2} u_{1R} X_P^{12} - \frac{g_s}{2} u_{1R} Y_P^{12}, \\
u_{3R;P} &= \partial_P u_{3R} - g_s u_{3R} U_P^3 + \frac{g_s}{2} u_{1R} X_P^{31} + \frac{g_s}{2} u_{1R} Y_P^{31} + \frac{g_s}{2} u_{2R} X_P^{23} - \frac{g_s}{2} u_{2R} Y_P^{23}.
\end{aligned}$$

Proof. Substitute Definition 7.2 into ρ_{mn} and consider Definition 7.3, then compute them, and then substitute $d'_{1L}, d'_{2L}, d'_{3L}, u'_{1L}, u'_{2L}, u'_{3L}$ into them, we finally obtain the results. $\square$

Remark 7.2. The above proposition shows a possible geometric origin of CKM mixing. We see that, in affine connection representation of gauge fields, $d'_{1L}, d'_{2L}, d'_{3L}, u'_{1L}, u'_{2L}, u'_{3L}$ spontaneously appear as geometric properties. There are still some problems about the detailed relation of mixing and they need to be further studied, but this may be solved in affine connection representation in prospect.

In addition, this proposition also shows that the new framework has an advantage that it never causes the problem that a proton decays into a lepton as do some GUTs.

Definition 7.4. If reference-system $\mathcal{F}$ satisfies

$$\begin{aligned}
\rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)} &= \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} = \rho_{\mathfrak{D}\mathfrak{D}} = \rho_{(\mathfrak{D}-2)(\mathfrak{D}-1)} = \rho_{(\mathfrak{D}-1)(\mathfrak{D}-2)} = \rho_{(\mathfrak{D}-1)\mathfrak{D}} = \rho_{\mathfrak{D}(\mathfrak{D}-1)} = \rho_{\mathfrak{D}(\mathfrak{D}-2)} \\
&= \rho_{(\mathfrak{D}-2)\mathfrak{D}} = 0, \\
\Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-2)P} &= \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-1)P} = \Gamma_{\mathfrak{D}\mathfrak{D}P} = \Gamma_{(\mathfrak{D}-2)(\mathfrak{D}-1)P} = \Gamma_{(\mathfrak{D}-1)(\mathfrak{D}-2)P} = \Gamma_{(\mathfrak{D}-1)\mathfrak{D}P} = \Gamma_{\mathfrak{D}(\mathfrak{D}-1)P} \\
&= \Gamma_{\mathfrak{D}(\mathfrak{D}-2)P} = \Gamma_{(\mathfrak{D}-2)\mathfrak{D}P} = 0,
\end{aligned}$$

we say $\mathcal{F}$ is a **lepton field**, otherwise $\mathcal{F}$ is a **hadron field**.

Suppose $\mathcal{F}$ is a hadron field. For $d_1, d_2, d_3, u_1, u_2, u_3$, if $\mathcal{F}$ satisfies that five of them are zero and the other one is non-zero, we say $\mathcal{F}$ is an **individual quark**.

Proposition 7.3. There does not exist an individual quark. In other words, if any five ones of $d_1, d_2, d_3, u_1, u_2, u_3$ are zero, then $d_1 = d_2 = d_3 = u_1 = u_2 = u_3 = 0$.

For an individual down-type quark, the above proposition is evidently true. Without loss of generality let $u_1 = u_2 = u_3 = 0$ and $d_1 = d_2 = 0$, thus $\rho_{(\mathfrak{D}-2)(\mathfrak{D}-2)} = \rho_{(\mathfrak{D}-1)(\mathfrak{D}-1)} = \rho_{\mathfrak{D}\mathfrak{D}} = 0$, hence we must have $d_3 = 0$.

For an individual up-type quark, this paper has not made progress on the proof yet. Nevertheless, Proposition 7.3 provides the color confinement with a new geometric interpretation, which is significant in itself. It involves a natural geometric constraint of the curvatures among different dimensions.

8 Conclusions

1. An affine connection representation of gauge fields is established in this paper. It has the following main points of view.

- (i) The holonomic connection Eq.(3) contains more geometric information than Levi-Civita connection. It can uniformly describe gauge field and gravitational field.
- (ii) Time is the total metric with respect to all dimensions of internal coordinate space and external coordinate space.
- (iii) On-shell evolution is described by gradient direction.
- (iv) Quantum theory is a geometric theory of distribution of gradient directions. It has a geometric meaning discussed in section 3.9.

2. In the affine connection representation of gauge fields, the following unified mathematical descriptions are achieved.

- (i) Gauge field and gravitational field can both be represented by affine connection. They have a unified coordinate description. Some parts of Γ_{NP}^M describe gauge fields such as electromagnetic, weak and strong interaction fields. The other parts of Γ_{NP}^M describe gravitational field.
 - (ii) Gauge field and elementary particle field have a unified geometric construction. They are both geometric entities constructed with semi-metric. The components ρ_{mn} of ρ_{MN} with $m, n \in \{4, 5, \dots, \mathfrak{D}\}$ describe leptons and quarks, the other components of ρ_{MN} may describe particle fields of dark matters.
 - (iii) Physical evolutions of gauge field and elementary particle field have a unified geometric description. Their on-shell evolution and quantum evolution both present as geometric properties about gradient direction.
 - (iv) Quantum theory and gravitational theory have a unified geometric interpretation and the same view of time and space. They both reflect intrinsic geometric properties of manifold.
3. Other advantages of the affine connection representation of gauge fields.

- (i) The geometric origination of coupling constants of interactions can be interpreted.
- (ii) Chiral asymmetry, MNS mixing and CKM mixing can appear spontaneously as geometric properties in affine connection representation, whereas in $U(1) \times SU(2) \times SU(3)$ principal bundle connection representation they can just only be artificially set up. Several postulates in the Standard Model are turned into geometric results that appear spontaneously in affine connection representation, so they are not necessary to be regarded as postulates anymore.
- (iii) It can avoid the problem that a proton decays into a lepton.
- (iv) Since the concept of point particle is thoroughly abandoned, this theory is not required to be renormalized.
- (v) There exists a possible geometric interpretation to the color confinement of quarks.

The Standard Model is not possessed of the above characteristics. In the affine connection representation, we can get a better interpretation of these physical properties. Therefore, to represent gauge fields by affine connection is probably a necessary step towards the ultimate theory of physics.

References

1. Kibble, T.W.B.: Lorentz invariance and the gravitational field. *Journal of Mathematical Physics* **2**(2), 212–221 (1961). DOI 10.1063/1.1703702
2. Sciama, D.: Recent developments in general relativity. *Festschrift for Infeld*, Pergamon, Oxford **415** (1962)
3. Harnard, J.P., Pettitt, R.B.: Gauge theories for space-time symmetries. i. *Journal of Mathematical Physics* **17**(10), 1827–1837 (1976). DOI 10.1063/1.522829
4. Heyde, P.v.d.: The field equations of the poincaré gauge theory of gravitation. *Physics Letters A* **58**(3), 141–143 (1976). DOI 10.1016/0375-9601(76)90266-8
5. Cho, Y.M.: Gauge theory, gravitation, and symmetry. *Physical Review D* **14**:12(12), 3341–3344 (1976). DOI 10.1103/PhysRevD.14.3341
6. Hehl, F.W., Kerlick, G.D., Nester, J.M.: General relativity with spin and torsion: Foundations and prospects. *Reviews of Modern Physics* **48**(3), 393–416 (1976). DOI 10.1103/RevModPhys.48.393
7. MacDowell, S.W., Mansouri, F.: Unified geometric theory of gravity and supergravity. *Physical Review Letters* **38**(23), 739–742 (1977). DOI 10.1103/PhysRevLett.38.1376
8. Chkareuli, J.L.: On the origin of poincaré gauge gravity. *Physics Letters B* **769**, 377–384 (2017). DOI 10.1016/j.physletb.2017.04.012
9. Chkareuli, J.L.: Poincaré gauge gravity an emergent scenario. *Physical Review D* **95**(8), 084051 (2017). DOI 10.1103/PhysRevD.95.084051
10. Obukhov, Y.N.: Poincaré gauge gravity: An overview. *International Journal of Geometric Methods in Modern Physics* **15**(supp01), 1840005–342 (2018). DOI 10.1142/S0219887818400054
11. Obukhov, Y.N., Hehl, F.W.: General relativity as a special case of poincaré gauge gravity. *Physical Review D* **102**(4), 044058 (2020). DOI 10.1103/PhysRevD.102.044058
12. Hehl, F.W., Kerlick, G.D., Heyde, P.v.d.: On hypermomentum in general relativity i. the notion of hypermomentum. *Ztschrift Fur Naturforschung A* **31**(2) (1976). DOI 10.1515/zna-1976-0201
13. Hehl, F.W., Kerlick, G.D., Heyde, P.v.d.: On hypermomentum in general relativity. ii. the geometry of space-time. *Zeitschrift Fur Naturforschung A* **31**(6), 524–527 (1976). DOI 10.1515/zna-1976-0602
14. Hehl, F.W., David, G., Kerlick, Heyde, P.v.d.: On hypermomentum in general relativity iii. coupling hypermomentum to geometry. *Zeitschrift Fur Naturforschung A* **31**(7), 823–827 (1976). DOI 10.1515/zna-1976-0724
15. Hehl, F.W.: On a new metric affine theory of gravitation. *Physics Letters B* **63**(4), 446–448 (1976). DOI 10.1016/0370-2693(76)90393-2
16. Hehl, F.W., Lord, E.A., Ne'Eman, Y.: Hadron dilation, shear and spin as components of the intrinsic hypermomentum current and metric-affine theory of gravitation. *Physics Letters B* **71**(2), 432–434 (1977). DOI 10.1016/0370-2693(77)90260-X
17. Lord, E.A.: The metric-affine gravitational theory as the gauge theory of the affine group. *Physics Letters A* **65**(1), 1–4 (1978). DOI 10.1016/0375-9601(78)90113-5
18. Hehl, F.W., Mccrea, J.D., Mielke, E.W., Ne'Eman, Y.: Metric-affine gauge theory of gravity: field equations, noether identities, world spinors, and breaking of dilation invariance. *Physics Reports-review Section of Physics Letters* **258** (1995). DOI 10.1016/0370-1573(94)00111-F
19. Sobreiro, R.F., Otoy, V.J.V.: Affine gauge theory of gravity and its reduction to the riemann-cartan geometry. *Journal of Physics Conference* **283**(1) (2010). DOI 10.1088/1742-6596/283/1/012032
20. Sobreiro, R.F., Otoy, V.J.V.: On the topological reduction from the affine to the orthogonal gauge theory of gravity. *Journal of Geometry and Physics* **61**(1), 137–150 (2011). DOI 10.1016/j.geomphys.2010.09.009
21. Cirilo-Lombardo, D.J.: Unified field theoretical models from generalized affine geometries. *International Journal of Theoretical Physics* **49**(6), 1288–1301 (2010). DOI 10.1007/s10773-010-0310-9
22. Cirilo-Lombardo, D.J.: Unified field theoretical models from generalized affine geometries ii. *International Journal of Theoretical Physics* **50**(6), 1699–1708 (2011). DOI 10.1007/s10773-011-0678-1
23. Cirilo-Lombardo, D.J.: Unified field theoretical models from generalized affine geometries iii. *International Journal of Theoretical Physics* **50**(11), 3621–3634 (2011). DOI 10.1007/s10773-011-0871-2
24. Kaluza, T.: Zum unitätsproblem der physik. *Sitzungsber Preuss Akad Wiss Berlin(Math. Phys.)* pp. 966–972 (1921). DOI 10.1142/S0218271818700017
25. Klein, O.: Quantentheorie und fünfdimensionale relativitätstheorie. *Zeitschrift für Physik A* **37**(12), 895–906 (1926). DOI 10.1080/01422418608228771
26. Klein, O.: The atomicity of electricity as a quantum theory law. *Nature* **118**, 516 (1926). DOI 10.1142/97898145716160007
27. Goddard, P., Goldstone, J., Rebbi, C., et al.: Quantum dynamics of a massless relativistic string. *Nuclear Physics* **56**(1), 109–135 (1973). DOI 10.1016/0550-3213(73)90223-X
28. Polchinski, J., Chaudhuri, S., Johnson, C.V.: Notes on d-branes. *arXiv:hep-th/9602052v2* (1996)
29. Taylor, W.: Lectures on d-branes, gauge theory and m(atrices). *arXiv:hep-th/9801182v2* (1998)
30. Bachas, C.P.: Lectures on d-branes. *arXiv:hep-th/9806199v2* (1999)
31. Giveon, A., Kutasov, D.: Brane dynamics and gauge theory. *Review of Modern Physics* **71**(4), 310–318 (1998). DOI 10.1103/RevModPhys.71.983

32. Wess, J., Zumino, B.: Supergauge invariant extension of quantum electrodynamics. *Nuclear Physics B* **78**(1), 1–13 (1974). DOI 10.1016/0550-3213(74)90112-6
33. Hanany, A., Witten, E.: Type iib superstrings, bps monopoles, and three-dimensional gauge dynamics. *Nuclear Physics B* **492**(1-2), 152–190 (1997). DOI 10.1016/S0550-3213(97)00157-0
34. Douglas, M.R., Govindarajan, S., Jayaraman, T., et al.: D-branes on calabiyau manifolds and superpotentials. *Communications in Mathematical Physics* **248**(1), 85–118 (2004). DOI 10.1007/s00220-004-1091-x
35. Haber, H.E., Kane, G.L.: The search for supersymmetry: Probing physics beyond the standard model. *Physics Reports* **117**(2-4), 75–263 (1985). DOI 10.1016/0370-1573(85)90051-1
36. Gunion, J.F., Haber, H.E.: Higgs bosons in supersymmetric models (i). *Nuclear Physics B, Particle Physics* **272**(1), 1–76 (1986). DOI 10.1016/0550-3213(86)90340-8
37. Gunion, J.F., Haber, H.E.: Higgs bosons in supersymmetric models (ii). implications for phenomenology. *Nuclear Physics B, Particle Physics* **278**(3), 449–492 (1986). DOI 10.1016/0550-3213(86)90050-7
38. Gunion, J.F., Haber, H.E.: Higgs bosons in supersymmetric models (iii). decays into neutralinos and charginos. *Nuclear Physics B, Particle Physics* **307**(3), 445–475 (1988). DOI 10.1016/0550-3213(88)90259-3
39. Gunion, J.F., Haber, H.E.: Errata for higgs bosons in supersymmetric models: I, ii and iii. arXiv:hep-ph/9301205v1 (1993)
40. Wu, T.T., Yang, C.N.: Concept of nonintegrable phase factors and global formulation of gauge fields. *Physical Review D* **12**(12), 3845–3857 (1975). DOI 10.1103/PhysRevD.12.3845
41. Chern, S.S., Chen, W.H., Lam, K.S.: *Lecture on Differential Geometry*. World Scientific Publishing Co. Pte. Ltd. (1999)
42. Nikodym, O.: Sur une generalisation des integrales de mj radon. *Fundamenta Mathematicae* **15**(1), 131–179 (1930)