

Comparison of Lidocaine, Huffing Maneuver and Combination of Both in Prevention Fentanyl Induced Cough Before Induction of Anesthesia: A Randomized Placebo-Controlled Trial

Laleh Dehghanpisheh

Shiraz University of Medical Sciences

Mohammadhossein Eghbal

Shiraz University of Medical Sciences

Fatemeh Bagheri Baravati

Shiraz University of Medical Sciences

Pooya Vatankhah (✉ Pvatankhah@sums.ac.ir)

Shiraz University of Medical Sciences

Research Article

Keywords: fentanyl, lidocaine, Huffing maneuver, cough, combination protocol

Posted Date: February 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-174791/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on March 27th, 2021. See the published version at <https://doi.org/10.1186/s12871-021-01313-w>.

Abstract

Background

Intravenous bolus injection of fentanyl has been frequently reported to be associated with cough reflex during patient anesthesia. However, the search for the most effective protocol continues. This study aimed to compare the effect of reducing cough reflex after injection of fentanyl in a fusion protocol by combining the injections of lidocaine and Huffing maneuver and comparing with a placebo control group, before anesthesia induction.

Methods

This prospective randomized controlled trial study was performed on 400 patients who were divided into four groups of combined protocol (group 1), lidocaine group (group 2), Huffing maneuver group (group 3), and the control receiving normal saline (group 4). Then patients were injected with 2.5 µg/kg fentanyl and monitored for two minutes regarding their cough reflex, as well as the severity.

Results

In group 1, 9 patients (9%), in group 2, 22 patients (22%), in the group 3 45 patients (45%) and in the group 4, 75 patients (75%), developed cough reflex following fentanyl injection. Also, 13 patients (13%) developed moderate and 4 (4%) developed severe coughs in the control group reported, while no reports of severe or moderate cough were among the intervention groups. There was a significant difference between the intervention group and the control group both in terms of the rate and severity of the fentanyl-induced cough.

Conclusion

By using a combination of lidocaine injection along and Huffing maneuver, better results can be obtained in reducing the frequency, and also the severity of cough followed by fentanyl injection.

Trial registration:

Trial Registration: The trial was registered with IRCT.IR (09/03/2018-No. IRCT20141009019470N74).

1. Introduction

Fentanyl-induced cough (FIC) is amongst the most common complications of anesthesia induction by intravenous bolus administration of fentanyl, which has been reported in 18 to 65% of patients. [1, 2]

Although this phenomenon is usually brief and self-limiting, during anesthesia induction, coughing is undesirable since it is associated with increased intracranial (ICP), intraocular, and intra-abdominal pressures. Also, multiple conjunctival and periorbital petechiae can be triggered by extreme FIC [3] and contribute to upper airway obstruction that may need urgent intervention [4]. Hence, prevention of FIC is of much clinical value.

Although multiple FIC-responsible mechanisms have been proposed, the exact mechanism remains unknown. Central sympathetic outflow could be blocked by fentanyl, thus stimulating the vagus nerve. This vagal activity change has been identified as a potential cause of cough and reflex bronchoconstriction [5, 6]

Lidocaine was successful in suppressing FIC in a meta-analysis by Kim et al, regardless of dose. [7] The mechanism by which lidocaine suppresses mechanically and chemically mediated cough reflex remains unclear, although it has been proposed that depression in brain-stem activity is a possibility [8].

The Huffing maneuver is a gentle, voluntary maneuver that opens the glottis following a high-pressure exhalation, and the air inside the upper airways is emptied along with secretions. This maneuver causes the opening of closed alveoli, reduces atelectasis, and increases the functional residual capacity of the lungs. [9, 10] Ambesh et al reported a significant decrease in the incidence and severity of FIC in patients who perform a huffing maneuver just before intravenous fentanyl injection. [11]

Various interventions have been performed to reduce the incidence of FIC, however none of which alone have demonstrated satisfactory reducing effect. In this study, the reducing effect of FIC was investigated in an integrated protocol with the simultaneous use of lidocaine injection along with the Huffing maneuvers before induction of anesthesia.

2. Material And Method

This study, which is a prospective randomized controlled trial, was performed on 400 patients who underwent fentanyl injection in hospitals affiliated to Shiraz University of Medical Sciences in 2018. Patients with non-emergency surgery aged 14 to 69 years were included in the study, while patients with a history of asthma, chronic obstructive pulmonary disease, chronic cough, upper respiratory tract infection in the two weeks before surgery, a history of smoking, steroid or bronchodilator treatment, or people taking ACE inhibitors were excluded.

The study was approved by the ethics committee of Shiraz University of Medical Sciences (IR.SUMS.MED.REC.1394.61), the institutional review board, and Iranian Registry of Clinical Trials (IRCT.ir; date: 09/03/2018-No. IRCT20141009019470N74) and conducted in compliance with local regulatory requirements, Good Clinical Practice (GCP), and the Declaration of Helsinki [12]. All stages of the research and the goals were explained by the researcher to all the patients and written informed consent was obtained from all patients or their legally authorized representatives. Also, it is ensured that the reluctance to participate in the study has no effect on their treatment process and they can leave our study whenever

they wish. Assuming 95% confidence level (first type alpha error 5%) and 80% power and considering the observation of at least 0.30 and expecting differences in treatment results between the intervention (0.25) and control (0.55) groups, we calculated that a total of 82 patients (i.e. 41 cases in the intervention group and 41 in the control group) would be required for the analysis (Fleiss with CC).

The Patients were then divided into four groups of 100 which were equivalent in sex and age, based on block randomization.

Group 1 (Integrated Protocol Group): In this group, one minute before fentanyl injection, 2 mg/kg lidocaine was injected into the patient and after 55 seconds, Huffing maneuver was performed after a deep inspiration for less than 5 seconds, and subsequently the patient was injected with 2.5 µg /kg fentanyl.

Group 2 (lidocaine group): In this group, one minute before fentanyl injection, 2 mg/kg lidocaine was injected into the patient and then fentanyl was injected into the patient at 2.5 µg/kg.

Group 3 (Huffing group): In this group, one minute before fentanyl injection, the Huffing maneuver was performed after a deep inspiration for less than 5 seconds and then fentanyl was injected at a rate of 2.5 µg /kg.

Group 4 (control group): In this group, one minute before the injection of fentanyl, 2 cc/kg of normal saline was injected into the patient and then fentanyl was injected at the rate of 2.5 µg /kg.

Then the patients were monitored by a neutral observer for two minutes, which was blinded from the patients grouping, and the presence or absence of reflex cough as well as the severity of cough (as 1–2 coughs assigned as first-degree cough, 3–4 as grade 2, while 5 coughs and more than 5 grade 3) was reported in each patient.

During these 2 minutes, all secondary changes related to fentanyl such as changes in heart rate or changes in blood pressure that required intervention were recorded. For the induction of anesthesia, propofol was injected for all patients if no coughs were recorded during the initial 2 minutes after fentanyl injection, while in cases where coughs persisted, propofol injection was delayed until the cessation of coughs. All data were collected and entered into SPSS statistical software for statistical analysis and analyzed using the chi-square test for qualitative variables and student test for quantitative variables. P-value < 0.05 was also considered significant.

3. Results

A total of 400 patients in four groups were included in the study during 2018. Patients were matched in terms of age and sex in groups (P > 0.05). The baseline demographic data of the patients is demonstrated in Table 1.

Table 1

Demographic and Fentanyl-induced cough frequency and severity based on the injection of Lidocaine, Huffing Maneuver, or combined therapy.

Variables	Groups n = 400				Pvalue*
	Integrated Protocol Group <i>n = 100</i>	lidocaine group <i>n = 100</i>	Huffing group <i>n = 100</i>	Control group <i>n = 100</i>	
Age	42.67 ± 12.73	40.99 ± 10.75	44.05 ± 14.84	41.24 ± 15.03	< 0.05
Sex					
<i>Male</i>	45 (45%)	50 (50%)	47 (47%)	47 (47%)	< 0.05
<i>Female</i>	55 (55%)	50 (50%)	53 (53%)	53 (53%)	
Fentanyl-induced cough frequency	9 (9%)	22 (22%)	45 (45%)	75 (75%)	< 0.001
Severity					
<i>Moderate</i>	0 (0%)	0 (0%)	0 (0%)	13 (13%)	< 0.001
<i>Severe</i>	0 (0%)	0 (0%)		4 (4%)	
* Chi-square test					

As demonstrated in Table 1, in the first group 9 patients (9%), in the second group 22 patients (22%), in the third group 45 patients (45%) and in the fourth group 75 patients (75%) had a reflex cough was following fentanyl injection. The frequency of FIC in intervention groups was significantly lower than the control group (P value < 0.001), also this symptom was significantly lower in the first group (integrated protocol) compared to the second groups (lidocaine injection alone; P < 0.001) And the third (Huffing maneuver alone; P = 0.009). Furthermore, comparing the second and third group (lidocaine injection alone vs. Huffing maneuver alone), the rates were significantly lower in the Huffing maneuver group (P value < 0.001).

In the control group, 13 cases (13%) had moderate coughs and 4 cases (4%) had severe coughs, while in the intervention groups, moderate and severe cough was not observed, while this difference between the control and intervention groups was also statistically significant (P-value < 0.001).

4. Discussion

During a cough, the pressure inside the chest reaches 300 mm Hg and the expiratory air velocity reaches 500 miles per hour [13]. This pressure and speed caused by coughing can cause adverse effects such as headache, dizziness, myalgia, hoarseness, urinary incontinence, and even rib fractures, especially in

patients with low bone density [14, 15]. Fentanyl is widely used in the operating room to anesthetize patients, but a typical complication of this valuable drug in patients during anesthesia is cough [16, 17].

In the present study, which aimed to combine the effects of Huffing maneuver and lidocaine injection before fentanyl administration to reduce the complication of FIC, it was found that FIC is prevalent after fentanyl injection in patients who no interventions in this regard was applied, which accounted for 75% of the patients in the control group. Our higher rates compared to most studies in this field, such as the study of Lin et al. [2], which reported a prevalence of 65%, and the study of Chandra et al., [18] which reported a prevalence of 34%, may have been due to the wider age range and the larger number of samples in this study.

Based on our data, the prevalence of cough was 45% in patients who received 2 µg / kg lidocaine before fentanyl injection, which was much higher compared to other studies such as Gecaj-Gashi et al. [19] who reported a prevalence of 16% with 1 µg / kg lidocaine injection and 22% with 0.5 µg / kg injection and Chandra et al. who reported a prevalence of 13%. [20] This could indicate that increasing the dose of lidocaine, as shown in another study by Chandra et al. [21], increased the prevalence of this complication. However, further studies in this regard are justified.

In our study, the incidence of cough was reported to be 22% in patients who underwent Huffing maneuvers before fentanyl injection. By applying a combination therapy, this rate was only observed in 9% of the patients, which was significantly lower compared to the control group. In this regard, a study by Ambesh et al. [11] showed that patients who underwent Huffing maneuvers before fentanyl injection had significantly lower incidences of drug-induced cough in the control group, however a study on the effect of the combination of huffing maneuver along with lidocaine injection has not been previously reported till date. Also in the present study, similar to the study by Ambesh et al. [11], the severity of drug-induced cough in the intervention groups was significantly lower than the control group, however, in other studies, no significant relationship was observed between the groups [2, 20].

5. Conclusion

By using a combination of lidocaine injection along and Huffing maneuver, better results can be obtained in reducing the frequency, and also the severity of cough followed by fentanyl injection. These results can be beneficial for anesthesiologists worldwide to avoid undesirable complications of FIC.

6. Declarations

6.1. Ethics approval and consent to participate

This study obtained ethics approval from the Institutional Review Board of Shiraz University of Medical Sciences (IR.SUMS.MED.REC.1394.61). The study was registered on IRCT.IR with the following identifier:

IRCT.ir; date: 09/03/2018-No. IRCT20141009019470N74. All participants were provided with written consent and were guaranteed that their information would be safeguarded by the researcher.

6.2. Consent for publication

Not applicable

6.3 Availability of data and materials

The dataset supporting the conclusions of this article is included within the article. Please contact the corresponding author in case of requiring any further information.

6.4. Competing interests

All authors declare that they have no competing interests.

6.5 Funding

No specific funding was obtained for this study.

6.6 Authors' contributions

L.D. and P.V. contributed to the conception or design of the work. M.E. contributed to the acquisition and analysis of the data. F.B. contributed to the interpretation of the data for the work. P.V. drafted the work. L.D. revised it critically for important intellectual content. All authors approved the final version to be published.

6.7. Acknowledgments

The authors would like to thank Reza Shahriarirad for assisting the manuscript preparation.

References

1. Lin J-A, Yeh C-C, Lee M-S, Wu C-T, Lin S-L, Wong C-S: **Prolonged injection time and light smoking decrease the incidence of fentanyl-induced cough.** *Anesthesia & Analgesia* 2005, **101**(3):670–674.
2. Lin C-S, Sun W-Z, Chan W-H, Lin C-J, Yeh H-M, Mok MS: **Intravenous lidocaine and ephedrine, but not propofol, suppress fentanyl-induced cough.** *Canadian Journal of Anesthesia* 2004, **51**(7):654–659.
3. Tweed WA, Dakin D: **Explosive coughing after bolus fentanyl injection.** *Anesthesia & Analgesia* 2001, **92**(6):1442–1443.

4. Ambesh S, Singh N, Srivastava K: **Fentanyl induced coughing caused life-threatening airway obstruction in a patient with arteriovenous malformation of tongue and hypopharynx.** *Int J Anesthesiol* 2009, **20**(20):7.
5. Ping-Wing L, Chung-Hsi H, Ya-Churn C: **Terbutaline inhalation suppresses fentanyl-induced coughing.** *Canadian journal of anaesthesia* 1996, **43**(12):1216.
6. Agarwal A, Azim A, Ambesh S, Bose N, Dhiraj S, Sahu D, Singh U: **Salbutamol, beclomethasone or sodium chromoglycate suppress coughing induced by iv fentanyl.** *Canadian journal of anaesthesia* 2003, **50**(3):297.
7. Kim JE, Min SK, Chae YJ, Lee YJ, Moon BK, Kim JY: **Pharmacological and nonpharmacological prevention of fentanyl-induced cough: a meta-analysis.** *Journal of anesthesia* 2014, **28**(2):257–266.
8. Poulton TJ, JAMES FM: **Cough suppression by lidocaine.** *Anesthesiology: The Journal of the American Society of Anesthesiologists* 1979, **50**(5):470–472.
9. Brauer S: **Nonpharmacologic airway clearance therapies.** *Australian Journal of Physiotherapy* 2007, **53**(1):67.
10. Deck D, Winston L, Katzung B, Masters S, Trevor A: **Basic & clinical pharmacology.** 2012.
11. Ambesh S, Singh N, Gupta D, Singh P, Singh U: **A huffing manoeuvre, immediately before induction of anaesthesia, prevents fentanyl-induced coughing: a prospective, randomized, and controlled study.** *British journal of anaesthesia* 2010, **104**(1):40–43.
12. Association WM: **World Medical Association Declaration of Helsinki: Ethical Principles for Medical Research Involving Human Subjects.** *JAMA* 2013, **310**(20):2191–2194.
13. Comroe J: **Special acts involving breathing.** *Physiology of respiration: an introductory text.* In.: Chicago, IL: Yearbook Medical Publishers; 1974.
14. Hanak V, Hartman TE, Ryu JH: **Cough-induced rib fractures.** In: *Mayo Clinic Proceedings: 2005.* Elsevier; 2005: 879–882.
15. Irwin RS, Baumann MH, Bolser DC, Boulet L-P, Braman SS, Brightling CE, Brown KK, Canning BJ, Chang AB, Dickey PV: **Diagnosis and management of cough executive summary: ACCP evidence-based clinical practice guidelines.** *CHEST Journal* 2006, **129**(1_suppl):1S-23S.
16. Kato R, Ross S, Foex P: **Fentanyl protects the heart against ischaemic injury via opioid receptors, adenosine A1 receptors and KATP channel linked mechanisms in rats.** *British journal of anaesthesia* 2000, **84**(2):204–214.
17. Weiskopf RB, Eger II EI, Noorani TM, Daniel M: **Fentanyl, Esmolol, and Clonidine Blunt the Transient Cardiovascular Stimulation Induced by Desflurane in Humans.** *Anesthesiology* 1994, **8**:15S0-1555.
18. Pandey CK, Raza M, Ranjan R, Singhal V, Kumar M, Lakra A, Navkar DV, Agarwal A, Singh RB, Singh U: **Intravenous lidocaine 0.5 mg·kg⁻¹ effectively suppresses fentanyl-induced cough.** *Canadian Journal of Anesthesia* 2005, **52**(2):172–175.
19. Gecaj-Gashi A, Nikolova-Todorova Z, Ismaili-Jaha V, Gashi M: **Intravenous lidocaine suppresses fentanyl-induced cough in Children.** *Cough* 2013, **15**(1):20.

20. Pandey CK, Raza M, Ranjan R, Lakra A, Agarwal A, Singh U, Singh RB, Singh PK: **Intravenous lidocaine suppresses fentanyl-induced coughing: a double-blind, prospective, randomized placebo-controlled study.** *Anesthesia & Analgesia* 2004, **99**(6):1696–1698.
21. Schlimp CJ, Wiedermann FJ: **Does fentanyl-induced cough justify pre-treatment with IV lidocaine 2 mg/kg-1.** *Can J Anaesth* 2005, **52**(2):207.