

Towards school-based mental health programs in Nigeria: The immediate impact of a depression-literacy program among school-going adolescents and their teachers

Olayinka Atilola (✉ draromedics@yahoo.com)

Lagos State University College of Medicine Ikeja

Olatunde Ayinde

University of Ibadan

Felix-Kingsley Obialo

Dominican University Ibadan

Sunday Oladotun Adeyemo

Olabisi Onabanjo University Ago-Iwoye

Adedapo Adegbaju

Federal Neuro-psychiatric Hospital Yaba

Robert Anthony

Adolescent Wellness, Inc

Research Article

Keywords: depression, depression literacy, adolescents, school mental-health

Posted Date: June 23rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1765925/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Depression-literacy, which is the foundational requirement for symptom recognition, positive attitude and help-seeking, is poor among adolescents in Nigeria. This study, therefore, aims to determine the impact of a school-based training program on depression-literacy among a cohort of high-school students and their teachers in South-West Nigeria.

Methods: An adapted version of the Break Free from Depression, a 4-module depression awareness curriculum for staff and students, was implemented among students and their teachers. Paired-sample T-test was used to assess the domain-specific (knowledge, attitude, and confidence) impact of the training by comparing the baseline and immediate (within the week of the training) post-scores.

Results: A total of 3,098 students and 294 teachers from 21 schools across three states in South-West Nigeria successfully completed the training. There was a significant positive difference ($p < .05$), at post-test, in the knowledge, attitude, and confidence among the students. The same was observed among teachers except for attitude where positive change did not reach significant level ($p = .06$). When statistically significant, the calculated effect size (eta squared) was highest for knowledge (students: $.07$, $p = .001$; teachers: $.08$, $p < .000$) and least for attitude (students: $.003$, $p = .002$ teachers: $.085$, $p = .06$). Multiple regression analyses result showed that the level of pre-scores predicted the magnitude of change in all domains of depression-literacy ($p < .05$) after controlling for age, gender, and type of school among the students, but not for teachers.

Conclusions: School-based depression-literacy programs can lead to significant positive change in knowledge, attitude, and confidence of students and teachers.

Background

Data from the Global Burden of Disease (GBD) studies have shown that mental disorders contributes significantly to disability among children and adolescents and, in fact, these disorders are likely to rank high among major child and adolescent public health challenges of the twenty-first century [1]. Among the common mental disorders that afflict children and adolescents, GBD data also establish depression as ranking highest in terms of burden [2]. Though prevalence rates vary from one region of the world to another, one cross-national study which sampled adolescents from more than ten countries across the world reported that one out of ten adolescents have clinically significant symptoms of depression [3]. In Nigeria, despite varying prevalence rates, several studies conducted in different parts of the country, have established that depression is common among children and adolescents. For instance, in South-West Nigeria, a study conducted among school-going adolescents in Ibadan identified mild to moderate depression in nearly a quarter (23.8%) and severe depression in 5.7% of respondents [4]. Oderinde et al also reported a prevalence rate of 16.3% among a sample of school-going adolescents in Ido-Ekiti [5]. Adewuya & Ologun reported a lower but equally high prevalence rate of 9.0% prevalence rate in a different sample of adolescents sampled from schools in Ile-Ife [6]. Studies conducted in other regions of the country also established a significantly high burden of adolescent depression [7, 8].

When left untreated, depression among children and adolescents is associated with significant short/medium-term risks such as missed school days and suicidal behaviours [9] and long-term negative consequences such as persistent recurrence, poorer physical health morbidity, and chronic alcohol abuse [10–12]. It is therefore appropriate that investment in strategies to reduce the burden of depression at key developmental phases across lifespan, particularly during adolescence, is being seen as a global public health goal [13]. Fortunately, there is robust evidence for safe, effective, and affordable pharmacologic and non-pharmacologic interventions for adolescent depression worldwide (14–16). However, for a variety of reasons, there has been significant barriers to mental health services, including treatment for depression, among adolescents worldwide, worse in low- and middle-income countries, and as such, creating a treatment-gap [17].

Interventions to reduce adolescent depression treatment-gap must acknowledge the fact that access to treatment of depression among adolescents is, among other factors, a function of the degree of recognition of the constellation of experiences which characterizes the syndrome of depression, as a health concern which can be understood, prevented, or treated [18]. In other words, adolescent depression-literacy must improve as a pre-condition for the success, in terms of uptake, of available intervention strategies. This is because poor depression-literacy, whereby depressive symptoms were interpreted as normal inconveniences of puberty or a personality trait, has been identified as one of the potential barriers to help-seeking among depressed adolescents [19]. Furthermore, there is preliminary evidence, from around the world, that school-based mental health awareness programs which targets the provisioning of accurate information about the nature of and services for mental health conditions to parents, teachers, and students, can and does lead to improved symptom recognition and uptake of mental-health services among school children [20–22]. Aside from being the foundational component upon which additional structures for promoting mental-health, such as symptom recognition, tolerant attitude, and improved service utilization, can be built [23]; improved mental-health literacy of students also come with additional benefits such as reduced prejudice and stigma, more positive emotional response towards peers who may be experiencing depression [24], and positive personal mental health in general [25, 26].

Recent research in Nigeria has shown that the health-education curriculum of Nigerian elementary and secondary schools focused on hygiene and environmental health with hardly any meaningful mental-health content [27]. Hence, knowledge of school-going adolescents, and indeed their teachers, about mental illness in general, is limited [28, 29]. By extension, attitude of school-going adolescents towards individuals with mental disorders has been shown to be negative in one study and reflected desire for social distance [30]. There have been very few school-based short-term training programs aimed at improving mental-health literacy of students in Nigeria in recent times. Such efforts have yielded immediate and sustained positive change in aspects of general mental-health literacy among small samples of children and adolescents attending elementary and secondary schools in the country [31, 32]. However, there has not been any documented intervention (school- or community-based) aimed at improving depression-literacy in particular among adolescents in Nigeria, and indeed in sub-Saharan Africa. This is despite the fact that more recent research, which examined depression-literacy specifically among school-going adolescents, found that only a paltry 4.8% of the sample correctly identified a depression vignette while another paltry 1.5% recommended professional help as a treatment option [33]. Furthermore, the few earlier intervention studies for general

mental-health literacy conducted in Nigeria did not involve teachers. Teachers are equally important members of the school environment and mental-health literacy has been shown to be equally poor among them [28, 29]. Improving the mental-health literacy of teachers alongside the students has been described as “critical” to the success of school-based mental-health promotion and intervention [34].

From the foregoing, it is obvious that more mental-health literacy programs are needed in Nigeria, and indeed Africa, especially large-scale intervention programs focussing on a specific mental-health problem such as depression. The present study, therefore, aimed to conduct an implementation of a standardized depression-literacy program among a large and representative sample of school-going adolescents and their teachers in Nigeria. In this paper, we report the immediate post-intervention impact, in terms of, satisfaction with the program, the knowledge gained and the socio-demographic factors that independently influenced it.

Methods

Design and Setting

The quasi-experimental, post-pre interventional study was conducted in secondary schools within the three states of Oyo, Ogun, and Osun in South-West Nigeria. The three states were picked based on convenience, though their combined population represents about 10% of the Nigerian population.

Sampling

The data analysed for this study came from an ongoing community project carried out in conjunction with the Rotary International, on which the authors served as technical and scientific partners. By the original design of the project, a convenient sample of schools from the three contiguous southwestern states of Oyo, Ogun, and Osun in Nigeria were selected, to reflect adequate representation of rural and urban, as well as private and public schools. In each selected school, school counsellors contacted all the students in each class to participate in the project. An explanatory note describing the project was sent to their parents. All contacted students whose parents provided written informed consent and who assented to participate were then included in the project. All the teachers in each of the selected schools were contacted to participate in the project through an explanatory note with an attached consent form. All teachers who returned signed consent forms were recruited into the project.

Procedures

The intervention

The intervention was adapted from the Break Free from Depression, a 4-module Depression Awareness Curriculum for Staff and Students developed by the Boston Children Hospital Neighborhood Partnerships [35]. The manual was developed from an initial depression tool-kit for schools [36], and later further enhanced through several years of working with school children with depression and getting feeds-back from

them. The present version has been used in several schools in the United States of America (USA) and in a few middle-income countries such as India. It is currently included in the best-practice registry of the Suicide Prevention Resource Center in the USA. The manual has three main components: The Staff (teachers and other administrative/ support staff) Training, Parent Events, and the implementation of the Student Curriculum. The teacher and parent components, with slight differences in the content and language to suit the audience, are both essentially geared towards educating the teachers/parents about the rationale for a depression awareness program in schools, nature and dynamics of depression and suicidal behaviors among adolescents, and ways to help and/or refer students/wards experiencing these symptoms. The student curriculum, on the other hand, has four modules originally. The first module (Module 1) is essentially an overview of the facts about the nature, sign, and symptoms of depression and suicidality, including efforts to dispel local myths about the condition. The second module (Module 2) is a film documentary aimed at 'bringing alive', in audio-visual format, the symptoms, stigma, risk factors, and coping strategies for depression. The third module (Module 3) is essentially a small-group debriefing session in which participants are encouraged to apply the knowledge gained from Module 1 to a set of pre-written case studies of depression and suicidality among hypothetical teenagers. The module also covers the key warning signs of suicide, basic creative problem-solving skills, and additional resources that can be accessed if further information is needed. The fourth module (Module 4) is an open discussion on self-help strategies for depression and the steps to be taken to access or help a friend to access additional mental-health services. It involves the use of presentations and role plays.

The pilot Implementation

Preparation

The entire school community, including students, teachers, parents' associations, and school authorities were briefed about the program and the potential benefits. A lead-staff, often the school counsellor, was identified to be the internal coordinator for the project in each school. The internal coordinator liaised between the program team and the school community, helped to provide answers to preliminary inquiries, addressed logistical issues in the course of the program, and served as the continuous link between the program team and the school during and after the program.

Adaptation

the original version of the depression awareness curriculum was adapted to suit the culture, resource-environment, and the available funds for the project. Firstly, the entire curriculum for the staff training, parents' event, and student curriculum was adapted to be more relevant to the culture of south-west Nigeria. This was in consideration of the observations from systematic review of qualitative studies around the world, which have established cultural differences and nuances in the perception of the nature of and explanatory model for mental disorders, including depression [37]. The adaptation was done by a panel, comprised of the investigators, who are mental-health practitioners with wide experience with the local realities of depression among adolescents in the region, particularly with reference to peculiarities in risk factors, coping strategies, idioms of expression of depressive psychopathology, and nuances in explanatory models of depression. The panel reviewed the entire curriculum and the case studies for appropriateness and

cultural relevance. Minor changes were made to the content of the curriculum and case studies based on a consensus of the experts. Secondly, we were not able to implement Module 2, that is, the film documentary part for the students. This was due to two main reasons. The first reason was the impracticability of culturally adapting the contents of the film documentary. The second reason was the logistics difficulties of airing the documentaries in some rural locations. Rather, we made sure to enrich the contents of Modules 1 and 3 to adequately cover the description of the symptoms and signs of depression with more illustrative examples.

Implementation Proper

All trainings were led by a member of the team supported by all other members of team. The training-lead is a doctoral candidate in clinical psychology (OA) with experience in conducting psycho-educational programs in schools. He is also fluent in both English and Yoruba, the local language of the region. All trainings were conducted in simple English, with occasional use of Yoruba, the local language to emphasize some points or concepts that were difficult to capture in English. The staff training was the first event, and it took place in batches in one of the school classrooms in each participating school. It involved teachers only. All available and consenting teachers took part. The parents' event was a single-day event. The parents were recruited through the schools' Parent Teacher Association. This was also a single-day event and all recruited parents who showed-up were asked to participate after giving their consent. The Module 1 and 4 of the student curriculum training was done in a single group of no more than 50 participants in one of the spacious school hall/classrooms. The Module 3 was implemented in small sub-groups of about 10 participants.

Evaluation

Pre- and post-survey was administered to both student- and teacher-participants as part of the training. The pre-survey was a 5-10-minute, 22-item survey which is part of the original Break Free from Depression Modules. The surveys measure respondents' knowledge (8 items) and attitude (5 items) about depression, as well as their level of confidence (9 items) in responding to a situation related to depression on a 4-point Likert scale ranging from strongly disagree to strongly agree. The pre-survey was administered before the start of the program. The post-survey, which was administered within the week of completing the training, was a replica of the pre-survey but, in addition, had extra sections assessing, on similar 4-point Likert scale, the overall satisfaction of respondents with the training. A single 'Yes' or 'No' question about whether respondent had referred someone, including themselves, to an adult or professional (as the case may be for students and teachers respectively), as a result of the training they had received from the program was also added to the post-survey. All pre-and post-surveys for each respondent were matched and linked with a code. In the final analysis, respondents who strongly agreed/ agreed were grouped together as one group while those who disagreed/ strongly disagreed were also grouped together. Items that were negative on the attitude scale were reversed scored in such a way that higher scores indicated better attitude.

Ethical Consideration

We followed the ethical guideline of ensuring voluntariness, confidentiality, and non-maleficence in the conduct of the research. The purpose of the research and any risk involved was explained to all would-be participants. Appropriate permissions were obtained from the local education authorities and the heads of each school. Written informed consent was obtained from parent and teachers, both for their own participation and that of their ward/students. The entire protocol of the project was also submitted to and approved by the local ethical review board. As additional benefit of participating in the program, a referral protocol for students who either sought help from the research team or were referred by friends or teachers was developed and implemented in a safe and confidential manner throughout the period of the project.

Statistical Analyses

Data was managed and analyzed using the Statistical Package for Social Sciences (SPSS) version 20.0. Descriptive statistics was used to present the age, gender, and other socio-demographic characteristics of participants across study sites. Paired-sample T-test was used to assess the overall and domain-specific (knowledge, attitude, and confidence) impact of the training within the same week of intervention by comparing the mean scores for the pre- and post-tests. A measure of effect sizes was obtained by dividing the overall mean difference in pre- and post-scores with the standard deviation. Multiple regression analysis, one model for the students and one model for the teachers, was used to determine the independent socio-demographic facilitators or barriers to improved literacy domains among the respondents. In the models, the dependent variable was the post intervention scores on the different domains as continuous variables. The independent variables were the socio-demographic data such as age, gender, type of school (public vs. private) and the baseline pre-test scores of the domain in question. The baseline pre-test score was included to correct for initial differences in the baseline scores of respondents. The model summary and the R-square for the model was presented after checking to make sure that standard assumptions such as linearity, collinearity, and homoscedasticity were not violated. Finally, the responses to the section on satisfaction with the program on the post-test was analyzed. The 4-point Likert responses was collapsed into a 'yes' (strongly agree and sort-of agree) and 'no' (strongly disagree and sort-of disagree) dichotomy. The proportion of respondents who reported satisfaction (yes response) for each of the satisfaction questions on the post-test were also recorded.

Results

The Break Free from Depression training modules, as modified, was successfully implemented in 21 public and private secondary schools across South-West Nigeria. The school authorities, teachers and students were mostly cooperative and supportive. The post-intervention results for the students and the teachers are presented separately.

Students

A total 3,098 secondary school students sampled from 21 public and private secondary schools in South-West Nigeria completed the training program. The mean age was 16.4 (\pm 1.58; range 11–19) years. There

were more females (n = 2049; 66.1%) than males. A slight majority of the students were from public secondary schools (n = 1769, 57.1%). As shown in Table 1, there was a significant positive difference ($p < .05$), at post-test, for each of the three domains of training, that is: knowledge of, attitude towards, and confidence in handling depression and suicidality among the students. The magnitude of the effect size (eta squared) was highest for knowledge (.07, medium effect) and least for attitude (.003, negligible). Multiple regression analyses result showed that the only statistically significant predictor of post intervention knowledge score was the level of pre-intervention knowledge scores among the students (see Table 2). Aside the pre intervention attitude scores which predicted post-intervention attitude score ($\beta = .256$, $p < .001$), students of private institutions had statistically significant better scores on attitude ($\beta = .170$, $p < .001$) compared with those in public schools. Age, sex, type of school, and pre-intervention level of self-reported confidence all predicted post-intervention confidence score in handling issues related to depression and suicidality. In the model, older respondents ($\beta = .083$, $p < .001$), females (.460, $p < .001$), private-school students ($\beta = .287$, $p = .002$), and those with higher pre-intervention confidence scores ($\beta = .286$, $p < .001$) all had significantly higher post-intervention confidence scores. Furthermore, at the completion of the program, almost three-quarters (71%, n = 2195) of the students who participated in the training agreed that the program was a good way to learn about depression and suicidality and would, therefore, recommend the program to a friend. Half of the participating students (50.9%, n = 1578) admitted to knowing a peer (including themselves) who sought help or advice about depression or suicidality from an adult, based on the lessons learnt from the training program. An overwhelming majority (82.5%, n=2552) expressed willingness to be a peer-trainer on the program if such opportunity was available in the future.

Teachers

A total 234 teachers from the 21 participating schools completed the training program. The mean age of the teachers was 39.8 years (SD: 9.97 years, range: 23 to 68 years). There were more females (n = 134;57.3%) than males. Majority of the teachers were from private secondary schools (n = 146; 62.4%). As shown in Table 1, self-reported knowledge of and confidence in handling issues related to depression and suicidality improved significantly ($p < .05$) and at roughly the same level of magnitude ($\eta^2 = .081$ and .080 respectively), which is of medium effect size, among the teachers at post-test. Though there was a demonstrable positive change in mean attitude scores post-intervention, the difference did not reach statistical significance ($p = .06$). Multiple regression analysis showed that pre-intervention attitude score was the sole predictor ($p < .001$) of the slight positive change in attitude (Table 3). Same regression models for knowledge and confidence did not show any statistically significant predictors of either outcome (Table 3)

Table 1
Paired Sample T-test for the Post and Pre-test comparison for both students and teachers

Paired differences									
	Mean difference	Std. Deviation	η^2 (Eta Square)	Std. Error Mean	95% CI of the difference		t	df	Sig. (2-tailed)
					Lower	Upper			
Students (n = 3,098)									
<i>Post-knowledge</i>	.458	1.663	.070	.030	.339	.517	15.327	3097	.001
<i>Post-attitude</i>	.081	1.449	.003	.026	.030	.132	3.113	3097	.002
<i>Post-confidence</i>	.514	2.822	.030	.051	.414	.613	10.129	3097	.001
Teachers (n = 234)									
<i>Post-knowledge</i>	.521	1.825	.081	.119	.286	.756	4.369	233	<.000
<i>Post-attitude</i>	.534	4.237	.085	.277	-.011	1.079	1.928	233	.06
<i>Post-confidence</i>	2.290	8.086	.802	.528	1.249	3.332	4.333	233	<.000

df: degree of freedom; t: Student T-test

Table 2
 Predictors of students' post-intervention knowledge of, attitude towards, and confidence about handling depression

Independent Variables	β	t	sig	Confidence interval	
				Lower bound	Upper bound
<i>Model 1 (dependent variable: post-knowledge)</i>					
Age	-.017	-.987	.324	-.042	.014
Sex (male coded 1, female 0)	-.002	-.111	.912	-.099	.089
School type (public coded 1 vs. private coded 0)	-.059	-1.289	.198	-.149	.031
Knowledge (pre-scores)	.183	10.342	.000	.140	.206
Constant	2.896	19.884	.000	4.607	5.614
<i>Model 2 (dependent variable: post-attitude)</i>					
Age	.002	-.168	.866	-.027	-.023
Sex (male coded 1, female coded 1)	.010	.231	.818	-.074	.094
School type (public coded 1 vs. private coded 0)	-.256	-6.230	.000	-.336	-.175
Attitude (pre-scores)	.170	9.437	.000	.336	-.175
Constant	2.383	10.85	.000	1.952	2.183
<i>Model 3 (dependent variable: post-confidence)</i>					
Age	.083	4.859	.000	.075	.177
Sex (male coded 1)	-.460	-5.204	.000	-.653	-.277
School type (public coded 1 vs. private coded 0)	-.287	-3.173	.002	-.433	-.102
Confidence (pre-scores)	.286	16.695	.000	.260	.330
Constant	3.092	6.966	.000	2.221	3.962
<i>Model 1: $R^2 = .035$, $F = 27.98$, $p < .001$; Model 2: $R^2 = .042$, $F = 34.55$, $p < .001$; Model 3: $R^2 = .10$, $F = 87.00$, $p < .001$</i>					

β : standardized co-efficient; t = Student T-test

Table 3
Predictors of teachers' post-intervention knowledge of, attitude towards, and confidence about handling depression

Independent Variables	β	t	sig	Confidence interval	
				Lower bound	Upper bound
<i>Model 1 (dependent variable: post-knowledge)</i>					
Age	.105	1.562	.120	-.004	.033
Sex (male coded 1, female 0)	.105	1.584	.114	-.073	.672
School type (public coded 1 vs. private coded 0)	.001	.002	.998	-.362	.363
Knowledge (pre-scores)	.100	1.434	.153	.153	.037
Constant	4.477	7.142	.000	3.242	5.712
<i>Model 2 (dependent variable: post-attitude)</i>					
Age	-.032	-.491	.624	-.018	.011
Sex (male coded 1, female coded 0)	.017	.260	.795	-.261	.341
School type (public coded 1 vs. private coded 0)	.234	1.573	.117	-.059	.527
Attitude (pre-scores)	.220	3.490	.001	.096	.345
Constant	2.174	5.031	.000	1.323	3.025
<i>Model 3 (dependent variable: post-confidence)</i>					
Age	.091	1.382	.168	-.099	.052
Sex (male coded 1)	.478	1.498	.135	-.151	1.107
School type (public coded 1 vs. private coded 0)	.111	.036	.971	-.599	.621
Confidence (pre-scores)	.131	1.961	.051	-.001	.263
Constant	5.187	5.569	.000	3.352	7.022
<i>Model 1: $R^2 = .050$, $F = 4.05$, $p = .003$; Model 2: $R^2 = .20$, $F = 2.20$, $p = .069$, Model 3: $R^2 = .34$, $F = 3.17$, $p = .085$</i>					

β : standardized co-efficient; t = Student T-test

Discussion

The successful execution of the modified version of the Break Free from Depression training module in Nigeria is a signal that such large-scale school-based mental-health programs is feasible and potentially

impactful. It also confirms earlier needs-assessments in Nigerian schools which have shown that teachers and school administrators recognized the need for and will welcome school-based training programmes that would improve mental-health literacy within the school ecosystem [38, 39]. The overall finding is that the intervention resulted in significant improvement in the three target domains of depression literacy (knowledge, attitude, and confidence) among the students and their teachers. The highest effect size, of moderate size, was seen in the knowledge domain followed by the confidence domain among the two groups. Other small-scale controlled studies conducted in Nigeria and other parts of Africa have demonstrated improvement in aspects of mental-health literacy after a short training program among students and teachers alike [31, 32, 40].

However, for both groups, that is; students and teachers in the present study, attitude was the least impacted, in terms of both the significance of change and the magnitude of impact. There have been mixed results on the impact of mental-health training programs on attitude among students and teachers. Some studies, including ones conducted in Africa, have found significantly improved attitude after mental-health literacy training programs [40, 41]. Other training programmes, however, have observed sustained poor attitude despite intervention [31, 32]. Recent systematic reviews showed that intervention program designs which involved extended contact-time spread over days or weeks and with multiple components (such as survivors lived-experiences and film/ documentaries) had better impact on attitudinal change compared with ultra-short and abridged training programs [42]. Future depression-literacy programs must be mindful of this design consideration. Furthermore, compared to the teachers' group, the students' group had an almost consistent significantly higher post intervention scores in more domains. This suggests that adolescence may be the golden period for achieving the best results when delivering mental health literacy interventions [43]. Adolescents are probably more likely than adults to find new knowledge interesting and be open minded enough for significant attitudinal change.

In terms of predictors, pre-intervention scores for the three domains of knowledge, attitude, and confidence were predictors of positive change in knowledge, attitude, and confidence, at least among the students. This finding is intuitive because it is expected that those with a pre-existing high-level of knowledge, attitude and confidence may also have better innate abilities to learn or are more likely to connect better with the training. The finding also suggests that mental health literacy interventions should be a continuing exercise, as those who are more likely to benefit from new information are those with better pre-existing information. Additionally, being a student in private school predicted improvements in attitude and confidence, while older adolescents and girls had better improvements on their post-intervention confidence in handling depression and suicidality issues compared with peers, all other factors kept constant. The sociodemographic nuances of impact of interventions are yet to be examined in the adolescent mental-health literacy literature and as such, we are unable to situate these observations. However, suffice to note that students of public and private schools in Nigeria often have wide differences in socio-economic background. Therefore, there might be certain socio-cognitive factors at play which are in favour of the demographics of the private-school students, but which were not measured in the present study. Likewise, there might be certain drivers of age and gender differences in the impact of the training program on attitude and confidence, which were not captured in our design.

Among teachers, improvement in knowledge and confidence was not dependent on age, gender, type of school where they taught, nor their pre-intervention assessment scores. Only their pre-intervention scores on attitude marginally predicted improvement in attitude. Aside from documenting the impact of the program on knowledge, attitude, and confidence, most of the mental-health literacy trainings that have been conducted hardly document the socio-demographic and other predictors of individual benefits. Therefore, there is hardly any previous study to compare these observations with. However, if these findings are replicated in future studies, it may have implications for design of similar interventions for maximal impact. These are areas for future exploration and replication.

There are a few other markers of success of the training program. Up to half of adolescents who participated in the training admitted to knowing someone (including themselves) who has sought help or advise from an adult for depression-related issues within the week of the training. While we did not determine the nature of the inquiry or the quality of the advice received, the finding indicates that the training stimulated conversation around depression and suicidality within the community. Such conversation may eventually lead to help-seeking. Furthermore, close to two-thirds of students who participated in the training program agreed that the program was a good way to learn about depression and suicidality. In the context of general mental-health literacy, school-going adolescents in Nigeria have found other school-based training programs acceptable [32]. This is another indicator of the acceptability of school-based mental-health literacy programs in Nigeria. Lastly, majority of the participating adolescents expressed willingness to be peer-trainers on the program if such opportunities were available. Though not yet implemented in the region of Africa, peer-led depression literacy programs has been shown to be acceptable and effective elsewhere [44]. Peer-led programs are viable alternatives to specialist-driven mental-health literacy programs in LMICs, such as Nigeria, where child mental-health specialist practitioners are in short supply and services are constrained [45]. For this reason, peer-led programs are more likely to be scalable in Nigeria than specialist-driven programs such as the present one and should be a future direction. The self-reported enthusiasm of the participating adolescents in the present study to serve as peer-educators is, therefore, encouraging.

There are a few design limitations of the present study that must be acknowledged. The first is that the schools which served as setting for the study were selected based on convenience. Therefore, the findings may not be easily generalizable. However, the schools were not selected based on any other bias that may have significantly influenced the results. Secondly, as acknowledged earlier, the training program was abridged to accommodate budget and logistics challenges. The results could have been different if the full training program was implemented. For the same reason, we report only the post-intervention outcomes within the week of intervention; we were unable to gather delayed outcome data. As such, we were not able to determine sustained impact. Other small-scale controlled trials of school-based mental-health literacy programs in Nigeria have documented sustained impact up to three months [31, 32]. Despite these limitations, the present study is still the largest school-based depression-literacy training program in Africa to date.

Abbreviations

SD: standard deviation

Declarations

Author contributions: All authors have participated in the concept, design, drafting or revising the manuscript. All authors read and approved the final manuscript.

Funding: The project from which the study was conceived was funded by the Rotary Foundation Global Grant (Grant number: GG1863183).

Availability of data and materials: The data and materials that support the findings of this study are available from the corresponding author upon reasonable request.

Declarations Ethics approval and consent to participate: The study protocol was approved by the review board of the Federal Neuropsychiatric Hospital Yaba Lagos, Nigeria. Participants were asked to sign an informed consent prior to their participation in the study.

Consent for publication: All authors agreed to publish this article in *Child and Adolescent Psychiatry and Mental Health*.

Competing interests: There are no competing interest to report.

References

1. Baranne ML, Falissard B. Global burden of mental disorders among children aged 5–14 years. *Child Adolesc Psychiatry Ment Health* [Internet]. 2018 Apr 12 [cited 2020 Jul 9];12. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5896103/>
2. Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012 Dec 15;380(9859):2197–223.
3. Atilola O, Balhara YPS, Stevanovic D, Avicenna M, Kandemir H. Self-reported mental health problems among adolescents in developing countries: results from an international pilot sample. *Journal of Developmental & Behavioral Pediatrics*. 2013;34(2):129–37.
4. Adeniyi AF, Okafor NC, Adeniyi CY. Depression and physical activity in a sample of nigerian adolescents: levels, relationships and predictors. *Child Adolesc Psychiatry Ment Health*. 2011 May 14;5:16.
5. Oderinde KO, Dada MU, Ogun OC, Awunor NS, Kundi BM, Ahmed HK, et al. Prevalence and Predictors of Depression among Adolescents in Ido Ekiti, South West Nigeria. *International Journal of Clinical Medicine*. 2018 Mar 6;9(3):187–202.
6. Adewuya AO, Ologun YA. Factors associated with depressive symptoms in Nigerian adolescents. *J Adolesc Health*. 2006 Jul;39(1):105–10.
7. Chinawa JM, Manyike PC, Obu HA, Aronu AE, Odutola O, Chinawa AT. Depression among adolescents attending secondary schools in South East Nigeria. *Ann Afr Med*. 2015 Mar;14(1):46–51.
8. Fatiregun A, Kumapayi TE. Prevalence and correlates of depressive symptoms among in-school adolescents in a rural district in southwest Nigeria. *Journal of adolescence*. 2014 Feb 1;37:197–203.

9. Glied S, Pine DS. Consequences and correlates of adolescent depression. *Arch Pediatr Adolesc Med*. 2002 Oct;156(10):1009–14.
10. Bohman H, Jonsson U, Päären A, Von Knorring AL, Olsson G, Von Knorring L. Long-term follow-up of adolescent depression. A population-based study. *Ups J Med Sci*. 2010 Feb;115(1):21–9.
11. Naicker K, Galambos NL, Zeng Y, Senthilselvan A, Colman I. Social, demographic, and health outcomes in the 10 years following adolescent depression. *J Adolesc Health*. 2013 May;52(5):533–8.
12. Thapar A, Collishaw S, Pine DS, Thapar AK. Depression in adolescence. *Lancet*. 2012 Mar 17;379(9820):1056–67.
13. Kieling C, Adewuya A, Fisher HL, Karmacharya R, Kohrt BA, Swartz JR, et al. Identifying depression early in adolescence. *Lancet Child Adolesc Health*. 2019 Apr;3(4):211–3.
14. Caelear AL, Christensen H. Review of internet-based prevention and treatment programs for anxiety and depression in children and adolescents. *Med J Aust*. 2010 Jun 7;192(S11):S12-14.
15. Cox GR, Fisher CA, De Silva S, Phelan M, Akinwale OP, Simmons MB, et al. Interventions for preventing relapse and recurrence of a depressive disorder in children and adolescents. *Cochrane Database Syst Rev*. 2012 Nov 14;11:CD007504.
16. Zhou X, Teng T, Zhang Y, Del Giovane C, Furukawa TA, Weisz JR, et al. Comparative efficacy and acceptability of antidepressants, psychotherapies, and their combination for acute treatment of children and adolescents with depressive disorder: a systematic review and network meta-analysis. *Lancet Psychiatry*. 2020 Jul;7(7):581–601.
17. Kieling C, Baker-Henningham H, Belfer M, Conti G, Ertem I, Omigbodun O, et al. Child and adolescent mental health worldwide: evidence for action. *Lancet*. 2011 Oct 22;378(9801):1515–25.
18. Patel V. Why adolescent depression is a global health priority and what we should do about it. *J Adolesc Health*. 2013 May;52(5):511–2.
19. Eigenhuis E, Waumans RC, Muntingh ADT, Westerman MJ, Meijel M van, Batelaan NM, et al. Facilitating factors and barriers in help-seeking behaviour in adolescents and young adults with depressive symptoms: A qualitative study. *PLOS ONE*. 2021 Mar 8;16(3):e0247516.
20. Hoven CW, Doan T, Musa GJ, Jaliashvili T, Duarte CS, Ovuga E, et al. Worldwide child and adolescent mental health begins with awareness: a preliminary assessment in nine countries. *Int Rev Psychiatry*. 2008 Jun;20(3):261–70.
21. Seedaket S, Turnbull N, Phajan T, Wanchai A. Improving mental health literacy in adolescents: systematic review of supporting intervention studies. *Tropical Medicine & International Health*. 2020;25(9):1055–64.
22. Wei Y, Hayden JA, Kutcher S, Zygmunt A, McGrath P. The effectiveness of school mental health literacy programs to address knowledge, attitudes and help seeking among youth. *Early Interv Psychiatry*. 2013 May;7(2):109–21.
23. Kutcher S, Bagnell A, Wei Y. Mental health literacy in secondary schools: a Canadian approach. *Child Adolesc Psychiatr Clin N Am*. 2015 Apr;24(2):233–44.
24. Skre I, Friborg O, Breivik C, Johnsen LI, Arnesen Y, Wang CEA. A school intervention for mental health literacy in adolescents: effects of a non-randomized cluster controlled trial. *BMC Public Health*. 2013

Sep 23;13(1):873.

25. Bjørnsen HN, Espnes GA, Eilertsen MEB, Ringdal R, Moksnes UK. The Relationship Between Positive Mental Health Literacy and Mental Well-Being Among Adolescents: Implications for School Health Services. *J Sch Nurs*. 2019 Apr;35(2):107–16.
26. Lam LT. Mental health literacy and mental health status in adolescents: a population-based survey. *Child and Adolescent Psychiatry and Mental Health*. 2014 Sep 25;8(1):26.
27. Atilola O, Ola B, Adewuya A. Towards school-based mental-health programs in Nigeria: Can the current student health-education curricula significantly advance mental health literacy? *Nigerian Journal of Psychiatry*. 2018;16:22–31.
28. Aluh DO, Dim OF, Anene-Okeke CG. Mental health literacy among Nigerian teachers. *Asia Pac Psychiatry*. 2018 Dec;10(4):e12329.
29. Bella T, Omigbodun O, Atilola O. Towards School Mental Health in Nigeria: Baseline Knowledge and Attitudes of Elementary School Teachers. *Advances in School Mental Health Promotion*. 2011 Jan 1;4(3):55–62.
30. Dogra N, Omigbodun O, Adedokun T, Bella T, Ronzoni P, Adesokan A. Nigerian secondary school children's knowledge of and attitudes to mental health and illness. *Clin Child Psychol Psychiatry*. 2012 Jul;17(3):336–53.
31. Bella-Awusah T, Adedokun B, Dogra N, Omigbodun O. The impact of a mental health teaching programme on rural and urban secondary school students' perceptions of mental illness in southwest Nigeria. *J Child Adolesc Ment Health*. 2014;26(3):207–15.
32. Oduguwa AO, Adedokun B, Omigbodun OO. Effect of a mental health training programme on Nigerian school pupils' perceptions of mental illness. *Child Adolesc Psychiatry Ment Health*. 2017;11:19.
33. Aluh DO, Anyachebelu OC, Anosike C, Anizoba EL. Mental health literacy: what do Nigerian adolescents know about depression? *Int J Ment Health Syst*. 2018 Feb 16;12:8.
34. Whitley J, Smith JD, Vaillancourt T. Promoting Mental Health Literacy Among Educators: Critical in School-Based Prevention and Intervention. *Canadian Journal of School Psychology*. 2013 Mar 1;28(1):56–70.
35. Reilly N, Prosper V, Jordan M, Feigenberg L, Capraro K. Break Free from Depression: A 4-Session Curriculum Addressing Adolescent Depression | Suicide Prevention Resource Center [Internet]. 2nd ed. Boston, MA: Boston Children Hospital; 2016 [cited 2021 Jul 4]. Available from: <https://www.sprc.org/resources-programs/break-free-depression-4-session-curriculum-addressing-adolescent-depression>
36. DeMaso D, Gold J. An adolescent mental health & wellness curriculum: A starter kit for schools | Suicide Prevention Resource Center [Internet]. 2006 [cited 2021 Jul 4]. Available from: <https://www.sprc.org/resources-programs/adolescent-mental-health-wellness-curriculum-starter-kit-schools>
37. Haroz EE, Ritchey M, Bass JK, Kohrt BA, Augustinavicius J, Michalopoulos L, et al. How is depression experienced around the world? A systematic review of qualitative literature. *Soc Sci Med*. 2017 Jun;183:151–62.

38. Ibeziako P, Bella T, Omigbodun O, Belfer M. Teachers' perspectives of mental health needs in Nigerian schools. *J Child Adolesc Ment Health*. 2009 Dec;21(2):147–56.
39. Ibeziako PI, Omigbodun OO, Bella TT. Assessment of need for a school-based mental health programme in Nigeria: perspectives of school administrators. *Int Rev Psychiatry*. 2008 Jun;20(3):271–80.
40. Kutcher S, Gilberds H, Morgan C, Greene R, Hamwaka K, Perkins K. Improving Malawian teachers' mental health knowledge and attitudes: an integrated school mental health literacy approach. *Glob Ment Health (Camb)*. 2015;2:e1.
41. Naylor PB, Cowie HA, Walters SJ, Talamelli L, Dawkins J. Impact of a mental health teaching programme on adolescents. *Br J Psychiatry*. 2009 Apr;194(4):365–70.
42. Ma KKY, Anderson JK, Burn AM. Review: School-based interventions to improve mental health literacy and reduce mental health stigma – a systematic review. *Child and Adolescent Mental Health* [Internet]. [cited 2022 Jun 1];n/a(n/a). Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/camh.12543>
43. Kutcher S, Wei Y, Costa S, Gusmão R, Skokauskas N, Sourander A. Enhancing mental health literacy in young people. *Eur Child Adolesc Psychiatry*. 2016 Jun;25(6):567–9.
44. Parikh SV, Taubman DS, Antoun C, Cranford J, Foster CE, Grambeau M, et al. The Michigan Peer-to-Peer Depression Awareness Program: School-Based Prevention to Address Depression Among Teens. *PS*. 2018 Apr;69(4):487–91.
45. Simelane SRN, de Vries PJ. Child and adolescent mental health services and systems in low and middle-income countries: from mapping to strengthening. *Curr Opin Psychiatry*. 2021 Nov 1;34(6):608–16.