

Effects of Biofilm Transfer and Electron Mediators Transfer on *Klebsiella quasipneumoniae* sp.203 Electricity Generation Performance in MFCs

CURRENT STATUS: UNDER REVIEW

Biotechnology for Biofuels  BMC

Yating Guo
Jiangsu Normal University

Guozhen Wang
Jiangsu Normal University

Hao Zhang
Jiangsu Normal University

Hongyu Wen
Jiangsu Normal University

✉ wenhy@jsnu.edu.cn Corresponding Author
ORCID: <https://orcid.org/0000-0001-5061-0569>

Wen Li
Jiangsu Normal University

DOI:

10.21203/rs.3.rs-17738/v1

SUBJECT AREAS

Applied & Industrial Microbiology

KEYWORDS

Microbial fuel cells, Electricity generation performance, Extracellular electron transfer, Biofilm, Electron mediators

Abstract

Background: Extracellular electron transfer (EET) is essential in improving the power generation performance of electrochemically active bacteria (EAB) in MFCs. *Klebsiella* has been proved to be an EAB capable of EET. Here, we cover the anode of MFC-1 with a layer of microfiltration membrane to block the effect of the biofilm mechanism, and then explore the EET of the electron mediator mechanism of *Klebsiella quasipneumoniae* sp.203 and electricity production performance of a *K. quasipneumoniae* sp.203-inoculated MFCs.

Results: Herein, we covered the anode of microbial fuel cells (MFCs) with a layer of microfiltration membrane to block the effect of the biofilm mechanism, and then explore the EET of the electron mediator mechanism of *K. quasipneumoniae* sp.203 and electricity production performance. In the absence of short-range electron transfer, we found that *K. quasipneumoniae* sp.203 can still produce certain power generation efficiency and redox activity. It was proved that in the case that EAB cannot attach to the growth anode, *K. quasipneumoniae* sp.203 can still perform EET through the electron mediator mechanism. To further verify the effect of electron mediators on electrochemical performance of MFCs, in the first cycle of well-functioning MFCs, the self-produced sterile supernatant was added to the different stages of electricity generation performance. We found that adding electron mediators during the rising phase, the MFCs can reach a maximum output voltage of 442mV in about 24 hours, which is about 70mV higher than MFC-Normal. Therefore, the addition of electron mediators can effectively improve the electrical performance in the stable growth stage of anode EAB in MFCs. Finally, we combined the CV analysis and HPLC-MS to analyze the anode supernatant of MFCs. It was speculated that *K. quasipneumoniae* sp.203 produced more than one electron mediator, which were 2,6-DTBHQ, 2,6-DTBBQ, ACNQ and DHNA.

Conclusions: To the best of our knowledge, the three modes of EET did not exist separately. *K. quasipneumoniae* sp.203 will adopt the corresponding electron transfer mode or multiple ways to realize EET according to the living environment to improve electricity generation performance.


Full Text

Due to technical limitations, full-text HTML conversion of this manuscript could not be completed.

However, the manuscript can be downloaded and accessed as a PDF.

Figures

A



B




Figure 1

(a) MFC-1 and MFC-2 were incubated for 680 hours of output voltage, and the anode medium was replaced when the output voltage dropped. (b) Polarization curve (hollow pattern) and power density curve (solid pattern) when the output voltage of the reaches its maximum value.




Figure 2

Nyquist plots from electrochemical impedance spectroscopy measurements of MFC-1 [black empty circle] and MFC-2 [red empty circle] scanned at 0.1~100kHz at open-circuit potential [the inset on the right is the charge transfer resistance in the high frequency region of MFC-2].




Figure 3

Cyclic voltammograms for MFC-1 and MFC-2




Figure 4

SEM images on anode carbon paper at different cycles of MFC-1 and MFC-2, [a]MFC-1: 1st cycle, [b] MFC-1: 3rd cycle, [c]MFC-1: 5th cycle [d]MFC-2: 1st cycle, [e] MFC-2: 3rd cycle, [f]MFC-2: 5th cycle.




Figure 5

Effect of electron mediator secreted by *Klebsiella quasipneumoniae* sp.203 on electricity generation performance of MFCs. [a]the rising phase, b]the gradual phase, [c]the falling phase

— 1st
— 2nd
— 3rd





Figure 6

(a) The CV of the supernatant of the anode on the 1-3 cycle (b) Quinones electron mediators secreted by *Klebsiella quasipneumoniae* sp.203




Figure 7

The transfer mechanism of the electron mediators in a *K. quasipneumoniae* sp.203-inoculated MFCs.