

# Income differences in screening, incidence, postoperative complications, and mortality of thyroid cancer in South Korea: a national population-based time trend study

**Hee-Yeon Kang**

Seoul National University College of Medicine <https://orcid.org/0000-0001-8530-8087>

**Ikhan Kim**

Seoul National University College of Medicine

**Yeon-Yong Kim**

National Health Insurance Service

**Jinwook Bahk**

Keimyung University

**Young-Ho Khang** (✉ [yhkhang@snu.ac.kr](mailto:yhkhang@snu.ac.kr))

Institute of Health Policy and Management, Seoul National University Medical Research Center, Seoul, Republic of Korea <https://orcid.org/0000-0002-9585-8266>

---

## Research article

**Keywords:** Incidence, Income, Medical overuse, Overdiagnosis, Postoperative complications, Thyroid cancer

**Posted Date:** March 24th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-17792/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on November 11th, 2020. See the published version at <https://doi.org/10.1186/s12885-020-07597-4>.

# Abstract

**Introduction:** The incidence of thyroid cancer (TC) has increased rapidly over the past few decades in Korea. This study investigated whether the TC epidemic has been driven by overdiagnosis.

**Methods:** We calculated the TC screening rate from mid-2008 through mid-2014, and the incidence, postoperative complications, and mortality rates of TC between 2006 and 2015, using data from the Korea Community Health Survey, the National Health Insurance Database, and the cause-of-death data of Statistics Korea. Trends in age-standardized rates of all indicators were examined, along with income gaps therein. Analyses were conducted for lung cancer and stroke as negative control outcomes.

**Results:** The incidence rate of TC increased from 46.6 per 100,000 to 115.0 per 100,000 between 2006 and 2012, and then decreased to 63.5 per 100,000 in 2015. Despite these remarkable changes in incidence, mortality did not fluctuate during the same period. High income was associated with high rates of screening, incidence, and postoperative complications, while low income showed an association with a high mortality rate. Analyses using negative control outcomes showed that high income was associated with low rates of both incidence and mortality, which contrasted with the patterns of TC. The recent decreases in TC incidence and postoperative complications, which reflect societal concerns about the overdiagnosis of TC, were more pronounced in high-income individuals than in low-income individuals.

**Conclusions:** The time trends in income gaps in screening, incidence, postoperative complications, and mortality of TC, as well as negative control outcomes, provided corroborating evidence of TC overdiagnosis in Korea.

## Introduction

Thyroid cancer (TC) incidence has increased in many countries around the world over the last few decades [1-3]. Its incidence in South Korea (hereafter 'Korea') has soared to an epidemic level [4-6]. In Korea, the incidence of TC in 2015 was 7 times higher than in 1999, and the incidence peaked in 2012, with a rate 12 times higher than was observed in 1999 [7]. Concerns about the overdiagnosis of TC have been raised in Korea. In 2010, the Korean Thyroid Association revised the management guidelines for patients with thyroid nodules and TC, including limitations on the use of fine needle aspiration cytology [8]. In 2014, the problem of the increase in TC incidence received media attention in Korea [9]. Some Korean doctors publicly pointed out the problems associated with screening for TC, arguing that it was urgent for the Korean government and medical community to restrict overdiagnosis and to take preventive measures against this issue [10]. Recently in Korea, the frequency of surgery for TC nosedived [11]. As a consequence, TC dropped from the most commonly diagnosed type of cancer to the third most commonly diagnosed type of cancer in 2015 in Korea [7]. However, a debate continues over the overdiagnosis of TC in Korea [12].

Overdiagnosis may occur more frequently in high-income strata than in low-income strata because individuals with high income are likely to use more medical services than those with low income, and they

are more exposed to the risk of overutilization than low-income individuals [13, 14]. More sensitive tests tend to be carried out in high-income patients in the screening process, and such measures may lead to an earlier diagnosis, thus increasing the possibility of overdiagnosis [14-16].

This study is based on the hypothesis that the rates of screening, incidence, and postoperative complications of TC would be higher in high-income individuals with greater access to medical care, whereas the mortality rate would show a different pattern. In particular, recently, rising concerns about the overdiagnosis of TC in Korea have led to decreases in both the incidence of TC and surgery for TC [10], and it was hypothesized that high-income individuals would respond more quickly to social concerns. Furthermore, it was assumed that major causes of death—lung cancer (as a representative of cancers) and stroke (as a representative of cardiovascular diseases)—have relatively a low probability of overdiagnosis, and that the incidence and mortality of lung cancer and stroke by income level would therefore show a different pattern from those of TC.

## **Material And Methods**

### **Data**

The study used data from the Korea Community Health Survey (KCHS) conducted in 2010, 2012, and 2014 by the Korea Centers for Disease Control and Prevention, data from the National Health Information Database (NHID) of NHIS between 2001 and 2015, and cause-of-death statistics from Statistics Korea between 2006 and 2015. The KCHS is a community-based cross-sectional health survey that targeted adults over 19 years of age nationwide [17]. The screening rate of TC was drawn from the KCHS items regarding TC screening. The incidence, mortality, and postoperative complication rates of TC, as well as the incidence and mortality rates of the negative control outcomes, were derived from the NHID and cause-of-death data. The NHID includes income-based insurance premiums, demographic variables, information on the date of death, and information on healthcare utilization, thereby making it possible to analyze the incidence of diseases, medical care utilization, and mortality [18]. The study analyzed the mortality rate using individual linkage between the NHID data and cause-of-death data. Individual linkage of the raw data was conducted within the NHIS, and researchers utilized aggregate data without personal identifiers for the analysis.

This study was approved by the National Health Insurance Service (NHIS) of Korea (NHIS-2018-1-191) and the Seoul National University Hospital Institutional Review Board (IRB No. E-1805-047-944). Informed consent was waived by the board.

### **Definitions of thyroid cancer screening, incidence, and mortality**

TC examinees were defined as those who responded “yes” to an item including “thyroid cancer” among questions regarding cancer screening in the most recent 2 years from the KCHS data. To calculate the incidence rate of TC in each year, the study targeted patients who were first diagnosed with TC (ICD-10 code: C73) as a primary or secondary diagnosis every year between 2006 and 2015. The first day of

treatment was considered as the date of diagnosis each year, and the study targeted patients who were admitted for TC once or more in a year or who received outpatient medical services for TC more than twice. However, patients with a history of any type of treatment for TC in the 5 years before the date of diagnosis (e.g., any history between January 1, 2001 and December 31, 2005 for an incident case on January 1, 2006) were considered to have a prior history of TC and were therefore excluded from the calculation of the incidence rate for that year. The number of cases and the age-standardized incidence rates between 2006 and 2015 derived from the NHID using the method described above were very close to those based on the Korea Central Cancer Registry (KCCR) (Additional file 1: Table S1). Since information on income was not available in the KCCR, we used the NHID for this study. Mortality of TC was defined as instances of people over 20 years of age with TC (ICD-10 code: C73) as the cause of death by year between 2006 and 2015.

## **Definition of thyroid cancer surgery-specific complications**

### **Negative control outcomes**

The negative control outcomes were lung cancer and stroke since they were assumed to be less susceptible to overdiagnosis. The incidence and mortality of both lung cancer and stroke were defined similarly to those of TC. In the annual healthcare utilization database of NHID, incident cases of lung cancer were defined as individuals 20 years old or over who were first diagnosed with lung cancer (ICD-10 code: C33-C34) between 2006 and 2015. This study targeted patients who were hospitalized for lung cancer at least once or had outpatient visits twice or more for a year from the diagnosis date, as for cases of thyroid cancer. Then, patients with a past history – those who have been treated for lung cancer during the five years before the diagnosis date – were excluded. The number of cases and the age-standardized incidence rates of lung cancer from the NHID were also very close to those based on the KCCR (Additional file 1: Table S2). Using the annual healthcare utilization database between 2006 and 2015, incident cases of stroke were defined by identifying patients 30 years of age or over who had received their first diagnosis of stroke (ICD-10 code: I60-I64) in that year and including those who were hospitalized with stroke more than once in each year. In addition, patients who had admitted for stroke the prior 5 years in each year between 2006 and 2015 were excluded because they were considered to have a past medical history. As with mortality from thyroid cancer, cases of mortality from lung cancer and stroke were defined based on linkage between the cause-of-death data from Statistics Korea and the NHID data. Those 20 years of age or older with an ICD-10 cause-of-death code of C33-C34 and those 30 years of age or older with an ICD-10 cause-of-death code of I60-I64 were defined as cases of mortality from lung cancer and stroke, respectively.

### **Income**

Household income from the KCHS data and national health insurance (NHI) premiums from the NHID data were used as an income index. Many previous studies in Korea have used the NHI premiums as an indicator of income [20, 21]. For the income index of TC examinees in the KCHS data, this study utilized converted values of equivalized income, in which items related to annual or monthly household income

were adjusted for household size. The income index of the incidence, mortality, and postoperative complications of TC and the negative control outcomes was calculated using equivalized income adjusted for household size using NHI premium data from the NHID. In addition, each equivalized income parameter drawn from the KCHS or NHID was divided into income quintiles by year, sex, and 5-year standardized age group [22].

## Statistical analysis

The changing patterns of the age-standardized screening, incidence, postoperative complication, and mortality rates of TC were analyzed by sex and income level between 2006 and 2015. For age standardization, subjects were divided into 5-year groups, and the 2010 mid-year population from Statistics Korea was used as the reference. Unlike other types of cancer, TC has a very low mortality rate; thus, the annual age-standardized mortality rate by income quintile was derived by combining the data between 2006 and 2010 and the data between 2011 and 2015, respectively, to ensure the stability of TC mortality rates. The incidence and mortality rates of lung cancer and stroke (the negative control outcomes) were derived using age-standardized rates for each income quintile by calendar year between 2006 and 2015. All analyses were performed using SAS version 9.4 (SAS Institute Inc., Cary, NC, USA).

## Results

Table 1 presents the numbers of study subjects aged 20 or over from the database that was used to derive the screening, incidence, postoperative complication, and mortality rates of TC. The number of study subjects in the KCHS data was relatively similar over the years: 225,116 in 2010, 225,188 in 2012, and 224,994 in 2014. The respondents who had received TC screening in the most recent 2 years gradually increased. The number of study subjects in the NHID between 2006 and 2015 has consistently increased, from 35,500,836 in 2006 to 39,696,326 in 2015. From 2006 to 2015, the number of patients with incident TC increased from 15,943 in 2006 to 44,174 in 2012, but then decreased to 25,165 in 2015. The number of patients with postoperative complications of TC surgery increased continually between 2006 and 2010, decreased in 2011, and then increased to its peak of 7,694 in 2013. Later, the number of patients gradually decreased, along with TC incidence. In contrast to the incidence and postoperative complication rate of TC, the mortality of TC remained between 300 and 400 every year, without any significant fluctuations between 2006 and 2015. The magnitudes of the screening, incident, postoperative complications, and deaths cases of TC were much higher for women than men. However, the trends were similar to both women and men. The numbers of patients with the negative control outcomes by year are presented in Additional file 1: Table S3.

Fig. 1. presents the age-standardized incidence and mortality rates of TC in Korea between 2006 and 2015. TC incidence in men and women combined soared from 46.6 to 115.0 per 100,000 from 2006 to 2012, but then plummeted through 2015. TC incidence among women in 2012 was 188.2 per 100,000, reflecting a nearly 2.4-fold increase from 79.1 in 2006. The incidence of TC in men was much lower than in women, but it increased by 3.2 times, from 13.9 per 100,000 in 2006 to 44.0 in 2013. The increasing

trend in TC incidence nosedived starting in 2012 among women and starting in 2013 among men. However, the mortality rate remained very low, at approximately 1.0 per 100,000 regardless of sex, and it showed no significant fluctuations between 2006 and 2015 (Additional file 1: Tables S4 and S5).

Table 2 presents the age-standardized screening, incidence, postoperative complication, and mortality rates of TC by calendar year and income level. In 2010, 2012, and 2014, from 12,562.0 to 18,717.5 adults aged 20 or over per 100,000 had undergone TC screening in the most recent 2 years. Between 2006 and 2015, TC incidence ranged from 46.6 to 115.0 per 100,000, and the postoperative complication rate was 8.9 to 19.7 per 100,000. This indicates that postoperative complications occurred in 14.1% to 20.6% of incident TC cases every year. In contrast, the mortality of TC was stable over time, ranging only from 1.0 per 100,000 between 2006 and 2010 and 0.8 per 100,000 between 2011 and 2015. Every year between 2006 and 2015, the highest income quintile showed high screening and incidence rates of TC. The highest income quintile also showed high rates of postoperative complications associated with TC in most years. However, the mortality rate showed an opposite pattern, as the highest mortality rate occurred in the lowest income quintile.

As shown in Fig. 2., every indicator, except the mortality rate, showed a higher age-standardized rate as income increased (Fig. 2a, b, and c). However, the mortality rate showed the opposite trend (Fig. 2d). The mortality rate of the lowest income quintile was higher than that of the other income quintiles. Furthermore, postoperative complications of TC showed a positive association with income between 2006 and 2013. In 2011, the complication rate decreased in every income group, and then it increased in 2012. The incidence rate began to decrease in 2013, and the complication rate of the highest income quintile nosedived, indicating that the highest-income participants responded more quickly to the decrease in TC incidence than other income quintiles (Fig. 2c).

Fig. 3. presents the age-standardized incidence and mortality rates of lung cancer and stroke by year and income level. Both lung cancer and stroke showed different patterns of incidence and mortality rates from TC. The incidence and mortality rates of lung cancer and stroke continually decreased. During the observation period, TC showed a rapid increase and decrease, but no similar fluctuation was observed for lung cancer (Fig. 3a) and stroke (Fig. 3b). The incidence of lung cancer showed no clear pattern by calendar year, although the mortality rate of lung cancer decreased (Fig. 3a). The incidence and mortality rates of stroke showed a consistent decrease between 2006 and 2015 (Fig. 3b). Every income group also presented a similar pattern. Furthermore, the patterns in incidence rates by income quintiles showed an opposite pattern from that of TC. Whereas TC had a high incidence and low mortality rate in the highest income quintile, for lung cancer and stroke, the lowest income quintile showed the highest rates of incidence and mortality (Fig. 3c, d, e, and f).

## Discussion

The study presented nationally representative rates of TC screening, incidence, postoperative complication, and mortality in Korea. In 2012, the year with the highest incidence of TC, the rate of TC

screening within the most recent 2 years was 17,449.9 per 100,000 adults aged 20 or over, while the incidence of TC was 115 per 100,000. In the same year, 17.9 per 100,000 had postoperative complications of TC and 0.9 per 100,000 died of TC. Incident cases of TC potentially included those who had not undergone TC screening, so these results show that TC screening led to a diagnosis of TC in fewer than 1 of 100 examinees. Furthermore, the complication rate in 2012 was 15.6% of incident TC cases ( $17.9/115 \times 100$ ). Thus, the complication rate was approximately 20 times higher than the mortality rate. It should be noted that this study only examined the rate of TC surgery-specific complications (hypoparathyroidism and vocal cord/fold paralysis) and excluded transient complications within 30 days after surgery. Active surgical interventions for TC are likely to decrease the mortality rate of TC, but this benefit should be balanced with the harm to the population by the postoperative complications of TC surgery. In most cases, the postoperative complications of TC surgery are temporary. However, permanent complications could lead to a low quality of life. Permanent vocal cord paralysis could change an individual's voice, and patients with permanent hypoparathyroidism need to receive calcium and vitamin D for their entire lifetime. In particular, TC surgery performed to vulnerable, high-risk patients, such as elderly patients with comorbidities, could lead to fairly negative results, including low quality of life, pneumonia, infection, heart complications, and death [13, 23]. The findings of this study, therefore, shed light on the magnitude of the side effects of TC surgery.

The incidence of TC showed very significant fluctuations, but the mortality rate showed no notable change. The incidence of TC between 2011 and 2013 increased more than 2-fold from 2006 in all income brackets. This increased incidence in such a short period is difficult to be explained by dietary and environmental risk factors. The increasing trend of TC incidence was more significant in women than in men (Fig. 1.), since ultrasonography for breast cancer screening is often performed jointly with thyroid ultrasonography in Korea. The increase in the incidence of TC and the concomitant lack of change in the mortality rate have been regarded as evidence of the overdiagnosis of TC [4, 24, 25]. In the study, the interpretation of the increase in the incidence of TC as being due to overdiagnosis was further corroborated by data showing that the incidence of TC decreased after 2012, but the mortality rate still showed no changes.

The screening, incidence, and postoperative complication rates of TC were higher in the high-income group than in other income groups. However, a higher mortality rate of TC was shown in the lowest income bracket. According to a previous review study, no clear relationship has been shown between the mortality rate of TC and socioeconomic status [26]. The high mortality rate in the lowest income group might be because this income group included Medical Aid recipients with underlying health problems [20, 21]. Previous studies in Korea have reported greater in-hospital mortality rates among Medical Aid recipients [27].

The incidence of TC decreased after 2012 as a result of concerns about TC overdiagnosis; in particular, the incidence decreased rapidly in the high-income group. Even the postoperative complication rate nosedived in the highest income group. In 2015, the incidence of TC postoperative complications in the highest income quintile had decreased to the level of the lowest income quintile. This could be interpreted

as indicating that high-income individuals responded more rapidly to concerns about the postoperative complications of TC. In 2011, the number of postoperative complications of TC decreased due to the 2010 revision of the management guidelines on thyroid nodules and TC by the Korean Thyroid Association [8]. Active surveillance associated with this 2010 revision might have contributed to the decline in postoperative complications of TC [28].

Lung cancer and stroke were used as negative control outcomes. They showed high incidence and mortality rates in low-income individuals. The high incidence and mortality rates of these diseases in low-income individuals could be explained through socioeconomic inequalities in risk factors, including gaps in the smoking rate across income levels [29] and similar gaps in blood pressure [30], as well as inequalities in the use of medical services and the quality thereof.

The study has strengths and limitations. This study employed large nationally representative data in Korea containing information on screening, healthcare utilization, and mortality. Several studies have been conducted on the incidence and mortality rates of TC, but very few studies have assessed the screening and complication rates. To the best of our knowledge, no study has investigated gaps across income levels in relation to the aforementioned indicators. Analyses by pathological subtype of TC were not conducted in this study. However, according to a medical records review of TC patients between 1962 and 2009 in Korea, the proportion of small tumors less than 1 cm in the total cases of TC increased from 6.1% to 43.1% [31]. In particular, papillary TC, for which overdiagnosis often occurs, accounted for over 90% of all cases of TC [4, 32]. A prior Korean study also showed that the increase in TC from 1999 to 2008 was mainly attributed to the increased incidence of small tumors [33]. In addition, a recent study demonstrated that the TC screening rate was highly associated with the papillary TC incidence rate between 2008 and 2010 across 16 administrative regions of Korea [34]. Even if papillary TC were concentrated in a particular income group, the results on the incidence of TC by income level found in this study would hardly change.

## Conclusions

This study suggests that the considerable fluctuations in the TC incidence over the past decade in Korea were driven by overdiagnosis. The study showed a high screening rate of TC in high-income individuals, leading to high incidence and postoperative complication rates of TC in that group. However, a similar pattern was not shown for the mortality rate. The incidence of lung cancer and stroke, negative control outcomes, did not show a similar pattern to TC by income level. In particular, the recent rapid decrease in the incidence and complication rates in high-income individuals is likely to have been a result of societal concerns over the TC overdiagnosis in Korea. These results provide corroborating evidence that overdiagnosis has caused a considerable proportion of the increase in TC incidence in Korea.

## Abbreviations

TC: Thyroid cancer; KCHS: Korea Community Health Survey; NHID: National Health Information Database; KCCR: Korea Central Cancer Registry

## **Declarations**

### **Ethics approval and consent to participate**

This study was approved by the Seoul National University Hospital Institutional Review Board (IRB No. E-1805-047-944). Informed consent was waived by the board.

### **Consent for publication**

Not applicable.

### **Availability of data and materials**

The Korea Community Health Survey (KCHS) data are publicly available. The National Health Information Database (NHID) that support the finding of this study are available from the Big Data Steering Department of the National Health Insurance Service, but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. The authors used the aggregate population and death data without personal identification numbers. De-identified aggregate data are available from the corresponding author after permissions are obtained from the Big Data Steering Department of the National Health Insurance Service in Korea.

### **Competing interests**

The authors declare that they have no competing interests.

### **Funding**

This study was supported by the Mid-career Researcher Program of the National Research Foundation (NRF) funded by the Ministry of Science & ICT (No. NRF-2014R1A2A1A11051392 and No. NRF-2017R1A2A2A05069746) and a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea [grant number: HI18C0446].

### **Authors' contributions**

YHK conceptualized the study, supervised the data analysis. HYK, IK, YYK contributed to data curation. HYK and IK analyzed and interpreted data. JB designed the analytic strategy. YHK and HYK drafted the manuscript. All authors read and approved the final manuscript.

### **Acknowledgements**

Not applicable

## References

1. Vaccarella S, Franceschi S, Bray F, Wild CP, Plummer M, Dal Maso L. Worldwide Thyroid-Cancer Epidemic? The Increasing Impact of Overdiagnosis. *N Engl J Med*. 2016;375(7):614-7.
2. Franceschi S, Vaccarella S. Thyroid cancer: An epidemic of disease or an epidemic of diagnosis? *Int J Cancer*. 2015;136(11):2738-9.
3. Lortet-Tieulent J, Franceschi S, Dal Maso L, Vaccarella S. Thyroid cancer “epidemic” also occurs in low- and middle-income countries. *Int J Cancer*. 2019;144(9):2082-7.
4. Ahn HS, Kim HJ, Welch HG. Korea's Thyroid-Cancer “Epidemic” – Screening and Overdiagnosis. *N Engl J Med*. 2014;371(19):1765-7.
5. Vaccarella S, Dal Maso L, Laversanne M, Bray F, Plummer M, Franceschi S. The Impact of Diagnostic Changes on the Rise in Thyroid Cancer Incidence: A Population-Based Study in Selected High-Resource Countries. *Thyroid*. 2015;25(10):1127-36.
6. Brito JP, Davies L. Is there really an increased incidence of thyroid cancer? *Curr Opin Endocrinol Diabetes Obes*. 2014;21(5):405-8.
7. Cancer registration statistics: cancer incident cases and incidence rates by site(24 items) and sex. Korean Statistical Information Service(KOSIS), Statistics Korea. 2018. [http://kosis.kr/statHtml/statHtml.do?orgId=117&tblId=DT\\_117N\\_A00022&conn\\_path=I2](http://kosis.kr/statHtml/statHtml.do?orgId=117&tblId=DT_117N_A00022&conn_path=I2). Accessed 31 Jan, 2018.
8. Yi KH, Park YJ, Koong SS, Kim JH, Na DG, Ryu JS, et al. Revised Korean Thyroid Association Management Guidelines for Patients with Thyroid Nodules and Thyroid Cancer. *Endocrinol Metab*. 2010;25(4):270-97.
9. Jung MH, Jung SE. What caused jump in thyroid cancer cases? *The Korea Times*. 2014 March 27, 2014. [http://www.koreatimes.co.kr/www/news/culture/2014/03/319\\_154183.html](http://www.koreatimes.co.kr/www/news/culture/2014/03/319_154183.html). Accessed 19 Jan, 2019.
10. Lee JH, Shin SW. Overdiagnosis and screening for thyroid cancer in Korea. *The Lancet*. 2014;384(9957):1848.
11. Ahn HS, Welch HG. South Korea's Thyroid-Cancer "Epidemic"--Turning the Tide. *N Engl J Med*. 2015;373(24):2389-90.
12. Jeon MJ, Kim HK, Kim EH, Kim ES, Yi HS, Kim TY, et al. Decreasing Disease-Specific Mortality of Differentiated Thyroid Cancer in Korea: A Multicenter Cohort Study. *Thyroid*. 2018;28(9):1121-7.
13. Welch HG, Fisher ES. Income and Cancer Overdiagnosis - When Too Much Care Is Harmful. *N Engl J Med*. 2017;376(23):2208-9.
14. Zagzag J, Kenigsberg A, Patel KN, Heller KS, Ogilvie JB. Thyroid cancer is more likely to be detected incidentally on imaging in private hospital patients. *J Surg Res*. 2017;215:239-44.
15. Brodersen J. How to conduct research on overdiagnosis. A keynote paper from the EGPRN May 2016, Tel Aviv. *Eur J Gen Pract*. 2017;23(1):78-82.

16. Welch HG, Schwartz L, Woloshin S. *Overdiagnosed : making people sick in the pursuit of health*. Boston, Massachusetts: Beacon Press; 2011.
17. Kang YW, Ko YS, Kim YJ, Sung KM, Kim HJ, Choi HY, et al. Korea community health survey data profiles. *Osong public health and research perspectives*. 2015;6(3):211-7.
18. Seong SC, Kim YY, Khang YH, Park JH, Kang HJ, Lee H, et al. Data Resource Profile: The National Health Information Database of the National Health Insurance Service in South Korea. *Int J Epidemiol*. 2017;46(3):799-800.
19. Papaleontiou M, Hughes DT, Guo C, Banerjee M, Haymart MR. Population-Based Assessment of Complications Following Surgery for Thyroid Cancer. *J Clin Endocrinol Metab*. 2017;102(7):2543-51.
20. Khang YH, Yang S, Cho HJ, Jung-Choi K, Yun SC. Decomposition of socio-economic differences in life expectancy at birth by age and cause of death among 4 million South Korean public servants and their dependents. *Int J Epidemiol*. 2010;39(6):1656-66.
21. Khang YH, Bahk J, Yi N, Yun SC. Age- and cause-specific contributions to income difference in life expectancy at birth: findings from nationally representative data on one million South Koreans. *Eur J Public Health*. 2016;26(2):242-8.
22. Kim I, Bahk J, Kim YY, Lee J, Kang HY, Lee J, et al. Comparison of District-level Smoking Prevalence and Their Income Gaps from Two National Databases: the National Health Screening Database and the Community Health Survey in Korea, 2009-2014. *J Korean Med Sci*. 2018;33(6):e44.
23. Grogan RH, Mitmaker EJ, Hwang J, Gosnell JE, Duh QY, Clark OH, et al. A population-based prospective cohort study of complications after thyroidectomy in the elderly. *J Clin Endocrinol Metab*. 2012;97(5):1645-53.
24. Davies L, Welch HG. Current thyroid cancer trends in the United States. *JAMA Otolaryngol Head Neck Surg*. 2014;140(4):317-22.
25. Davies L, Morris LG, Haymart M, Chen AY, Goldenberg D, Morris J, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Disease State Clinical Review: The Increasing Incidence of Thyroid Cancer. *Endocr Pract*. 2015;21(6):686-96.
26. Faggiano F, Partanen T, Kogevinas M, Boffetta P. Socioeconomic differences in cancer incidence and mortality. *IARC Sci Publ*. 1997(138):65-176.
27. Kim Y, Oh J, Jha A. Contribution of hospital mortality variations to socioeconomic disparities in in-hospital mortality. *BMJ Qual Saf*. 2014;23(9):741-8.
28. Haymart MR, Miller DC, Hawley ST. Active Surveillance for Low-Risk Cancers - A Viable Solution to Overtreatment? *N Engl J Med*. 2017;377(3):203-6.
29. Kim I, Bahk J, Yoon TH, Yun SC, Khang YH. Income Differences in Smoking Prevalences in 245 Districts of South Korea: Patterns by Area Deprivation and Urbanity, 2008-2014. *J Prev Med Public Health*. 2017;50(2):100-26.
30. Khang YH, Lynch JW, Jung-Choi K, Cho HJ. Explaining age-specific inequalities in mortality from all causes, cardiovascular disease and ischaemic heart disease among South Korean male public servants: relative and absolute perspectives. *Heart*. 2008;94(1):75-82.

31. Cho BY, Choi HS, Park YJ, Lim JA, Ahn HY, Lee EK, et al. Changes in the clinicopathological characteristics and outcomes of thyroid cancer in Korea over the past four decades. *Thyroid*. 2013;23(7):797-804.
32. Oh CM, Jung KW, Won YJ, Shin A, Kong HJ, Lee JS. Age-Period-Cohort Analysis of Thyroid Cancer Incidence in Korea. *Cancer Res Treat*. 2015;47(3):362-9.
33. Park S, Oh CM, Cho H, Lee JY, Jung KW, Jun JK, et al. Association between screening and the thyroid cancer “epidemic” in South Korea: evidence from a nationwide study. *BMJ*. 2016;355:i5745.
34. Ahn HS, Kim HJ, Kim KH, Lee YS, Han SJ, Kim Y, et al. Thyroid Cancer Screening in South Korea Increases Detection of Papillary Cancers with No Impact on Other Subtypes or Thyroid Cancer Mortality. *Thyroid*. 2016;26(11):1535-40.

## Tables

**Table 1** Study subjects (aged 20 or over) from KCHS and NHID in Korea, 2006-2015

Sex	Year	KCHS			NHID		
		No. of study subjects	No. of subjects with TC screening	No. of total population	No. of TC incident cases	No. of subjects with postoperative complications	No. of TC deaths
All	2006	-	-	35500836	15943	3043	322
	2007	-	-	35881719	20564	3565	358
	2008	-	-	36504109	26474	4450	332
	2009	-	-	36925146	32480	5654	342
	2010	225116	31701	36669946	34981	6946	344
	2011	-	-	37725424	40081	5640	382
	2012	225188	42249	38139862	44174	6937	358
	2013	-	-	38574788	42818	7694	379
	2014	224994	44942	38941467	31923	6610	339
	2015	-	-	39696326	25165	4294	331
Women	2006	-	-	18007925	13629	2631	234
	2007	-	-	18180997	17464	3111	253
	2008	-	-	18507966	22294	3851	230
	2009	-	-	18717880	27307	4924	233
	2010	122432	20011	18589856	28956	5889	254
	2011	-	-	19114446	33125	4754	269
	2012	123971	27322	19326295	36161	5860	235
	2013	-	-	19541003	34362	6269	266
	2014	123226	29248	19717940	25529	5431	256
	2015	-	-	20095936	19768	3449	228
Men	2006	-	-	17492911	2314	412	88
	2007	-	-	17700722	3100	454	105
	2008	-	-	17996143	4180	599	102
	2009	-	-	18207266	5173	730	109
	2010	102684	11690	18080090	6025	1057	90
	2011	-	-	18610978	6956	886	113
	2012	101217	14927	18813567	8013	1077	123
	2013	-	-	19033785	8456	1425	113
	2014	101768	15694	19223527	6394	1179	83
	2015	-	-	19600390	5397	845	103


TC thyroid cancer; KCHS Korea Community Health Survey; NHID National Health Insurance Database

**Table 2** Age-standardized screening, incidence, postoperative complications, and mortality rates of TC by income quintiles in Korea, 2006-2015

Age-standardized rate <sup>a</sup>	Year	Overall	Income Q1 (lowest)	Income Q2	Income Q3	Income Q4	Income Q5 (highest)
Screening prevalence	Mid 2008-mid 2010	12562.0 (12415.9-12708.0)	8824.2 (8515.1-9133.3)	10084.2 (9781.0-10387.5)	11487.7 (11180.1-11795.2)	13057.9 (12734.2-13381.6)	17675.7 (17317.5-18033.9)
	Mid 2010-mid 2012	17449.9 (17279.2-17620.6)	11506.9 (11161.8-11852.0)	13612.7 (13263.2-13962.3)	16305.0 (15933.0-16677.0)	18874.5 (18494.1-19254.8)	24439.6 (24020.1-24859.1)
	Mid 2012-mid 2014	18717.5 (18540.9-18894.2)	11977.5 (11606.0-12349.0)	14934.3 (14574.5-15294.0)	17118.8 (16741.1-17496.4)	20138.9 (19748.2-20529.6)	26110.0 (25675.6-26544.5)
Incidence rate	2006	46.6 (45.8-47.3)	36.4 (35-37.8)	37.8 (36.4-39.3)	42.7 (41.2-44.3)	50.2 (48.5-51.9)	66.0 (64.1-68.0)
	2007	58.7 (57.9-59.6)	43.9 (42.3-45.5)	47.3 (45.7-48.9)	55.0 (53.2-56.7)	66.3 (64.3-68.2)	81.6 (79.4-83.7)
	2008	73.7 (72.8-74.6)	53.7 (52.0-55.4)	59.1 (57.3-60.9)	68.0 (66.1-70.0)	83.6 (81.5-85.7)	104.2 (101.9-106.6)
	2009	88.7 (87.7-89.6)	63.6 (61.8-65.4)	73.5 (71.6-75.5)	82.1 (80.0-84.2)	99.8 (97.5-102.1)	124.4 (121.9-127.0)
	2010	95.4 (94.4-96.4)	69.5 (67.6-71.4)	79.6 (77.5-81.6)	90.1 (88.0-92.3)	106.4 (104.0-108.8)	131.4 (128.8-134.1)
	2011	105.8 (104.8-106.9)	78.0 (76.1-80.0)	91.3 (89.2-93.5)	104.0 (101.7-106.3)	119.1 (116.6-121.5)	136.8 (134.2-139.4)
	2012	115.0 (113.9-116)	84.8 (82.8-86.9)	100.1 (97.8-102.3)	113.3 (110.9-115.7)	128.3 (125.8-130.9)	148.3 (145.6-151.0)
	2013	110.5 (109.4-111.5)	81.0 (79.0-83.0)	97.8 (95.6-100.1)	108.0 (105.7-110.4)	125.9 (123.4-128.4)	139.6 (136.9-142.2)
	2014	81.7 (80.8-82.6)	63.4 (61.6-65.1)	75.5 (73.6-77.5)	81.5 (79.5-83.5)	91.2 (89.1-93.3)	97.2 (95.0-99.4)
	2015	63.5 (62.7-64.2)	50.5 (49.0-52.1)	58.4 (56.7-60.1)	63.5 (61.8-65.3)	71.3 (69.5-73.2)	73.4 (71.5-75.3)
Postoperative complications rate	2006	8.9 (8.6-9.2)	6.5 (5.9-7.1)	7.7 (7.0-8.3)	8.4 (7.7-9.1)	9.8 (9.1-10.6)	12.0 (11.2-12.9)
	2007	10.1 (9.8-10.5)	7.8 (7.2-8.5)	8.1 (7.4-8.7)	10.3 (9.6-11.1)	11.7 (10.9-12.5)	12.9 (12.1-13.8)
	2008	12.4 (12.0-12.8)	10.0 (9.3-10.8)	10.6 (9.8-11.3)	12.3 (11.4-13.1)	13.4 (12.5-14.2)	15.8 (14.9-16.7)
	2009	15.4 (15.0-15.8)	12.3 (11.5-13.1)	13.7 (12.8-14.5)	15.4 (14.5-16.3)	16.5 (15.6-17.4)	19.2 (18.2-20.2)
	2010	18.9 (18.5-19.4)	15.1 (14.2-16.0)	17.3 (16.4-18.3)	18.7 (17.8-19.7)	20.2 (19.2-21.2)	23.2 (22.1-24.3)
	2011	14.8 (14.4-15.2)	11.6 (10.8-12.4)	13.8 (13.0-14.7)	14.8 (13.9-15.6)	16.1 (15.2-17.0)	17.8 (16.8-18.7)
	2012	17.9 (17.5-18.4)	15.3 (14.4-16.2)	16.4 (15.5-17.3)	18.1 (17.2-19.1)	19.3 (18.3-20.3)	20.6 (19.6-21.6)
	2013	19.6 (19.2-20.1)	15.9 (15.1-16.8)	18.0 (17.1-19.0)	20.0 (19.0-21.0)	21.7 (20.6-22.7)	22.5 (21.5-23.6)
	2014	16.8 (16.4-17.2)	14.2 (13.4-15.1)	16.4 (15.5-17.3)	17.7 (16.8-18.6)	18.4 (17.4-19.4)	17.1 (16.2-18.0)
	2015	10.7 (10.4-11.0)	9.6 (8.9-10.3)	9.8 (9.2-10.5)	11.3 (10.5-12.0)	12.1 (11.4-12.9)	10.5 (9.8-11.3)
Mortality rate	2006-2010	1.0 (1.0-1.1)	1.3 (1.1-1.4)	0.9 (0.8-1.0)	1.0 (0.9-1.1)	1.0 (0.9-1.1)	1.0 (0.9-1.1)
	2011-2015	0.8 (0.8-0.9)	1.0 (0.9-1.1)	0.8 (0.7-0.9)	0.8 (0.7-0.9)	0.8 (0.7-0.9)	0.7 (0.7-0.8)


<sup>a</sup>Age-standardized rates per 100,000 were presented with 95% confidence intervals.

## Figures




**Figure 1**

the age-standardized incidence and mortality rates of TC in Korea between 2006 and 2015



## Figure 2

Every indicator, except the mortality rate, showed a higher age-standardized rate as income increased (Fig. 2a, b, and c). However, the mortality rate showed the opposite trend (Fig. 2d). The mortality rate of the lowest income quintile was higher than that of the other income quintiles. Furthermore, postoperative complications of TC showed a positive association with income between 2006 and 2013. In 2011, the complication rate decreased in every income group, and then it increased in 2012. The incidence rate began to decrease in 2013, and the complication rate of the highest income quintile nosedived, indicating that the highest-income participants responded more quickly to the decrease in TC incidence than other income quintiles (Fig. 2c).



**Figure 3**

The age-standardized incidence and mortality rates of lung cancer and stroke by year and income level. Both lung cancer and stroke showed different patterns of incidence and mortality rates from TC. The incidence and mortality rates of lung cancer and stroke continually decreased. During the observation period, TC showed a rapid increase and decrease, but no similar fluctuation was observed for lung cancer (Fig. 3a) and stroke (Fig. 3b). The incidence of lung cancer showed no clear pattern by calendar year,

although the mortality rate of lung cancer decreased (Fig. 3a). The incidence and mortality rates of stroke showed a consistent decrease between 2006 and 2015 (Fig. 3b). Every income group also presented a similar pattern. Furthermore, the patterns in incidence rates by income quintiles showed an opposite pattern from that of TC. Whereas TC had a high incidence and low mortality rate in the highest income quintile, for lung cancer and stroke, the lowest income quintile showed the highest rates of incidence and mortality (Fig. 3c, d, e, and f).

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.docx](#)