

# On the electrochemical NADH regeneration: Influence of electrode potential, pH and NAD<sup>+</sup> concentration

**Emad Aamer**

Hochschule Bremerhaven

**Jorg Thöming**

University of Bremen

**Michael Baune**

University of Bremen

**Nicholas Reimer**

University of Bremen

**Ralf Dringen**

University of Bremen

**Manuela Romero**

University of Bremen

**Ingmar Bösing** (✉ [ingmarboesing@uni-bremen.de](mailto:ingmarboesing@uni-bremen.de))

University of Bremen

---

## Article

### Keywords:

**Posted Date:** July 8th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1792009/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---




# Abstract

Electrochemical  $\text{NAD}^+$  reduction is a promising method to regenerate NADH for enzymatic reactions. Many different electrocatalysts have been tested in the search for high yields of the 1,4-isomer of NADH, the active NADH, but aside from electrode material, other system parameters such as pH, electrode potential and educt concentration also play a role in NADH regeneration. The effect of these last three parameters and the mechanisms behind their influence on NADH regeneration was systematically studied and presented in this paper. With percentages of active NADH ranging from 10% - to 70% and faradaic efficiencies between 1% and 30%, it is clear that all three system parameters drastically affect the reaction outcome. The optimised reaction conditions were applied to test the consumption of the regenerated NADH by an enzymatic reaction. Electrochemical  $\text{NAD}^+$  reduction in the presence of pyruvate and lactate dehydrogenase revealed that the 1,4-isomer of NADH, the physiological substrate of oxidoreductases, accounted for around 45% of the detectable electrochemically regenerated NADH.

## 1. Introduction

Nicotinamide adenine dinucleotide (NADH) is an enzymatic cofactor that plays an essential role in many biochemical reactions [1]. Enzymatic reactions catalysed by many dehydrogenases rely on NADH as a cofactor for electrons and hydrogen shuttle. In nature, NADH is found in all living cellular organisms and is oxidised during cellular respiration [2]. Industrially it is used in the preparation of organic chiral compounds, for the aerobic transformation of chromium (VI), which is toxic, to chromium (III) [3]. There are numerous other uses of NADH which makes it an extremely important compound [4]. However, the industrial use of NADH is limited due to its high cost and low availability. Therefore, it is important to regenerate NADH from its oxidised form  $\text{NAD}^+$  to make its industrial use feasible [5]. Several methods of NADH regeneration have been tested so far including chemical, photochemical, biological, and electrochemical processes [6]. The electrochemical process for regenerating NADH is of great importance as the operational costs are low, chemical reducing agents are unnecessary, the reaction can be monitored and controlled easily by applying a potential and measuring current, and operational windows can be found, to reduce byproducts during the synthesis, that must be later separated from the product itself [7].

The electrochemical reduction of  $\text{NAD}^+$  to NADH proceeds in two steps [2,8]:



The electrochemical reduction of  $\text{NAD}^+$  on bare metal electrodes at significantly high cathodic overpotentials takes place through one-electron transfer followed by a rapid reaction of the  $\text{NAD}^{\bullet}$  radicals

to form the inactive dimer  $\text{NAD}_2$  <sup>[7,9]</sup> (step 2b). The NADH regeneration-reaction crucial step consists of a second electron transfer followed by protonation (step 2a). Reaction step 2a can not only yield the desired active 1,4-NADH but also enzymatically inactive isomers 1,2-NADH and 1,6-NADH. This reaction step is slow compared to reaction step 2b which is mainly why the direct electrochemical regeneration of NADH on bare metal electrodes often yields high amounts of the enzymatically inactive dimer  $\text{NAD}_2$  <sup>[2,10-12]</sup>. In another study <sup>[13]</sup>, Jaegfeldt observed that the second electron transfer occurs only at more (-1.7V) cathodic potentials compared to the first (-1.1V).

It could already be shown that the choice of the electro-catalysts for the electrochemical NADH regeneration plays an important role: on bare copper electrodes, Barin et al. achieved around 50 % and on copper foam around 80 % active 1,4-NADH <sup>[10]</sup>, while Damian could regenerate up to 80 % active NADH on bare gold electrodes <sup>[8]</sup>. The electrochemical regeneration of NADH on titanium <sup>[2]</sup> or modified glassy carbon electrodes <sup>[5]</sup> was reported to yield up to 98 % active NADH. Unfortunately, no information on the faradaic efficiency and the mole yield was given.

Not only the catalyst material plays a significant role in the NADH regeneration but also the electrode potential. As mentioned above, Jaegfeldt reported more negative potentials necessary for the second electron transfer step, which favours reaction step 2a and the protonation. On the other hand, the hydrogen evolution reaction (HER), which can be a competing reaction to the  $\text{NAD}^+$  reduction reaction, is also favoured at more cathodic potentials. It can be seen from the reaction equation that the reaction kinetics of step 2a depends upon the availability of  $\text{H}^+$  on the reaction site. If the availability of  $\text{H}^+$  on the surface of the electrode is decreased by consumption due to HER, it might diminish the overall NADH formed.

Therefore, it is not surprising that many authors found a dependency of the ratio of active NADH on the electrode potential. Ali et al. reported surprisingly high amounts of 1,4-NADH yield (98 %) at a high negative potential of -1.8 V vs  $\text{Ag}|\text{AgCl}|\text{KCl}_{3\text{M}}$  <sup>[14]</sup> with glassy carbon electrodes (our own experiments could not reproduce this yield). Contrary to their observations, Jaegfeldt et al. reported around 50 % 1,4-NADH, 30 % 1,6-NADH and 20% dimers at -1.7 V vs  $\text{Ag}|\text{AgCl}|\text{KCl}_{3\text{M}}$  <sup>[13]</sup>. Damian reported a decreasing yield of active NADH with increasing electrode potential <sup>[8,9]</sup> and explained this observation with the higher amount of  $\text{NAD}^\bullet$  radicals due to the enhanced reaction kinetics and transport at more cathodic potentials favouring dimerization. Barin et al. showed an initial increment in the yield of active NADH with decreasing potential and with further decreasing potential eventually a decreased yield of active NADH <sup>[11]</sup>.

Unfortunately, most of the studies do not discuss the faradaic efficiency of the process or quantify the concentrations of the regenerated active NADH, even though both are relevant parameters to evaluate the applicability of the process. Furthermore, many studies investigate the electrochemical NADH regeneration in phosphate buffer, even though it has been shown that NADH is unstable in this buffer solution <sup>[15]</sup>.

An overview of the current literature and the reaction scheme suggests that there are (next to the catalysts material) at least three system parameters that can affect the yield of active NADH: the electrode potential, the availability of  $H^+$ , and the concentration of  $NAD^{\bullet}$  radicals at the electrode, which is (among others) determined by the  $NAD^+$  concentration and the kinetic of the first electron transfer (Reaction 1). The latter, which is favoured by a high concentration of  $NAD^+$  or a high material transport to the electrode (caused by – among others – a high cathodic potential), might enhance the dimerization (as can be seen from reaction step 2b. The availability of  $H^+$  is essential for the protonation of the radicals (reaction step 2a), while the electrode potential controls the reaction kinetics of the NADH regeneration and the HER, which in turn might also influence the concentration of  $NAD^{\bullet}$  radicals at the electrode surface and the  $H^+$  availability.

In the study presented here, these assumptions have been tested by systematically changing the solution's pH and  $NAD^+$  concentration while studying the yield of active NADH for different electrode potentials on a bare copper electrode. Furthermore, the faradaic efficiency of the process has been calculated and the molar yield is presented as well as the dependency on all three system parameters – electrode potential, pH and substrate concentration, on the electrochemical NADH regeneration.


Additionally, we propose that it can be advantageous to combine the electrochemical regeneration of NADH with enzyme-catalysed reactions, using the regenerated cofactor directly to produce valuable chemical compounds, e.g. for industrial use. This has been done for reducing reactions to facilitate the production of chemicals in an NADH-dependent manner, for example by reducing  $CO_2$  to formate with the formate dehydrogenase<sup>[16]</sup>. To test, if enzyme activity can be maintained under the applied potential and during the NADH regeneration, lactate dehydrogenase (LDH) and pyruvate are added to the reaction chamber.

## 2. Experimental

All electrochemical measurements were carried out in a three-electrode H-cell containing 0.1 M Tris buffer at room temperature, where anodic and cathodic half cells were separated by a glass frit. A copper sheet (99.9 % purity) of size 1.5 cm × 1.5 cm was used as the working electrode. Before each measurement, it was ground and polished with diamond paste (6  $\mu m$ ) and cleaned with ethanol and deionized water. An Ag/AgCl (3 mM KCl) electrode served as reference electrode while a platinum sheet electrode was used as counter electrode. The electrochemical cell was purged with nitrogen for at least 15 min before each measurement and the subsequent measurements were carried out in a nitrogen atmosphere. The potentiostat used was a Metrohm Autolab PGSTAT204 with the Nova software package. The Linear Sweep Measurements were performed with a scan rate of 0.1 V/s between 0 V and -1.5 V (vs Ag/AgCl). All potentials in this paper refer to an Ag/AgCl electrode, where cathodic potentials are negatively drawn. The regeneration measurements were carried out at constant potentials for 1800 seconds.

The determination of active NADH was done with an enzyme test involving lactate dehydrogenase (LDH), pyruvate, and consequent measurements of the extinction at 340 nm with a UV/VIS spectrometer.

The addition of pyruvate and LDH leads to the reaction of enzymatically active 1,4-NADH as follows:



Three possible products of the  $\text{NAD}^+$  reduction, 1,4-NADH, 1,6-NADH and  $\text{NAD}_2$  show extinction at 340 nm [6,9] making it impossible to determine the percentage of active 1,4-NADH by UV/VIS measurements alone. By measuring the extinction after the  $\text{NAD}^+$  reduction (while the solution contains 1,4-NADH, 1,6-NADH and  $\text{NAD}_2$ ) and after the addition of LDH and pyruvate, when all 1,4-NADH is consumed by the reaction with pyruvate, and calculating the difference of both extinctions (before addition of LDH – including 1,4-NADH and after addition of LDH – without 1,4-NADH), the percentage of active NADH can be calculated. It should be mentioned that the addition of pyruvate also increases the extinction at 340 nm, which must be then subtracted from the results.

Enzymatic utilization of regenerated NADH was determined during electrochemical NADH regeneration under a potential of -1.3 V at a solution pH of 7 with 0.75 mM  $\text{NAD}^+$ , and 5 mM pyruvate with and without 2 U/mL LDH. The total volume in the reaction chamber was 15 mL. During regeneration, samples of 300  $\mu\text{L}$  were taken from the reaction chamber at regular time intervals. Each measurement was repeated at least three times.

## 3. Results And Discussion

### 3.1 Buffer system

Many research groups working on the electrochemical reduction of  $\text{NAD}^+$  have used phosphate as the buffer system [5,7,10–12]. This is not advisable, as it was shown that NADH is unstable in phosphate buffer [15]. An increased degeneration of NADH in combination with an enzyme assay to test the ratio of active NADH will lead to an overestimation of active NADH since the degeneration of NADH and  $\text{NAD}_2$  due to the buffer system will lead to a decreasing signal in the UV/VIS at 340 nm. Furthermore, LSV measurements in phosphate buffer show a reduction peak due to the reduction of phosphate groups at the same potential as the  $\text{NAD}^+$  reduction occurs (Fig 1), which will lead to a worse faradaic efficiency.

To test if NADH shows better stability in Tris buffer, phosphate and Tris buffer solutions at different pH (6.8, 7.2, 7.6) values containing 1.0 mM NADH were prepared and the extinction at 340 nm was measured over time by UV/VIS measurements. While the extinction decreases drastically in phosphate buffer (nearly 60 % of start extinction after 1300 min), the extinction remained constant in Tris buffer (Fig. 2), indicating better stability of NADH in Tris buffer over time.

### 3.2 Reduction of $\text{NAD}^+$

Linear sweep voltammetry (LSV) was performed to identify the electrochemical behaviour of NAD<sup>+</sup> reduction at different pH values and to identify the potential where hydrogen evolution reaction starts. Fig. 3a and Fig. 3b show the LSV taken on a copper electrode in the absence and the presence of NAD<sup>+</sup>, respectively. The LSV was performed in the potential region -0 V to -1.5 V vs Ag/AgCl (3 M KCl). An increase in current density at higher cathodic potential is due to hydrogen evolution reaction (HER), which is kinetically favoured at higher cathodic potential, and at lower pH values as observed in Fig 3a. The current density at a certain pH is lower (less cathodic) when NAD<sup>+</sup> is not present in the electrolyte. However, when NAD<sup>+</sup> is added to the electrolyte, a significant increase in current density results due to the reduction of NAD<sup>+</sup>. The reduction potential of NAD<sup>+</sup> as seen from the increase of cathodic current was in agreement with prior research [5,9,10,17]. In the presence of NAD<sup>+</sup>, reduction starts at around -0.9 V. However, Fig. 3 suggests that to achieve the NAD<sup>+</sup> reduction at an appreciable rate, higher cathodic potentials are required.

### 3.3 Influence of pH

The electrochemical behaviour of NAD<sup>+</sup> at different potentials was previously studied [2,9]. It has been discussed that the yield of enzymatically active NADH depends upon the electrode potential and the electrode material [9]. Both the electrode's material and potential directly affect the amount of adsorbed hydrogen H<sub>ads</sub> at the surface of the electrode which is available for NAD<sup>•</sup> radicals to form NADH, and the hydrogen evolution reaction. However, the availability of H<sup>+</sup> and thus the amount of adsorbed H<sub>ads</sub> can also be affected by changing the pH of the electrolyte. Thus, if the above-stated hypothesis is true – the adsorbed H<sub>ads</sub> affect the NADH regeneration - one should observe a dependency of active 1,4-NADH on the solution's pH.

To study the influence of electrolyte's pH several experiments were carried out in the pH range between 6.5 and 9.0 in a batch electrochemical cell using Cu as the working electrode. The percentage of enzymatically active 1,4-NADH formed was determined by an enzyme assay as described in the experimental section.

The percentage of enzymatically active 1,4-NADH regenerated strongly depends upon the pH of the electrolyte as well as the electric potential (Fig. 4). At the highest pH (pH 9.0) increasing the cathodic potential from -0.9 V to -1.5 V results in an increment in the average amount of active 1,4-NADH. At low cathodic potentials, the first electron transfer occurs drastically faster compared to the second electron transfer, thus NAD<sup>•</sup> radicals form at the electrode surface and their dimerization is faster than the second electron transfer. Furthermore, at low cathodic potentials (and high pH values) the adsorption of hydrogen to the electrode surface by the Volmer reaction [18] is very low, thus the availability of adsorbed hydrogen for the formation of NADH is limited. With increasing electrode potential, the reaction kinetics of the second electron transfer step and the hydrogen adsorption at the electrode surface increase. Therefore, with increasing cathodic potential the percentage of active 1,4-NADH increases at pH 9.

With decreasing pH and potential until -1.1 V (in case of pH 6.5) or -1.3 V (in case of pH 7 and pH 8) there is a visible increase of active NADH. But by further decreasing the potential a decrease in active NADH occurs. This is due to an increase in the HER, which limits the amount of  $H_{ads}$  necessary for the NADH regeneration. At lower pH and potential values, the reaction kinetics of the HER are faster than reaction step 2a.

As an overall trend from pH 9 to pH 7, with decreasing pH, an increasing percentage of active NADH can be seen. With further decreasing pH to 6.5 the percentage of active NADH decreases due to the accelerated HER. This observation is further supported by the course of the faradaic efficiency (FE) for the 1,4-NADH regeneration at different potentials and pH values (Figure 4b): with increasing cathodic potential from -0.9 V to -1.5 V the faradaic efficiency drastically drops for all pH values, due to the accelerated HER and due to a large number of electrons being used for the non-desirable HER and formation of  $NAD_2$ . At -1.5 V the highest faradaic efficiency can be observed for pH 9 (slowest HER) and the lowest FE can be observed at pH 6.5 (fastest HER). At an electrode potential of -0.9 V, one can observe the lowest FE at pH 9 (lowest availability of  $H_{ads}$  and thus favoured dimerization), while the highest FE can be observed at pH 6.5 (high  $H_{ads}$  availability).

Due to the interaction of both competing reactions, HER and NADH regeneration, which are influenced by the electrode potential, for each pH value a specific potential depending peak of active NADH can be found (Figure 5). This tipping potential – the potential at which the highest percentage of active 1,4-NADH can be found and at which a further decrease of electrode potential will lead to a worse yield of active NADH, is strongly pH-dependent. The higher the pH, the more cathodic the tipping potential. Accordingly, the most cathodic tipping potential (-1.5 V) can be found for pH 9 and the least cathodic tipping potential (-1.1 V) for pH 6.5.

The total amount of NADx (all NADH isomers and  $NAD_2$  dimer) generated during 30 minutes of electrochemical reaction depends on pH as well as the electrode potential (Figure 6a). Lowering the electrode potential from -0.9 V to more cathodic potentials increases the number of moles generated at all different electrolyte pH values, due to the enhanced reaction kinetics. However, it is also shown that if the pH is increased, while the electrode potential is kept constant, the moles of NADx first increase from pH 6.5 to pH 7, and later decrease between pH 7.0 and pH 9.0. With decreasing pH, the HER is favoured over the  $NAD^+$  reduction, which leads to a drop of regenerated NADx and pH 7.0 seems to be the ideal combination of available  $H_{ads}$  and low HER.

The data obtained for the amounts of active 1,4-NADH formed within 30 min at different pH-values are similar to that observed for the sum of all NADx species (Figure 6b). But in contrast to the overall amounts produced, the dependency on the electrode potential shows a difference. A maximum of active 1,4-NADH was observed at -1.3 V, since more cathodic potentials (-1.5 V) lead to faster Hydrogen Evolution Reaction and result in higher dimerization due to the lack of adsorbed  $H_{ads}$ . At any given potential, pH 7.0 regenerated the maximum amount of 1,4-NADH. At lower pH than 7.0, HER reaction is

favoured over reaction step 2a, while at pH values higher than 7.0 dimerization is kinetically favoured due to less availability of  $H^+$ .

### 3.4 Influence of $NAD^+$ concentration

Damian et al. showed that increasing the concentration of  $NAD^+$  in the bulk solution increases the surface coverage of the  $NAD^+$  on the electrode rapidly [9]. Increasing the concentration further would level off indicating that the saturated monolayer  $NAD^+$  coverage is reached [8]. But this is only true for less cathodic potentials where the competing reactions are not taking place. However, if the cathodic potential is higher or near the  $NAD^+$  reduction potential, then the reduction reaction is faster than the adsorption of  $NAD^+$  molecules on the electrode surface [19]. Hence, the reduction of  $NAD^+$  will be diffusion controlled at higher cathodic potentials.

The more cathodic the potential, the faster the formation of radicals and the higher the mass transport towards the electrode's surface. This may lead to an accumulation of  $NAD^\bullet$  radicals on the electrode's surface, which would lead to an accelerated formation of  $NAD_2$  dimers since reaction step 2b depends on the dimer concentration. If this – next to the accelerated HER and the depletion of adsorbed hydrogen – is a further reason for the decrease of active 1,4-NADH at high cathodic potentials, this must be a concentration-dependent phenomenon. As a consequence, NAD dimerization should be decreased with lower  $NAD^+$  concentration which should lead to a percental enrichment of  $NAD_x$  in active 1,4-NADH.

By decreasing the  $NAD^+$  concentration from 3 mM stepwise down to 0.1 mM, the percentage of active 1,4-NADH increases significantly for all electrode potentials (Fig 7a). The tipping potential – the potential at which a more cathodic potential will lead to a decreasing 1,4-NADH percentage – of the 3 mM solution is -1.1 V whereas the tipping potential for the remaining concentrations is -1.3 V. At the lowest concentration, 0.1 mM  $NAD^+$ , the difference of active NADH between -1.3 V and -1.5 V is smallest. These observations support the hypothesis that the reaction in step 2b, the dimerization, is a concentration-dependent reaction.

The effect of concentration on faradaic efficiency displays a less clear trend (Fig. 7b). The lowest concentration of  $NAD^+$  shows the lowest faradaic efficiency, which might be due to a low  $NAD^+$  concentration at the electrode's surface and thus elevated HER compared to  $NAD^+$  reduction. In addition, only for a concentration of 1.5 mM, the FE decreases steadily with higher cathodic potentials. In the case of the remaining concentrations, the highest FE can be found at -1.1 V, which might be due to the combination of relatively high 1,4-NADH yield and low HER.

### 3.5 Enzymatic utilisation of produced NADH during regeneration



To test whether optimised reaction conditions for NADH regeneration provide sufficient amounts of 1,4-NADH (active NADH) to continuously allow an enzymatic reduction reaction, LDH was either added or not to the reaction chamber, containing 0.75 mM  $\text{NAD}^+$  and 5 mM pyruvate, before the electrochemical NADH regeneration was started. For different regeneration times, samples were taken from the reaction chamber and their absorption at 340 nm was recorded (Fig. 8).

During the NADH regeneration, a clear increase in absorption over time was observed for both reactions with and without LDH. Nevertheless, for each time point investigated, the absorbance values for LDH- and pyruvate-containing reactions were significantly lower, representing only around 55 % of the values determined for the reactions without the enzyme. This suggests that around 45 % of the regenerated  $\text{NADx}$  is consumed by the LDH and thus is active 1,4-NADH. This shows that a coupling of the electrochemical regeneration and enzymatic reaction in the same system is possible.

An amount of 45 % active NADH is significantly lower compared to around 60 % achieved under similar conditions without the presence of LDH and pyruvate. On the one hand, this could be explained by the fact that enzymatic reactions need time; and that the electrochemically regenerated 1,4-NADH, continuously produced at the electrode, would require additional time to get coupled with the LDH for lactate formation. On the other hand, the enzyme may not work optimally in the presence of the copper electrode with applied potential. It is known that copper ions, which can be released by the copper electrode<sup>[20]</sup>, can reduce the activity of enzymes. Whether an applied potential and an ongoing electrochemical reaction affect or not the catalytic behaviour of enzymes will be explored by us in the future.

## 4. Conclusion

Three different parameters influencing the electrochemical NADH regeneration and the yield of active 1,4-NADH from this reaction were tested: The solution's pH was changed, which affected the availability of protons and adsorbed  $\text{H}_{\text{ads}}$  at the electrode's surface, necessary for the formation of  $\text{NAD}^{\bullet}$  radicals (reaction step 2a). The  $\text{NAD}^+$  concentration was changed, which affected the concentration of newly formed radicals at the electrode's surface and thus the possibility of radical dimerization to  $\text{NAD}_2$  according to reaction step 2b. As for the third parameter, the electrode potential was changed for all tested pH values and concentrations. The electrode potential controls the reaction kinetics and delivers the energy for the first electron transfer (reaction step 1), which affects the  $\text{NAD}^{\bullet}$  radical concentration and the second electron transfer (reaction step 2a) for the production of NADH. In addition, the potential controls the HER which affects the availability of protons and adsorbed hydrogen  $\text{H}_{\text{ads}}$ .

Changing the pH of the solution shows a significant effect on the percentage of 1,4-NADH. The combination of electric potential and the solution's pH controls the HER which affects the NADH regeneration. Thus, the tipping potential for maximum active NADH shifts with increasing pH to more cathodic potentials.

By lowering the  $\text{NAD}^+$  concentration a higher percentage of active NADH can be achieved. This is due to an overall lower  $\text{NAD}^\bullet$  radical concentration at the electrode surface and thus, a less favoured dimerization.

Even on a bare metal electrode the range of active NADH could yield between approx. 10 % and 70 % and the faradaic efficiency between approx. 1 % and 30 % simply by adjusting different system parameters. The understanding of the parameters affecting the electrochemical NADH regeneration, presented here, will help to optimise conditions which may allow finally to bring the process from laboratory to application scale.

Lastly, LDH and pyruvate were added to the reaction chamber during the regeneration of NADH from  $\text{NAD}^+$ . The data from this experimental setup suggests that the enzyme could use the 1,4-isomer formed continuously during the NADH regeneration and catalyse the reaction converting pyruvate to lactate. Future experiments should further explore how to improve the yield of active 1,4-NADH during electrochemical  $\text{NAD}^+$  reduction in the presence of enzymes to enhance the online use of regenerated, active NADH for enzymatic reduction reactions.

## Declarations

Data availability

All data will be available on reasonable request from the corresponding author.

Acknowledgments

This project was funded by the European Regional Development Fund (ERDF). Furthermore, we thank Sven Kerzenmacher for fruitful scientific discussions on our project.

## References

1. Sellés Vidal, L., Kelly, C. L., Mordaka, P. M. & Heap, J. T. Review of NAD(P)H-dependent oxidoreductases: Properties, engineering and application. *Biochimica et Biophysica Acta - Proteins and Proteomics* vol. 1866 327–347 (2018).
2. Ali, I., Khan, T. & Omanovic, S. Direct electrochemical regeneration of the cofactor NADH on bare Ti, Ni, Co and Cd electrodes: The influence of electrode potential and electrode material. *Journal of Molecular Catalysis A: Chemical* **387**, 86–91 (2014).
3. Albo, J., Perfecto-Irigaray, M., Beobide, G. & Irabien, A. Cu/Bi metal-organic framework-based systems for an enhanced electrochemical transformation of  $\text{CO}_2$  to alcohols. *Journal of  $\text{CO}_2$  Utilization* **33**, 157–165 (2019).
4. Karyakin, A. A., Ivanova, Y. N. & Karyakina, E. E. Equilibrium ( $\text{NAD}^+/\text{NADH}$ ) potential on poly(Neutral Red) modified electrode. *Electrochemistry Communications* **5**, 677–680 (2003).

5. Ali, I., Gill, A. & Omanovic, S. Direct electrochemical regeneration of the enzymatic cofactor 1,4-NADH employing nano-patterned glassy carbon / Pt and glassy carbon / Ni electrodes. *Chemical Engineering Journal* **188**, 173–180 (2012).
6. Saba, T., Burnett, J. W. H., Li, J., Kechagiopoulos, P. N. & Wang, X. A facile analytical method for reliable selectivity examination in cofactor NADH regeneration. *Chemical Communications* **56**, 1231–1234 (2020).
7. Ho Baik, S., Kang, C., Cheol Jeon, I. & Eok Yun, S. *Direct electrochemical regeneration of NADH from NAD<sup>+</sup> using cholesterol-modified gold amalgam electrode. Biotechnology Techniques* vol. 13 (1999).
8. Damian, A. & Omanovic, S. Electrochemical reduction of NAD<sup>+</sup> on a polycrystalline gold electrode. *Journal of Molecular Catalysis A: Chemical* **253**, 222–233 (2006).
9. Damian, A., Maloo, K. & Omanovic, S. Direct electrochemical regeneration of NADH on Au, Cu and Pt-Au electrodes. *Chemical and Biochemical Engineering Quarterly* **21**, 21–32 (2007).
10. Barin, R., Rashid-Nadimi, S., Biria, D. & Asadollahi, M. A. Direct electrochemical regeneration of 1,4-NADH at the copper foam and bimetallic copper foam. *Electrochimica Acta* **247**, 1095–1102 (2017).
11. Azem, A., Man, F. & Omanovic, S. Direct regeneration of NADH on a ruthenium modified glassy carbon electrode. *Journal of Molecular Catalysis A: Chemical* **219**, 283–299 (2004).
12. Moiroux, J. & Elving, P. J. Adsorption phenomena in the NAD<sup>+</sup>/NADH system at glassy carbon electrodes. *Journal of Electroanalytical Chemistry* **102**, 93–108 (1979).
13. Jaegfeldt, H. A study of the products formed in the electrochemical reduction of nicotinamide-adenine-dinucleotide. *Journal Electroanalytical Chemistry* **128**, 355–370 (1981).
14. Ali, I., Soomro, B. & Omanovic, S. Electrochemical regeneration of NADH on a glassy carbon electrode surface: The influence of electrolysis potential. *Electrochemistry Communications* **13**, 562–565 (2011).
15. Wu, J. T., Wu, L. H. & Knight, J. A. Stability of NADPH: Effect of Various Factors on the Kinetics of Degradation Materials and Methods. *Clin. Chem. Clinical Chemistry* **32232**, 314–319 (1986).
16. Barin, R., Biria, D., Rashid-Nadimi, S. & Asadollahi, M. A. Enzymatic CO<sub>2</sub> reduction to formate by formate dehydrogenase from *Candida boidinii* coupling with direct electrochemical regeneration of NADH. *Journal of CO<sub>2</sub> Utilization* **28**, 117–125 (2018).
17. Kim, S. *et al.* Electrochimica Acta Electrochemical NADH regeneration and electroenzymatic CO<sub>2</sub> reduction on Cu nanorods / glassy carbon electrode prepared by cyclic deposition. *Electrochimica Acta* **210**, 837–845 (2016).
18. M. R. Gennero, de C. & Cesar, A. C. Hydrogen evolution reaction: Analysis of the Volmer-Heyrovsky-Tafel mechanism with a generalized adsorption model. *Journal of Electroanalytical Chemistry*, **372**, 209–223 (1994).
19. Damian, A. & Omanovic, S. Interactive adsorption behavior of NAD<sup>+</sup> at a gold electrode surface. *Langmuir* **23**, 3162–3171 (2007).

20. van Assche, F. & Clijsters, H. Effects of metals on enzyme activity in plants. *Plant, Cell and Environment* **13**, 195–206 (1990).

## Figures




Figure 1

Linear potential sweep from -0.4 V to -1.4 V with the scan rate of 0.1 V/s vs. Ag|AgCl|KCl3M in 0.1 M phosphate buffer pH 7 with and without  $\text{NAD}^+$ .




Figure 2

Extinction at 340 nm in the percentage of start extinction over time for 0.1 mM NADH in Tris buffer (  $\Delta$  ) and Phosphate buffer ( O ) (both pH 6.8).






Figure 3

LSV taken using a copper electrode in 0.1 M Tris buffer flushed with  $N_2$  at pH 6.5, pH 7, pH 8 and pH 9 with a scan rate of 0.1 V/s. **a)** without  $NAD^+$  **b)** with 1.5 mM  $NAD^+$ .



**Figure 4**

Influence of pH and electrode potential on the NADH regeneration. **a**) Percentage of active 1,4-NADH produced with a copper electrode at potential (-0.9 V to -1.5 V) vs. Ag|AgCl|KCl<sub>3M</sub> and using 1.5 mM NAD<sup>+</sup> in Tris buffer for 1800 sec. **b**) Faradaic efficiency of the NADH regeneration.



**Figure 5**

Normalised amount of active NADH over the electrode potential for different solution pHs. Arrows indicate the tipping potential for the different solution pHs. The normalised active NADH was calculated by dividing the active NADH by the maximum active NADH of each pH.




Figure 6

Moles of reaction products after 30 min of  $\text{NAD}^+$  reduction depending on the electrode potential for different pH. **a)** NADx (NADH isomers and  $\text{NAD}_2$ ); **b)** 1,4 NADH.





Figure 7

Concentration depending on NADH regeneration for different electrode potentials. **a)** Percentage of active NADH depending on  $\text{NAD}^+$  concentration and electrode potential; **b)** Faradaic efficiency of 1,4-NADH

regeneration depending on  $\text{NAD}^+$  concentration and electrode potential.



**Figure 8**

See image above for figure legend