Measurement of Phthalate Acid Esters in Non-alcoholic Malt Beverages by MSPE-GC/MS Method in Tehran City: With the Approach of Chemometric

Hana Rezaei

TUMS: Tehran University of Medical Sciences

Mojtaba Moazzen

Shahid Beheshti University of Medical Sciences School of Nutrition Sciences and Food Technology

Nabi Shariatifar (✉ nshariati@tums.ac.ir)

Tehran University Of Medical Sciences <https://orcid.org/0000-0002-2861-5758>

Gholamreza Jahed Khaniki

Tehran University of Medical Sciences Dental School

Mohammad Hadi Dehghani

Tehran University of Medical Sciences

Majid Arabameri

Semnan University of Medical Sciences and Health Services

Mahsa Alikord

Tehran University of Medical Sciences

Research Article

Keywords: Magnetic solid-phase extraction (MSPE), GC/MS, Phthalate acid esters (PAEs), Non-alcoholic malt beverages

Posted Date: February 9th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-179959/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

The goal of this research was to assess the phthalic acid esters (DEP, DMP, BBP, DBP, DEHP, DnOP and total PAE) in non-alcoholic malt beverages bottled by multi-walled carbon nanotubes were magnetized with iron (MWCNT-Fe₃O₄) using Mass spectrometry is gas chromatography (GC-MS) in Tehran .

The results showed that the mean of DEHP (which has also been found to be carcinogenic) in all samples was lower than the standard levels (5944.73 ng/l). The maximum concentration of DEHP in four samples was upper than the standard levels (8957.87 ng/l). Also, scanning electron microscopy (SEM), scattered energy X-ray (EDX) and X-ray diffraction (XRD) analysis were used to describe the generated MWCNT-Fe₃O₄. Multivariate techniques and heat-map visualization were used to assess the correlation among the type and levels of PAEs with brand, color, product date, pH, sugar, volume and gas pressure. Therefore, based on heat-map and principal component analysis (PCA) results, the Bis (2- ethyl hexyl) phthalate (DEHP) and total PAEs were the closest accessions, indicating that these variables had similar trends. Based on the results, it can be stated that due to the low average of total phthalate esters in non-alcoholic malt beverages, there is no serious health hazard of these compounds for humans.

1. Introduction

Large amounts of phthalic acid esters (PAEs) are used annually to produce a wide variety of common consumer goods around the world (Arfaeinia et al. 2020, Harunarashid et al. 2017). PAEs are used mainly as plasticizers to increase durability, workability, and flexibility of polymeric materials in polyethylene terephthalate (PET) packaging or other plastic packaging, But, in addition, they can be present in many products such as inks, paints, cosmetics and adhesives. (Rafiei Nazari et al. 2018). These compounds bind to polymers through frail subaltern molecular interactions with the chains of polymer. Also, PAEs have employed in physically attached to the polymer, so easily released from the products and then transported to the environment (with migration way), and finally can cause contaminate air, soil, water, and food products (Dobaradaran et al. 2020, Jarošová 2006, Kouhpayeh et al. 2017). In recent years, food contamination with PAEs has become a problem of public concern. PAEs have been reported in beverages such as alcoholic and non-alcoholic malt beverages, soft drinks, meat and meat products, fruits, vegetables, dairy products, cereals, vegetable edibe oils and other food products (Kiani et al. 2018, Liu et al. 2020). PAEs are also widespread pollutants in the environment and appear to be the cause of cancer in humans and animals, as well as endocrine disorders. The USEPA (Environmental Protection Agency of USA) has listed PAEs as the important contaminant (Jarošová 2006, Moazzen et al. 2019, Shen et al. 2019). The most important PAEs include bis(2-ethylhexyl) phthalate (DEHP), di-n-octyl phthalate (DnOP), butylbenzyl phthalate (BBP), dimethyl phthalate (DMP), diethyl phthalate (DEP) and dibutyl phthalate (DBP). DEHP is one of the most important these compounds, which causes liver cancer in rodents (Liu et al. 2020, Sinha et al. 2019). DBP, BBP, DEHP and some phthalate esters metabolites in animals have shown teratogenic effects. The phthalates are involved in the pathogenesis of asthma, allergic symptoms, contribute to airway remodeling and the premature puberty in young girls (Harunarashid et al. 2017, Kiani et al. 2018). Several analytical techniques have been developed to detect PAE in various samples packaged in plastics (especially PET), such as drinking water, beverages, and cosmetics. High-performance liquid chromatography (HPLC), Gas chromatography (GC) and GC combined with mass spectrometry (GC-MS) were commonly used (Dobaradaran et al. 2020, Moazzen et al. 2019, Wu et al. 2014). Numerous preparation methods can be used to measure these compounds in different matrixes, including single-drop microextraction (SDME), liquid-liquid extraction (LLE), solvent extraction (SE), liquid-phase microextraction (LPME), LPME method based on the solidification of a floating organic micro drop (LPME-SFO), accelerated solvent extraction (ASE), stir-bar sorptive extraction (SBSE), solid-phase extraction (SPE) and column chromatographic cleanup (CC), but these methods are expensive, insensitive and time-consuming (Kiani et al. 2019, Roudbari et al. 2020, Shariatifar et al. 2020). Recently, a different type for solid-phase extraction (SPE), that called MSPE or magnetic solid-phase extraction has been established (this method based on the using of magnetic adsorbents). Magnetic adsorbents compared with non-magnetic adsorbents, can make separation process faster and easier without any additional procedures such as filtration or centrifugation. The MSPE method can prevent the time-consuming column operations seen in SPE (Kouhpayeh et al. 2017, Moazzen et al. 2018, Shen et al. 2019). Principal component analysis is one of the most important statistical technique used to characterize the interrelationships between variables and visualization of the data patterns (Ghelichkhani et al. 2019, Samiee et al. 2020). Heat-map used to represent the similar or vastly different expression status characteristics values (Arabameri et al. 2019b, Heydarieh et al. 2020). Considering the high consumption of these products in the daily diet, it is necessary to investigate the PAEs contaminants in alcoholic malt beverages, since so far, no research has been performed on PAEs contaminants in non-alcoholic malt beverages in Iran. Therefore, this research was conducted to investigate of PAEs in non-alcoholic malt beverages based on MSPE-GC/MS method by using MWCNT-Fe₃O₄ as magnetic adsorbent offered in Tehran, Iran. Also the PCA was applied to characterize the interrelationships between variables and visualization of the data patterns and Heat-map was applied to represent similar or vastly different expression characteristics values

2. Materials And Methods

2.1. Reagents and Chemicals

Phthalate acid esters (DEHP, BBP, DBP, DEP, DMP and DNOP) and other used chemicals including ethanol, methanol, NaCl and n-hexane were obtained from Sigma-Aldrich Company (St. Louis, MO). $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$, Benzyl benzoate (internal standard (IS)), NaBH_4 , HCl, NaOH and 3,5-dinitrosalicylic acid (DNSA) were purchased from Merck Company (Germany). The diameter and length of multi walled carbon nanotube (MWCNTs, Panchkula, India) were 30–60 nm and 5.0–30 mm, respectively. The grade of all the other used chemical and solvents were analytical reagent. The stock solution of analyzed contaminant was ready in methanol (100 mg/mL). Afterward, the solutions of phthalate acid esters working standard were ready by consecutive dilutions of the stock solution with distilled water-methanol (50: 50 v/ v). To prepare I.S., benzyl benzoate (10 mg) was dissolved in ethanol (10 mL), and at the end for preparing samples, 100 μL of prepared I.S. was added. On the same day of the study, the QC (quality control) were prepared from diluted solutions of stock standard. The prepared solutions were retained at 4°C and in the dark place, until analysis. In this research, all the laboratory glass dishes (before using) were washed with an $\text{Al}(\text{O})_3$ solution, and then they were immersed in acetone for 40min, and were washed with n-hexane as well as finally were dried (in the oven) for 5 h at 150°C.

2.2. Determination of sugar

In the first, five samples with different dilutions were prepared from sucrose stock solution (1000 mg/dL) as standard. Afterward, 10mL of non-alcoholic malt beverages samples diluted with distilled water up to the 100 mL as samples. Insert 2 mL of each standard solution into a test tube and insert two mL of distilled water into a separate test tube (blank solution). Then, two mL of HCl hydrochloric acid (six M) solution was added and placed in Bain-marie for 10min and then 8 mL of 2.5 M NaOH solution was added to neutralize. Afterward, add two mL of 0.05 M 3,5-dinitrosalicylic acid solution and cover the test tube with cover (film) and shake well to mix. Then was placed in Bain-marie for five min followed by so cold water for ten min (The time between 3,5-dinitrosalicylic acid solution addition and measurements should be same for all test solutions.). Finally, the absorbance of the five standards and references at 580 nm was measured using UV/Vis Spectrophotometer (Roig & Thomas 2003).

2.3. Gas measurement

The gas pressure of the non-alcoholic malt beverages samples were determined using a digital gas meter at room temperature set at 25°C.

2.4. The pH measurement

The pH of the non-alcoholic malt beverages samples were determined using using a digital pH meter at room temperature and pressure of 1atm (Godwill et al. 2015).

2.5. Preparation of adsorbent

Magnetic adsorbents ($\text{MWCNTS-Fe}_3\text{O}_4$) were prepared according our previous study (Kiani et al. 2018, Kouhpayeh et al. 2017, Moazzen et al. 2018).

2.6. Evaluate the properties of the prepared adsorbent

Phase identification of adsorbent was assessed by XRD (X-ray Diffraction) and characterization of elemental was assessed by EDX (energy dispersive X-ray) and morphological analysis was assessed by SEM (scanning electron microscope) (Moazzen et al. 2019). The models of our devices were SEM: PHILIPS, and S360 Mv2300, EDX: PHILIPS, S360, and Mv2300 and XRD: Philips, X'PertPro 2002.

2.7. Sampling, sample preparation, and instrumentation

In the beginning, 120 non-alcoholic malt beverages samples were bought with five brand of most commonly used from chain store in Tehran, Iran and all samples were degassed (with a bath of ultrasonic) at temperature of room for 20 minute. Afterward, 10mg of prepared adsorbent ($\text{MWCNT-Fe}_3\text{O}_4$) were activated with solutions of methanol and water. To the ten milliliter of each degassed non-alcoholic malt beverage samples, 10 mg the activated adsorbent, 0.5 gr of NaCl, and 100 μL I.S. were added. The mixture was mixed strongly (to extract the contaminant compounds) for four minute, after that adsorbent was gathered (with an exterior magnet) to the side of the laboratory dish (within 90 seconds) and the other compounds in the mixture were discarded. Then, 2 mL of n-hexane was added to the adsorbent and was mixed with blender (2 min) vigorously, to elute contaminant compounds (PAEs) from the adsorbent. After that, by using an external magnet, the adsorbent was collected to the side of the laboratory dish, and the supernatant was moved to a vial. Desorption solvent was

dehydrated with a mild stream of N₂ gas (at room temperature) and was maintained in the cold place such as refrigerator. Finally, the dehydrated contents of the previous step was dissolved in one mL solvent (n-hexane), and one µL of the mentioned solution was injected into the gas chromatography (Agilent 7890 N) –mass spectrometry (5975) device. The column of chromatographic was DB-5–J & W Scientific (30m, 250µm, 0.5µm). Helium (He) was selected as the gas of carrier at one mL/min (ratio of split of 50:1). Splitless was the mode of injection with an inlet temperature of 290-centigrade degree. The program of GC-MS temperature was: 80-centigrade degree, retained for two minute, 80–285 centigrade degree at seven °C/min, retained for ten minute. For the quantitative determination of PAEs compounds, the mode of selective ion monitoring was used. The device outputs (the retention times (RT), quantitative and qualitative ions) of six phthalate acid esters and I.S. are presented in Table 1.

Table 1

The RT of each analyst, quantitative and qualitative ions for the GC/MS outputs of 6 phthalate acid esters and internal standard.

Group of ions	Compounds	RT (minute)	Quantitative ions (abundance) (m/z)	Qualitative ions (abundance) (m/z)
1	Di methyl phthalate (DMP)	12.8–13.2	163	76, 134, 162, 195
2	Di ethyl phthalate (DEP)	14.2–14.8	149	121, 149, 177, 222
3	Di butyl phthalate (DBP)	20.1–20.4	149	121, 149, 205, 223
4	Butyl benzyl phthalate (BBP)	25.2–25.6	149	91, 149, 206, 238
5	Benzyl benzoate (IS)	26.1–26.4	105	212 (40), 194 (35)
6	Bis(2- ethyl hexyl) phthalate (DEHP)	27.1–27.6	149	113, 149, 167, 279
7	Di-n-octyl phthalate (DNOP)	29.3–29.5	149	149, 179, 261, 279

2.8. Optimization of method of the extraction process

The optimization of method was performed according to the 1 factor at a time method. (Kiani et al. 2018, Moazzen et al. 2019).

2.9. Validation of method

The validation of method was performed based on the currently established Guideline of FDA (U.S. Food and Drug Administration) for industries (Kouhpayeh et al. 2017, Moazzen et al. 2018).

2.10. Statistical analysis

The outcomes were statistically analyzed by SPSS version 18 (SPSS Inc, Chicago, IL) for Windows. Data analysis done by test of Kolmogorov–Smirnov and tests of Kruskal-Wallis. Statistical significance was a p-value of <0.05. For a better understanding of the most significant contribution to distribution PAEs levels among the different samples, the PCA was done by the software of SPSS (Arabameri et al. 2019a, Heydarieh et al. 2020). Multivariate techniques were used to assess the correlation between the amount and type of PAEs levels with properties samples. Heat-map analysis was used to analyze the correlation between samples online at <https://biit.cs.ut.ee/clustvis/>.

3. Results And Discussion

3.1. Quantitative analysis

To draw the calibration curves were prepared five different concentration levels of phthalate acid esters (DEHP, DBP, DEP, DNOP, BBP and DMP) at the range of 10 to 12000 ng/L (including 10, 100, 1000 and 5000 ng/L). The correlation coefficient ranged from 0.9979 to 0.9997 and the LODs (detections limit) and LOQs (quantifications limit) for the target analytes were 13 to 30 ng/L and 39 to 90 ng/L, respectively (Table 2). The recovery values of the 6 PAEs were 94.2–104.3% with the RSDs less than 7.6%. To control the quality three levels (50, 500 and 5000 ng/L) from mix of PAEs were prepared and were analyzed duplicate in several days. The inter- and intra- day precision measured for three consecutive days in triplicate analyzes and they were lower than 7.8% and 8%, respectively (Table 3). The selectivity of method was examined by analyzing 25 non-alcoholic malt beverages.

Table 2

The linear range, LOD, LOQ and coefficient of estimation of the developed MSPE technique for measurement of phthalate acid esters

Target compound	Linear range (ng/L)	Detections limit (LOD) (ng/L)	quantifications limit (LOQ) (ng/L)	Coefficient of estimation (r ²)
Di methyl phthalate (DMP)	10–12000	23	69	0.9979
Di ethyl phthalate (DEP)	10–12000	15	45	0.9984
Di butyl phthalate (DBP)	10–12000	13	39	0.9991
Butyl benzyl phthalate (BBP)	10–12000	30	90	0.9990
Di-n-octyl phthalate (DNOP)	10–12000	18	54	0.9988
Bis(2-ethyl hexyl) phthalate (DEHP)	10–12000	26	78	0.9997

Table 3

Estimated recoveries, precisions and accuracies for determination of the PAEs compounds at three different concentrations (n = 6) in QC samples.

Target compound	Sample	Nominal concentration (ng/L)	Mean of calculated concentration (ng/L)	RSD(%) of calculated concentration (intraday)	RSD(%) of calculated concentration (interday)	RE(%) of calculated concentration	Estimated recoveries (%)	RSD(%) of calculated recovery
Di methyl phthalate (DMP)	QCI	50	52	6.4	7.3	4	95.3	7.2
	QCII	500	503	7.1	6.6	0.6	96.0	6.4
	QCIII	5000	5050	7.8	7.9	1	97.3	7.0
Di ethyl phthalate (DEP)	QCI	50	47	5.3	7.1	-6	99.7	6.9
	QCII	500	510	6.4	8.0	2	96.3	7.6
	QCIII	5000	5100	6.2	5.8	2	94.2	5.7
Di butyl phthalate (DBP)	QCI	50	53	6.3	7.2	6	98.4	7.0
	QCII	500	505	6.8	6.7	1	98.4	5.6
	QCIII	5000	5045	7.0	6.9	0.9	99.8	6.3
Butyl benzyl phthalate (BBP)	QCI	50	56	5.6	5.9	12	104.3	5.8
	QCII	500	512	6.4	7.7	2.4	98.3	7.2
	QCIII	5000	5039	6.5	7.0	0.78	98.6	6.0
Di-n-octyl phthalate (DNOP)	QCI	50	56	5.2	6.3	12	98.2	5.7
	QCII	500	504	7.3	7.2	0.8	98.9	6.9
	QCIII	5000	5105	6.6	5.9	2.1	97.1	5.8
Bis(2-ethyl hexyl) phthalate (DEHP)	QCI	50	49	6.9	7.8	-2	101.3	7.0
	QCII	500	515	6.0	6.7	3	98.8	5.6
	QCIII	5000	5070	6.8	7.3	1.4	98.4	6.9

3.3. Images of SEM and analysis of EDX

The SEM image and EDX analysis of the magnetic adsorbents are presented in Figs. 1 and 2. The graphs show that the placement of magnetic particle on the surface of multi wall carbon nanotubes is comparatively uniform and after filling with magnetic particles, the surface of the multi wall carbon nanotubes became rougher (Fig. 1). Besides, there's no significant modification is detected in the surface of construction of the magnetic adsorbents after extraction procedure. A growth in adsorbent diameter is apparent. In addition, we

concluded that the adsorbent exhibited a chain-like morphology without apparent collection. This conducts to a high ability of adsorption of the adsorbent.

The chemical compounds of the prepared adsorbent were assessed by using EDX analysis. The spectrum of EDX displayed oxygen (O), iron (Fe) and carbon (C). The atomic C, Fe, and O ratio (64.6, 22.7 and 12.7, respectively) as the principal elements in the structure of prepared adsorbent confirmed the quantitative representation of the existence of Fe_3O_4 nanoparticles on the MWCNT surface (Fig. 2).

3.4. XRD images

By XRD analysis, the construction of multi wallet carbon nanotubes and MWCNT- Fe_3O_4 composites were more confirmed. The patterns of XRD of adsorbent were exhibited in Fig. 3. The strong diffraction peaks at $2\theta = 31/77^\circ$ and $2\theta = 45/52^\circ$ were shown MWCNTs and Fe_3O_4 , respectively. The get XRD outcomes displayed that the Fe_3O_4 were efficaciously coated on the surface and texture of multi wallet carbon nanotubes using a co-precipitation technique.

3.5. Evaluation of phthalate esters in non-alcoholic malt beverages bottled in PET bottle

Mean concentrations and other statistical analyses of PAEs in all samples were shown in Table 4. There was a significant difference between the groups in terms of phthalate esters ($P < 0/05$). According to Table 4 from the research mean of all compounds in all the samples were less than the standard defined by the EPA (6000 ng/L) and WHO-EU (8000 ng/L) in drinking water, but maximum of DEHP in samples (4 samples) was upper than the EPA and WHO-EU standards (8957.87 ng/L).

Table 4
Statistical analysis of phthalate esters in non-alcoholic malt beverages(ng/L)

Target compound	mean	SD	min	max
DMP	18.31	8.75	8.6	98.1
DEP	151.33	31.73	74.83	268.9
DBP	496.73	108.99	300.76	803.21
BBP	nd	—	nd	nd
DEHP	5944.73	2518.14	1897.12	8957.87
DnOP	46.16	13.24	15.43	97.13
Total	6657.28	1600.9	2412.5	9483.93

The mean value and range of pH in samples was 3.3 (3.18–3.53) and sugar (per 100g) was 8.7 (4.4–11.9) and gas pressure was 1140 (830–1398) millimeters of mercury (mmHg). There was a significant difference between the groups in terms of sugar content and gas pressure ($P < 0/05$).

M oazzen ,M et al. (2018) showed the highest concentration of PAEs in carbonated soft drinks was DEHP, that it was upper than the standard level (STL) in the 4 samples (14,008, 9301.6, 9201 and 6766.6 ng/L) and other PAEs compounds were lower than the STL, which was somewhat similar to our study (Moazzen et al. 2018).

Xu,X et al. (2019) showed 3 PAEs compounds (DBP,DMP and DEP) were measured in ten common brands of bottled water (Made of PET) in Beijing (China), ranging from 101.97 $\mu\text{g/kg}$ to 709.87 $\mu\text{g/kg}$ (Xu et al. 2020).

Vincenzo Russo, M et al in 2014 with the determination of PAEs in soft drinks and alcoholic drinks (light) showed that DEHP (3.6–101 ng/mL), DBP (1.9–4.4 ng/mL), DiBP (0.2–2.5 ng/mL), DEP (0.1–1.0 ng/mL) and BBP (0.08–0.8 ng/mL), are present in all samples, while iBcEP (0.08 ng/mL) and DMP (1.9 ng/mL) are present only in 1 beer sample, that was higher than compare our study (Russo et al. 2014).

Rafiei Nazari, R et al. (2017) with determine the migration modelling of PAEs from non-alcoholic beer bottles showed that storage duration increased and temperature resulted in an increase in migration level ranging from 0.6 $\mu\text{g/L}$ to 2.9 $\mu\text{g/L}$, which was lower than our study (Rafiei Nazari et al. 2018).

Carnol, L et al. (2017) identified 6 PAEs of Luxembourgish beer stored in various containers (aluminum, glass and can bottle) and total PAEs were found in all samples at levels of 61.56 µg / L, which was upper than the present study (Carnol et al. 2017).

Victor E. Balderas-Hernández et al. (2020) identified PAEs in tequila beverage and showed that 22% of samples (65 samples) lacked PAE. DINP (1.64–3.43 mg/kg), BBP (0.05–2.91 mg/kg) and DEP (0.13–0.27 mg/kg) were found in 5 (1.69%), 37 (12.54%) and 11 (3.73%) samples, respectively. However, these levels weren't higher than the maximum standard level of PAEs for alcoholic beverages. DEHP (0.03–4.64 mg/kg) and DBP (0.01–2.20 mg/kg) were found in 224 (75.93%) and 96 (32.54%) samples, from them just 15 (5.08%) and 10 (3.39%) samples, respectively, exceeded the maximum standard levels for alcoholic drinks. Bis(2-ethylhexyl) phthalate was the most repetitious PAEs detected in tequila and detected concentrations of DEHP were two-times higher in ultra-aged tequilas compared to those in white tequilas (Balderas-Hernández et al. 2020).

Yang, JF et al. (2016) measured the PAEs in non-alcoholic drinks bottled in PET from China and showed Bis(2-ethylhexyl) phthalate contained the highest median and mean contents (0.62 ng/g and 1.60 ng/g), followed by DBP (0.27 ng/g and 1.34 ng/g). (Yang et al. 2017).

Rodríguez-Ramos, R et al (2020) by evaluating plastic migrants in non-alcoholic and alcoholic drinks showed that there are 4 PAEs in the range of 0.14–1.1 µg / L in some beers, 1 PAE in the range of 1.2–1.5 µg / L in 3 grape juices and 6 PAE in several cider samples, in the range of 0.3–2.1 µg / L, which was less than the present study (Rodríguez-Ramos et al. 2020).

Wu, PG et al. (2014) with determination of PAEs in non-alcoholic beverages showed a wide variety of PAEs contents was detected in 48 non-alcoholic beverages. Bis(2-ethylhexyl) phthalate was the most abundant PAEs compound followed by DOP, DPP and DBP. Bis(2-ethylhexyl) phthalate was detected in fruit juice samples (0.022–0.126 mg/L), sport beverages (0.015–0.098 mg/L), coffee (0.028–0.159 mg/L) and tea (0.016–0.123 mg/L), that was higher than our study (Wu et al. 2014).

March, JG et al. (2015) showed low-alcohol beer had DEHP 0.4 ± 0.2 and DBP 2.2 ± 0.4 µg/L, that DEHP was lower than our study and DBP was higher our study (March & Cerdà 2015).

Heinemeyer, G et al. (2013) showed that the DEHP in Beer (alcoholic) was 0.022 and in nonalcoholic Beverages was 0.020 µg/g (Heinemeyer et al. 2013).

Wang, F et al. (2017) showed DEHP in alcoholic beverages (liquor) was ranged from 0.6182 to 1.0890 µg/mL, that was higher than our study (Wang et al. 2017).

Pang, YH et al (2019) showed DEHP in alcoholic carbonated beverage and beer was not found, that was lower than our study (Pang et al. 2020).

The data comparison of this research with other articles shows differences that can be due to reasons such as: alcoholic or non-alcoholic beverage, use of plastic or other containers, contamination of raw materials or secondary contaminants during the production process, carbonated or non-carbonated beverage and also the amount of gas in carbonated beverage (pH), duration Keep in the package as well as the amount of other ingredients in the drink.

3.6. Structural relationship of parameters

Multivariate techniques and heat-map visualization were applied to evaluate the correlation between the type and levels of PAEs with brand, color, product date, pH, sugar, volume and gas pressure. Consequently, based on heat-map and PCA results, the Bis(2- ethyl hexyl) phthalate (DEHP) and total PAEs were the closest accessions, indicating that this variables had similar trends. Heat-map clustered the 40 non-alcoholic malt beverages using correlation distance and average linkage, reflecting similarities and relationship among the type and levels of PAEs samples. The heat-map clearly grouped samples into two major clusters and two sub clusters (Fig. 4). First cluster includes DEHP and total PAEs, Second cluster contains two sub-groups with brand, color, product date, pH, sugar, volume, gas pressure, DMP, DEP, DBP, BBP and DNOP.

Quantitative results obtained for the type and levels of PAEs were used to PCA to investigate the most significant contribution among non-alcoholic malt beverages. The compounds included samples brand, color, product date, pH, sugar, volume, gas pressure and levels of PAEs (DEHP, DMP, DNOP, BBP, DBP and DEP).

The dependence relations between different PAEs can be found from Fig. 4, the graph subset illustrates a visual representation about the relations among different PAEs. The nearest neighbor among the properties, the greater significant relationship existed among dependent

variable.

As shown in Fig. 5, the first five principal components accounted for 77.86% of the data variance in all samples, and their contribution rates were 26.80%, 15.87%, 14.27%, 11.46% and 9.43%, respectively. The DEHP and total PAEs were the closest accessions, indicating which these variables had similar trends. The Total phthalate, DEP, DBP, BEHP had a high positive correlation with PC1, while had negative correlation with sugar content and gas pressure. The results showed that the sugar, pH, brand had a positive correlation with PC2, while had negative correlation with gas pressure and DOP.

4. Conclusion

In first magnetic adsorbent was ready using a simple, sensitive and cost-effective method for measuring 6 PAEs (DEHP, DMP, BBP, DnOP, DEP and DBP) from non-alcoholic malt beverages. Moreover, SEM, EDX, XRD, VSM and TEM analysis were used to describe the generated MWCNT-Fe₃O₄. Analysis of non-alcoholic malt beverage samples evaluated in Tehran and showed that the concentration of none of the PAEs released from the bottles was upper than the standard levels (6 µg/L by USEPA). The correlation analysis focuses on the numerical relationship among types and levels of PAEs in all samples. Comparing relationships between PAEs levels from different non-alcoholic malt beverages showed that the Bis (2- ethyl hexyl) phthalate (DEHP) and total PAEs were the closest accessions. Furthermore, the results of heat-map visualization clearly indicated that relationship between PAEs levels under different samples was properly diagnosed. Based on the results, it can be stated that due to the low average of total phthalate esters in non-alcoholic malt beverages, there is no serious health hazard of these compounds for humans.

Declarations

Acknowledgments

This work was performed in Department of Environmental Health laboratory of Health School of Tehran University of Medical Sciences.

Funding

This study was supported by Research Grant of Health School of Tehran University of Medical Sciences.

Competing interests

The authors have no conflicts of interest to declare in this research.

Ethical Approval

"This study does not involve any human or animal testing" or "This study was approved by the School of Public Health of Tehran University of Medical Sciences.

Consent to participate

All authors participated in this work.

Consent to publish

All authors agree to publish.

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Author's contributions

Nabi Shariatifar, Gholamreza Jahed: Conceptualization, Supervision, Design of study, Writing- Reviewing and Editing. **Hana Rezaei and Mojtaba Moazzen:** Data curation, Writing- Original draft preparation. **Mohammad Hadi Dehghani:** Visualization, Investigation. **Mahsa Alikord:** Software, Methodology. **Majd Arabameri :** Software, Validation:

References

1. Arabameri M, Nazari RR, Abdolshahi A, Abdollahzadeh M, Mirzamohammadi S, Shariatifar N, Barba FJ, Mousavi Khaneghah A (2019a): Oxidative stability of virgin olive oil: evaluation and prediction with an adaptive neuro-fuzzy inference system (ANFIS). *Journal of the Science of Food and Agriculture* 99, 5358-5367
2. Arabameri M, Nazari RR, Abdolshahi A, Abdollahzadeh M, Mirzamohammadi S, Shariatifar N, Barba FJ, Mousavi Khaneghah A (2019b): Oxidative stability of virgin olive oil: evaluation and prediction with an adaptive neuro-fuzzy inference system (ANFIS). *Journal of the Science of Food and Agriculture*
3. Arfaeina L, Dobaradaran S, Nasrzadeh F, Shamsi S, Poureshgh Y, Arfaeina H (2020): Phthalate acid esters (PAEs) in highly acidic juice packaged in polyethylene terephthalate (PET) container: Occurrence, migration and estrogenic activity-associated risk assessment. *Microchemical Journal* 155, 104719
4. Balderas-Hernández VE, Ornelas-Salas JT, Barba de la Rosa AP, De Leon-Rodriguez A (2020): Diminution of migration of phthalic acid esters in tequila beverage by the year of production. *Journal of Environmental Science and Health, Part B* 55, 148-154
5. Carnol L, Schummer C, Moris G (2017): Quantification of six phthalates and one adipate in Luxembourgish beer using HS-SPME-GC/MS. *Food Analytical Methods* 10, 298-309
6. Dobaradaran S, Akhbarizadeh R, Mohammadi MJ, Izadi A, Keshtkar M, Tangestani M, Moazzen M, Shariatifar N, Mahmoodi M (2020): Determination of phthalates in bottled milk by a modified nano adsorbent: Presence, effects of fat and storage time, and implications for human health. *Microchemical Journal* 159, 105516
7. Ghelichkhani G, Modaresi MH, Rashidi L, Shariatifar N, Homapour M, Arabameri M (2019): Effect of the spray and freeze dryers on the bioactive compounds of olive leaf aqueous extract by chemometrics of HCA and PCA. *Journal of Food Measurement and Characterization*, 1-13
8. Godwill EA, Jane IC, Scholastica IU, Marcellus U, Eugene AL, Gloria OA (2015): Determination of some soft drink constituents and contamination by some heavy metals in Nigeria. *Toxicology Reports* 2, 384-390
9. Harunarashid NZIH, Lim LH, Harunsani MH (2017): Phthalate sample preparation methods and analysis in food and food packaging: a review. *Food Analytical Methods* 10, 3790-3814
10. Heinemeyer G, Sommerfeld C, Springer A, Heiland A, Lindtner O, Greiner M, Heuer T, Krems C, Conrad A (2013): Estimation of dietary intake of bis (2-ethylhexyl) phthalate (DEHP) by consumption of food in the German population. *International journal of hygiene and environmental health* 216, 472-480
11. Heydari A, Arabameri M, Ebrahimi A, Ashabi A, Monjazeab Marvdashti L, Shokrollahi Yancheshmeh B, Abdolshahi A (2020): Determination of Magnesium, Calcium and Sulphate Ion Impurities in Commercial Edible Salt. *Journal of Chemical Health Risks* 10, 93-102
12. Jarošová A (2006): Phthalic acid esters (PAEs) in the food chain. *Czech J. Food Sci* 24, 223-231
13. Kiani A, Ahmadloo M, Shariatifar N, Moazzen M, Baghani AN, Khaniki GJ, Taghinezhad A, Kouhpayeh A, Khaneghah AM, Ghajarbeygi P (2018): Method development for determination of migrated phthalate acid esters from polyethylene terephthalate (PET) packaging into traditional Iranian drinking beverage (Doogh) samples: a novel approach of MSPE-GC/MS technique. *Environmental Science and Pollution Research* 25, 12728-12738
14. Kiani A, Shariatifar N, Shahsavari S, Ahmadloo M, Moazzen M (2019): Investigating the Presence of Polycyclic Aromatic Hydrocarbons in Doogh. *Journal of Mazandaran University of Medical Sciences* 29, 10-23
15. Kouhpayeh A, Moazzen M, Jahed Khaniki GR, Dobaradaran S, Shariatifar N, Ahmadloo M, Azari A, Nazmara S, Kiani A, Salari M (2017): Extraction and determination of phthalate esters (PAEs) in Doogh. *Journal of Mazandaran University of medical sciences* 26, 257-267
16. Liu JM, Li CY, Zhao N, Wang ZH, Lv SW, Liu JC, Chen LJ, Wang J, Zhang Y, Wang S (2020): Migration regularity of phthalates in polyethylene wrap film of food packaging. *Journal of Food Science*
17. March JG, Cerdà V (2015): An innovative arrangement for in-vial membrane-assisted liquid-liquid microextraction: application to the determination of esters of phthalic acid in alcoholic beverages by gas chromatography-mass spectrometry. *Analytical and Bioanalytical Chemistry* 407, 4213-4217
18. Moazzen M, Mahvi AH, Shariatifar N, Jahed Khaniki G, Nazmara S, Alimohammadi M, Ahmadkhaniha R, Rastkari N, Ahmadloo M, Akbarzadeh A (2018): Determination of phthalate acid esters (PAEs) in carbonated soft drinks with MSPE/GC-MS method. *Toxin Reviews* 37, 319-326
19. Moazzen M, Khaneghah AM, Shariatifar N, Ahmadloo M, Eş I, Baghani AN, Yousefinejad S, Alimohammadi M, Azari A, Dobaradaran S (2019): Multi-walled carbon nanotubes modified with iron oxide and silver nanoparticles (MWCNT-Fe₃O₄/Ag) as a novel adsorbent for determining PAEs in carbonated soft drinks using magnetic SPE-GC/MS method. *Arabian Journal of Chemistry* 12, 476-488

20. Pang Y-H, Yue Q, Huang Y-y, Yang C, Shen X-F (2020): Facile magnetization of covalent organic framework for solid-phase extraction of 15 phthalate esters in beverage samples. *Talanta* 206, 120194
21. Rafiei Nazari R, Noorian S, Arabameri M (2018): Migration modelling of phthalate from non-alcoholic beer bottles by adaptive neuro-fuzzy inference system. *Journal of the science of food and agriculture* 98, 2113-2120
22. Rodríguez-Ramos R, Socas-Rodríguez B, Santana-Mayor Á, Rodríguez-Delgado MÁ (2020): A simple, fast and easy methodology for the monitoring of plastic migrants in alcoholic and non-alcoholic beverages using the QuEChERS method prior to gas chromatography tandem mass spectrometry. *Analytical and Bioanalytical Chemistry* 412, 1551-1561
23. Roig B, Thomas O (2003): Rapid estimation of global sugars by UV photodegradation and UV spectrophotometry. *Analytica Chimica Acta* 477, 325-329
24. Roudbari A, Nazari RR, Shariatifar N, Moazzen M, Abdolshahi A, Mirzamohammadi S, Madani-Tonekaboni M, Delvarianzadeh M, Arabameri M (2020): Concentration and health risk assessment of polycyclic aromatic hydrocarbons in commercial tea and coffee samples marketed in Iran. *Environmental Science and Pollution Research*, 1-13
25. Russo MV, Notardonato I, Avino P, Cinelli G (2014): Determination of phthalate esters at trace levels in light alcoholic drinks and soft drinks by XAD-2 adsorbent and gas chromatography coupled with ion trap-mass spectrometry detection. *Analytical Methods* 6, 7030-7037
26. Samiee S, Fakhri Y, Sadighara P, Arabameri M, Rezaei M, Nabizadeh R, Shariatifar N, Mousavi Khaneghah A (2020): The concentration of polycyclic aromatic hydrocarbons (PAHs) in the processed meat samples collected from Iran's market: a probabilistic health risk assessment study. *Environmental Science and Pollution Research*
27. Shariatifar N, Dadgar M, Fakhri Y, Shahsavari S, Moazzen M, Ahmadloo M, Kiani A, Aenehvand S, Nazmara S, Khanegah AM (2020): Levels of polycyclic aromatic hydrocarbons in milk and milk powder samples and their likely risk assessment in Iranian population. *Journal of Food Composition and Analysis* 85, 103331
28. Shen C, Wu T, Zang X (2019): Hollow fiber stir bar sorptive extraction combined with GC–MS for the determination of phthalate esters from children's food. *Chromatographia* 82, 683-693
29. Sinha P, Anshu AK, Tiwari M, Ahmad MY (2019): Determination of PAEs in soil of municipal wards of Patna, India by microwave assisted extraction and LC-MS/MS. *Soil and Sediment Contamination: An International Journal* 28, 162-170
30. Wang F, Liu Y, Tang Z, Hou M, Wang C, Wang X, Wang Q, Xiao Q (2017): Simultaneous determination of 15 phthalate esters in commercial beverages using dispersive liquid–liquid microextraction coupled to gas chromatography-mass spectrometry. *Analytical Methods* 9, 1912-1919
31. Wu P-G, Pan X-D, Ma B-J, Wang L-Y, Zhang J (2014): Determination of phthalate esters in non-alcoholic beverages by GC–MS and optimization of the extraction conditions. *European Food Research and Technology* 238, 607-612
32. Xu X, Zhou G, Lei K, LeBlanc GA, An L (2020): Phthalate Esters and Their Potential Risk in PET Bottled Water Stored under Common Conditions. *International Journal of Environmental Research and Public Health* 17, 141
33. Yang J-F, Yang L-M, Zheng L-Y, Ying G-G, Liu C-B, Luo S-L (2017): Phthalates in plastic bottled non-alcoholic beverages from China and estimated dietary exposure in adults. *Food Additives & Contaminants: Part B* 10, 44-50

Figures

Figure 1

SEM of magnetic adsorbent (A: MWCNTs and B:MWCNTs-Fe3O4)

Figure 2

EDX of magnetic MWCNTs/Fe3O4

Figure 3

The XRD image of MWCNTs/Fe₃O₄

Figure 4

Heat map of PAEs in non-alcoholic malt beverages

Figure 5

Principal component analysis plot of PAEs in non-alcoholic malt beverages samples