

Effect of acupuncture on insomnia in menopausal women: A Study protocol for a Randomized Controlled Trial

Shanshan Li

Shanghai Municipal Hospital of Traditional Chinese Medicine, Shanghai University of Traditional Chinese Medicine

Ping Yin

Shanghai Municipal Hospital of Traditional Chinese Medicine, Shanghai University of Traditional Chinese Medicine

Xuan Yin

Shanghai Municipal Hospital of Traditional Chinese Medicine, Shanghai University of Traditional Chinese Medicine

Anna Bogachko

Shanghai Municipal Hospital of Traditional Chinese Medicine, Shanghai University of Traditional Chinese Medicine

Tingting Liang

Shanghai Municipal Hospital of Traditional Chinese Medicine, Shanghai University of Traditional Chinese Medicine

Lixing Lao

School of Chinese Medicine, The University of Hong Kong and University of Maryland School of Medicine

Shifen Xu

Shanghai Municipal Hospital of Traditional Chinese Medicine, Shanghai University of Traditional Chinese Medicine <https://orcid.org/0000-0001-5595-3480>

Method Article

Keywords:

Posted Date: November 7th, 2018

DOI: <https://doi.org/10.21203/rs.1.11/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Trials on May 30th, 2019. See the published version at <https://doi.org/10.1186/s13063-019-3374-8>.

Abstract

Background: China today has high occurrences of insomnia in perimenopausal women. Some studies have shown that acupuncture might be effective in treating primary insomnia and insomnia related to depression and stroke. Although there are some programs supporting insomnia during the menopausal transition, there are few randomized controlled trials to provide evidence regarding their effectiveness. We design a randomized controlled trial of suitable sample size to verify the effectiveness of acupuncture in patients with insomnia during the menopausal transition and to form an optimized acupuncture treatment protocol. **Method/Design:** In this randomized, single-site, single-blind, placebo-controlled trial, 84 eligible patients will be recruited and randomly assigned to either the acupuncture group (n = 42) or the sham control group (n = 42) in a 1:1 ratio. Participants will receive a total of 18 sessions of treatment for eight consecutive weeks. Treatments will be given three times per week in the first four weeks, twice a week for the next two weeks and finally treatment once weekly for the final two weeks. Treatment will utilize eight main acupoints (GV20, GV24, GV29, RN6, RN4, SP6, HT7, EX-HN22), and extra two acupoints based on syndrome differentiation. The primary outcome will be assessed using the Pittsburgh Sleep Quality Index (PSQI). The secondary outcomes will be measured by sleep parameters recorded in the Actigraphy (SE, TST, SA), Insomnia Severity Index (ISI), Self-Rating Anxiety Scale (SAS), Self-Rating Depression Scale (SDS) and Menopause Quality of Life (Men-QOL). The primary outcomes will be assessed at baseline, week four, week eight, and first and the third month following the end of treatment. **Discussion:** If the results confirm that acupuncture is effective and safe for the treatment on insomnia in menopausal women, this positive outcome could provide evidence for clinical application.

Background

Background

The menopausal transition (MT) is often accompanied by a myriad of symptoms, the most common of which being insomnia[1-3]. Insomnia is characterized by having difficulty falling asleep, frequent nocturnal wakings with difficulty falling back asleep[4] and unexplained pain. A study conducted by the National Institutes of Health estimates the prevalence of insomnia in menopausal women at 40-50%[5-8]. Women who experience poor sleep quality during the MT are more likely to have critical problems[9] such as depression, anxiety, memory and cognitive deprivation, cardiovascular disease and reduced quality of life[3, 4]. Yet, there are few effective therapies available for treating sleep problems during the MT. Yoga, therapeutic massage and exercise may be helpful for sleep disorders, but they have not yet been proven as an effective independent treatment[10]. Hormone therapy (HT) is currently the most effective treatment, but is also associated with a number of risks[11, 12]. HT can reduce hot flashes at night, and some studies have reported that it can reduce sleep disorders[13]. However, increasing numbers of women are refusing HT due to the risks (myocardial infarction, stroke, venous thromboembolism and breast cancer) outlined by trials published in the Women's Health Initiative[14]. Moreover, these trials reported that HT has a weaker effect on insomnia than previously thought[15].

Acupuncture, an important part of Traditional Chinese Medicine (TCM), has been used for thousands of years in China to treat various diseases including sleep disorders[5]. Many studies have confirmed the efficacy of acupuncture for primary insomnia and related depressive disorders[16-18]. Based on the results that the efficacy of acupuncture for depression-related insomnia [19] and insomnia following stroke [20], we believe that acupuncture might be an effective treatment for insomnia during the MT. However, effects of acupuncture on sleep disorders remain uncertain because of flawed research methodology, such as improper sample size, treatment times and follow-up problems[21].

The main objective of this study is to determine whether acupuncture is effective as compared to a sham acupuncture control on insomnia in menopausal women. The findings of this trial will provide useful information in forming an optimal acupuncture treatment protocol.

Methods/design

Methods/Design

Hypotheses: Acupuncture treatment helps menopausal transition insomnia patients improve their sleep quality and other related symptoms.

Objectives:

1. To compare the differences in improvement of insomnia, assessed by the Pittsburgh Sleep Quality Index (PSQI) and Insomnia Severity Index (ISI), between the intervention group and control group.
2. To compare the differences in improvement of mood, measured by the Self-Rating Anxiety Scale (SAS) and Self-Rating Depression Scale (SDS), between the intervention group and control group.
3. To compare the differences in improvement of Menopause Quality of Life (MenQOL), between the intervention group and control group.

Design: This is a single-site, single-blind, randomized, placebo-controlled clinical trial that will be carried out in the Shanghai Municipal Hospital of Traditional Chinese Medicine. Eligible patients will be randomly divided into the acupuncture group and the sham acupuncture group in a 1:1 allocation ratio. All participants will sign the informed consent before proceeding with the trial.

The flow chart of the study process is as follows in Fig. 1. Timing of treatment assessments and data collection are as follows in Table 1.

Table1. Timing of treatment assessments and data collection

	STUDY PERIOD					
	Enrolment	Baseline	Treatment phase		Follow-up phase	
	<i>-1week</i>	<i>0week</i>	<i>4week</i>	<i>8week</i>	<i>4week</i>	<i>12week</i>
ENROLMENT:						
Eligibility screen	X					
Informed consent	X					
Medical history	X					
Allocation		X				
INTERVENTIONS:						
Acupuncture		X	X	X		
Sham Acupuncture		X	X	X		
ASSESSMENTS:						
Primary outcome						
PSQI		X	X	X	X	X
Secondary outcomes						
ISI		X		X		
Actigraphy		X		X		
SDS		X		X		
SAS		X		X		
Men-QoL		X	X	X	X	X
Others						
Estazolam dose		X	X	X	X	X
Adverse events		X	X	X	X	X
Patients's satisfaction						X
Success of blinding						X

Recruitment: Participants of the study will be recruited through the outpatient clinic, hospital-based Wechat advertising, and posters in Shanghai Municipal Hospital of Traditional Chinese Medicine in Shanghai, China. Those who are interested in participating in the study will be screened either via telephone or on-site. They will receive eight-week treatment of acupuncture and a series of assessments on sleep disorders free of charge. During the study, participants will be required to wear a wrist Actigraphy to record their sleep quality. The participants who meet the inclusion criteria will be informed of the research details and asked to sign the informed consent before the treatment commence.

Inclusion Criteria: Participants with the following conditions will be included:

- (1) Menopausal women aged 45~60;
- (2) Participants who meets the diagnostic criteria of traditional Chinese medicine in accordance with the above perimenopausal and insomnia[22];
- (3) Participants in line with the international classification of sleep disorders (ICSD-3) for the diagnosis of insomnia;
- (4) Participants whose PSQI score is higher than five;

(5) Participants who voluntarily agree to join the study and to sign the written informed consent form for the RCT.

Exclusion Criteria: Participants with the following conditions will be excluded:

(1) Participants who have severe mental illness such as depression, anxiety, schizophrenia and so on;

(2) Participants with serious heart, brain, kidney or liver disease;

(3) Participants with secondary insomnia caused by systemic diseases such as pain, fever, cough, surgery and external environment disturbance;

(4) Participants who have taken hormones for 6 or more months;

(5) Participants who are pregnant or currently lactating.

Intervention:

Participants in each group will receive 18 real or sham treatments within an eight-week period. In each group, they will receive either acupuncture or sham acupuncture treatment three times per week for the first four weeks, twice per week for the next two weeks and once per week for the last two weeks. Each treatment will last for 30 minutes. The each patient will be placed in a separate quiet space and lying supine. The acupuncturists will be registered practitioners with more than three years of experience in acupuncture practice. In order to increase the patients' compliance, they will accept the intervention and assessment by reservation, and they will receive some financial subsidies (200 RMB) after the follow-up assessments.

In the treatment group, acupuncture needles will be standard stainless steel, sterile, and disposable (0.25 × 40 mm and 0.30×40mm in length; Jia Jian, China). The control group will use the Streitberger Placebo-needle at the same acupoints. The treatment methods of acupuncture and acupoints are shown in Table 2.

Table 2. Details of intervention

	Intervention group	Control group
Main Acupoints	GV20, GV24, GV29, RN6, RN4, SP6, HT7, EX-HN22	GV20, GV24, GV29, RN6, RN4, SP6, HT7, EX-HN22
Added Acupoints	Kidney yang deficiency GV4, BL23	Kidney yang deficiency GV4, BL23
	Kidney yin deficiency KI3, KI7	Kidney yin deficiency KI3, KI7
Depth of insertion	GV20, GV24, GV29, RN6, RN4, EX-HN22 10mm	No insertion
	HT7, SP6, GV4, BL23, KI7, KI3 15mm	
Needle type	Steel needle (Wuxi Jiajian Medical Co. Ltd. Wuxi, China)	Blunt-tip needle (Streitberger Placebo-needle)
Needle sensation	With de-qi sensation	Without de-qi sensation
Electric stimulation	Needle on GV20, GV29 connected to G6805-2 Multi-Purpose Health Device (Shanghai Medical Instruments High-Techno, China). with electric pulse at a frequency of 2.5 Hz and an intensity of 45 mA	Needle on GV20, GV29, connected to G6805-2 Multi-Purpose Health Device (Shanghai Medical Instruments High-Techno, China). without electric pulse
Frequency and duration	Three times per week for the first four weeks	Three times per week for the first four weeks
	Twice per week for the next two weeks	Twice per week for the next two weeks
	Once per week for the final two weeks	Once per week for the final two weeks

The acupuncture group:

Participants in this group will receive acupuncture treatment during the first eight weeks. According to our in depth research on insomnia and acupuncture experience, Baihui (GV20), Shenting (GV24), Yintang (GV29), Qihai (RN6), Guanyuan (RN4) and bilateral Anmian (EX-HN22), Sanyinjiao (SP6), Shenmen (HT7) will be used as main acupoints[17]. Acupuncturists will be permitted to add acupuncture points based on TCM syndrome differentiation [23]. According to our own clinical practice, potentially relevant extraneous points include Mingmen (GV4) and Shenshu (BL23) for kidney yang deficiency, and TaiXi (KI3) and Fuliu (KI7) for kidney yin deficiency. The needles will be inserted into the skin to the depth of 10–30 mm and manipulated manually (including lifting, thrusting, and rotating) until the patient reports needling sensations (Deqi sensation). The needles on GV20 and GV29 will be connected to a G6805-2 Multi-Purpose Health Device (Huayi company), using continuous wave type, frequency at 2.5HZ, and intensity of 45 mA. Needles will be retained for 30 minutes before removal.

The control group:

Participants in this group will receive sham acupuncture treatment. We will use a non-invasive placebo control, the Streitberger Placebo-needle. Acupoints are the same as with acupuncture group, without

insertion. An electroacupuncture apparatus (G6805-2 Multi-Purpose Health Device) will be set beside the patients and connected to the GV20 and GV29, without electrical pulse.

Outcome measures

We will make assessments at baseline, week four, week eight, post-treatment, and with follow-ups at the first and third month after the end of treatment.

Primary outcome

Pittsburgh Sleep Quality Index (PSQI):

The PSQI is a self-rated questionnaire used to measure general sleep quality. It comprises 19 self-rated items and five other-rated items[20]. It assesses sleep based on seven domains in the past month: sleep duration, sleep disorders, sleep-onset latency, daytime dysfunction, sleep efficiency, use of medications to sleep, and overall sleep quality[24]. Each domain is rated from 0 to 3, and the accumulated scores of the seven domains constitute the total score of the PSQI (0–21). A total score of more than 5 indicates poor quality of sleep[25].

Secondary outcomes:

Insomnia Severity Index (ISI):

ISI is designed to assess the nighttime and daytime symptoms of insomnia. It includes seven items rated on a 0- to 4-point scale, total score ranges from 0–28[26]. All self-report questionnaires are conducted in Chinese. A higher score indicates a more severe insomnia. ISI classification: clinical insomnia (score 0-7), mild insomnia (8-14), moderate insomnia (score 15-21), and severe insomnia (score 22-28).

Actigraphy assessments:

The wActiSleep-BT actigraph (Actigraph LLC, Pensacola, USA), will be worn on the patients' wrist, which can record the sleep quality by noting sleep awakenings (SA), total sleep time (TST), sleep onset, sleep latency as well as the efficiency of sleep. The analysis of sleep condition and sleep quality will be performed by the software ActiLife6 (Version 6.8.1, Actigraph LLC) [17, 20].

Self-Rating Anxiety Scale (SAS):

We will use SAS to assess the degree of anxiety in patients. The questionnaire includes 20 items, as follows: 15 negative and 5 positive descriptions, the opposite of the score[27]. The total score of the 20 items more than 40 points, indicating the state of anxiety. A higher score indicates more severe anxiety.

Self-Rating Depression Scale (SDS):

The SDS is a self-rated scale for evaluating the severity of depression [28]. The main statistical index consists of 20 questions. We first calculate the standard score by ranking the participant's responses to

20 questions. From this figure the total score determined as is 1.25 times of the standard score, making use of the integral part [17]. The demarcation point of depression is 50 points, the higher the score, the more significant the depression tendency.

Menopause Quality of Life (MenQOL):

MenQOL is a questionnaire used to assess the quality of life in menopause, including 29 items, which are divided into four domains, Vasomotor (Items 1 to 3), Psychosocial (Items 4 to 10), Physical (Items 11 to 26), and Sexual (Items 27 to 29). For analyses, we convert the item scores to a score ranging from 1 to 8 (NO, 0-6). Each domain mean ranges from 1 to 8. The overall questionnaire score is the mean of the domain means. A higher score indicates a worse quality of life [29].

Estazolam dose:

Patients will be allowed to take Estazolam (0.5mg-2.0mg) when they have difficulty in falling asleep for more than three consecutive days during the trial. They will be asked to record the dose and the time of taking Estazolam on the case report form (CRF).

Sample size:

The sample size calculation was based on the change of PSQI scores. Systematic review[21] shows that at least 2.70 point difference in PSQI scores between acupuncture and sham-acupuncture has clinical significance. We expected that acupuncture would out-perform sham-acupuncture by three points, therefore a sample size of 35 participants should be recruited in each group. Considering about a 20% dropout rate, each group will take 42 cases. So a total of 84 participants should be recruited for this RCT.

$N = \dots$

Randomization and Blinding:

We will use the block randomization method. We plan to use SPSS version 23.0 software to generate a random number table. The participants who meet the criteria will be randomly assigned to acupuncture group or control group with 1:1 ratio. The random allocation sequence will be generated in a block. The block size will be randomized to six. Then the independent researcher will pre-pack envelopes with the corresponding number according to the randomization schedule. Each patient will receive an envelope labeled with their name containing their numbered group assignment.

Patients will be asked to wear an eye-patch when they receive treatment. Only the acupuncturists will know the group assignments. The participants and other researchers (including the data analysts, outcome assessors, statistician will be blinded to group allocation. In order to ensure the successful implementation of the blinding method, all researchers will be trained before the trial begins.

Safety and undesirable effects:

Patients will be advised in the event of any undesirable effect after the application of acupuncture that they discontinue acupuncture and contact their doctors, the doctor will diagnose and treat the adverse reactions. They will also be asked to contact the researcher who will have completed an undesirable effect form as well as the undesirable effect record log. Adverse events such as fainting, allergies and pain may occur in acupuncture clinical trials[30]. If the participant has the reaction of fainting, the needle will be immediately withdrawn. The patient will be returned to a supine position and given warm water or sugar water. The patient will be allowed to rest until they have made a full recovery. In the case of pain and allergies, the needle should be taken out immediately and treated according to the specific circumstances.

Any adverse event (include any discomfort, symptoms, or diseases occurring during the trial) will be reported by the patients and doctors after acupuncture treatment. All details of adverse events will be reported in the Case Report Form. At the end of the study we will analyze the influence of all events.

Data management:

Patient characteristics will be recorded in CRFs, which will be stored in the researcher's work office. Quality of CRF completion will be monitored by the specified researcher. All the original data of the RCT will be entered into the ResMan research Manager of the Clinical Trial Management Public Platform. The relevant users will be trained and the system will be tested before it is officially launched to ensure that the system meets the trial requirements. Only relevant personnel will receive the account number and password once the system is officially launched. The clinical supervisor will monitor the work of the clinical trial center at least once a month.

Statistical analysis:

All data will be analyzed by Intention-to-Treat (ITT), including data from any participants who have dropped out of the RCT during the trial. All data will be entered twice by two different researchers to ensure the accuracy. The statistical analysis of data will be carried out by SPSS 23.0 software. Descriptive characteristics of baseline statistics will be reported as the mean \pm standard deviation or median. The two study groups will be compared using t test for continuous measures and chi-square tests for categorical measures. To compare changes in sleep quality between two groups as the primary analysis, a repeated measures ANOVA (General Linear Model) will be used. P-Values ≤ 0.05 will be considered statistically significant.

Monitoring:

To improve the quality of this RCT, the whole process will be conducted under the supervision of a qualified clinical trial expert and be carried out by Shanghai Municipal Hospital of Traditional Chinese Medicine. The Clinical Research Center of Drugs of the Shanghai University of Traditional Chinese Medicine will provide data monitoring with access to any interim results and will make the final decision to terminate the trial if necessary.

Clinical trial registration: This RCT was registered in the Chinese Clinical Trial Registry (ChiCTR1800018645), Date: 2018-10-01. URL: <http://www.chictr.org.cn/showproj.aspx?proj=31482>

Discussion

Acupuncture treatment has been practiced in China for more than two millennia[31] and in recent years has been widely used in western countries [32]. Recent studies have found that acupuncture can be used to relieve vasomotor symptoms during the MT [23], but have not yet established the effects of acupuncture on insomnia during the MT. This is a valuable clinical RCT aimed at treating insomnia in menopausal women. We use acupuncture and sham acupuncture[33] to evaluate the efficacy of acupuncture on improving sleep quality. In addition, we use a strictly randomized controlled trial, utilizing the control group method to reduce test bias and ensure the reliability of the RCT.

The improvement on previous trials offered in this RCT is the method of point selection. Individualized treatment upon syndrome differentiation is the basis of TCM theory[23], and the selection of points is a question for the world acupuncture researchers [34]. To give some evidence for the effects of treatment upon syndrome differentiation, we set eight points as main points, and add two acupoints based on individual differences in the syndromes. Another advantage is the use of Actigraphy. This wrist-worn sleep monitor can provide an objective assessment for the treatment effect [35]. This is very useful for us to compare and evaluate the change of patients' sleep quality before and after treatment.

Another innovation in this trial is the design of the treatment frequency. In the first month we will give full dose (three times per week) treatment. After the initial month we will decrease the frequency of the acupuncture treatment. This will be beneficial for us to observe the acupuncture therapeutic effect, and to achieve the purpose of maintaining the minimum therapeutic dose for the long-term.

However, this RCT still faces some challenges. (1) The single-blinded method. For clinical trials of acupuncture, it is inevitable to know about the acupuncture treatment or sham acupuncture treatment for the acupuncturists. All the participants will be asked to wear the eye-patch and arranged in a separate quiet space during the treatment. (2) The application of the sham acupuncture method. The acupuncturists will receive several sessions of strict training before the study begins to ensure the proper sham acupuncture techniques. (3) The challenge of participant compliance. We will provide patient outreach over the phone to arrange the treatment time reasonably and to improve the attendance rate of the patients. The researchers will conduct follow-up interviews by phone to record patient outcomes.

To achieve our clinical goals, we will strive to standardize every step of the study, including acupoints selection, acupuncture operation, needle apparatus, and therapists' clinical experience. We expect that this trial will provide strong positive evidence for acupuncture treatment of sleep disorders in the perimenopausal period.

Trial status:

The first investigators' meeting took place on 31 August 2018. The RCT is in preparation now and will launch in November 2018. Recruitment is expected to end in late 2020.

Declarations

Ethics approval and consent to participate:

This RCT was approved by the Ethics Committee of Shanghai Municipal Hospital of Traditional Chinese Medicine on September 18, 2018 (certificate number 2018SHL-KY-09). The purpose, procedures, and potential risks of the RCT will be explained clearly to the participants. All participants shall give their written informed consent before joining the RCT.

Consent for publication:

Not applicable.

Availability of data and material:

The trial data will be available from the corresponding author upon reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding:

The study is partly supported by grants from Shanghai Committee of Science and Technology, China (No 18401905200). Shanghai Hospital Development Center, China (No SHDC12016124), Shanghai Municipal Commission of Health and Family Planning, China (No 201640026) and State Administration of Traditional Chinese Medicine, China (Grant No. JDZX2015024).

Authors' contributions:

The trial was designed and developed by SFX and LXL. The manuscript was drafted by SSL. The protocol was carefully revised, edited by SSL and PY. XY, AB, and TTL contributed the discussion. All authors read and approved the final manuscript.

Acknowledgements:

Shifen Xu provided general support as the head of the Acupuncture Department in Shanghai Municipal Hospital of Traditional Chinese Medicine, and was responsible for the design of this RCT and for communicating important protocol modifications to relevant parties. We would like to thank Dr. Philippa Hazlewood from the International Education College, Shanghai University of Traditional Chinese Medicine, for her editorial support.

Publisher's note:

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors' information:

1Shanghai Municipal Hospital of Traditional Chinese Medicine, Shanghai University of Traditional Chinese Medicine, Shanghai 200071, China.

2School of Chinese Medicine, University of Hong Kong, Hong Kong, China.

3University of Maryland School of Medicine, Baltimore, MD, USA 21201

References

1. Prairie BA, Wisniewski SR, Luther J, Hess R, Thurston RC, Wisner KL, Bromberger JT: **Symptoms of depressed mood, disturbed sleep, and sexual problems in midlife women: cross-sectional data from the Study of Women's Health Across the Nation.** *Journal of women's health* 2015, **24**(2):119-126.
2. Chiu HY, Hsieh YJ, Tsai PS: **Acupuncture to Reduce Sleep Disturbances in Perimenopausal and Postmenopausal Women: A Systematic Review and Meta-analysis.** *Obstetrics and gynecology* 2016, **127**(3):507-515.
3. Yazdi Z, Sadeghniiat-Haghighi K, Ziaee A, Elmizadeh K, Ziaeeha M: **Influence of sleep disturbances on quality of life of Iranian menopausal women.** *Psychiatry journal* 2013, **2013**:907068.
4. Sun D, Shao H, Li C, Tao M: **Sleep disturbance and correlates in menopausal women in Shanghai.** *Journal of psychosomatic research* 2014, **76**(3):237-241.
5. Cong Fu NZ, Zhen Liu, Lu-hua Yuan, Chen Xie, Wen-jia Yang, et al.: **Acupuncture Improves Perimenopausal Insomnia: A Randomized Controlled Trial.** *SLEEP* 2017, **40**:1-8.
6. Slopian R, Wichniak A, Pawlak M, Slopian A, Warenik-Szymankiewicz A, Sajdak S: **Disturbances of sleep continuity in women during the menopausal transition.** *Psychiatria polska* 2015, **49**(3):615-623.
7. Berecki-Gisolf J, Begum N, Dobson AJ: **Symptoms reported by women in midlife: menopausal transition or aging?** *Menopause* 2009, **16**(5):1021-1029.
8. Jin F, Tao M, Teng Y, Shao H, Li C, Mills E: **Knowledge and attitude towards menopause and hormone replacement therapy in Chinese women.** *Gynecologic and obstetric investigation* 2015, **79**(1):40-45.
9. Philip S. Eichling JS: **Menopause Related Sleep Disorders.** *Journal of Clinical Sleep Medicine* 2005, **1**:291-300.

10. Joshua z SA, Claire L, Sara Nowakowski: **Treatment of Insomnia, Insomnia Symptoms, and Obstructive Sleep Apnea During and After Menopause:Therapeutic Approaches.** *HHS Public Access* 2015, **1**:63-83.
11. H.Hachul TKPG, A. L. Maciel, F. Yagibara, S.Tufik, L. bittencourt: **Acupuncture improves sleep in postmenopause in a randomized, double-blind, placebo-controliied study.** *CLIMACTERIC* 2013, **16**:36-40.
12. Avis NE, Coeytaux RR, Levine B, Isom S, Morgan T: **Trajectories of response to acupuncture for menopausal vasomotor symptoms.** *Menopause* 2017, **24**(2):171-179.
13. Huang MI, Nir Y, Chen B, Schnyer R, Manber R: **A randomized controlled pilot study of acupuncture for postmenopausal hot flashes: effect on nocturnal hot flashes and sleep quality.** *Fertility and sterility* 2006, **86**(3):700-710.
14. Jennifer Hays JK, Ockene, Robert, Jane M, Kotchenm, et al.: **Effects of estrogen plus progestin on health-related quality of life.** *The NEW ENGLAND JONAL of MEDICINE* 2003, **348**:1839-1845.
15. Fredi K, Adriane Fugh-Berman: **Complementary and alternative medicine for menopausal symptoms- a review of randomized, controlled trials.** *ACADEMIA AND CLININ* 2002:805-814.
16. Yin X, Xu J, Dong B, Ma J, Chen Z, Yin P, Wu J, Zhu B, Cao Y, Zheng H *et al.*: **Efficacy and Safety of Electroacupuncture on Treating Depression Related Sleep Disorders: Study Protocol of a Randomized Controlled Trial.** *Evidence-based complementary and alternative medicine : eCAM* 2016, **2016**:1069597.
17. Yin X, Gou M, Xu J, Dong B, Yin P, Masquelin F, Wu J, Lao L, Xu S: **Efficacy and safety of acupuncture treatment on primary insomnia: a randomized controlled trial.** *Sleep medicine* 2017, **37**:193-200.
18. Wing-Fai Yeung K-FC, Kwok-Chu Tso,Shi-Ping,Zhang-jin Zhang,Lai-Ming Ho: **Electroacupuncture for residual insomnia associated with major depressive disorder- a randomized controlled trial.** *SLEEP* 2011, **34**:807-815.
19. Dong B, Chen Z, Yin X, Li D, Ma J, Yin P, Cao Y, Lao L, Xu S: **The Efficacy of Acupuncture for Treating Depression-Related Insomnia Compared with a Control Group: A Systematic Review and Meta-Analysis.** *BioMed research international* 2017, **2017**:9614810.
20. Cao Y, Yin X, Soto-Aguilar F, Liu Y, Yin P, Wu J, Zhu B, Li W, Lao L, Xu S: **Effect of acupuncture on insomnia following stroke: study protocol for a randomized controlled trial.** *Trials* 2016, **17**(1):546.
21. Cheuk DK, Yeung WF, Chung KF, Wong V: **Acupuncture for insomnia.** *The Cochrane database of systematic reviews* 2012(9):CD005472.
22. Sateia MJ: **International classification of sleep disorders-third edition: highlights and modifications.** *Chest* 2014, **146**(5):1387-1394.

23. Avis NE, Coeytaux RR, Isom S, Prevette K, Morgan T: **Acupuncture in Menopause (AIM) study: a pragmatic, randomized controlled trial.** *Menopause* 2016, **23**(6):626-637.
24. Jutta Backhaus KJ, Andreas Brocks, Dieter Riemann, Fritz Hohhagen: **Test-retest reliability and validity of the Pittsburgh Sleep Quality Index in primary insomnia.** *Journal of Psychosomatic Research* 2002, **53**:737-740.
25. Daniel J. Buysse CFRI, Timmothy H. Monk, Susan R. Berman, David J. Kupfer: **The Pittsburgh Sleep Quality Index- a new instrument for psychiatric practice and research.** *Psychiatry Research* 1988, **28**:193-213.
26. Célyne H. Bastien AV, Charles M. Morin: **Validation of the Insomnia Severity Index as an outcome measure for insomnia research.** *Sleep medicine* 2001, **2**:297-307.
27. Li RX, Ma M, Xiao XR, Xu Y, Chen XY, Li B: **Perimenopausal syndrome and mood disorders in perimenopause: prevalence, severity, relationships, and risk factors.** *Medicine* 2016, **95**(32):e4466.
28. WILLIAM W. K. ZUNG D: **A Self-Rating Depression Scale.** *ARCHIVES OF GENERAL PSYCHIATRY* 1965, **12**:63-70.
29. Julie Lemieux MD, Wendy R, Paul E., James N. Kathleen I, et al.: **Quality of Life From Canadian Cancer Trials Group MA.17R: A Randomized Trial of Extending Adjuvant Letrozole to 10 years.** *JOURNAL OF CLINICAL ONCOLOGY* 2018, **36**:563-571.
30. Li YQ, Lu YY, Zhang J, Guo Y: **Analysis on the situation of adverse reaction to acupuncture and acupuncture risk.** *Chinese Acupuncture & Moxibustion* 2011, **31**(8):764-768.
31. ADAM BURKE DM, CLAIRE DYE, LAURA CHYU: **Acupuncture use in the United States- findings from the National Health Interview Survey.** *THE JOURNAL OF ALTERNATIVE AND COMPLEMENTARY MEDICINE* 2006, **12**:639-648.
32. Yeung WF, Chung KF, Yung KP, Ho FY, Ho LM, Yu YM, Kwok CW: **The use of conventional and complementary therapies for insomnia among Hong Kong Chinese: a telephone survey.** *Complementary therapies in medicine* 2014, **22**(5):894-902.
33. Lao L, Bergman S, Hamilton GR, Langenberg P, Berman B: **Evaluation of acupuncture for pain control after oral surgery: a placebo-controlled trial.** *Arch Otolaryngol Head Neck Surg* 1999, **125**(5):567-572.
34. Albert Yeung DM, Jonathan Alpert, Yantao Ma, Maurizio Fava: **Acupuncture for Major Depressive Disorder: Has Its Efficacy Been Disproved?** *LETTERS TO THE EDITOR* 2007:1617-1622.
35. Short MA, Gradisar M, Lack LC, Wright H, Carskadon MA: **The discrepancy between actigraphic and sleep diary measures of sleep in adolescents.** *Sleep medicine* 2012, **13**(4):378-384.

Figures




Figure 1

Flowchart of the trial

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SPIRITchecklist.doc](#)