

# Ocular biomarkers for cognitive impairment in nonagenarians; a prospective cross-sectional study

Jacoba Alida van de Kreeke (✉ [ja.vandekreeke@amsterdamumc.nl](mailto:ja.vandekreeke@amsterdamumc.nl))

Amsterdam UMC, location VUmc <https://orcid.org/0000-0002-0165-1839>

**Nienke Legdeur**

Alzheimer Center Amsterdam, Dept. of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, location VUmc, Amsterdam

**Maryam Badissi**

Alzheimer Center Amsterdam, Dept. of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, location VUmc, Amsterdam

**H. Ton Nguyen**

Ophthalmology Dept., Amsterdam UMC, location VUmc, Amsterdam

**Elles Konijnenberg**

Alzheimer Center Amsterdam, Dept. of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam

**Jori Tomassen**

Alzheimer Center Amsterdam, Dept. of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam

**Mara ten Kate**

Alzheimer Center Amsterdam, Dept. of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam

**Anouk den Braber**

Dept. of Biological Psychology, VU University Amsterdam, Amsterdam

**Andrea B. Maier**

Dept. of Medicine and Aged Care, @AgeMelbourne, Royal Melbourne Hospital, University of Melbourne, Melbourne / Dept. of Human Movement Sciences, @AgeAmsterdam, Amsterdam Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam

**H. Stevie Tan**

Ophthalmology Dept., Amsterdam UMC, Location VUmc, Amsterdam

**Frank D. Verbraak**

Ophthalmology Dept., Amsterdam UMC, location VUmc, Amsterdam

**Pieter Jelle Visser**

Alzheimer Center Amsterdam, Dept. of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam

## Research article

**Keywords:** retina, optical coherence tomography, fundus photography, Singapore I Vessel Assessment, nonagenarians, ocular biomarkers, cognitive impairment

**Posted Date:** July 3rd, 2019

**DOI:** <https://doi.org/10.21203/rs.2.10871/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published at BMC Geriatrics on April 28th, 2020. See the published version at <https://doi.org/10.1186/s12877-020-01556-1>.

# Abstract

**Background:** Ocular imaging receives much attention as a source of potential biomarkers for dementia. In the present study, we analyze these ocular biomarkers in cognitively impaired and healthy participants in a population aged over 90 years (= nonagenarian), and elucidate the effects of age on these biomarkers. **Methods:** For this prospective cross-sectional study, we included individuals from the EMIF-AD 90+ study, consisting of a cognitively healthy (N=67) and cognitively impaired group (N=33), and the EMIF-AD PreclinAD study, consisting of cognitively healthy controls aged  $\geq 60$  (N=198). Participants underwent Optical Coherence Tomography (OCT) and fundus photography of both eyes. OCT was used to assess total and individual inner retinal layer thickness in the macular region (Early Treatment Diabetic Retinopathy Study circles) as well as peripapillary retinal nerve fiber layer thickness, fundus images were analyzed with Singapore I Vessel Assessment to obtain 7 retinal vascular parameters. Values for both eyes were averaged. Differences in ocular biomarkers between the 2 nonagenarian groups were analyzed using linear regression, differences between the individual nonagenarian groups and controls were analyzed using generalized estimating equations. **Results:** Ocular biomarkers did not differ between the healthy and cognitively impaired nonagenarian groups. Both nonagenarian groups differed in most ocular biomarkers from the younger controls. **Conclusion:** Ocular biomarkers were not associated with cognitive impairment in nonagenarians, making their use as a screening tool for dementing disorders in this group limited. However, ocular biomarkers were significantly associated with chronological age, which were very similar to those ascribed to occur in Alzheimer's Disease.

## Introduction

The eye, and more specifically the retina, shares many similarities with the brain. Both are derived from the same embryological tissue and consist of a complex combination of neuronal tissue and glial cells (1, 2). One could consider the retina to be an extension of the brain (1, 2). Assessment of the retina may provide information on diseases causing cognitive impairment, such as Alzheimer's disease (AD). Therefore, studies have been performed on the diagnostic value of optical techniques used to examine the retina in AD, as these ocular markers may be less invasive and cheaper than most established biomarkers for brain diseases (3-8).

There is ample evidence of retinal changes in dementia (3-8). In AD, retinal thinning has been observed on Optical Coherence Tomography (OCT), especially of the inner layers (3, 5). Studies have also shown differences between individuals with and without dementia in the (micro)vascular state of their retinal vessels (such as vessel width, tortuosity and fractal dimension) through fundus photography, although these results were sometimes contradicting (8-11).

It is estimated that up to 40% of individuals aged 90 and over suffer from dementia (12). In nonagenarians (individuals between the ages of 90-100), dementia is mostly caused by mixed pathologies, including AD and vascular brain damage (13, 14). This may be reflected in the retinal biomarkers, causing the expected changes due to for example AD (retinal thinning and changes in microvasculature) to be less discriminative in this elderly population compared to younger individuals. Furthermore, many of the

changes occurring in the retina of individuals with dementia also occur naturally with the aging process (15, 16).

In this cross-sectional study, we aimed to 1) investigate whether retinal (layer) thickness and retinal vascular parameters differ between cognitively healthy and cognitively impaired nonagenarians and 2) elucidate the effects of age on these ocular biomarkers.

## Methods

### Participants

This study consists of combined data from 2 Amsterdam UMC sub-studies of the European Medical Information Framework for Alzheimer's Disease (EMIF-AD): the EMIF-AD 90+ study and the EMIF-AD PreclinAD cohort. The 90+ study consists of cognitively healthy and cognitively impaired subjects, aged  $\geq 90$  or over. For extensive recruitment information we refer to our set-up paper of this study by Legdeur et al (17). The PreclinAD cohort (18) is a cohort consisting of cognitively healthy participants (monozygotic twins) aged  $\geq 60$ , recruited from the Netherlands Twin Registry (19). The studies adhered the Tenets of the Declaration of Helsinki and written informed consent was obtained from all participants. The Medical Ethics Committee of the Amsterdam UMC approved both studies.

For complete in- and exclusion criteria of the EMIF-AD 90+ study, we refer to our set-up paper of this study by Legdeur et al (17). In short, inclusion criteria for the cognitively normal group of the 90+ study were: age  $\geq 90$  years and cognitively healthy. This group is referred to as 'healthy nonagenarian' group.

Inclusion criteria for individuals with cognitive impairment (CI) of the 90+ study were: a diagnosis of amnesic Mild Cognitive Impairment (aMCI) (20) or a diagnosis of probable or possible AD (21). As during the study we had difficulties identifying subjects of 90 years and older with aMCI or probable or possible AD, we broadened the inclusion criteria in this group to subjects older than 85 years. Six individuals from this group were aged 85-90 years. This group is referred to as 'CI nonagenarian' group.

Inclusion criteria for the PreclinAD study were: age  $\geq 60$  years, monozygosity and cognitively healthy. For complete in-/exclusion criteria we refer to the set-up paper of this study by Konijnenberg et al (18). This group is referred to as the 'control' group.

### Medical history

Data about the medical and family history and medication use, in particular on the presence of diabetes mellitus, hypertension and coronary disease, were collected through a structured interview, in combination with information provided by the study partner (if available), general practitioner and/or medical specialist.

### Ophthalmological examination

All participants underwent the following ophthalmological examinations: best corrected visual acuity, intra-ocular pressure, refraction data, slit lamp examination, indirect funduscopy, fundus photography and OCT. Controls received tropicamide 0.5% to enable these examinations, nonagenarians both tropicamide 0.5% and phenylephrine 5% (as mydriasis was harder to achieve in these very elderly patients). If a nonagenarian suffered from coronary stenosis, only tropicamide was given, due to the slight risk of phenylephrine inducing a coronary spasm. All photographs/OCT images were assessed by an experienced ophthalmologist (HTN or FDV) for unexpected pathology. Participants suffering from ophthalmological conditions severely interfering with the (neuro)retina or image quality were excluded from analyses (severe cataract, macular degeneration, glaucoma, diabetic retinopathy, vascular occlusions). Eyes with diseases considered to interfere with the OCT measurements excluded from OCT analyses could still be included in the SIVA analyses and vice versa (e.g. AMD with geographical atrophy interfered with OCT, but not fundus image analyses). This resulted in a slightly different study population for the OCT and SIVA analyses.

## **Optical Coherence Tomography**

Using spectral domain OCT (Spectralis, Heidelberg), dense macular scans (49 B-scans) and axonal ring scans around the optic nerve head (ONH) were acquired. Total retinal thickness and individual layer thickness was measured in the macular region. The following individual retinal layers were analyzed: retinal nerve fiber layer (RNFL), ganglion cell layer (GCL) and inner plexiform layer (IPL). A distinction was made between the inner and outer macular ring according to the standard Early Treatment and Diabetic Retinopathy Study (ETDRS) macular grid (1-3mm around the fovea for inner ring and 3-6mm around the fovea for outer ring). For further details on the acquiring of OCT data we refer to our earlier paper by van de Kreeke et al (22).

## **Fundus photography and quantitative assessment of retinal vasculature**

Digital fundus images were made of the fundus of both eyes in all participants (Topcon TRC 50DX type IA). All images were graded by a trained grader (JAvdK) using the Singapore I Vessel Assessment (SIVA) software (version 3.0, National University of Singapore, Singapore) (9-11). The following 7 retinal vascular parameters were analyzed: central retinal artery equivalent (CRAE), central retinal vein equivalent (CRVE), arteriole-venular ratio (AVR), fractal dimension of the arteriolar network (FDa), fractal dimension of the venular network (FDv), curvature tortuosity of the arterioles (cTORTa) and curvature tortuosity of the venules (cTORTv). All values for retinal vascular parameters were measured within zone C (i.e. 0.5 – 2 disc diameters around the optic nerve head). For further information on the analyses of fundus images we refer to our earlier paper by van de Kreeke et al (23).

## **Statistical analysis**

First, we compared group means of all ocular outcome measures of the healthy and CI nonagenarian groups using linear regression, corrected for age, sex and diabetes. Additionally, mean differences between both nonagenarian groups and younger healthy controls were obtained using Generalized Estimating Equations (GEE). GEE was used to correct for clustering in the data from twin pairs in the control group. It also allowed us to correct for confounders such as sex and a diagnosis of diabetes. We deliberately did not correct this analysis for age, to illustrate the differences based on aging effects. Curvature tortuosity (cTORT) values for arteries and veins were log-transformed to normalize their distribution. All statistical analyses were performed using SPSS (IBM, version 22).

## Results

From the total 298 participants included in the cohorts, 51 (17.1%) participants were excluded for both the OCT and SIVA analyses, but these were not necessarily the same participants, although there was a high overlap. For the OCT analyses, 9 were excluded due to low quality scans/failed imaging and 42 due to ophthalmological pathology. For the SIVA analyses, 24 were excluded due to low quality images/failed imaging and 27 due to ophthalmological pathology. Interfering ophthalmological pathology consisted mostly of glaucoma and (severe) AMD. Additional file 1 shows the reasons for exclusion in more detail, categorized per group (control, healthy nonagenarian and CI nonagenarian). Although the two study populations (i.e. for OCT and SIVA analyses) were slightly different from each other in terms of included individuals, they were very similar in their demographics, and statistical analyses revealed no significant differences. As such, table 1 shows the demographics of the combined study populations included for analyses.

Table 1: Demographics of the study population.

When comparing the two nonagenarian groups, no significant differences in any of the ocular biomarkers were found (table 2).

Table 2: Differences between means for ocular biomarkers in both the healthy and the CI nonagenarian groups.

Both the healthy and the CI nonagenarians differed significantly in multiple ocular biomarkers when compared to the younger control group, with most ocular parameters being lower in the nonagenarian groups. Figures 1 and 2 show boxplots for the 3 groups, table 3 shows the mean differences of both nonagenarian groups compared to the control group, corrected for twin dependencies, sex and diabetes.

<Figure 1>

Figure 1 title: Boxplots for retinal layer thicknesses as measured with Optical Coherence Tomography in the 3 groups.

Figure 1 legend: Dotted line represents the mean. CI = Cognitively Impaired, ONH = Optic Nerve Head.

<Figure 2>

Figure 2 title: Boxplots for retinal vascular parameters obtained with Singapore I Vessel Assessment in the 3 groups.

Figure 2 legend: Dotted line represents the mean. CI = Cognitively Impaired, ONH = Optic Nerve Head.

Table 3: Differences in ocular biomarkers of both nonagenarian groups compared to the younger control group.

## Discussion

Ocular biomarkers showed no differences between cognitively healthy and cognitively impaired nonagenarians. However, nonagenarians did show extensive differences in their ocular biomarkers when compared to a younger control group, which were very similar to changes often attributed to AD in existing literature.

Our analyses showed no statistical difference in any of the studied ocular biomarkers between healthy and CI nonagenarians (table 2), suggesting that ocular biomarkers have a limited role in the detection of cognitive impairment in nonagenarians. A possible explanation for the lack of differences may be that other disorders that can have a profound effect on ocular biomarkers (such as hypertension, diabetes and coronary/cardiovascular disease), are also more prevalent in nonagenarians, and may obscure a possible neurodegenerative effect (24-26). Another explanation can be that other aging related processes affect ocular biomarkers, again obscuring differences between the nonagenarian groups. Indeed, we found that both nonagenarian groups showed large differences in ocular biomarkers, including total/GCL/IPL thickness in the macula, average and most individual segments pRNFL thickness and all retinal vascular parameters except tortuosity (figures 1 and 2, table 3), when compared to younger healthy controls. Several studies showed that aging is associated with thinning of the retina, as well as retinal vascular changes (15, 16, 26-30). Normal aging unavoidably causes wearing of the body, as is seen in many other organ systems (31). Especially in nonagenarians, compensatory mechanisms start failing, and such wear and tear effects become more and more pronounced, resulting in multimorbidity (32). It is likely that this process also occurs in the retina. Blood vessels gradually start to decline due to aging effects like atherosclerosis, causing damage to (micro)vessels, leading to changes in retinal vascular parameters such as thinner vessel calibers and a lower fractal dimension of the vascular network (26). This in turn may lead to a lower or insufficient blood and oxygen supply, causing damage to the neuronal tissue of the retina, resulting in the thinning of its layers (33, 34).

A problem we noticed when performing this study was the difficulty to make reliable images in nonagenarians. In this population there was a very high prevalence of ophthalmological pathology. Some of the diseases were already known at the time of the study and under treatment of an ophthalmologist, but several participants suffered, without their knowing, from ophthalmological pathology requiring medical attention. This resulted in a high exclusion of nonagenarian participants due to bilateral

pathology, as illustrated in supplemental figure 1. Approximately double the percentage of participants had to be excluded for OCT or SIVA analyses when comparing them with the younger control group. Additionally, acquiring good quality scans/images was harder in the nonagenarian groups, due to their lower endurance, physical impairments and, in the case of CI nonagenarians, cognitive functioning. This suggests that in a substantial (~25-30%) percentage of nonagenarians, ocular biomarkers for the diagnosis of dementia cannot be used, due to an inability to obtain reliable images/scans or bilateral ophthalmological pathology being present which interferes with the interpretation of the imaging.

A strength of the study was the extensive characterization of all participants. Participants were comprehensively screened for possible confounding pathologies, and were excluded (e.g. glaucoma, severe AMD, vascular occlusions etc.) or controlled for (diabetes).

Both nonagenarian groups were relatively small, limiting statistical power. Furthermore, as we included nonagenarians that were able to perform the study, which included 2-3 days of assessment (2 days in a hospital setting), this could also have introduced a bias, as only relatively healthy individuals of this elderly population will have been included in this study.

## Conclusion

We found no significant differences between the healthy and CI nonagenarian groups for any of the studied ocular biomarkers. We did find significant differences in many of the ocular biomarkers when comparing the two nonagenarian groups to a healthy younger control group, suggesting a large effect of age on these biomarkers. The combination of these findings, along with the difficulties we encountered in obtaining suitable images in our nonagenarian population, leads to the conclusion that the use of ocular biomarkers in a population of the oldest-old is very limited.

## List Of Abbreviations:

AD = Alzheimer's Disease

aMCI = amnesic Mild Cognitive Impairment

AMD = Age-related Macular Degeneration

AVR = Arteriole-Venular Ratio

BCVA = Best Corrected Visual Acuity

CDR = Clinical Dementia Rating

CERAD = Consortium to Establish a Registry for Alzheimer's Disease

CI = Cognitively Impaired

CRAE = Central Retinal Artery Equivalent

CRVE = Central Retinal Vein Equivalent

cTORTa = curvature Tortuosity of arteries

cTORTv = curvature Tortuosity of veins

ETDRS = Early Treatment and Diabetic Retinopathy Study

FDa = Fractal Dimension of arteries

FDv = Fractal Dimension of veins

GCL = Ganglion Cell Layer

GDS = Geriatric Depression Scale

GEE = Generalized Estimating Equations

IPL = Inner Plexiform Layer

MMSE = Mini Mental State Exam

OCT = Optical Coherence Tomography

ONH = Optic Nerve Head

pRNFL = peripapillary Retinal Nerve Fiber Layer

RNFL = Retinal Nerve Fiber Layer

RT = Retinal Thickness

SIVA = Singapore I Vessel Assessment

TICS-m = Telephone Interview for Cognitive Status modified

## **Declarations**

### **Ethics approval and consent to participate:**

The study adhered the Tenets of the Declaration of Helsinki and written informed consent was obtained from all participants. The Medical Ethics Committee of the Amsterdam UMC approved the study.

### **Consent for publication:**

Not applicable.

## Availability of data and material:

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Competing interest:

Pieter Jelle Visser is a member of the editorial board (section editor) of BMC Geriatrics. None of the other authors have competing interests.

## Funding:

This work has received support from the EU/EFPIA Innovative Medicines Initiative Joint Undertaking (EMIF grant number 115372).

These funding sources had no involvement in the collection, analysis and interpretation of data; in the writing of the report; or in the decision to submit the article for publication.

## Authors' contribution:

JAvdK wrote the manuscript, acquired data, analyzed the results, interpreted the data and contributed to the conception and design of the work. NL and HTN acquired data, contributed to the conception and design of the work and critically revised the manuscript. MB, EK, JT and MtK acquired data and critically revised the manuscript. AdB and ABM contributed to the conception and design of the work and critically revised the manuscript. HST critically revised the manuscript. FDV and PJV interpreted the data, contributed to the conception and design of the work and critically revised the manuscript. All authors read and approved the final manuscript.

## Acknowledgements:

Not applicable.

## References

1. London A, Benhar I, Schwartz M. The retina as a window to the brain-from eye research to CNS disorders. *Nat Rev Neurol*. 2013;9(1):44-53.
2. Jindal V. Interconnection between brain and retinal neurodegenerations. *Mol Neurobiol*. 2015;51(3):885-92.
3. Chan VTT, Sun Z, Tang S, Chen LJ, Wong A, Tham CC, et al. Spectral-Domain OCT Measurements in Alzheimer's Disease: A Systematic Review and Meta-analysis. *Ophthalmology*. 2018.
4. Chhablani PP, Ambiya V, Nair AG, Bondalapati S, Chhablani J. Retinal Findings on OCT in Systemic Conditions. *Semin Ophthalmol*. 2018;33(4):525-46.

5. den Haan J, Verbraak FD, Visser PJ, Bouwman FH. Retinal thickness in Alzheimer's disease: A systematic review and meta-analysis. *Alzheimers Dement (Amst)*. 2017;6:162-70.
6. Frost S, Martins RN, Kanagasingam Y. Ocular biomarkers for early detection of Alzheimer's disease. *J Alzheimers Dis*. 2010;22(1):1-16.
7. Lim JK, Li QX, He Z, Vingrys AJ, Wong VH, Currier N, et al. The Eye As a Biomarker for Alzheimer's Disease. *Front Neurosci*. 2016;10:536.
8. McGrory S, Cameron JR, Pellegrini E, Warren C, Doubal FN, Deary IJ, et al. The application of retinal fundus camera imaging in dementia: A systematic review. *Alzheimers Dement (Amst)*. 2017;6:91-107.
9. Cheung CY, Hsu W, Lee ML, Wang JJ, Mitchell P, Lau QP, et al. A new method to measure peripheral retinal vascular caliber over an extended area. *Microcirculation*. 2010;17(7):495-503.
10. Cheung CY, Tay WT, Mitchell P, Wang JJ, Hsu W, Lee ML, et al. Quantitative and qualitative retinal microvascular characteristics and blood pressure. *J Hypertens*. 2011;29(7):1380-91.
11. Cheung CY, Zheng Y, Hsu W, Lee ML, Lau QP, Mitchell P, et al. Retinal vascular tortuosity, blood pressure, and cardiovascular risk factors. *Ophthalmology*. 2011;118(5):812-8.
12. Corrada MM, Brookmeyer R, Berlau D, Paganini-Hill A, Kawas CH. Prevalence of dementia after age 90: results from the 90+ study. *Neurology*. 2008;71(5):337-43.
13. LoGiudice D, Watson R. Dementia in older people: an update. *Intern Med J*. 2014;44(11):1066-73.
14. Yang Z, Slavin MJ, Sachdev PS. Dementia in the oldest old. *Nat Rev Neurol*. 2013;9(7):382-93.
15. Wei Y, Jiang H, Shi Y, Qu D, Gregori G, Zheng F, et al. Age-Related Alterations in the Retinal Microvasculature, Microcirculation, and Microstructure. *Invest Ophthalmol Vis Sci*. 2017;58(9):3804-17.
16. Demirkaya N, van Dijk HW, van Schuppen SM, Abramoff MD, Garvin MK, Sonka M, et al. Effect of age on individual retinal layer thickness in normal eyes as measured with spectral-domain optical coherence tomography. *Invest Ophthalmol Vis Sci*. 2013;54(7):4934-40.
17. Legdeur N, Badissi M, Carter SF, de Crom S, van de Kreeke A, Vreeswijk R, et al. Resilience to cognitive impairment in the oldest-old: design of the EMIF-AD 90+ study. *BMC Geriatr*. 2018;18(1):289.
18. Konijnenberg E, Carter SF, Ten Kate M, den Braber A, Tomassen J, Amadi C, et al. The EMIF-AD PreclinAD study: study design and baseline cohort overview. *Alzheimers Res Ther*. 2018;10(1):75.
19. Boomsma DI, de Geus EJ, Vink JM, Stubbe JH, Distel MA, Hottenga JJ, et al. Netherlands Twin Register: from twins to twin families. *Twin Res Hum Genet*. 2006;9(6):849-57.
20. Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med*. 2004;256(3):183-94.

21. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology*. 1984;34(7):939-44.
22. van de Kreeke JA, Nguyen HT, den Haan J, Konijnenberg E, Tomassen J, den Braber A, et al. Retinal layer thickness in preclinical Alzheimer's disease. *Acta Ophthalmol*. 2019.
23. van de Kreeke JA, Nguyen HT, Konijnenberg E, Tomassen J, den Braber A, Ten Kate M, et al. Retinal and Cerebral Microvasculopathy: Relationships and Their Genetic Contributions. *Invest Ophthalmol Vis Sci*. 2018;59(12):5025-31.
24. De Clerck EE, Schouten JS, Berendschot TT, Kessels AG, Nuijts RM, Beckers HJ, et al. New ophthalmologic imaging techniques for detection and monitoring of neurodegenerative changes in diabetes: a systematic review. *Lancet Diabetes Endocrinol*. 2015;3(8):653-63.
25. Ikram MK, Ong YT, Cheung CY, Wong TY. Retinal vascular caliber measurements: clinical significance, current knowledge and future perspectives. *Ophthalmologica*. 2013;229(3):125-36.
26. Wang SB, Mitchell P, Liew G, Wong TY, Phan K, Thiagalingam A, et al. A spectrum of retinal vasculature measures and coronary artery disease. *Atherosclerosis*. 2018;268:215-24.
27. Hoffmann EM, Schmidtman I, Siouli A, Schuster AK, Beutel ME, Pfeiffer N, et al. The distribution of retinal nerve fiber layer thickness and associations with age, refraction, and axial length: the Gutenberg health study. *Graefes Arch Clin Exp Ophthalmol*. 2018;256(9):1685-93.
28. Won JY, Kim SE, Park YH. Effect of age and sex on retinal layer thickness and volume in normal eyes. *Medicine (Baltimore)*. 2016;95(46):e5441.
29. Huang QF, Wei FF, Zhang ZY, Raaijmakers A, Asayama K, Thijs L, et al. Reproducibility of Retinal Microvascular Traits Decoded by the Singapore I Vessel Assessment Software Across the Human Age Range. *Am J Hypertens*. 2018;31(4):438-49.
30. Leung H, Wang JJ, Rochtchina E, Tan AG, Wong TY, Klein R, et al. Relationships between age, blood pressure, and retinal vessel diameters in an older population. *Invest Ophthalmol Vis Sci*. 2003;44(7):2900-4.
31. Wyss-Coray T. Ageing, neurodegeneration and brain rejuvenation. *Nature*. 2016;539(7628):180-6.
32. Formiga F, Ferrer A, Sanz H, Marengoni A, Alburquerque J, Pujol R, et al. Patterns of comorbidity and multimorbidity in the oldest old: the Octabaix study. *Eur J Intern Med*. 2013;24(1):40-4.
33. Frydkjaer-Olsen U, Soegaard Hansen R, Pedersen K, Peto T, Grauslund J. Retinal Vascular Fractals Correlate With Early Neurodegeneration in Patients With Type 2 Diabetes Mellitus. *Invest Ophthalmol Vis Sci*. 2015;56(12):7438-43.

34. Lasta M, Pemp B, Schmidl D, Boltz A, Kaya S, Palkovits S, et al. Neurovascular dysfunction precedes neural dysfunction in the retina of patients with type 1 diabetes. *Invest Ophthalmol Vis Sci*. 2013;54(1):842-7.

## Tables

**Table 1:** Demographics of the study population.

	<b>Controls</b>	<b>Healthy nonagenarians</b>	<b>CI nonagenarians</b>
OCT analyses N	172	52	23
SIVA analyses N	173	50	24
Age (years)	70.4 ( $\pm 7.5$ )	92.4 ( $\pm 1.9$ )	91.9 ( $\pm 2.9$ )
Sex, female N (%)	105 (57.4%)	30 (52.6%)	20 (74.1%)
BCVA (LogMAR)	0.02 ( $\pm 0.10$ )	0.14 ( $\pm 0.21$ )	0.13 ( $\pm 0.22$ )
Intra-ocular pressure (mmHg)	14.4 ( $\pm 2.8$ )	15.9 ( $\pm 2.0$ )	15.6 ( $\pm 2.2$ )
Spherical equivalent	0.28 ( $\pm 1.86$ )	-0.02 ( $\pm 1.41$ )	0.05 ( $\pm 0.13$ )
MMSE (median, IQR)	29.0 (29.0-30.0)	29.0 (28.0-30.0)	24.0 (20.0-26.0)
Hypertension N (%)	75 (41.0%)	23 (40.4%)	16 (59.3%)
Diabetes mellitus N (%)	8 (4.4%)	3 (5.3%)	2 (7.4%)
Coronary disease N (%)	18 (9.8%)	14 (24.6%)	7 (25.9%)

Data are means from the groups combined unless otherwise specified. CI = Cognitively Impaired, N = Number of participants, OCT = Optical Coherence Tomography, SIVA = Singapore I Vessel Assessment, BCVA = Best Corrected Visual Acuity, MMSE = Mini-Mental State Exam, IQR = Inter-Quartile Range, CERAD = Consortium to Establish a Registry for Alzheimer's Disease. Coronary disease in this case means a history of myocardial infarction or angina pectoris.

**Table 2:** Differences between means for ocular biomarkers in both the healthy and the CI nonagenarian groups.

	Healthy nonagenarians	CI nonagenarians	SE	p- value
Macular retinal layer thickness:				
Total RT inner ring (µm)	323.5	317.3	5.0	0.218
Total RT outer ring (µm)	282.2	276.1	4.0	0.133
Total RT outer ring (µm)	23.0	22.2	0.8	0.353
RNFL inner ring (µm)	36.4	35.5	1.4	0.518
RNFL outer ring (µm)	43.9	42.9	1.6	0.550
GCL inner ring (µm)	30.0	29.9	1.1	0.940
GCL outer ring (µm)	36.6	35.5	1.0	0.275
IPL inner ring (µm)	25.5	24.9	0.8	0.449
IPL outer ring (µm)				
Peripapillary RNFL:				
Average (µm)	88.3	89.1	2.7	0.762
Nasal superior (µm)	90.6	96.6	4.9	0.229
Nasal (µm)	67.1	69.9	3.5	0.428
Nasal inferior (µm)	101.9	102.3	5.3	0.940
Temporal inferior (µm)	123.0	122.1	5.2	0.868
Temporal (µm)	69.1	66.9	3.2	0.471
Temporal superior (µm)	118.9	118.0	4.5	0.854
Retinal vascular parameters:				
CRAE	123.6	120.3	2.3	0.156
CRVE	184.3	179.7	4.6	0.325
AVR	0.676	0.673	0.014	0.818
FDa	1.152	1.152	0.011	0.977
FDv	1.137	1.137	0.011	0.987
cTORTa*	-9.75	-9.80	0.062	0.407
cTORTv*	-9.73	-9.69	0.054	0.472

Linear regression, corrected for age, sex and diabetes. CI = Cognitively Impaired, SE = Standard Error of difference, RT = Retinal Thickness, RNFL = Retinal Nerve Fiber Layer, GCL = Ganglion Cell Layer, IPL = Inner Plexiform Layer, CRAE = Central Retinal Artery Equivalent, CRVE = Central Retinal Vein Equivalent, AVR = Arteriole-Venular Ratio, FDa = Fractal Dimension of arteries, FDv = Fractal Dimension of veins, cTORTa = curvature Tortuosity of arteries, cTORTv = curvature Tortuosity of veins. \* Log transformation applied.

**Table 3:** Differences in ocular biomarkers of both nonagenarian groups compared to the younger control group.

	Estimated marginal mean of controls	Mean difference compared to controls					
		Healthy nonagenarians	SE	p-value	CI nonagenarians	SE	p-value
Macular retinal layer thickness:	<i>340.1</i>	<b>-18.0</b>	<b>3.0</b>	<b>&lt;0.001</b>	<b>-22.5</b>	<b>5.0</b>	<b>&lt;0.001</b>
Total RT	<i>294.8</i>	<b>-12.7</b>	<b>2.4</b>	<b>&lt;0.001</b>	<b>-18.1</b>	<b>3.7</b>	<b>&lt;0.001</b>
inner ring (µm)	<i>21.9</i>	<b>1.0</b>	<b>0.5</b>	<b>0.032</b>	<b>0.4</b>	<b>0.7</b>	<b>0.559</b>
Total RT	<i>36.3</i>	<b>0.2</b>	<b>0.9</b>	<b>0.829</b>	<b>-0.7</b>	<b>1.2</b>	<b>0.549</b>
outer ring (µm)	<i>49.8</i>	<b>-6.1</b>	<b>0.9</b>	<b>&lt;0.001</b>	<b>-6.5</b>	<b>1.7</b>	<b>&lt;0.001</b>
Total RT	<i>34.2</i>	<b>-4.2</b>	<b>0.6</b>	<b>&lt;0.001</b>	<b>-4.1</b>	<b>1.2</b>	<b>0.001</b>
outer ring (µm)	<i>41.0</i>	<b>-4.5</b>	<b>0.6</b>	<b>&lt;0.001</b>	<b>-5.2</b>	<b>0.9</b>	<b>&lt;0.001</b>
RNFL inner ring (µm)	<i>28.4</i>	<b>-2.9</b>	<b>0.4</b>	<b>&lt;0.001</b>	<b>-3.5</b>	<b>0.8</b>	<b>&lt;0.001</b>
RNFL outer ring (µm)							
GCL inner ring (µm)							
GCL outer ring (µm)							
IPL inner ring (µm)							
IPL outer ring (µm)							
Peripapillary RNFL:	<i>96.9</i>	<b>-8.6</b>	<b>1.6</b>	<b>&lt;0.001</b>	<b>-7.7</b>	<b>2.5</b>	<b>0.002</b>
Average (µm)	<i>103.7</i>	<b>-13.1</b>	<b>3.0</b>	<b>&lt;0.001</b>	<b>-7.1</b>	<b>4.9</b>	<b>0.145</b>
Nasal superior (µm)	<i>73.3</i>	<b>-6.5</b>	<b>2.3</b>	<b>0.004</b>	<b>-3.1</b>	<b>3.3</b>	<b>0.349</b>
Nasal inferior (µm)	<i>110.4</i>	<b>-8.7</b>	<b>3.7</b>	<b>0.018</b>	<b>-8.1</b>	<b>4.4</b>	<b>0.066</b>
Temporal inferior (µm)	<i>138.6</i>	<b>-15.4</b>	<b>3.3</b>	<b>&lt;0.001</b>	<b>-16.7</b>	<b>4.1</b>	<b>&lt;0.001</b>
Temporal superior (µm)	<i>70.7</i>	<b>-1.4</b>	<b>2.0</b>	<b>0.486</b>	<b>-4.1</b>	<b>2.5</b>	<b>0.108</b>
Temporal superior (µm)	<i>132.2</i>	<b>-13.5</b>	<b>2.9</b>	<b>&lt;0.001</b>	<b>-13.8</b>	<b>4.1</b>	<b>0.001</b>
Retinal vascular parameters:	<i>126.8</i>	<b>-3.1</b>	<b>1.5</b>	<b>0.038</b>	<b>-6.4</b>	<b>2.4</b>	<b>0.009</b>
CRAE	<i>195.3</i>	<b>-10.9</b>	<b>3.0</b>	<b>&lt;0.001</b>	<b>-14.4</b>	<b>4.2</b>	<b>0.001</b>
CRVE	<i>0.652</i>	<b>0.024</b>	<b>0.010</b>	<b>0.013</b>	<b>0.017</b>	<b>0.009</b>	<b>0.045</b>
AVR	<i>1.173</i>	<b>-0.021</b>	<b>0.007</b>	<b>0.003</b>	<b>-0.021</b>	<b>0.010</b>	<b>0.035</b>
FDa	<i>1.156</i>	<b>-0.019</b>	<b>0.007</b>	<b>0.009</b>	<b>-0.018</b>	<b>0.009</b>	<b>0.040</b>
FDv	<i>-9.690</i>	<b>-0.057</b>	<b>0.041</b>	<b>0.171</b>	<b>-0.109</b>	<b>0.054</b>	<b>0.045</b>
cTORTa*	<i>-9.670</i>	<b>-0.054</b>	<b>0.038</b>	<b>0.153</b>	<b>-0.031</b>	<b>0.041</b>	<b>0.445</b>

GEE, corrected for sex and a diabetes. SE = Standard Error, CI = Cognitively Impaired, RT = Retinal Thickness, RNFL = Retinal Nerve Fiber Layer, GCL = Ganglion Cell Layer, IPL = Inner Plexiform Layer, CRAE = Central Retinal Artery Equivalent, CRVE = Central Retinal Vein Equivalent, AVR = Arteriole-Venular Ratio, FDa = Fractal Dimension of arteries, FDv = Fractal Dimension of veins, cTORTa = curvature Tortuosity of arteries, cTORTv = curvature Tortuosity of veins. \* Log transformation applied. Values in **bold** are significant at  $p < 0.05$ .

## Additional File Information:

*File name:* additional file 1


*File format:* .pdf

*Title of data:* Additional file 1: Reasons for excluding participants, categorized per group.

*Description of data:* Note that participants were only excluded if bilateral problems were present, explaining why the total N may be lower than the subdivided numbers taken together (i.e. 1 eye of a participant may fall in 1 category, and the contralateral eye in another). CI = Cognitively Impaired, OCT = Optical Coherence Tomography, SIVA = Singapore I Vessel Assessment, AMD = Age-related Macular Degeneration, ERM = Epiretinal Membrane, PPA = Peripapillary Atrophy, CSC = Central Serous Chorioretinopathy.

## Figures


# Macular Area



# ONH

Figure 1

Title: Boxplots for retinal layer thicknesses as measured with Optical Coherence Tomography in the 3 groups. Legend: Dotted line represents the mean. CI = Cognitively Impaired, ONH = Optic Nerve Head.



**Figure 2**

Title: Boxplots for retinal vascular parameters obtained with Singapore I Vessel Assessment in the 3 groups. Legend: Dotted line represents the mean. CI = Cognitively Impaired, ONH = Optic Nerve Head.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.pdf](#)