

# Low impact of regular PCR testing on presence at work site during the COVID-19 pandemic: Experiences during an open observational study in Lower Saxony 2020-21

**Lisa K. Seiler**

Leibniz University Hannover

**Susanne Stolpe**

University Hospital Essen

**Nils Stanislawski**

Leibniz University Hannover

**Frank Stahl**

Leibniz University Hannover

**Martin Witt**

Leibniz University Hannover

**Rebecca Jonczyk**

Leibniz University Hannover

**Stefanie Heiden**

Leibniz University Hannover

**Holger Blume**

Leibniz University Hannover

**Bernd Kowall**

University Hospital Essen

**Cornelia Blume** (✉ [blume@iftc.uni-hannover.de](mailto:blume@iftc.uni-hannover.de))

Leibniz University Hannover

---

## Research Article

**Keywords:** Workplace safety, COVID-19, PCR test, Home Office

**Posted Date:** July 21st, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1822640/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

**Additional Declarations:** No competing interests reported.

---

**Version of Record:** A version of this preprint was published at BMC Public Health on February 3rd, 2023. See the published version at <https://doi.org/10.1186/s12889-023-15036-9>.

## Abstract

Since social distancing during the COVID-19-pandemic had a profound impact on professional life, this study investigated the effect of PCR testing on on-site work.

PCR screening, antibody testing, and questionnaires offered to 4,890 working adults in Lower Saxony were accompanied by data collection on demographics, family status, comorbidities, psychosocial situation, health-related behavior, and the number of work-related contacts. Relative risks (RR) with 95% confidence intervals were estimated for the associations between regular PCR testing and other work and health-related variables, respectively, and working on-site. Analyses were stratified by the suitability of work tasks for mobile office.

Between April 2020 and February 2021, 1,643 employees underwent PCR testing. Whether mobile working was possible strongly influenced the work behavior. Persons whose work was suitable for mobile office (mobile workers) had a lower probability of working on-site than persons whose work was not suitable for mobile office (RR = 0.09 (95%CI: 0.07–0.12)). In mobile workers, regular PCR-testing was slightly associated with working on-site (RR = 1.19 (0.66; 2.14)). In those whose working place was unsuitable for mobile office, the corresponding RR was 0.94 (0.80; 1.09). Compared to persons without chronic diseases, chronically ill persons worked less often on-site if their workplace was suitable for mobile office (RR = 0.73 (0.40; 1.33)), but even more often if their workplace was not suitable for mobile office (RR = 1.17 (1.04; 1.33)).

If work was suitable for mobile office, regular PCR testing did not have a strong effect on presence at the work site.

Trial registration: An ethics vote of the responsible medical association (Lower Saxony, Germany) retrospectively approved the evaluation of the collected subject data in a pseudonymized form in the context of medical studies (No. Bo/30/2020; Bo/31/2020; Bo/32/2020).

## Background

A significant problem with preventive measures such as social distancing against the spread of a pandemic is the social and economic collateral damage caused by the strict isolation of groups of people. In Italy, strict isolation of significant parts of the population led to a considerable reduction in COVID-19-positive patients and those severely affected [1, 2] but was associated with significant economic and social consequences. As soon as more targeted information was available on how widespread positivity for the pandemic-causing pathogen (i.e., SARS-CoV-2) is in different population groups, more targeted measures were taken that led to a definition of SARS-CoV-2 risk groups [3]. In an early population-based study on SARS-CoV-2 prevalence in a large German municipality not affected by a super spreading event, only one in four SARS-CoV-2 infections in private households was reported and known to the health authorities [4]. Social isolation of the general population caused unsustainable social and economic conditions in the long term. Others show, for example, that specific employees, such as teachers in a similar situation as university docents with frequent social contacts, had a particular load of psychic stress due to the necessity of social distancing during the lockdown, preventing them from their educational mission [5]. Remote working was also a question of gender, as has recently been shown by Bezak et al.: Females were more stressed professionally, socially, and personally than males, which was partially caused by employers' high expectations. Working from home here was shown to be a challenge, complicated by a lack of preparedness with the rapid start of the lockdown measures [6]. Gibbs et al. documented with a study on 112 desk workers that working from home was often associated with non-work sedentary resulting in declining physical functionality and worsened sleep quality [7].

In this project, PCR screening for SARS-CoV-2 infectivity was carried out in groups of populations with an increased risk for infection based on their medical risk factors or workplace (e.g., due to frequent contact with other people). Untargeted testing for SARS-CoV-2 infectivity was carried out to enable more movement freedom for individuals, especially for more

presence at their workplace. Untargeted PCR testing at the time of the study start was not recommended by the Robert Koch Institute (RKI) [8] since resources for this diagnostic measure were insufficient at the beginning of the pandemic and should be reserved for symptomatic individuals with suspicion of COVID-19. PCR testing requires expensive laboratory instrumentation and highly skilled laboratory personnel. In addition, due to logistic reasons, several days were initially needed until the results were available [5].

The study presented here comprised selected groups of participants from several occupational groups in Lower Saxony from April 2020 to February 2021. We hypothesized that for the working world, in particular, rapid and highly precise detection of infectiousness is helpful to prevent the spreading of the virus during working in presence. Without testing, we presumed that many employees would have to work from home for preventive reasons or an increased infection risk caused by frequent unavoidable professional contacts on-site. Particularly in the case of work processes that can only be carried out poorly in the "home office", this would also be economically significant. By testing, the employer could thus continuously operate or reopen essential sectors of the company's work. The study aimed to show that free and voluntary tests offered to employees at a workplace could help to avert economic damage due to missing workforce on-site, despite the additional costs associated with PCR testing. Of note, there was an increase from 15% of employees working from home pre-pandemic compared to 66% since the pandemic's start [9].

We primarily aimed to address whether repetitive testing of non-symptomatic subjects in the context of the SARS-CoV-2 pandemic positively affected working on-site. Moreover, we aimed to examine the impact of further factors like health status, smoking, psychosocial problems, and risk perception on presence at the work site.

## **Materials And Methods**

### **Study design**

In the context of infection prevention, mobile screening facilities for offering voluntary COVID-19 PCR tests for employees were positioned close to the workplaces of participating institutions [10, 11]. These institutions were, among others, a company located in the southern part of Lower Saxony, the Leibniz University Hannover (LUH), a theatre and, several nursing homes and schools in Hannover (Tab. S1). Since not all members of the announced occupational groups could be tested, each institution's respective employer or head invited those persons to be entitled to free testing. Employees with frequent personal contacts or comorbidities were the first asked to participate. No feedback was given to the respective employer on whether a selected person used the offer for regular testing. Altogether, 4,890 test persons in Lower Saxony were recorded concerning their infectivity and contamination with SARSCoV2 during April 2020 and February 2021. References to gender, age, and risk factors were drawn in combination with the data of the test persons, and a subgroup of test persons (N = 1,643) was retrospectively questioned concerning their working and health behavior and perception of risk during work. Furthermore, employees could also volunteer for testing.

### **Questionnaire campaign**

For each test, participants answered a questionnaire regarding contacts with COVID19-positive persons, symptoms of a COVID-19-infection, feeling strongly impaired in the last three months due to worries about health or due to problems related to family and friends or work or due to financial difficulties, chronic diseases, changed behavior regarding health service utilization (refraining from seeing a doctor in case of feeling sick, canceling of appointments or use of telephone consulting) during the last six months, and self-assessed risk for corona infection (0 = no risk, 5 = high risk). Starting in August 2020, questions on subjective health, emotional stress, and health behavior were included in the questionnaire and answered by those participants receiving antibody testing (N = 929). After the test period, all participants with known email-address received an additional online questionnaire in August 2021 (N = 3,846). These participants were asked about workplace characteristics before the pandemic and during the testing period about suitability for mobile work and

risk of infection due to frequent contact with coworkers, customers, or pupils. Response rate was 4 % (N = 1,643). The questionnaires can be found in the supplement.

## Variable definitions

Working primarily on-site was coded with 'yes' if the participants worked two days or less at home. The type of workplace was coded according to the workplace at which participants worked most of the week. Suitability of work for mobile working was coded 'yes' if it was rated as suitable or primarily suitable by participants. Regular testing was defined as at least three tests with a maximal time lag of 14 days each. The minimum length of a test period with regular testing was 28 days. Chronic diseases (diabetes, obesity, heart disease, chronic lung disease, chronic bowel disease) or COVID-19-symptoms were coded as present if ever mentioned during the test period. Feeling strongly impaired during the last six months was coded as present if participants responded accordingly to at least one out of seven questions relating to worries about health, difficulties with partner, family or friends, strain due to care for children, parents or other family, stress at work or school, financial problems, or missing close contacts. Social problems in the last three months were coded present if participants reported either feeling lonely, isolated, or excluded from the community. A change in health-related behavior (such as avoided, reduced, or canceled visits to a doctor or cancelled preventive or rehabilitation therapy or telephone consultation) was coded as present if ever mentioned during the test period.

## Missing values

Some questions we analyzed were only asked during an antibody test. As not all participants underwent antibody testing, the questions on subjective health, emotional stressors, and health behavior had more missings (N = 714).

We did a complete case analysis.

## Statistical analyses

We estimated the prevalence of participant and workplace characteristics stratified by the suitability of the workplace for mobile office. Fitting log-binomial regression models, we estimated univariate crude relative risks for working primarily on-site with 95 % confidence intervals (CI) for variables related to workplace conditions, regular testing, and personal characteristics. Further, we stratified these analyses by the suitability of the work for mobile work. We did not adjust for potential confounders [12] as we did not want to assess causal effects. All analyses were done using SAS9.4.

## Results

### Study participants

Among the 1,643 participants filling out the online questionnaire, 53% were male. The mean age was 42.7 years (standard deviation (sd) 12.2) (Table 1). 34% of the participants worked in university departments, 54% at a biotechnological company, and 7% in nursing homes or schools. Three out of four participants (N = 1,224) rated their work suitable for a mobile office. 43% (N = 714) estimated that 80% 10 % of their work could be fulfilled in a mobile office.

Table 1

Results of online questionnaire (N = 1.643): characteristics of work places not/only limited suitable for mobile office and suitable for mobile office.

	Not/only limited suitable for mobile office (N = 419)	Suitable for mobile office (N = 1,224)	Total (N = 1,643)
Sex			
Male	203 (48.5)	660 (53.9)	863 (52.5)
Female	216 (51.5)	564 (40.1)	780 (47.5)
Age (mean, (Std))	42.6 (12.9)	42.7 (12.0)	42.7 (12.2)
<30	87 (20.8)	213 (17.4)	300 (18.3)
30-<50	177 (42.2)	594 (48.5)	771 (46.9)
50-65	155(37.0)	417 (34.1)	572 (34.8)
Living in a single household	92 (22.0)	213 (17.4)	305 (18.6)
Self-rated risk for corona infection (0-5) (mean (std)) (N = 927))	2.58 (1.19)	2.31 (1.00)	2.39 (1.07)
Low (0-<2)	51 (19.8)	153 (22.8)	204 (22.0)
Medium (2-<3)	63 (24.5)	230 (34.3)	293 (31.6)
High (3-5)	143 (55.6)	287 (42.8)	430 (46.4)
Number of tests in test period (mean (std))	8.8 (6.6)	8.5 (6.0)	8.6 (6.2)
1-2 tests	83 (19.8)	228 (18.6)	311 (18.9)
3-5	86 (20.5)	204 (16.7)	290 (17.7)

<sup>1</sup> Regular testing: at least 3 tests with a maximum lag time of 14 days

<sup>2</sup> Includes diabetes, adiposity, heart disease, chronic lung disease, chronic bowel disease

<sup>3</sup> includes avoided, reduced or cancelled visits to a doctor or cancellation of rehabilitation/preventive therapy or use of telephone consultation

<sup>4</sup> feeling impaired due to worries about health, difficulties with partner, family or friends, strain due to care for children, parents or other family, stress at work or school, financial problems or missing social close contacts

<sup>5</sup> Sometimes or often feeling lonely in a community or feeling excluded or isolated

	Not/only limited suitable for mobile office (N = 419)	Suitable for mobile office (N = 1,224)	Total (N = 1,643)
6-9	60 (14.3)	256 (20.9)	316 (19.2)
10-14	113 (27.0)	399 (32.6))	512 (31.2)
15 and more tests	77 (18.4)	136 (11.1)	213 (13.0)
Employer			
University	117 (27.9)	448 (36.6)	565 (34.4)
Nursery home/School	69 (16.5)	53 (4.3)	122 (7.4)
Sartorius	186 (44.4)	696 (56.9)	882 (53.7)
Other	47 (11.2)	27 (2.2)	74 (4.5)
Regular Tests <sup>1</sup> no	322 (76.9)	1,069 (87.4)	1,391 (84.7)
Yes	97 (23.1)	154 (12.6)	251 (15.3)
Medical/social occupation (N = 1.054)	224 (71.1)	456 (61.7)	680 (64.5)
Working mostly on site before pandemic - no	27 (6.4)	332 (27.1)	359 (21.9)
Yes	392 (93.6)	892 (72.9)	1,284 (78.1)
Working mostly on site during test period	86 (20.5)	1,048 (85.6)	1,134 (69.0)
no			
yes	333 (79.5)	176 (14.4)	509 (31.0)
Type of working place (before pandemic)			

<sup>1</sup> Regular testing: at least 3 tests with a maximum lag time of 14 days

<sup>2</sup> Includes diabetes, adiposity, heart disease, chronic lung disease, chronic bowel disease

<sup>3</sup> includes avoided, reduced or cancelled visits to a doctor or cancellation of rehabilitation/preventive therapy or use of telephone consultation

<sup>4</sup> feeling impaired due to worries about health, difficulties with partner, family or friends, strain due to care for children, parents or other family, stress at work or school, financial problems or missing social close contacts

<sup>5</sup> Sometimes or often feeling lonely in a community or feeling excluded or isolated

	Not/only limited suitable for mobile office (N = 419)	Suitable for mobile office (N = 1,224)	Total (N = 1,643)
Single room	73 (17.4)	310 (25.3)	383 (23.3)
Multi-person room (2–4)	62 (14.8)	352 (28.8)	414 (25.2)
Room with 5 persons or more	60 (14.3)	471 (38.5)	531 (32.4)
Production/Laboratory	178 (42.5)	59 (4.8)	237 (14.4)
Sales/outdoor/logistic	46 (11.0)	32 (2.6)	78 (4.8)
Type of working place (during pandemic)			
Single room	96 (22.9)	519 (42.4)	615 (37.4)
Multi-person room (2–4)	56 (13.4)	263 (21.5)	319 (19.4)
Room with 5 persons or more	43(10.3)	302 (24.7)	345 (21.0)
Production/Laboratory	185 (44.2)	64 (5.2)	249 (15.2)
Sales/outdoor/logistic	15 (3.6)	11 (0.9)	26 (1.6)
Part of work that can be fulfilled in home office			
0-<50%	405 (96.7)	168 (13.7)	573 (34.9)
50%-<80%	7 (1.7)	349 (28.5)	356 (21.7)
80%-100%	7 (1.7)	707 (57.8)	714 (43.4)
Chronic diseases <sup>2</sup> (N = 1,618) no	323 (78.0)	970 (80.6)	1,293 (79.9)

<sup>1</sup> Regular testing: at least 3 tests with a maximum lag time of 14 days

<sup>2</sup> Includes diabetes, adiposity, heart disease, chronic lung disease, chronic bowel disease

<sup>3</sup> includes avoided, reduced or cancelled visits to a doctor or cancellation of rehabilitation/preventive therapy or use of telephone consultation

<sup>4</sup> feeling impaired due to worries about health, difficulties with partner, family or friends, strain due to care for children, parents or other family, stress at work or school, financial problems or missing social close contacts

<sup>5</sup> Sometimes or often feeling lonely in a community or feeling excluded or isolated

	Not/only limited suitable for mobile office (N = 419)	Suitable for mobile office (N = 1,224)	Total (N = 1,643)
yes	91 (22.0)	234 (19.4)	325 (20.1)
Smoking (N = 1,604) no	314 (76.0)	1,029 (86.4)	1,343 (83.7)
yes	99 (24.0)	162 (13.6)	261 (16.3)
Covid symptoms in last 14 days – no	272 (64.9)	699 (57.2)	971 (59.1)
yes	147 (35.1)	524 (42.8)	671 (40.9)
Change in behaviour in regard to health service utilization <sup>3</sup> (N = 924) no	133 (52.2)	294 (44.0)	427 (46.2)
Yes	122 (47.8)	375 (56.0)	497 (53.8)
Feeling strongly impaired due to personal problems <sup>4</sup> in last 3 months (N = 929)–	132 (51.4)	311 (46.3)	443 (47.7)
no			
yes	125 (48.6)	361 (53.7)	486 (52.3)
Social problems in last 3 months <sup>5</sup> (N = 929)	187 (72.8)	463 (68.9)	650 (70.0)
- no			
yes	70 (27.2)	209 (31.1)	279 (30.0)
<sup>1</sup> Regular testing: at least 3 tests with a maximum lag time of 14 days			
<sup>2</sup> Includes diabetes, adiposity, heart disease, chronic lung disease, chronic bowel disease			
<sup>3</sup> includes avoided, reduced or cancelled visits to a doctor or cancellation of rehabilitation/preventive therapy or use of telephone consultation			
<sup>4</sup> feeling impaired due to worries about health, difficulties with partner, family or friends, strain due to care for children, parents or other family, stress at work or school, financial problems or missing social close contacts			
<sup>5</sup> Sometimes or often feeling lonely in a community or feeling excluded or isolated			

## Test frequency

About 40% of the participants in the online survey had a maximum of five tests; 13% were tested 15 times or more. The mean number of tests was 8.6 (sd 6.2). Participants with workplaces unsuitable for mobile office were more often tested regularly (23% vs. 13%) (Table 2).



Table 2

Working mostly on site – by demographic characteristics, job characteristics, comorbidities and other factors (N, % and relative risk with 95% confidence interval). N = 1,643

	working mostly on site			Crude relative risk (95%-Confidence interval)
	(max. 2 days/month at home)			
	Yes	No	% yes	
Gender male	185	678	21.4	Ref.
female	197	583	25.3	1.18 (0.99; 1.40)
Age (10 yrs)				1.02 (1.00; 1.04)
<30	66	234	22.0	Ref.
30-<50	161	610	20.9	0.95 (0.74; 1.22)
50-65	155	417	27.1	1.23 (0.96; 1.58)
Part of work that can be done effectively at home (every 10% increase)				0.64 (0.62; 0.68)
<50%	317	256	55.3	Ref.
50-70%	32	324	9.0	0.16 (0.12; 0.23)
80-100%	33	681	4.6	0.08 (0.06; 0.12)
Work suited for mobile work No	300	119	71.6	Ref.
Yes	82	1,142	6.7	0.09 (0.07; 0.12)
Employer				
Nurse home/school	55	67	45.1	Ref.
University	100	465	17.7	0.39 (0.30; 0.51)
Sartorius	191	691	21.7	0.48 (0.38; 0.61)
Other	36	38	48.7	1.08 (0.80; 1.46)
Working place during test period				
Single room	83	532	13.5	Ref.
Multiple persons room	108	556	16.3	1.21 (0.93; 1.57)
Laboratory/production	158	91	63.5	4.70 (3.77; 5.87)
Sales / field service	33	82	28.7	2.13 (1.50; 3.02)
Number of job related contacts > 15min before testing				
None	12	20	37.5	Ref.
1-5 persons	119	508	19.0	0.51 (0.31; 0.81)
6 or more	251	733	25.5	0.68 (0.43; 1.08)
Regular tests * no	304	1,087	21.9	Ref.
Yes	78	173	31.1	1.42 (1.15; 1.75)

	working mostly on site (max. 2 days/month at home)			Crude relative risk (95%-Confidence interval)
	Yes	No	% yes	
	Smoker No	277	1,066	
Yes	96	165	36.8	1.78 (1.47; 2.16)
Medical or social occupation no	93	281	24.9	Ref.
Yes	179	501	26.3	1.06 (0.85; 1.31)
Chronic disease No	292	1,001	22.6	Ref.
Yes	86	239	26.5	1.17 (0.95; 1.44)
Diabetes No	361	1,205	23.1	Ref.
Yes	15	29	34.1	1.48 (0.97; 2.25)
Obesity No	350	1,191	22.7	Ref.
Yes	26	44	37.1	1.64 (1.19; 2.25)
Chronic lung disease No	348	1,137	23.4	Ref.
Yes	30	102	22.7	0.97 (0.70; 1.35)

<b>Ever Contact with Covid Case No</b>	<b>340</b>	<b>1113</b>	<b>23.4</b>	<b>Ref</b>
Yes	41	145	22.0	0.94 (0.71; 1.25)
Covid symptoms in last 14 days No	251	720	25.9	Ref
Yes	131	540	19.5	0.76 (0.63; 0.91)
Self-rated risk of corona infection (0–5) increase by 1 <sup>3</sup> (N = 927)				1.12 (1.01; 1.25)
Low Self-rated risk (0-<2)	48	156	23.5	Ref
Medium self-rated risk (2-<3)	62	231	21.2	0.90 (0.65; 1.25)
Higher self-rated risk (3+)	115	315	26.7	1.14 (0.85; 1.52)
Change in behaviour in regard to health service utilization <sup>4</sup> (N = 924)				
No	120	307	28.1	Ref
Yes	105	392	21.1	0.75 (0.60; 0.94)
Feeling strongly impaired due to personal problems <sup>5</sup> in last 3 months (N = 929)				
No	112	331	25.3	Ref
Yes	114	372	23.5	0.93 (0.74; 1.16)
Social problems <sup>6</sup> in last 3 months (N = 929)				
No	63	163	31.7	Ref
Yes	136	567	22.3	0.71 (0.55; 0.90)
<sup>1</sup> Regular testing: at least 3 tests with a maximum lag time of 14 days				
<sup>2</sup> Includes diabetes, obesity, heart disease, chronic lung disease, chronic bowel disease				
<sup>3</sup> A rating of 0 indicates no self-assessed risk, a rating of 5 indicates a very high self-assessed risk of corona infection				
<sup>4</sup> includes avoided, reduced or cancelled visits to a doctor or cancellation of rehabilitation/preventive therapy or use of telephone consultation				
<sup>5</sup> feeling impaired due to worries about health, difficulties with partner, family or friends, strain due to care for children, parents or other family, stress at work or school, financial problems or missing social close contacts				
<sup>6</sup> Sometimes/often feeling lonely in a community or feeling excluded or feeling isolated				

## Testing and working on-site

Working on-site was reduced during the pandemic compared to pre-pandemic times, from 94–80% in the group with work unsuitable for mobile office and from 73–14% in those with suitable work (Table 1). Suitability of the work for mobile office was most strongly associated with presence at work (RR = 0.09 (0.07; 0.12)) (Table 2). In mobile workers, regular

PCR-testing was slightly associated with working on-site (RR = 1.19 (0.66; 2.14)). In those whose workplace was not suitable for mobile office, the corresponding RR was 0.94 (0.80; 1.09) (Table 3).

Table 3

Working mostly on site – by demographic characteristics, job characteristics, comorbidities and other factors (N, % and relative risk with 95% confidence interval) - stratified by suitability for mobile work. N = 1,643.

	Job suitable for mobile office (N = 1,224)		Job not suitable for mobile office (N = 419)	
	Working mostly on site		Working mostly on site	
	N (%)	RR (95%-CI)	N (%)	RR (95%-CI)
Gender male	32 (4.9)	Ref.	153(75.4)	Ref.
female	50 (8.9)	1.83 (1.19; 2.81)	147 (68.1)	0.90 (0.80; 1.02)
Age (years)				
< 30	15 (7.0)	Ref.	51 (58.6)	Ref.
30<50	31 (5.2)	0.74 (0.41; 1.35)	130 (73.5)	1.25 (1.03; 1.53)
50–65	36 (8.6)	1.23 (0.69; 2.19)	119 (76.8)	1.31 (1.08; 1.59)
Part of work that can be done effectively at home (every 10% increase)	-	0.77 (0.71; 0.84)	-	0.73 (0.67; 0.80)
<50%	26 (15.5)	Ref.	291 (71.9)	Ref.
50–70%	29 (8.3)	0.54 (0.33; 0.88)	3 (42.9)	0.60 (0.25; 1.41)
80–100%	27 (23.8)	2.18 (1.21; 3.62)	6 (85.7)	1.19 (0.88; 1.62)
Employer				
Nurse home/school	12 (22.6)	Ref.	43 (62.3)	Ref.
University	34 (7.6)	0.34 (0.19; 0.61)	66 (56.4)	0–91 (0.71; 1.15)
Sartorius	31 (4.5)	0.20 (0.11; 0.57)	160 (86.0)	1.38 (1.14; 1.67)
Other	5 (18.5)	0.82 (0.32; 2.08)	16 (66.0)	1.06 (0.80; 1.39)
Number of job related contacts > 15min duration - before testing period				
None	2 (10.0)	Ref.	10 (83.3)	Ref.
1–5 persons	34 (6.7)	0.67 (0.17; 2.59)	85 (71.4)	0.86 (0.65; 1.13)

<sup>1</sup> Regular testing: at least 3 tests with a maximum lag time of 14 days

<sup>2</sup> Includes diabetes, adiposity, heart disease, chronic lung disease, chronic bowel disease

<sup>3</sup> A rating of 0 indicates no self-assessed risk, a rating of 5 indicates a very high self-assessed risk of corona infection

<sup>4</sup> includes avoided, reduced or cancelled visits to a doctor or cancellation of rehabilitation/preventive therapy or use of telephone consultation

<sup>5</sup> Sometimes or often feeling lonely in a community or feeling excluded or feeling isolated

	Job suitable for mobile office (N = 1,224)		Job not suitable for mobile office (N = 419)	
	Working mostly on site		Working mostly on site	
6 or more	46 (6.6)	0.66 (0.17;2.53)	205 (17.2)	0.85 (0.66; 1.11)
Regular tested <sup>1</sup> No	70 (6.6)	Ref.	234 (72.7)	Ref.
Yes	12 (7.8)	1.19 (0.66; 2.14)	66 (68.0)	0.94 (0.80; 1.09)
Living alone No	62 (6.1)	Ref.	235 (71.9)	Ref.
Yes	20 (9.4)	1.53 (0.94; 2.47)	65 (70.7)	0.98 (0.85; 1.14)
Current Smoker No	62 (6.0)	Ref.	80 (80.8)	Ref.
Yes	16 (9.9)	1.64 (0.97; 2.77)	215 (68.5)	1.18 (1.04; 1.33)
Medical or social occupation no	20 (7.1)	Ref.	73 (80.2)	Ref.
Yes	41 (9.0)	1.27 (0.76; 2.13)	138 (61.6)	0.77 (0.66; 0.89)
Chronic disease <sup>2</sup> No	68 (7.0)	Ref.	224 (69.4)	Ref.
Yes	12 (5.1)	0.73 (0.40; 1.33)	74 (81.3)	1.17 (1.04; 1.33)
Diabetes No	77 (6.6)	Ref.	284 (71.5)	Ref.
Yes	2 (7.1)	1.08 (0.28; 4.19)	13 (81.3)	1.14 (0.89; 1.44)
Obesity No	76 (6.6)	Ref.	274 (70.3)	Ref.
Yes	3 (6.4)	0.97 (0.32; 2.95)	23 (100)	1.42 (1.33; 1.52)
Chronic lung disease No	75 (6.8)	Ref.	273 (71.5)	Ref.
Yes	5 (5.0)	0.74 (0.30; 1.78)	25 (78.1)	1.09 (0.90; 1.33)
Self-rated risk of corona infection <sup>3</sup>				
0-<2	11 (7.2)	Ref.	37 (72.6)	Ref.
3-	19 (8.3)	1.15 (0.56;2.35)	43 (68.3)	0.94 (0.74; 1.19)

<sup>1</sup> Regular testing: at least 3 tests with a maximum lag time of 14 days

<sup>2</sup> Includes diabetes, adiposity, heart disease, chronic lung disease, chronic bowel disease

<sup>3</sup> A rating of 0 indicates no self-assessed risk, a rating of 5 indicates a very high self-assessed risk of corona infection

<sup>4</sup> includes avoided, reduced or cancelled visits to a doctor or cancellation of rehabilitation/preventive therapy or use of telephone consultation

<sup>5</sup> Sometimes or often feeling lonely in a community or feeling excluded or feeling isolated

	Job suitable for mobile office (N = 1,224)		Job not suitable for mobile office (N = 419)	
	Working mostly on site		Working mostly on site	
4–5	27 (9.4)	1.31 (0.67; 2.57)	88 (61.5)	0.84 (0.69; 1,05)
Contact with Corona Case no	73 (6.7)	Ref.	267 (72.8)	Ref.
Yes	8 (6.0)	0.88 (0.44; 1.80)	33 (63.5)	0.87 (0.70; 1.08)
Change in behaviour in regard to health service utilization <sup>4</sup> (N = 924) no	25 (8.5)	Ref.	95 (56.9)	Ref.
Yes	33 (8.8)	1.03 (0.63; 1.70)	72 (43.1)	0.83 (0.69; 0.99)
Social problems <sup>5</sup> in last 3 months(N = 929)	17 (12.1)	Ref.	46 (79.3)	Ref.
No				
Yes	41 (7.7)	0.64 (0.38; 1.09)	122 (61.3)	0.77 (0.65; 0.92)
<sup>1</sup> Regular testing: at least 3 tests with a maximum lag time of 14 days				
<sup>2</sup> Includes diabetes, adiposity, heart disease, chronic lung disease, chronic bowel disease				
<sup>3</sup> A rating of 0 indicates no self-assessed risk, a rating of 5 indicates a very high self-assessed risk of corona infection				
<sup>4</sup> includes avoided, reduced or cancelled visits to a doctor or cancellation of rehabilitation/preventive therapy or use of telephone consultation				
<sup>5</sup> Sometimes or often feeling lonely in a community or feeling excluded or feeling isolated				

## Demographics and working on-site

Women were more often present at their workplace than men (RR = 1.18 (0.99; 1.40)), and participants aged 50–65 years were more often present at their workplace than younger persons (RR = 1.23 (0.96; 1.58)) (Table 2). Generally, associations between participants` and work characteristics, respectively, and working onsite often differed in strata of work suitability for mobile office.

## Family status, lifestyle factors, comorbidity, and working on-site

Current smokers were more likely to be present on-site regardless of the suitability of their workplace for a mobile office (RR = 1.64 (0.97; 2.77) for jobs suitable for mobile office, RR = 1.18 (1.04; 1.33) for jobs not suitable for mobile office) (Table 3). In participants with jobs unsuitable for mobile office, the prevalence of chronic conditions was associated with higher workplace presence (RR = 1.17 (1.04; 1.33)) than in participants with a workplace suitable for a mobile office (RR = 0.73 (0.40; 1.33)) (Table 3). In addition, participants living alone were more often present on-site even if their job was suitable for mobile office than those who reported not living alone (RR = 1.53 (0.94; 2.47)) (Table 3).

## Risk assessment and working on-site

In persons with jobs suitable for mobile office, presence at work was more frequent the higher the self-rated risk of corona infection was (RR = 1.15 (0.56; 2.35) for medium self-rated risk, and RR = 1.31 (0.67; 2.57) for high self-rated risk compared to low self-rated risk) (Table 3). Furthermore, among participants with a job unsuitable for mobile office, persons who reported a change in their health-related behavior caused by the pandemic were more seldom on-site (RR = 0.87 (0.70; 1.08)).

## Discussion

The strongest predictor for working on-site was the lack of suitability of the workplace for mobile office. Regular PCR testing did not substantially impact on presence at the work site. In those with jobs suitable for mobile office, regular tests only led to a slight increase in work site presence.

On-site workplace absenteeism was associated with reporting more frequent contact with others lasting longer than 15 minutes at work even before the pandemic. In addition, absence from work was associated with altered health-related behaviors, clustered social problems, and experienced contact with COVID19 cases. The above factors were further related to actual or perceived psychological distress from the pandemic. Participants who feel more stressed seem to prefer to work at home, regardless of whether their workplace allows them to perform their job from home.

People with lower social status may not be able to set up a workplace at home because they do not have enough space or the financial resources to do so. They may also be less likely to have the opportunity in their work context to decide for themselves whether to work at home. Current smoking was associated with a higher prevalence at work in all strata. One reason for this could be a relatively lower social status of smokers compared to nonsmokers [13]. However, smokers may also be more interested in social contacts. Interestingly, having an existing chronic disease was not associated with a lower prevalence at work. Because chronic diseases are more prevalent among people with lower social status [14], this factor may be the leading cause of the association. For those living alone, the higher likelihood of working locally may reflect a need for social contact otherwise limited by the official nationwide lockdown regulation.

In this work, a higher self-assessed risk of infection correlates with a higher prevalence of on-site work, with the on-site presence most likely being the cause of the self-assessed increased risk. It is worth noting that this relationship was only present among those with the option of mobile working and not among those who did not have the option of moving their workplace home. Social stress and changes in health-related behaviors due to the pandemic were associated with lower prevalence at work. People experiencing these stresses would want to avoid additional potential viral exposure at work.

## Limitations

Tested participants in the online survey do not represent the entire target group of this study. In some workplaces, such as the university, participants took care of the registration to participate in the voluntary test offer themselves. However, in some workplaces, such as the participating company and nursing homes, the respective supervisor or manager gave a recommendation to some individuals to be tested in addition to independent voluntary participation. Specific encouragement by employers to use the test may have harmed motivation to be tested over a more extended period. Only about 50% of all tested participants took up to five tests. We have no information on the educational background of those tested, although it can be assumed that some of the associations presented reflect a person's underlying educational background [13]. This is particularly true for the association between current smoking or obesity and on-site attendance.

The online questionnaire could not be delivered to all tested individuals because some had no valid e-mail address. A comparison between the tested population and the population that answered the online questionnaire showed that university employees were overrepresented in the online survey. People working in nursing homes or schools were underrepresented.

In future pandemic situations, testing should be mandatory, especially for people who cannot work on a mobile basis with a great risk for infection, as shown for German health workers during April 2020 and April 2021 [15]. This was later practiced to some extent during the pandemic in Germany - mainly based on the SARS-CoV-2 rapid tests - albeit with comparatively lower test reliability than PCR testing [16]. Even PCR testing is not a universal remedy due to a relatively moderate sensitivity at best, as shown by a study on inpatients in Finland [17]. Identification of vulnerable working groups



with a specific need for frequent testing thus represents an essential contribution to occupational safety and health services necessary for the future [18]. It may be additionally meaningful in older populations and populations at risk, which were threatened by prolonged grief disorders upon long-lasting social isolation [19]. Consequent PCR testing against virus spreading was proven successful by Luxembourg's mass screening program [20] and is urgently postulated by others [21].

## Conclusion

Programs aimed at increasing workplace prevalence in situations such as the COVID19 pandemic were primarily intended to cover workplaces with little discretion about allowing workers to work at home. Offering voluntary testing to the workforce did not affect workplace attendance among individuals whose jobs were not conducive to mobile work. Nevertheless, because the tests were used more frequently by individuals in such a workplace, they could help increase workplace safety, particularly at production sites with an exceptionally high risk of transmission through the goods produced or in service sectors with particularly vulnerable customers, such as the health sector. Prioritization of such work areas recommended by public bodies would be desirable. It may be mentioned that the company supported by PCR testing in this project was able to expand its production unhindered by labor shortages throughout the pandemic [22]. In contrast, publications from the Netherlands and Morocco document a lack of productivity and dissatisfaction during the lockdown in working society[23, 24].

Based on the results of this study with voluntary participants, we cannot recommend whether the costs required for mass prophylactic testing justify the effort to keep certain occupational groups in attendance during future pandemics. We cannot decide if testing projects would thus replace the lockdown imposed in the SARS-CoV-2 crisis as a preventive measure against the pandemic spread in the workforce. In agreement with other groups, we value the effectiveness of testing as a means to combat pandemic scenarios. To be more effective as a preventive measure, PCR testing probably would have had to be controlled more strictly.

## List Of Abbreviations

CI	Confidence interval
Fig.	Figure
LUH	Leibniz University Hannover
MHH	Hannover Medical School
PCR	Polymerase chain reaction
RR	Relative risk
RKI	Robert Koch Institute
Sd	Standard deviation
Tab.	Table

## Declarations

Ethics approval and consent to participate

For these studies, the "Aerztekammer Niedersachsen" issued an ethical approval in August 2020 (No. Bo/30/2020; Bo/31/2020; Bo/32/2020). All methods were carried out in accordance with relevant guidelines and regulations. Study

subjects belonged to three different groups located in the southern part of Lower Saxony, Germany: The first group, defined as “education/culture,” included employees from universities, theater, and administration in and close to Hannover, Germany. The second group, defined as “company,” consisted of employees of a biotechnological company in Goettingen, Germany. The third group included both employees and residents of retirement and nursing homes as well as schools in and close to Hannover (designated as “nursing homes”). Informed consent was obtained from all subjects and/or their legal guardian(s).

#### Consent for publication

Not applicable.

#### Availability of data and materials

De-identified subject data sets will be available upon written request to the corresponding author following publication.

Competing interests: The authors declare no conflict of interest. The sponsor did not exert any influence or make any recommendation as to which groups of people should be tested. The offer of testing was requested by various institutions or groups of persons themselves.

#### Funding

The study was partially financed by state funds from the Ministry of Economics of Lower Saxony. The Ministry of Economics had no role in the design and conduct of the project or related studies; data collection, management, analysis, or interpretation; or writing of this article.

#### Authors' contributions

Conceptualization: Cornelia Blume, Nils Stanislawski, Holger Blume and Susanne Stolpe; Methodology: Frank Stahl, Martin Witt, Rebecca Jonczyk and Cornelia Blume; Software: Nils Stanislawski and Holger Blume; Validation: Susanne Stolpe and Bernd Kowall; Formal Analysis: Martin Witt and Bernd Kowall; Investigation: Nils Stanislawski and Cornelia Blume; Resources: Stefanie Heiden; Data Curation: Nils Stanislawski; Writing – Original Draft Preparation: Lisa Seiler, Susanne Stolpe and Cornelia Blume; Writing – Review & Editing: Bernd Kowall, Frank Stahl, Rebecca Jonczyk and Nils Stanislawski; Visualization: Lisa Seiler; Supervision: Cornelia Blume; Project Administration: Stefanie Heiden and Cornelia Blume; Funding Acquisition: Stefanie Heiden

#### Acknowledgements

The project group would like to thank Dr. Corinna Hauß (MVZ Labor Limbach Hannover GbR) and Dipl.-Ing. Jan Vöckler for their support and advice.

We further thank several PhD students at the Institutes of Technical Chemistry and Microelectronic Systems (Leibniz University Hannover, Germany) for intense work on both PCR and antibody testing as well as technical and organizational support in this project. Furthermore, we thank the medical and dental students from the Hannover Medical School (MHH) and from the university in Goettingen for their support registering subjects and taking swabs. We thank Nils Hoppe and his team for their support of the study's ethical vote and procedures with regard to data protection (Centre of Ethics and Law in the Life Sciences, Leibniz University, Hannover, Germany).

## References

1. Chu DKW, Pan Y, Cheng SMS, Hui KPY, Krishnan P, Liu Y, et al. Molecular Diagnosis of a Novel Coronavirus (2019-nCoV) Causing an Outbreak of Pneumonia. *Clinical chemistry*. 2020;66:549–55. doi:10.1093/clinchem/hvaa029.
2. Sebastiani G, Massa M, Riboli E. Covid-19 epidemic in Italy: evolution, projections and impact of government measures. *Eur J Epidemiol*. 2020;35:341–5. doi:10.1007/s10654-020-00631-6.
3. Standl F, Jöckel K-H, Stang A. COVID-19 and the need of targeted inverse quarantine. *Eur J Epidemiol*. 2020;35:339–40. doi:10.1007/s10654-020-00629-0.
4. Pritsch M, Radon K, Bakuli A, Le Gleut R, Olbrich L, Guggenbühl Noller JM, et al. Prevalence and Risk Factors of Infection in the Representative COVID-19 Cohort Munich. *Int J Environ Res Public Health* 2021. doi:10.3390/ijerph18073572.
5. Silva NSSE, Rose Elizabeth Cabral B, Leão LL, Pena GdG, Pinho L de, Magalhães TA de, et al. Working conditions, lifestyle and mental health of Brazilian public-school teachers during the COVID-19 pandemic. *Psychiatriki*. 2021;32:282–9. doi:10.22365/jpsych.2021.045.
6. Bezak E, Carson-Chahhoud KV, Marcu LG, Stoeva M, Lhotska L, Barabino GA, et al. The Biggest Challenges Resulting from the COVID-19 Pandemic on Gender-Related Work from Home in Biomedical Fields-World-Wide Qualitative Survey Analysis. *Int J Environ Res Public Health* 2022. doi:10.3390/ijerph19053109.
7. Barone Gibbs B, Kline CE, Huber KA, Paley JL, Perera S. Covid-19 shelter-at-home and work, lifestyle and well-being in desk workers. *Occup Med (Lond)*. 2021;71:86–94. doi:10.1093/occmed/kqab011.
8. RKI. Neuartiges\_Coronavirus/Massnahmen\_Verdachtsfall. 2022. [https://www.rki.de/DE/Content/InfAZ/N/Neuartiges\\_Coronavirus/Massnahmen\\_Verdachtsfall\\_Infografik\\_DINA3.pdf?\\_\\_blob=publicationFile](https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Massnahmen_Verdachtsfall_Infografik_DINA3.pdf?__blob=publicationFile).
9. Sandra Vogel and Marc Breitenbroich (IW). Germany:working life in the Covid19-pandemic. 2022. [file:///C:/Users/blume/Desktop/lit\\_work\\_behavior/wpef21016.pdf](file:///C:/Users/blume/Desktop/lit_work_behavior/wpef21016.pdf).
10. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* 2020. doi:10.2807/1560-7917.ES.2020.25.3.2000045.
11. Jonczyk R, Stanislawski N, Seiler LK, Blume H, Heiden S, Lucas H, et al. Combined Prospective Seroconversion and PCR Data of Selected Cohorts Indicate a High Rate of Subclinical SARS-CoV-2 Infections—an Open Observational Study in Lower Saxony, Germany. *Microbiology Spectrum*. 2022;10:e01512-21.
12. Hernán MA. The C-Word: Scientific Euphemisms Do Not Improve Causal Inference From Observational Data. *Am J Public Health*. 2018;108:616–9. doi:10.2105/AJPH.2018.304337.
13. Lampert T. Smoking, physical inactivity, and obesity: associations with social status. *Dtsch Arztebl Int*. 2010;107:1–7. doi:10.3238/arztebl.2010.0001.
14. Geyer S. Soziale Ungleichheiten beim Auftreten chronischer Krankheiten. [Social inequalities in the occurrence of chronic diseases]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2016;59:181–7. doi:10.1007/s00103-015-2277-4.
15. Platten M, Nienhaus A, Peters C, Cranen R, Wisplinghoff H, Kersten JF, et al. Cumulative Incidence of SARS-CoV-2 in Healthcare Workers at a General Hospital in Germany during the Pandemic-A Longitudinal Analysis. *Int J Environ Res Public Health* 2022. doi:10.3390/ijerph19042429.
16. Šterbenc A, Tomič V, Bidovec Stojković U, Vrankar K, Rozman A, Zidarn M. Usefulness of rapid antigen testing for SARS-CoV-2 screening of healthcare workers: a pilot study. *Clin Exp Med*. 2022;22:157–60. doi:10.1007/s10238-021-00722-y.
17. Kortela E, Kirjavainen V, Ahava MJ, Jokiranta ST, But A, Lindahl A, et al. Real-life clinical sensitivity of SARS-CoV-2 RT-PCR test in symptomatic patients. *PLoS One*. 2021;16:e0251661. doi:10.1371/journal.pone.0251661.

18. Felkner SA, Streit JMK, McDaniel M, Schulte PA, Chosewood LC, Delclos GL, On BOTWPAP. How Will the Future of Work Shape OSH Research and Practice? A Workshop Summary. *Int J Environ Res Public Health* 2021. doi:10.3390/ijerph18115696.
19. Goveas JS, Shear MK. Grief and the COVID-19 Pandemic in Older Adults. *Am J Geriatr Psychiatry*. 2020;28:1119–25. doi:10.1016/j.jagp.2020.06.021.
20. Wilmes P, Zimmer J, Schulz J, Glod F, Veiber L, Mombaerts L, et al. SARS-CoV-2 transmission risk from asymptomatic carriers: Results from a mass screening programme in Luxembourg. *Lancet Reg Health Eur*. 2021;4:100056. doi:10.1016/j.lanepe.2021.100056.
21. Johnson-León M, Caplan AL, Kenny L, Buchan I, Fesi L, Olhava P, et al. Executive summary: It's wrong not to test: The case for universal, frequent rapid COVID-19 testing. *EClinicalMedicine*. 2021;33:100759. doi:10.1016/j.eclinm.2021.100759.
22. Handelsblatt. Sartorius boomt. 2022. <https://www.handelsblatt.com/unternehmen/industrie/quartalszahlen-corona-pandemie-befluegelt-labora-ausruester-sartorius-bis-ins-naechste-jahr-hinein/27720694.html>.
23. Azizi A, Achak D, Aboudi K, Saad E, Nejjari C, Noura Y, et al. Health-related quality of life and behavior-related lifestyle changes due to the COVID-19 home confinement: Dataset from a Moroccan sample. *Data Brief*. 2020;32:106239. doi:10.1016/j.dib.2020.106239.
24. Huls SPI, Sajjad A, Kanters TA, Hakkaart-van Roijen L, Brouwer WBF, van Exel J. Productivity of Working at Home and Time Allocation Between Paid Work, Unpaid Work and Leisure Activities During a Pandemic. *Pharmacoeconomics*. 2022;40:77–90. doi:10.1007/s40273-021-01078-7.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterial.pdf](#)