

A new family of eukaryotic and bacterial deubiquitinases spreading and diversifying through transposons and horizontal gene transfer

Ilka Erven

Institute for Genetics, University of Cologne <https://orcid.org/0000-0002-1095-6861>

Elena Abraham

Institute of Biochemistry, University of Cologne <https://orcid.org/0000-0002-3255-8565>

Thomas Hermanns

Institute for Genetics, University of Cologne

Ulrich Baumann

Institute of Biochemistry, University of Cologne

Kay Hofmann (✉ Kay.Hofmann@uni-koeln.de)

Institute for Genetics, University of Cologne <https://orcid.org/0000-0002-2289-9083>

Article

Keywords:

Posted Date: July 15th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1842489/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Communications on December 10th, 2022. See the published version at <https://doi.org/10.1038/s41467-022-35244-y>.

Abstract

Several distinct families of eukaryotic deubiquitinases (DUBs) regulate various aspects of ubiquitin signaling. Apart from one metalloenzyme family, six classes of eukaryotic papain-fold cysteine proteases are known to have DUB activity. A seventh cysteine-based DUB class is found in intracellular bacteria, which use DUB effectors to evade the ubiquitin-based defense response. Here, we report on the presence of an additional DUB class broadly distributed in eukaryotes and several bacteria. So far, the only known members of this family are the large tegument proteins of $\alpha/\beta/\gamma$ herpesviruses, which act as relatively unspecific deubiquitinases and deneddylases attached to the outer face of the viral capsid. By using a bioinformatical screen, we identified distant homologs of this family in transposons of vertebrates and invertebrates, in agaricomycete fungi, insects, nematodes, cnidaria, protists and selected intracellular bacteria. Members of this VTD (Viral tegument-like DUB) family use an atypical Cys-Asp-His catalytic triad to cleave ubiquitin chains. Some activities resemble the viral tegument DUBs in preferring K48-linked ubiquitin chains and also cleaving NEDD8, while other members are highly specific for K6- or K63-linked ubiquitin chains. Judging by the crystal structures of K48- and K6-specific members, the subfamilies differ considerably in their ubiquitin recognition mode. Our analyses suggest that this deubiquitinase family evolved from YopT/HopN-like non-DUB proteases and was co-opted early by DNA transposons, helping the spread of this family. In several independent events, the deubiquitinase domain of these transposons was 'domesticated' by various taxa, giving rise e.g. to the *Drosophila* male sterile (3)76Ca gene and several nematode genes with male-specific expression.

Introduction

Ubiquitination, the posttranslational modifications of proteins by covalent attachment of ubiquitin, is an important cellular regulatory mechanism. Unlike protein phosphorylation, the modification by ubiquitin rarely activates or inactivates proteins directly, but rather regulates other aspects such as protein degradation, intracellular transport, and recruitment to particular sites¹. An important contributor to the diversity of ubiquitin-based signals is the ability of multiple ubiquitin residues to be subject to further ubiquitination, leading to the formation of ubiquitin chains of different linkage types^{2,3}. While ubiquitin chains linked through lysine residues K48 and K11 often lead to proteasomal degradation of the modified substrates, K63-linked ubiquitin chains rather affect protein localization events. Besides the abundant K48, K63 and K11-linked chains, several minor linkage types are currently attracting attention due to their highly-regulated occurrence under cellular stress situations⁴⁻⁶. Of particular interest are K6-linked chains, which have been first observed after DNA damage, where they are probably generated by the ubiquitin ligase BRCA1⁷. K6 chains also play a role in the regulation of mitophagy, the removal of defective mitochondria by selective autophagy. Parkin, the mitophagy-associated ubiquitin ligase, can generate K6 chains and is itself modified and regulated by K6 ubiquitin^{8,9}. Furthermore, K6 chains generated by the ubiquitin ligase LRSAM1 are important for the defense against intracellular bacteria via ubiquitin-dependent autophagy.¹⁰

Deubiquitinating enzymes (DUBs) cleave peptide- and isopeptide-bonds downstream of ubiquitin. While some DUBs are required for ubiquitin activation and recycling, most DUB enzymes regulate ubiquitin signaling by undoing the activity of ubiquitin ligases - either by cleaving within ubiquitin chains or by completely removing ubiquitin from the substrate. Typical eukaryotes harbor many DUB types with different specificities for the substrates and/or the linkage of the attached ubiquitin chains¹¹. Apart from a few metalloproteases belonging to the JAMM/MPN+ family, most deubiquitinating enzymes are cysteine proteases, which – despite a high degree of structural variability – all assume a papain-like fold¹². In accordance with sequence relationship and structural criteria, eukaryotic cysteine-DUBs are usually categorized in six different classes: USP (ubiquitin-specific proteases), UCH (ubiquitin carboxyl-terminal hydrolases), OTU (ovarian tumor domain proteins), Josephins (Ataxin-3-like proteins), MINDY (MIU-containing new DUB family), and ZUP1/ZUFSP (zinc-finger ubiquitin protease 1). Among those, the highly-abundant USP class comprises mostly substrate-specific DUBs, while members of the OTU class are known for their pronounced linkage specificity¹³.

Many bacteria and viruses also code for deubiquitinating enzymes, which work within the host cell to counteract ubiquitin-based cellular defense responses. Most bacterial DUBs either belong to the OTU class or are members of the 'CE-clan', a class of papain-fold enzymes that includes proteases for several ubiquitin-related modifiers such as NEDD8 or SUMO. Most DUB effectors secreted by intracellular bacteria prefer K63-linked chains or have no strong linkage preference; it is assumed that these enzymes remove ubiquitin from the bacterial surface or from the cytoplasmic face of bacteria-containing vacuoles, thereby helping the bacteria to evade clearance by ubiquitin-mediated targeting to the lysosome. For *Legionella pneumophila*, two additional DUB activities have been identified: The effector RavD specifically cleaves linear (M1-linked) ubiquitin chains, thereby inhibiting inflammation and preventing M1-based xenophagy signals. The *Legionella* effector LotA, which belongs to the OTU class, has a K6-specific DUB activity of unknown biological significance¹⁴. Viral deubiquitinases are typically divergent members of the USP, OTU or CE-clan families and often fulfill multiple proteolytic functions¹⁵. Besides removing ubiquitin, many viral DUBs can also cleave the ubiquitin-related modifier ISG15, an important interferon-stimulated component of the antiviral response^{16,17}. In addition, viral DUBs are often tasked with processing viral polyproteins at multiple sites. This latter cleavage does not involve the hydrolysis of isopeptide bonds – a hallmark of DUB activity – but the cleaved peptide bonds are typically flanked by a di-glycine motif resembling the C-terminus of ubiquitin. Despite the structural differences between the cysteine DUB classes found in eukaryotes, their viruses, and prokaryotic pathogens, these enzymes not only share a common papain fold and a conserved active site topology; there are also detectable sequence similarities between different DUB classes, which argue for a common evolutionary ancestry and have been used for predicting novel and unusual DUB families^{18,19}.

The protease domain found in the large tegument protein of α , β and γ herpesviruses forms a small DUB family^{20–22}, which differs in several respects from other deubiquitinase classes: While these proteases share the papain-like fold and the ability to cleave ubiquitin chains of various linkage types, the unusual Cys-Asp-His active site has a different architecture²³. Moreover, extensive bioinformatical analyses did

not detect any sequence relationship to other deubiquitinase classes, suggesting a different evolutionary origin¹⁸. A number of biological roles for the tegument-associated deubiquitinases in different herpesviruses have been proposed, including the inhibition of NF- κ B and interferon signaling, and the prevention of selective autophagy^{22,24-26}. BPLF1, the tegument-associated DUB from Epstein-Barr virus, was shown to also possess deneddylase activity and thereby to modulate host-encoded cullin-based ubiquitin ligases to the virus advantage²⁷. The importance of the DUB domain of herpesvirus large tegument proteins for virion stability and virus entry could be separated from contributions of the unstructured C-terminal portion of the protein²⁸, which is required for attaching the protein to the outside of the viral capsid²⁹.

Based on the unusual active site architecture and the general dissimilarity to other deubiquitinases, the authors of the M48 crystal structure suggested that this tegument protein from human cytomegalovirus is the founding member of a new deubiquitinase family²³. To this day, the scope of this family has been limited to a set of closely related proteins from vertebrate herpesviruses. By performing a comprehensive bioinformatical screen, followed by biochemical and structural characterization of interesting candidates, we were able to identify active members of this family in a wide range of eukaryotic species and a few host-associated bacteria. Due to the similarity to the herpesvirus proteins, we refer to this extended enzyme family as VTD (for Viral Tegument-like Deubiquitinases). Our data suggest an origin of the VTD family by branching off from a non-DUB protease family, followed by a change in active site topology. Several pieces of evidence suggest that VTD proteins have spread to other taxa by becoming incorporated into DNA transposons and later 'domesticated' in multiple independent events. During this process, subfamilies of VTDs evolved to assume different linkage specificities, highlighting the versatility of this unusual deubiquitinase class.

Results

Bioinformatical discovery of VTD homologs

In order to learn more about the origins and evolution of the enigmatic herpesviral tegument DUBs, we used bioinformatical screens to identify distant homologs. In the first approach, a multiple alignment of all recognizable herpesviral homologs of UL36 (Herpes Simplex Virus), UL48 (Human Cytomegalovirus) and BPLF1 (Epstein-Barr Virus) was generated by using MAFFT³⁰. The resulting alignment was converted into a generalized sequence profile³¹ and used to search current protein databases. After two cycles of iterative profile refinement, a set of 2951 significantly matching sequences was retrieved, among them 1730 sequences from herpesviruses and 1221 non-viral sequences. Sequence analysis of the latter group revealed that the majority of non-viral homologs are found within the open reading frame of Helitrons, a class of rolling-circle DNA transposons present in all eukaryotic kingdoms³². Autonomous Helitron elements lack the classical transposase/integrase found in other DNA transposons, but rather encode a so-called 'RepHel' protein with both HUH-endonuclease and 5' to 3' helicase activity.³³ The Helitron members hosting the recognizable VTD domains carry an additional C-terminal AP-type

endonuclease and thus belong to the Helitron subclass of rolling-circle transposons³³. Previously, Helitrons have been described to sometimes include OTU-like domains, although their potential deubiquitinase activity has never been addressed³³. In fact, 121 of the VTD-containing Helitrons were found to also contain OTU domains. With the exception of several pseudogenized sequences, the tegument-like domains of presumed autonomous Helitrons have all catalytic residues conserved, suggesting that these transposon-associated domains have DUB activity. VTD-containing Helitrons were identified in vertebrates, insects, cnidarians, molluscs and plants. For further biochemical and structural characterization, we selected two Helitrons from zebrafish: DrT1 with two VTD domains but without OTU (LOC108180207 on chromosome 20) and DrT2 with a single active VTD and an additional OTU domain (AL772241 on chromosome 25) (Fig. 1a,b).

A second large group of non-viral VTD domains were identified in insects outside of a recognizable transposon context. Although these VTD domains are closely related to the Helitron-encoded VTDs (Supplementary Fig. 1), they are likely devoid of enzymatic activity as they lack one or more of the active site residues. Many of the identified insect sequences carry multiple copies of the inactive VTD (iVTD) domains, including three of the five identified proteins from *Drosophila melanogaster*: CG32436 (5 copies), CG939 (3 copies), CG4669 (2 copies), CG14402 (1 copy), CG32462 (1 copy). Interestingly, all five drosophila proteins are annotated to have a testis-specific expression pattern³⁴.

Besides the helitrons and inactive insect proteins, the list of identified VTD homologs also contains a few bacterial sequences, including Wc-VTD1 (wcv_1294) from *Waddlia chondrophila*, an emerging pathogen linked to miscarriages and infertility^{35,36}. While the bacterial sequences reach a significant similarity score, they are far more divergent from the viral VTDs (Fig. 1a, Supplementary Fig. 1). Their relationship to the established VTDs could be corroborated by Hidden-Markov-Model (HMM) searches starting from the bacterial members, which also uncovered a second, particularly divergent VTD in the *Waddlia chondrophila* gene Wc-VTD2 (wcv_1327) – a protein that did not reach significant scores in the initial search. Both *Waddlia* VTDs were selected for further biochemical analysis, their domain architecture is shown in Fig. 1b. The inclusion of the bacterial sequences into HMM searches revealed a further VTD subfamily with members in the fungal class *Agaricomycetes*, as well as in several protist classes. The fungal and protist VTD domains are rather similar to each other, but quite different from the viral and transposon-encoded VTDs (Supplementary Fig. 1). All members show active site conservation, and we selected two fungal members from *Serendipita indica* (PIIN_06337) and *Serpula lacrymans* (LOC18815922), and one protist member from the pathogenic amoeba *Naegleria fowleri* (FDP41_009013) for further analysis. Their domain structure is shown in Fig. 1b.

Finally, an HMM constructed from all VTDs found so far was run against protein family databases of various species, using the HHsearch algorithm³⁷. In this search, the *Drosophila melanogaster* male sterile gene ms(3)76Ca (CG14101) and five uncharacterized genes from the model nematode *Caenorhabditis elegans* (C09H10.9, C18G1.9, C14C11.1, ZC317.6, Y53F4B.36) were identified, which are broadly conserved in other insects and nematodes and do not form part of transposons. The coding sequences

of these genes are restricted to a single VTD domain (Fig. 1a,b), which contains all predicted catalytic residues. As shown in Supplementary Fig. 1, this invertebrate subfamily is quite divergent and shows no particular relationship to the insect-encoded inactive VTDs. We selected the *Drosophila* ms(3)76 Ca sequence for further biochemical analysis.

Helitron-encoded VTDs have similar activities as herpesviral deubiquitinases.

The three intact VTD domains encoded by the two selected *Danio rerio* helitrons were expressed in *E. coli*, purified, and tested against a set of model substrates comprising C-terminally propargylated ubiquitin and ubiquitin-like modifiers. These activity-based probes use a C-terminal PA-group to react covalently with functional thiol proteases of the correct specificity, resulting in an adduct of increased molecular weight³⁸. The zebrafish VTDs DrT1-VTD1 and DrT1-VTD2, both derived from the same Chr.20 helitron copy, reacted readily with Ub-PA and NEDD8-PA, while DrT2-VTD from the Chr.25 helitron copy only reacted with the propargylated ubiquitin probe (Fig. 2a-c). No reaction with the other three tested probes SUMO1-PA, SUMO3-PA and ISG15^{CTD}-PA (C-terminal domain of ISG15) could be detected. To directly compare the ubiquitin- and NEDD8-directed activities of the helitron-VTDs, they were additionally incubated with fluorogenic model substrates Ub-AMC and NEDD8-AMC, both carrying a C-terminal cleavable 7-amino-4-methylcoumarin (AMC) group. DrT1-VTD1 and DrT1-VTD2 were similarly efficient in removing ubiquitin and NEDD8 from the C-terminally fused fluorophore, as indicated by an increase in fluorescence. In accordance with these results, DrT2-VTD was active against the ubiquitin-AMC substrate, albeit with markedly lower reactivity compared to the other two helitron VTDs, while its ability to cleave NEDD8-AMC was negligible (Fig. 2d-i). To further investigate the DUB activity and linkage preferences of the three newly discovered deubiquitinases, the three enzymes were tested against a panel of di-ubiquitin species comprising all eight homotypic chain types. All three DrT-VTDs preferentially cleave K48- and K63-linked di-ubiquitin chains, with DrT1-VTD1 and DrT1-VTD2 also showing substantial activity towards K11-linked chains (Fig. 2j-l). These initial results show that the three helitron-derived VTDs cleave ubiquitin chains with a similar preference to that published for the viral tegument deubiquitinase UL36 from HSV-1³⁹.

A Helitron-VTD structure

To further elucidate the structural basis for VTD activity, we solved the crystal structure of DrT1-VTD2 (residues 300 to 520 of Uniprot entry A0A2R8QBC4) at a resolution of 1.90 Å and compared it to the ubiquitin complex of mouse cytomegalovirus M48, the only available structure of a herpesviral tegument deubiquitinase. As shown in Fig. 3a, the asymmetric unit contained one DrT1-VTD2 domain, which was completely resolved with the exception of residues 107 to 109 located in a protruding loop where Leu108 and Gly109 have rather weak density. In total, the structure contains four α -helices and nine β -strands, which adopt the form of an α - β - α sandwich, in which a bifurcated antiparallel β -sheet is flanked by α -helices. The helitron-VTD structure could be superimposed with the M48 catalytic domain (pdb:2J7Q) with an RMS distance of 2.5 Å over 192 residues, demonstrating an almost identical fold (Fig. 3b). The only major difference was observed for the aforementioned unstructured loop of DrT1-VTD2, which in

M48 corresponds to a well-folded β -hairpin contacting the outgoing S1-ubiquitin²³. Since the DrT1-VTD2 structure does not contain ubiquitin, the β -hairpin structure might be induced by substrate binding rather than representing a real difference between viral and transposon-encoded VTD domains. While viral and transposon-encoded VTDs adopt the papain-like fold shared with all other thiol deubiquitinases, they have a highly unusual active site architecture (Fig. 3c). In VTDs, the catalytic histidine and aspartate residues form an invariable Asp-x-His motif, while in typical DUBs the catalytic Asp/Asn residue follows the His residue at a variable distance. As a consequence, the catalytic histidine (His453 in DrT1-VTD2 and His158 in M48) resides on the opposite site of the catalytic triad, creating a mirror image to the classic papain fold active site (Supplementary Fig. 2a,b). In the herpesviral M48 structure, a second histidine (His141) is positioned close to the catalytic site and was found to act as a partial replacement for the main catalytic residue His-158²³. By contrast, the helitron VTD structure does not contain a second histidine residue with the potential to be involved in catalysis.

While in terms of sequence similarity and active site architecture, the VTDs are quite distinct from other deubiquitinases and even other cysteine proteases, a structure similarity search of the DrT1-VTD2 using the DALI method⁴⁰ revealed an interesting relationship to bacterial YopT-like enzymes. This family contains non-DUB proteases – often targeting small G-proteins – as demonstrated for *Pseudomonas syringae* AvrPphB and its homolog YopT from *Yersinia pestis*⁴¹. The structural alignment of DrT1-VTD2 with AvrPphB (pdb:1UKF) gave an RMS of 5.0 Å over 160 residues and shows an overall similar fold, while highlighting the different active site architecture (Fig. 3d,e), since the YopT family uses the conventional Cys-His-Asp catalytic triad. To confirm the predicted role of Cys322, Asp451 and His453 as the catalytic triad of DrT1-VTD2, these residues were mutated to alanine and tested for their catalytic activity. Ubiquitin-AMC assays as well as incubation of K48-linked di-ubiquitin chains with C322A, D451A or H453A mutants demonstrated a complete loss of activity (Fig. 3f,g).

Since all attempts to crystallize DrT-VTD2 in complex with ubiquitin failed, the structural correspondence to M48 was used to identify potential ubiquitin-binding interfaces by superposition of the M48 ~ Ubiquitin complex and DrT1-VTD2 (Supplementary Fig. 2c). In M48, a protruding β -hairpin consisting of residues 108 to 115 was found to be important for ubiquitin binding, with Leu110 and Tyr113 making important contacts with His68 and Val70 of ubiquitin's Ile44 patch²³. This β -hairpin corresponds to the flexible 402–413 loop in DrT-VTD2. However, the β -hairpin region is not well conserved among VTD-type enzymes, making it difficult to identify corresponding residues. When the DrT1-VTD2 residues 403–411 were replaced by a short Gly-Gly-Ser linker, the enzymatic activity against Ub-AMC and di-ubiquitin was completely lost (Fig. 3h,i). By contrast, single point mutations of three potential contact residues within the loop (P405A, A406G and L407A) and one residue in the adjacent β 4 strand (F398A) showed little effect on the chain-cleaving activity (Fig. 3i), suggesting a complex multi-factorial recognition of the ubiquitin Ile44 patch. To further assess the importance of Ile44 recognition, DrT1-VTD2 was incubated with a ubiquitin activity-based probe, in which Ile44 was mutated to alanine (Ub^{I44A}-PA). Interestingly, DrT1-VTD2 reacted with this mutant probe as well as with wildtype Ub-PA (Fig. 3j), suggesting a lack of direct Ile44 recognition by the DUB. As a second substrate recognition surface, typical DUBs bind the C-

terminal R-x-R motif of the S1 ubiquitin by one or two salt bridges using acidic residues of the DUB. Based on their conservation pattern in the VTD family and their positioning within the structure, Asp376 and Glu377 might be suitable candidates for this interaction. Individual mutations of both residues (D376A and E377A) resulted in a strongly reduced chain cleaving activity (Fig. 3i), and also a reduced cleavage of Ub-AMC, the latter effect being more pronounced for E377A (Fig. 3h).

Bacterial VTD domains show linkage-specific ubiquitin cleavage

While the viral and helitron-encoded VTD domains resemble each other in sequence, structure and specificity, the bacterial VTDs, of which Wc-VTD1 is a typical example, are more divergent. The second Waddlia-encoded Wc-VTD2 is particularly divergent and forms an outlier even to the other bacterial VTDs (Supplementary Fig. 1). We analyzed the catalytic domain (aa279-494) of Wc-VTD1 and the full-length Wc-VTD2 for their ability to cleave model substrates and ubiquitin chains. Both Waddlia VTDs reacted with the activity-based Ub-PA probe, but not with the analogous UBL probes SUMO1-PA, SUMO3-PA, NEDD8-PA or ISG15CTD-PA (Fig. 4a,b). When tested for their catalytic activity against AMC substrates, neither Wc-VTD1 nor Wc-VTD2 were able to cleave Ub-AMC or NEDD8-AMC (Fig. 4c,d). In the di-ubiquitin cleavage assay, Wc-VTD1 showed a strong preference for K6-linked chains. Apart from some trace activity against K11-diUb, none of the other linkages showed any cleavage in this assay (Fig. 4e). By contrast, Wc-VTD2 preferentially hydrolyzed K63-linked chains, with minor activity towards K11-linked di-ubiquitin (Fig. 4f). The strong linkage specificity of the two Waddlia VTDs suggests an important contribution of the proximal (S1') ubiquitin binding site and might thus explain the lack of activity against the mono-Ub based AMC substrate (Fig. 4c). Summarizing, the bacterial VTDs do not only diverge in sequence conservation from their viral counterparts, they have also acquired a different specificity, possibly accompanied by a different ubiquitin-recognition mode.

Structures of Wc-VTD1 reveal an unusual Ub binding mode

In order to gain insight into the functional differences of bacterial VTDs in comparison to their relatives from viruses and transposons, the structure of Wc-VTD1 was solved in the unbound form at 1.8 Å resolution as well as in a covalent complex with ubiquitin at 1.9 Å resolution. The latter was achieved by reacting the protease with ubiquitin-PA, which traps the Wc-VTD1 in an intermediate state with the ubiquitin being bound to the S1 site in a covalent linkage to the active cysteine. The tertiary structure contains nine α -helices and seven β -strands, adopting a fold with overall similarity to the viral and transposon-encoded VTDs (Fig. 5a, Fig SXXX). The catalytic triad is formed by Cys319, Asp440, and His442 (oxyanion hole?). As previously observed in the M48 crystal structure²³, the conformation of the catalytic histidine appears as non-productive for catalysis due to the large distance to the cysteine. However, considering the sequence alignment (Fig. 1a) and the site-directed mutagenesis experiments (Fig. 5h), the assignment of these residues to the catalytic triad is safe. The structures in the unbound and complexed state can be superimposed with an RMS distance of 0.538 Å over 144 C α -atoms excluding some 40 residues, thus indicating little overall conformational changes upon binding to the Ub-PA substrate. However, some parts of Wc-VTD1 undergo relatively large movements upon substrate

binding. Helix $\alpha 9$ (residues 474 to 486) and its preceding loop (residues 469 to 476) are pushed about 9 Å toward the active site by ubiquitin's loop consisting of residues 7 to 11, where it offers an interaction surface for the Ile-36 patch of the S1 ubiquitin (Fig. 5a,b,c). On the opposite side, helix $\alpha 1$ (residues 302 to 305) together with the following amino acids 306 to 311 is also shifted by a distance of about 8 Å. However, helix $\alpha 1$ itself does not seem to be involved in direct interactions with the bound substrate. It is located on the surface rather distant from the active site and is involved in the crystal contacts between the two crystallographically independent molecules in the apo structure. Roughly at this position, the proximal (S1') ubiquitin would be placed and thus helix $\alpha 1$ may be important for linkage specificity. The β -hairpin turn consisting of residues 421 to 423 is pushed away by about 2 Å from the active site cleft by ubiquitin's four very C-terminal residues (Fig. 5b). To compare the ubiquitin interaction of Wc-VTD1 to viral VTDs, a superposition of the bacterial protease with M48 was created, yielding an RMSD of 1.76 Å over 1113 atoms (Supplementary Fig. 3). Wc-VTD1 does not contain the ubiquitin-interacting β -hairpin of its viral relatives, but rather an irregular loop comprising residues 380–386 (colored yellow Fig. 5a,d and Supplementary Fig. 3b). This loop makes extensive contacts mainly via Pro383 and Phe384 to Leu8, Ile44, His68 and Val70 of the bound ubiquitin. The entire Ile44 patch of ubiquitin is localized within a hydrophobic pocket of the enzyme, which contributes multiple interactions between modifier and protease (Fig. 5d, Supplementary Fig. 3b).

Three residues were chosen to investigate their importance for ubiquitin recognition, based on their structural position: Phe384 located on the P383/F384 loop and Ala392 on the adjacent $\beta 2$ strand both contact the Ile44 patch, while Leu481 is part of the mobile $\alpha 9$ helix and contacts the Ile36 patch of ubiquitin. All three tested mutations (F384A, A392G and L481A) strongly reduce the activity of Wc-VTD1 on K6-linked ubiquitin chains (Fig. 5e). While the F384A and A392G mutants showed some residual activity, cleavage was completely abrogated by the L481A mutation, demonstrating the importance of the Ile36-patch recognition. To test if Wc-VTD1 shares with the helitron-VTD the property that the Ile44 patch of ubiquitin is contacted, while Ile44 itself is not crucial, the Wc-VTD1 domain was also incubated with the Ile44-mutated activity-based probe Ub^{I44A}-PA. Unlike the transposon-encoded enzyme, Wc-VTD1 only reacted with the wildtype Ub-PA, showing that this enzyme depends on a recognition of Ile44 itself (Fig. 5f). The alignment of Wc-VTD1 with M48 and other VTDs (Fig. 1a) suggested Glu363 and Asn364 as potential salt bridge partners for the basic RxR motif at the ubiquitin C-terminus. Mutation of either residue indeed had a strong impact on Wc-VTD1 activity against K6-linked chains: The E363A mutant appeared to be as inactive as the active site mutants, while the N364A mutation severely reduced K6 linked di-ubiquitin cleavage. (Fig. 5h). However, in the experimental crystal structure the side-chains of Asn364 of Wc-VTD1 and Arg74 of ubiquitin are completely disordered and there is no evidence for a significant interaction visible in the electron density map; the same is true for the moderately well-defined Arg72 and Glu363 side chains (Fig. 5g). A possible explanation for this is the very high ionic strength of the crystallization buffer (2 M citrate), which may interfere with the electrostatic interactions like salt bridges⁴².

Fungal and Naegleria VTDs are K6-directed deubiquitinases

Besides the transposon-derived and bacterial VTDs, potential candidates were also discovered in agaricomycete fungi and various protists. As fungal representatives, the VTD domain of *Serendipita indica* (Si-VTD) and *Serpula lacrimans* (Sl-VTD) were analyzed, while the short protein Nf-VTD from the pathogenic amoeba *Naegleria fowleri* was analyzed as protist representative. Since bacterially expressed Si-VTD and Sl-VTD become unstable upon tag removal, both proteins were purified with an N-terminal 6His-Smt3 tag. When incubated with a panel of UBL-derived activity-based probes, none of the three VTD candidates showed any reactivity (Fig. 6a-c). Similarly, activity against the fluorogenic substrates Ub-AMC and NEDD8-AMC was mostly absent, except for a weak cleavage of Ub-AMC by Nf-VTD (Fig. 6d-i). By contrast to their poor activity against mono-Ub based substrates, all three tested VTD enzymes were able to cleave di-ubiquitin chains, with a marked preference for K6-linkages (Fig. 6j-l). Nf-VTD at 500 nM cleaved about half of the K6 chains within 3 hours, with minor activity against K48 chains. (Fig. 6j). Si-VTD required a higher concentration of 5 μ M for reaching a similar level of cleavage (Fig. 6k), while Sl-VTD was more active at 500 nM and cleaved most of the K6-diUb after 1h (Fig. 6l). Unlike the *Naegleria* enzyme, the two fungal VTDs were truly K6-specific and showed no reactivity against any other linkage type. Thus, in accordance with their sequence similarity to bacterial VTDs, the fungal and protist homologs share the K6-specificity observed for *Waddlia* Wc-VTD1 (Fig. 4e).

Drosophila male-sterile VTD is a K48-directed deubiquitinase

As a representative for the divergent invertebrate-type VTDs found in insects and vertebrates, the protein encoded by the *Drosophila* Male sterile (3) 76Ca gene (Dm-VTD) was tested. As shown in Fig. 7a, the enzyme reacts with the Ub-PA probe, but also with a probe based on the C-terminal moiety of ISG15 – the latter being without biological relevance, since ISG15 is a vertebrate-specific modifier. No reactivity was observed towards NEDD8-PA or SUMO-PA probes (Fig. 7a). Dm-VTD was also shown to cleave the fluorogenic substrates Ub-AMC, but not NEDD8-AMC (Fig. 7b,c). When testing Dm-VTD against a panel of di-ubiquitin substrates, a preferential cleavage of K48-linked chains was observed, with some reactivity towards K63 and K11 chains. Despite the large evolutionary distance (Supplementary Fig. 1), the enzymatic properties of Dm-VTD resemble those of viral and helitron-encoded VTDs.

Discussion

VTD-type deubiquitinases are found in the genomes of animals, fungi, protists and bacteria, but their phyletic distribution is rather patchy. Horizontal gene transfer and recurrent gene losses are known to cause unusual phyletic distributions with different degrees of sparseness⁴³. In the evolution of the VTD family, both mechanisms appear to be at play. There are several discontinuous taxa with broadly observed VTD genes: One of them are the *Herpesviridae*, which evolved 180–220 million years ago (mya)⁴⁴. VTDs are found in α -, β -, and γ -herpesviruses, which infect mammals and birds, while no VTDs are found in *Alloherpesviridae*, a sister group that comprises fish and amphibian herpesviruses. Interestingly, *Alloherpesviridae* contain a large tegument protein with an OTU domain⁴⁵, suggesting that these viruses too might benefit from a ubiquitin-cleaving activity attached to the capsid. A second VTD-containing

taxon is formed by the *Agaricomycetes*, a mushroom-forming class of *Basidiomycete* fungi. Most sequenced *Agaricomycete* genomes encode a single protein with a VTD-domain fused to an RBR-type ubiquitin ligase; in other genomes (e.g. the eponymous *Agaricus bisporus*) this gene is unannotated, but still present. The VTD-containing *Agaricomycetes* evolved approx. 185 mya, while the *Dacrymycetes*, which diverged ~ 40 million years earlier^{46,47}, encode the corresponding RBR gene without a VTD domain. Apparently, the acquisition of viral and fungal VTDs happened roughly simultaneously in the mesozoic era, around the advent of the first mammals.

Older VTD genes are found in the invertebrates. Almost all nematode genomes of clade III (*Ascaridomorpha*, *Spirumorpha*) and clade V (*Rhabditids*, *Strongylids*) contain VTD genes, while other nematode clades lack VTDs. A recent phylogenetic analysis joins clade III and V nematodes in one taxon, suggesting that a VTD gene was acquired by a common precursor of these clades – possibly 400–800 mya⁴⁸. VTD-containing genes are also found in many arthropod clades, including *Hexapoda* (several flies, wasps, butterflies, beetles), *Arachnomorpha* (several ticks), *Crustacea* (several ostracods), and *Myriapoda* (*Strigamia*). However, only very few species within these highly populated taxa do contain VTD genes, raising the question if this sparsity is the result of sweeping gene losses in most lineages, or if it is caused by multiple independent acquisitions – possibly from an arthropod-associated transposon source. The former idea is supported by dendrogram analysis of extant arthropod VTDs, showing that their similarity roughly follows the phylogenetic tree of the host organism. Other invertebrate VTD genes might have originated from transposons: Several *Cnidaria* (corals, sea anemones) contain multiple genes with VTD domains fused to various other protein regions – among them several transposon-associated domains. While VTD proteins are widespread in eukaryotes, bacterial VTDs are currently limited to two members of *Chlamydiales* and one *Myxobacterium*, arguing against a bacterial origin of this family.

Judging by the similarity of VTD structures to members of the YopT/HopN family, while also considering the unusual VTD active site architecture and lack of detectable sequence relationship to other DUB classes, it is likely that the first VTD proteases branched off a YopT-like precursor gene in some eukaryotic species. Since all characterized VTD proteins cleave ubiquitin, while known YopT/HopN proteases have other targets⁴¹, this specificity shift might be a consequence of the ‘flipped’ active site topology. Interestingly, a similar change of active site topology with associated specificity shift has recently been described for members of the ZUFSP deubiquitinase family: ZUP1 and other members of the ZUFSP family use the canonical active site architecture found in papain and YopT/HopN proteases, while the related UBL-specific proteases UFSP2 and ATG4 use a VTD-like catalytic triad including the D-x-H motif^{49–52}. Apart from VTDs, members of the ZUFSP family are the only known deubiquitinases lacking the aromatic gatekeeper motif¹⁸, suggesting a mechanistic similarity between these enzyme families.

There is insufficient data to fully elucidate the evolutionary history of the extant VTD family, but the following hypothesis is plausible and compatible with the experimental data: After the ‘birth’ of the VTD family from a non-DUB source, it appears to have been co-opted relatively early by one or more transposon classes. Some of these transposons spread among invertebrates, leading to the introduction

of VTDs into clade III/V nematodes and an early arthropod precursor. Judging by the extant *Drosophila* VTD, these early enzymes probably preferred K48 chains, suggesting a role in protecting proteins from proteasomal degradation. Later, the K6-specific VTD form evolved – possibly in some protist or associated transposon – and was subsequently introduced into the precursor of *Agaricomycetes* and several protist lineages, including *Amoebozoa*. The latter are the most likely source for the bacterial K6-specific VTDs, since both *Waddlia* and *Simkania* are found in free-living amoebae^{35,53}. At approximately the same time, the precursor of *Herpesviridae* co-opted a K48-selective VTD – probably from a Helitron-type rolling circle transposon, which became incorporated into the large tegument protein of vertebrate herpesviruses.

Due to our failure to co-crystallize VTD domains complexed with di-ubiquitin chains, we cannot pinpoint the structural basis for the different linkage-specificities of the K6-preferring VTDs from fungi, protists and bacteria and the K48/K63/K11-preferring VTDs from viruses and transposons. The latter group does not only efficiently cleave three different linkage types, but is also highly active towards Ub-AMC, suggesting that a binding of the proximal (S1') ubiquitin is not important for catalysis. By contrast, the K6-specific VTDs are nearly inactive against Ub-AMC and might therefore harbor a specific recognition surface for the proximal (S1') ubiquitin in K6-orientation. The α 1-helix and flanking regions of Wc-VTD1 are well positioned to fulfill such a role. Since the α 1-region is shifted markedly between free and ubiquitin-bound structures of Wc-VTD1 (Fig. 5b), a crosstalk between S1- and S1'-recognition appears possible. Despite the requirement of a S1'-ubiquitin for efficient cleavage, the K6-specific Wc-VTD1 also recognizes multiple surfaces of the outgoing S1-ubiquitin. The poorly structured P383/F384 loop contacts the Ile44-patch of ubiquitin and appears to functionally replace the hairpin loop of viral and helitron-encoded VTDs. In addition, the Ile36 patch of ubiquitin is contacted by the C-terminal α 9-helix of Wc-VTD1 (Fig. 5a), which is completely absent in the viral and helitron VTD structures. As highlighted in Fig. 1a, this helix shows little sequence conservation but is present in all VTDs with K6-specificity (Wc-VTD1, Nf-VTD, Si-VTD, SI-VTD) while lacking from VTDs with other specificities. Thus, a possibly indirect role of the α 9-helix in K6-recognition cannot be excluded. In a recent study on the K6-specific DUB LotA, an interaction between the Ile36 and Ile44 patches of two adjacent K6-linked ubiquitin moieties was reported, which had to be dissolved by the enzyme before cleaving the isopeptide bond⁵⁴. It is possible that the α 9-helix of Wc-VTD1 fulfills a similar task.

Another open question concerns the functional role of VTD enzymes in their host species – and why many species groups apparently don't need these activities. Given that ubiquitination by K48-, K63- and K6-linked chains usually leads to very different outcomes, it is unlikely that all VTD deubiquitinases fulfil a common task. For the K48-cleaving herpesviral tegument DUBs, a number of biological effects have been published, often targeting the ubiquitin-mediated degradation of signaling components^{22,24-26} or using the deneddylase activity to target cullin-based ubiquitin ligases of the host²⁷. The closely related helitron-encoded VTDs might have a similar function by preventing the proteasomal degradation of proteins required for transposition. Interestingly, *Alloherpesviridae* – a virus group closely related to *Herpesviridae* – have replaced the VTD-domain in their tegument proteins by OTU-domains⁴⁵. Similarly,

several helitron-type rolling circle transposons also contain OTU domains besides – or instead of – VTD domains. Although neither viral nor transposon OTU domains have been tested for catalytic activity, those observations suggest that herpesviruses and helitrons benefit from a type of deubiquitination activity, which can be encoded by different DUB classes. Another intriguing observation is the male-specific expression of different animal VTD classes: There are five *Drosophila* genes that encode proteins with inactive VTD domains closely related to the helitron-VTDs, all of which are expressed exclusively in the male testis³⁴. The same is true for the *Drosophila* male sterile gene *ms(3)76Ca* (CG14101), which encodes a very distantly related but enzymatically active VTD domain. This gene, like its homologs from the nematode *C. elegans*, is also expressed exclusively in the male germline⁵⁵. Although the male-specific VTD types are not closely related, they both appear to be derived from domesticated transposons and either cleave K48-chains or are inactivated versions of K48-cleaving enzymes. It is therefore tempting to speculate that their transposon-encoded ancestors might have performed K48-linked deubiquitination in the male germline – possibly helping to overcome a mechanism for limiting transposon inheritance.

For the K6-specific VTDs, their biological role is equally enigmatic. In eukaryotes, no K6-specific DUBs are known; OTUD3 prefers K6 chains but also cleaves K11 linkages¹³. K6-linked ubiquitin chains have been implicated in various biological processes including DNA damage and mitophagy^{7–9}, but the particular role of this linkage type is insufficiently understood⁶. The only known example of a truly K6-specific deubiquitinase is the bacterial effector LotA from *Legionella pneumophila*, whose N-terminal OTU domain harbors this activity^{14,54}. *Waddlia chondrophila* is unrelated to *Legionella*, but has a similar intracellular lifestyle within bacteria-containing vacuoles. Thus, an analogous role for the K6-specific LotA and Wc-VTD1 appears plausible – possibly through the evasion of ubiquitin-dependent autophagy after modification by the K6-specific ubiquitin ligase LRSAM1¹⁰. Besides the aforementioned herpesviral and helitron-related domain displacements, the LotA/Wc-VTD1 dichotomy might be a third case of co-opting non-homologous DUB types for fulfilling analogous tasks. It thus appears that VTD-type deubiquitinases are widespread and versatile enzymes, which can functionally replace OTUs – a DUB class known for their linkage-selective chain cleavage.

Materials & Methods

Sequence Analysis

Sequence alignments were generated using the L-INS-I algorithm of the MAFFT package³⁰. The multiple alignments were used for the generation of generalized profiles using pftools³¹, and Hidden-Markov-Models using the HHSEARCH³⁷. Generalized profile searches were performed using all proteins from the Uniprot database (<https://www.uniprot.org>). Protein clustering was performed using the CLANS software⁵⁶. Structure comparisons were performed using the DALI method⁴⁰.

Cloning & Mutagenesis

The DrT-VTDs were amplified from *Danio rerio* genomic DNA or, in the case of DrT1-VTD2, from *Danio rerio* cDNA (kind gifts from Sigrun Korsching, University of Cologne), Wc-VTD2 was amplified from *Waddlia chondrophila* genomic DNA (kind gift from Carole Kebbi Beghdadi, University of Lausanne), Si-VTD was amplified from *Serendipita indica* cDNA (kind gift from Alga Zuccaro, University of Cologne), and Dm-VTD was amplified from *Drosophila melanogaster* cDNA (kind gift from Mirka Uhlirova, University of Cologne). All amplifications were done by PCR, using Phusion High Fidelity Kit (New England Biolabs). Wc-VTD1, Nf-VTD and SI-VTD coding regions were obtained by gene synthesis (IDT). The PCR fragments and gBlocks were cloned into pOPIN-S and pOPIN-K vectors⁵⁷ using the In-Fusion HD Cloning Kit (Takara Clontech). Point mutations were introduced using the QuikChange Lightning kit (Agilent Technologies).

Constructs for ubiquitin-PA purification (pTXB1-ubiquitin¹⁻⁷⁵) were a kind gift of David Komander (WEHI, Melbourne). SUMO1¹⁻⁹⁶, SUMO3¹⁻⁹¹ and ISG15⁷⁹⁻¹⁵⁶ were amplified by PCR with an N-terminal 3xFlag tag and cloned into the pTXB1 vector (New England Biolabs) by restriction cloning according to the manufacturers protocol. The Ub^{I44A} activity-based probe was created by introducing the mutation using the QuikChange Lightning kit (Agilent Technologies).

Protein expression & purification

All VTD candidates including all truncations and mutants were expressed from the pOPIN-S (in case of the bacterial and eukaryotic proteases) with an N-terminal 6His-Smt3-tag or pOPIN-K vector (in case of the transposon-derived candidates) with an N-terminal 6His-GST-tag. *Escherichia coli* (Strain: Rosetta (DE3) pLysS) were transformed with constructs expressing DUBs and 2-6 l cultures were grown in LB medium at 37 °C until the OD600 of 0.8 was reached. The cultures were cooled down to 18 °C and protein expression was induced by addition of 0.2 mM isopropyl β -d-1-thiogalactopyranoside (IPTG).

The expression of selenomethionine substituted proteins was carried out as described previously by⁵⁸: In brief, the expression cultures were grown in minimal medium until the OD600 of 0.8 was reached. The cultures were cooled down to 18 °C, mixed with feed-back inhibition amino acid mix (0.5 g/l final concentration), metal trace elements (0.1% final concentration) and vitamins (0.01% final concentration) and induced with 0.2 mM IPTG. After 16 h, the cultures were harvested by centrifugation at 5000 $\times g$ for 15 min. After freeze thaw, the pellets were resuspended in binding buffer (300 mM NaCl, 20 mM TRIS pH 7.5, 20 mM imidazole, 2 mM β -mercaptoethanol) containing DNase and Lysozyme, and lysed by sonication using 10 s pulses with 50 W for a total time of 10 min. Lysates were clarified by centrifugation at 50,000 $\times g$ for 1 h at 4 °C and the supernatant was used for affinity purification on HisTrap FF columns (GE Healthcare) according to the manufacturer's instructions. The 6His-Smt3 tag was removed by incubation with SENP1⁴¹⁵⁻⁶⁴⁴, the 6His-GST tag was removed by incubation with 3C protease. Si-VTD and SI-VTD were purified including the N-terminal 6His-Smt3 tag. The proteins were simultaneously dialyzed in binding buffer. The liberated affinity-tag and the His-tagged SENP1 and 3C proteases were removed by a second round of affinity purification with HisTrap FF columns (GE Healthcare). All proteins were purified

with a final size exclusion chromatography (HiLoad 16/600 Superdex 75 or 200 pg) in 20 mM TRIS pH 7.5, 150 mM NaCl, 2 mM dithiothreitol (DTT), concentrated using VIVASPIN 20 Columns (Sartorius), flash frozen in liquid nitrogen, and stored at -80°C . Protein concentrations were determined using the absorption at 280 nm (A_{280}) using the proteins' extinction coefficients derived from their sequences.

Synthesis of activity-based probes

All activity-based probes used in this study were expressed as C-terminal intein fusion proteins as described previously⁵⁹: In brief, the fusion proteins were affinity purified in buffer A (20 mM HEPES, 50 mM sodium acetate pH 6.5, 75 mM NaCl) from clarified lysates using Chitin Resin (New England Biolabs) following the manufacturer's protocol. On-bead cleavage was performed by incubation with cleavage buffer (buffer A containing 100 mM MesNa (sodium 2-mercaptoethanesulfonate)) for 24 h at room temperature (RT). The resin was washed extensively with buffer A and the pooled fractions were concentrated and subjected to size exclusion chromatography (HiLoad 16/600 Superdex 75pg) with buffer A. To synthesize the propargylated probe, 300 μM Ub/Ubl-MesNa were reacted with 600 μM propargylamine hydrochloride (Sigma Aldrich) in buffer A containing 150 mM NaOH for 3 h at RT. Unreacted propargylamine was removed by size exclusion chromatography and the probe was concentrated using VIVASPIN 20 Columns (3 kDa cutoff, Sartorius), flash frozen and stored at -80°C . The NEDD8-PA was a kind gift from David Pérez Berrocal and Monique Mulder (Department of Cell and Chemical Biology, Leiden University).

Chain generation

Met1-linked di-ubiquitin was expressed as a linear fusion protein and purified by ion exchange chromatography and size exclusion chromatography. K6-, K11-, K48-, and K63-linked ubiquitin chains were enzymatically assembled using UBE2S Δ C (K11), CDC34 (K48), and Ubc13/UBE2V1 (K63) as previously described^{60,61}. In brief, ubiquitin chains were generated by incubation of 1 μM E1, 25 μM of the respective E2, and 2 mM ubiquitin in reaction buffer (10 mM ATP, 40 mM TRIS (pH 7.5), 10 mM MgCl_2 , 1 mM DTT) for 18 h at RT. K6-linked ubiquitin chains were assembled by incubation of 1 μM E1, 25 μM E2 (UbCH7), 25 μM E3 (Nle1) and 2 mM ubiquitin in reaction buffer. The generated mixture of K6- and K48-linked chains was treated with 10 μM OTUB1. The respective reactions were stopped by 20-fold dilution in 50 mM sodium acetate (pH 4.5) and chains of different lengths were separated by cation exchange using a Resource S column (GE Healthcare). Elution of different chain lengths was achieved with a gradient from 0 to 600 mM NaCl.

Crystallization

DrT1-VTD2 (selenomethionine substituted), Wc-VTD1 (selenomethionine substituted) and Wc-VTD1~Ub were crystallized using sitting drop vapor diffusion with commercially available sparse matrix screens. 96

well iQ crystallization plates containing 30 μ l of the respective screening conditions were mixed with 10 mg/ml protein in the ratios 1:2, 1:1 and 2:1 in 300 nl drops. For DrT1-VTD2 and Wc-VTD1, the initial conditions containing the most promising crystals were optimized by gradually changing the chemical components included in the respective screening condition. 80 μ l reservoir solution were pipetted into 48-well MRC plates and sitting drop vapour diffusion was performed by mixing 10 mg/ml protein in the ratios 1:2, 1:1 and 2:1 in drops of 3 μ l. Initial DrT1-VTD2 crystals were detected in Proplex D12 (0.1 M Tris pH 8.5, 20% w/v PEG 6000) and several other conditions after 3 days at 20°C. Optimization was carried out with 3 μ l drops (protein/precipitant ratios: 2:1, 1:1 and 1:2) and precipitant solutions varying in pH or PEG 6000 concentration respectively. Optimized crystals were harvested and cryoprotected with reservoir containing 10% glycerol. For unbound Wc-VTD1, initial crystals were detected in JCSG H8 (0.2 M sodium chloride, 0.1 M BisTris pH 5.5, 25 % w/v PEG 3350) and several other conditions after 3 days at 20°C. Optimization was carried out with 3 μ l drops (protein/precipitant ratios: 2:1, 1:1 and 1:2) and precipitant solutions varying in sodium chloride or PEG 3350 concentration respectively. Optimized crystals were harvested and cryoprotected with reservoir containing 20% glycerol. 1.2 mM Wc-VTD1 were incubated with 760 μ M ubiquitin-PA for 18 h at 4 °C. Unreacted Wc-VTD1 and Ub-PA were removed by size exclusion chromatography and the complex concentrated to a concentration of 7 mg/ml. The covalent Wc-VTD1~Ub crystals were detected in Salt RX B6 (0.1 M BisTris propane pH 7.0, 2.0 M Ammonium citrate tribasic) after 2 days at 20 °C. Crystals were harvested without further optimization and cryoprotected with the addition of 10% glycerol.

Data collection, phasing, model building, and refinement

Diffraction data for DrT1-VTD2, Wc-VTD1 and the Wc-VTD1~Ub-PA complex was collected at the Deutsches Elektronen-Synchrotron (DESY), Hamburg, Germany at beamline P13 at the EMBL outstation ⁶². All datasets were processed using XDS ⁶³. For DrT1-VTD2 and Wc-VTD1, initial phases were determined using selenomethionine SAD experiments and SHELXC/D/E ⁶⁴, which were driven by HKL2MAP ⁶⁵. Both protein structures were subsequently built automatically using the ARP/wARP Web Service ⁶⁶ or the buccaneer pipeline implemented in CCP4 ⁶⁷. The Wc-VTD1~Ub-PA complex was solved by molecular replacement using PHASER ⁶⁸ as implemented in the phenix package ⁶⁹ with a single molecule of Wc-VTD1 and a full-length model of ubiquitin as search models. For further refinement, necessary restraints for the propargyl moiety were calculated using AceDRG implemented in the CCP4 package ⁷⁰. Initial models were refined using iterative cycles of phenix.refine (DrT1-VTD2, Wc-VTD1) or RefMac (Wc-VTD1~Ub-PA) and manually rebuilt using COOT ⁷¹. For structural analysis, the PyMOL (<http://www.pymol.org>) and ChimeraX Graphics Systems⁷² were used.

AMC assays

Activity assays of DUBs against AMC-labeled substrates were performed using reaction buffer (150 mM NaCl, 20 mM TRIS pH 7.5, 10 mM DTT), 1 μ M DUBs (deviating concentrations are indicated in the

respective figures and the corresponding legends), 5 μ M Ub-AMC (UbiQ-Bio, The Netherlands) or 5 μ M NEDD8-AMC (Enzo Life Science). The reaction was performed in black 96-well plates (Corning) at 30 °C and fluorescence was measured using the Infinite F200 Pro plate reader (Tecan) equipped for excitation wavelength of 360 nm and an emission wavelength of 465 nm. The presented results are means of three independent cleavage assays.

Activity based probe assays

DUBs were prediluted to 2 \times concentration (10 μ M) in reaction buffer (20 mM TRIS pH 7.5, 150 mM NaCl and 10 mM DTT) and 1:1 combined with 100 μ M Ub-, Ub^{I44A}-, SUMO1, SUMO3, ISG15^{CTD} or NEDD8-PA for 18 hours at 4°C. The reaction was stopped by the addition of 2x Laemmli buffer, resolved by SDS-PAGE, and Coomassie stained.

Ubiquitin chain cleavage

DUBs were preincubated in 150 mM NaCl, 20 mM TRIS pH 7.5 and 10 mM DTT for 10 min. The cleavage was performed for the indicated time points with 25 nM up to 500 nM DUBs (as indicated in the respective figure legends) and 25 μ M di-ubiquitin (K11, K48, K63, M1, K6 synthesized as described above, others purchased from Boston Biochem) at RT. The reactions were stopped with 2x Laemmli buffer, resolved by SDS-PAGE, and Coomassie stained.

Tables

Table 1. Data collection and refinement statistics.

	DrT1-VTD2	Wc-VTD1	Wc-VTD1~Ubiquitin
Wavelength (Å)	0.9747	0.9763	1.0000
Resolution range (Å) ¹	42.37 - 1.90 (1.97 - 1.90)	51.06 - 1.70 (1.80 - 1.70)	50.88 - 1.73 (1.84 - 1.73)
Space group	C 2 2 2 ₁	P 1 2 ₁ 1	P 3 ₂ 2 1
Unit cell (Å, deg)	119.01 120.67 34.17 90 90 90	37.12 70.30 75.89 90 101.84 90	101.77 101.77 63.20 90 90 120
Total reflections	481,803 (14,406)	279,246 (41,211)	690,794 (107,527)
Unique reflections ²	37,458 (2724)	81,659 (13,147)	39,674 (6,344)
Multiplicity ²	12.9 (5.3)	3.4 (3.2)	17.4 (16.9)
Completeness (%)	100.0 (99.9)	98.8 (98.3)	100.0 (100.0)
Mean I/sigma(I)	11.5 (1.7)	8.5 (1.0)	11.0 (1.1)
Wilson B-factor	23.3	25.9	30.6
R-merge	0.198 (1.02)	0.079 (1.18)	0.146 (2.57)
R-meas	0.209 (1.09)	0.106 (1.6)	0.148 (2.78)
CC1/2	0.998 (0.63)	0.995 (0.63)	0.998 (0.50)
Refinement resolution range	42.37 - 1.90 (1.97 - 1.90)	51.06 - 1.70 (1.76 - 1.70)	50.88 - 1.73 (1.79 - 1.73)
Reflections used in refinement	19941 (1930)	41742 (4124)	39633 (3914)
Reflections used for R-free	1494 (144)	2075 (208)	1985 (196)
R-work	0.167 (0.257)	0.205 (0.382)	0.167 (0.320)
R-free	0.213 (0.283)	0.244 (0.393)	0.192 (0.373)
CC (work)	0.96 (0.87)	0.96 (0.80)	0.968 (0.74)
CC (free)	0.93 (0.89)	0.96 (0.71)	0.96 (0.68)
Number of non-hydrogen atoms	1832	3382	2402
macromolecules	1674	3201	2246
ligands	0	0	28
solvent	158	181	139
Protein residues	212	406	284
RMS (bonds)	0.005	0.015	0.011
RMS (angles)	0.85	1.18	1.22
Ramachandran favored (%)	96.14	99.24	98.92
Ramachandran allowed (%)	3.86	0.76	1.08
Ramachandran outliers (%)	0.00	0.00	0.00
Rotamer outliers (%)	0.00	0.57	0.40
Clashscore	3.62	6.17	4.66
Average B-factor	31.06	43.90	39.46
macromolecules	30.42	43.92	38.91
ligands			66.66
solvent	37.87	43.41	45.07
Number of TLS groups	5	2	2
PDB entry	8ADD	8ADC	8ADB

¹ Statistics for the highest-resolution shell are shown in parentheses

² Friedel pairs are counted as different reflections for the anomalous data sets DrT1-VTD2 and Wc-VTD1

Declarations

Acknowledgments

We thank Christiane Horst and Claudia Poschner for expert technical assistance. We are also grateful to David Pérez Berrocal, Monique Mulder, Sigrun Korsching, Mirka Uhlirova and Alga Zuccaro for reagents and constructs. The synchrotron data was collected at beamline P13 operated by EMBL Hamburg at the PETRA III storage ring (DESY, Hamburg, Germany). We would like to thank David von Stetten, Michael Agthe, Gleb Bourenkov, and Isabel Bento for the assistance in using the beamline. Deubiquitinase research in the lab of KH is supported by DFG grant HO 3783/3-1. Crystals were grown using equipment of the Cologne Crystallization facility (C₂f), which is supported by DFG grant INST 216/949-1 FUGG.

Author contributions

IE performed most biochemical experiments and crystallization, EA solved the X-ray structures, TH contributed to the experiments, UB supervised the crystallography, KH initiated and supervised the study and contributed bioinformatical analyses. All authors contributed to data analysis and the writing of the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Data availability

The X-ray structures of DrT1-VTD1 and Wc-VTD in its apo and ubiquitin-bound form have been deposited at the PDB database under the accession numbers 8ADD, 8ADC and 8ADB, respectively. The data underlying the findings of this study are available in this article and its Supplementary Information or are available from the corresponding author upon reasonable request.

References

1. Komander, D. & Rape, M. The ubiquitin code. *Annu Rev Biochem* **81**, 203–229, (2012).
2. Swatek, K. N. & Komander, D. Ubiquitin modifications. *Cell Res* **26**, 399–422, (2016).
3. Yau, R. & Rape, M. The increasing complexity of the ubiquitin code. *Nat Cell Biol* **18**, 579–586, (2016).
4. Tracz, M. & Bialek, W. Beyond K48 and K63: non-canonical protein ubiquitination. *Cell Mol Biol Lett* **26**, 1, (2021).
5. van Huizen, M. & Kikkert, M. The Role of Atypical Ubiquitin Chains in the Regulation of the Antiviral Innate Immune Response. *Front Cell Dev Biol* **7**, 392, (2019).
6. Michel, M. A., Swatek, K. N., Hospenthal, M. K. & Komander, D. Ubiquitin Linkage-Specific Affimers Reveal Insights into K6-Linked Ubiquitin Signaling. *Mol Cell* **68**, 233–246 e235, (2017).

7. Wu-Baer, F., Lagazon, K., Yuan, W. & Baer, R. The BRCA1/BARD1 heterodimer assembles polyubiquitin chains through an unconventional linkage involving lysine residue K6 of ubiquitin. *J Biol Chem* **278**, 34743–34746, (2003).
8. Durcan, T. M. *et al.* USP8 regulates mitophagy by removing K6-linked ubiquitin conjugates from parkin. *EMBO J* **33**, 2473–2491, (2014).
9. Ordureau, A. *et al.* Quantitative proteomics reveal a feedforward mechanism for mitochondrial PARKIN translocation and ubiquitin chain synthesis. *Mol Cell* **56**, 360–375, (2014).
10. Huett, A. *et al.* The LRR and RING domain protein LRSAM1 is an E3 ligase crucial for ubiquitin-dependent autophagy of intracellular *Salmonella Typhimurium*. *Cell Host Microbe* **12**, 778–790, (2012).
11. Clague, M. J., Urbe, S. & Komander, D. Breaking the chains: deubiquitylating enzyme specificity begets function. *Nat Rev Mol Cell Biol* **20**, 338–352, (2019).
12. Pruneda, J. N. & Komander, D. Evaluating enzyme activities and structures of DUBs. *Methods Enzymol* **618**, 321–341, (2019).
13. Mevissen, T. E. *et al.* OTU deubiquitinases reveal mechanisms of linkage specificity and enable ubiquitin chain restriction analysis. *Cell* **154**, 169–184, (2013).
14. Kubori, T., Kitao, T., Ando, H. & Nagai, H. LotA, a *Legionella* deubiquitinase, has dual catalytic activity and contributes to intracellular growth. *Cell Microbiol* **20**, e12840, (2018).
15. Bailey-Elkin, B. A., Knaap, R. C. M., Kikkert, M. & Mark, B. L. Structure and Function of Viral Deubiquitinating Enzymes. *J Mol Biol* **429**, 3441–3470, (2017).
16. Proulx, J., Borgmann, K. & Park, I. W. Role of Virally-Encoded Deubiquitinating Enzymes in Regulation of the Virus Life Cycle. *Int J Mol Sci* **22**, (2021).
17. Zhang, Q., Jia, Q., Gao, W. & Zhang, W. The Role of Deubiquitinases in Virus Replication and Host Innate Immune Response. *Front Microbiol* **13**, 839624, (2022).
18. Hermanns, T. *et al.* An evolutionary approach to systematic discovery of novel deubiquitinases, applied to *Legionella*. *Life Sci Alliance* **3**, (2020).
19. Scheel, H., Tomiuk, S. & Hofmann, K. Elucidation of ataxin-3 and ataxin-7 function by integrative bioinformatics. *Hum Mol Genet* **12**, 2845–2852, (2003).
20. Kattenhorn, L. M., Korb, G. A., Kessler, B. M., Spooner, E. & Ploegh, H. L. A deubiquitinating enzyme encoded by HSV-1 belongs to a family of cysteine proteases that is conserved across the family Herpesviridae. *Mol Cell* **19**, 547–557, (2005).
21. Kim, E. T., Oh, S. E., Lee, Y. O., Gibson, W. & Ahn, J. H. Cleavage specificity of the UL48 deubiquitinating protease activity of human cytomegalovirus and the growth of an active-site mutant virus in cultured cells. *J Virol* **83**, 12046–12056, (2009).
22. Yla-Anttila, P., Gupta, S. & Masucci, M. G. The Epstein-Barr virus deubiquitinase BPLF1 targets SQSTM1/p62 to inhibit selective autophagy. *Autophagy* **17**, 3461–3474, (2021).

23. Schlieker, C. *et al.* Structure of a herpesvirus-encoded cysteine protease reveals a unique class of deubiquitinating enzymes. *Mol Cell* **25**, 677–687, (2007).
24. Ye, R., Su, C., Xu, H. & Zheng, C. Herpes Simplex Virus 1 Ubiquitin-Specific Protease UL36 Abrogates NF-kappaB Activation in DNA Sensing Signal Pathway. *J Virol* **91**, (2017).
25. Kumari, P. *et al.* Essential role of HCMV deubiquitinase in promoting oncogenesis by targeting anti-viral innate immune signaling pathways. *Cell Death Dis* **8**, e3078, (2017).
26. Gupta, S., Yla-Anttila, P. & Masucci, M. G. A novel mechanism for regulation of the type I IFN response by herpesvirus deconjugases. *Microb Cell* **5**, 259–261, (2018).
27. Gastaldello, S. *et al.* A deneddylase encoded by Epstein-Barr virus promotes viral DNA replication by regulating the activity of cullin-RING ligases. *Nat Cell Biol* **12**, 351–361, (2010).
28. Kim, Y. E., Oh, S. E., Kwon, K. M., Lee, C. H. & Ahn, J. H. Involvement of the N-Terminal Deubiquitinating Protease Domain of Human Cytomegalovirus UL48 Tegument Protein in Autoubiquitination, Virion Stability, and Virus Entry. *J Virol* **90**, 3229–3242, (2016).
29. Newcomb, W. W. & Brown, J. C. Structure and capsid association of the herpesvirus large tegument protein UL36. *J Virol* **84**, 9408–9414, (2010).
30. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular biology and evolution* **30**, 772–780, (2013).
31. Bucher, P., Karplus, K., Moeri, N. & Hofmann, K. A flexible motif search technique based on generalized profiles. *Comput Chem* **20**, 3–23, (1996).
32. Kapitonov, V. V. & Jurka, J. Helitrons on a roll: eukaryotic rolling-circle transposons. *Trends Genet* **23**, 521–529, (2007).
33. Thomas, J., Phillips, C. D., Baker, R. J. & Pritham, E. J. Rolling-circle transposons catalyze genomic innovation in a mammalian lineage. *Genome Biol Evol* **6**, 2595–2610, (2014).
34. Leader, D. P., Krause, S. A., Pandit, A., Davies, S. A. & Dow, J. A. T. FlyAtlas 2: a new version of the *Drosophila melanogaster* expression atlas with RNA-Seq, miRNA-Seq and sex-specific data. *Nucleic Acids Res* **46**, D809-D815, (2018).
35. de Barys, M. & Greub, G. *Waddlia chondrophila*: from biology to pathogenicity. *Microbes Infect* **15**, 1033–1041, (2013).
36. Baud, D. *et al.* *Waddlia chondrophila* and Male Infertility. *Microorganisms* **8**, (2020).
37. Soding, J. Protein homology detection by HMM-HMM comparison. *Bioinformatics* **21**, 951–960, (2005).
38. Ekkebus, R. *et al.* On terminal alkynes that can react with active-site cysteine nucleophiles in proteases. *J Am Chem Soc* **135**, 2867–2870, (2013).
39. Lin, J. *et al.* UL36 Encoded by Marek's Disease Virus Exhibits Linkage-Specific Deubiquitinase Activity. *Int J Mol Sci* **21**, (2020).
40. Holm, L. Using Dali for Protein Structure Comparison. *Methods Mol Biol* **2112**, 29–42, (2020).

41. Shao, F., Merritt, P. M., Bao, Z., Innes, R. W. & Dixon, J. E. A *Yersinia* effector and a *Pseudomonas* avirulence protein define a family of cysteine proteases functioning in bacterial pathogenesis. *Cell* **109**, 575–588, (2002).
42. Horovitz, A., Serrano, L., Avron, B., Bycroft, M. & Fersht, A. R. Strength and co-operativity of contributions of surface salt bridges to protein stability. *J Mol Biol* **216**, 1031–1044, (1990).
43. Keeling, P. J. & Palmer, J. D. Horizontal gene transfer in eukaryotic evolution. *Nat Rev Genet* **9**, 605–618, (2008).
44. Davison, A. J. Evolution of the herpesviruses. *Vet Microbiol* **86**, 69–88, (2002).
45. Michel, B. *et al.* The genome of cyprinid herpesvirus 3 encodes 40 proteins incorporated in mature virions. *J Gen Virol* **91**, 452–462, (2010).
46. Varga, T. *et al.* Megaphylogeny resolves global patterns of mushroom evolution. *Nat Ecol Evol* **3**, 668–678, (2019).
47. Floudas, D. *et al.* The Paleozoic origin of enzymatic lignin decomposition reconstructed from 31 fungal genomes. *Science* **336**, 1715–1719, (2012).
48. International Helminth Genomes, C. Comparative genomics of the major parasitic worms. *Nat Genet* **51**, 163–174, (2019).
49. Hermanns, T. *et al.* A family of unconventional deubiquitinases with modular chain specificity determinants. *Nat Commun* **9**, 799, (2018).
50. Kwasna, D. *et al.* Discovery and Characterization of ZUFSP/ZUP1, a Distinct Deubiquitinase Class Important for Genome Stability. *Mol Cell* **70**, 150–164 e156, (2018).
51. Hermanns, T., Pichlo, C., Baumann, U. & Hofmann, K. A structural basis for the diverse linkage specificities within the ZUFSP deubiquitinase family. *Nat Commun* **13**, 401, (2022).
52. Satoo, K. *et al.* The structure of Atg4B-LC3 complex reveals the mechanism of LC3 processing and delipidation during autophagy. *EMBO J* **28**, 1341–1350, (2009).
53. Lamoth, F. & Greub, G. Amoebal pathogens as emerging causal agents of pneumonia. *FEMS Microbiol Rev* **34**, 260–280, (2010).
54. Warren, G. D. *et al.* Mechanism of Lys6 poly-ubiquitin specificity by the *L. pneumophila* deubiquitinase LotA. *bioRxiv*, 2022.2005.2011.491541, (2022).
55. Ebbing, A. *et al.* Spatial Transcriptomics of *C. elegans* Males and Hermaphrodites Identifies Sex-Specific Differences in Gene Expression Patterns. *Dev Cell* **47**, 801–813 e806, (2018).
56. Frickey, T. & Lupas, A. CLANS: a Java application for visualizing protein families based on pairwise similarity. *Bioinformatics* **20**, 3702–3704, (2004).
57. Berrow, N. S. *et al.* A versatile ligation-independent cloning method suitable for high-throughput expression screening applications. *Nucleic Acids Res* **35**, e45, (2007).
58. Van Duyne, G. D., Standaert, R. F., Karplus, P. A., Schreiber, S. L. & Clardy, J. Atomic structures of the human immunophilin FKBP-12 complexes with FK506 and rapamycin. *J Mol Biol* **229**, 105–124, (1993).

59. Borodovsky, A. *et al.* Chemistry-based functional proteomics reveals novel members of the deubiquitinating enzyme family. *Chem Biol* **9**, 1149–1159, (2002).
60. Bremm, A., Freund, S. M. & Komander, D. Lys11-linked ubiquitin chains adopt compact conformations and are preferentially hydrolyzed by the deubiquitinase Cezanne. *Nat Struct Mol Biol* **17**, 939–947, (2010).
61. Komander, D. & Barford, D. Structure of the A20 OTU domain and mechanistic insights into deubiquitination. *Biochem J* **409**, 77–85, (2008).
62. Cianci, M. *et al.* P13, the EMBL macromolecular crystallography beamline at the low-emittance PETRA III ring for high- and low-energy phasing with variable beam focusing. *J Synchrotron Radiat* **24**, 323–332, (2017).
63. Kabsch, W. Xds. *Acta crystallographica. Section D, Biological crystallography* **66**, 125–132, (2010).
64. Sheldrick, G. M. *et al.* Shelx. (2012).
65. Pape, T. & Schneider, T. R. HKL2MAP: a graphical user interface for macromolecular phasing with SHELX programs. *Journal of Applied Crystallography* **37**, 843–844, (2004).
66. Langer, G., Cohen, S. X., Lamzin, V. S. & Perrakis, A. Automated macromolecular model building for X-ray crystallography using ARP/wARP version 7. *Nat Protoc* **3**, 1171–1179, (2008).
67. Cowtan, K. The Buccaneer software for automated model building. 1. Tracing protein chains. *Acta crystallographica. Section D, Biological crystallography* **62**, 1002–1011, (2006).
68. McCoy, A. J. *et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658–674, (2007).
69. Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta crystallographica. Section D, Biological crystallography* **66**, 213–221, (2010).
70. Long, F. *et al.* AceDRG: a stereochemical description generator for ligands. *Acta Crystallogr D Struct Biol* **73**, 112–122, (2017).
71. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta crystallographica. Section D, Biological crystallography* **66**, 486–501, (2010).
72. Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci* **30**, 70–82, (2021).

Figures

in red and other domain types in grey. The position of the VDT domains as aligned in part a) of the figure are indicated by numbers.




Figure 2

Activity of Helitron-encoded VTDs. a-c) Reaction of DrT1-VTD1 (a), DrT1-VTD2 (b) or DrT2-VTD with Ub and Ubl activity-based probes. Asterisks mark the shifted bands after reaction. **d-e)** Activity of 5 nM DrT1-

VTD1 against Ubiquitin-AMC (d) and 1 nM DrT1-VTD1 against NEDD8-AMC (e). The RFU values are the means of triplicates. **f-g)** Activity of 5 nM DrT1-VTD2 against Ubiquitin-AMC (f) and NEDD8-AMC (g). The RFU values are the means of triplicates. **h-i)** Activity of 100 nM DrT2-VTD against Ubiquitin-AMC (d) and NEDD8-AMC (e). The RFU values are the means of triplicates. **j-l)** Linkage specificity analysis of DrT-VTDs. A panel of homotypic di-ubiquitin chains was incubated with DrT1-VTD1 (j), DrT1-VTD2 (k) or DrT2-VTD (l) for the indicated time points.




Figure 3

Crystal structure of a Helitron-encoded VTD. **a)** Crystal structure of DrT1-VTD2 in cartoon representation. The catalytic triad and residues potentially important for ubiquitin interaction are shown as sticks. The unstructured loop is coloured purple. **b)** Structural superposition of DrT1-VTD2 (blue) with M48 (PDB: 2J7Q, green). The RMSD is 2.5 Å for 192 C_α atoms. **c)** Active site architecture of DrT1-VTD2 (green sticks) compared to M48 (PDB: 2J7Q, purple sticks). **d)** Structural superposition of DrT1-VTD2 (blue) with AvrPphB (PDB: 1UKF, pink). The RMSD is 5.0 Å over 160 residues. **e)** Active site architecture of DrT1-VTD2 (green sticks) compared to AvrPphB (PDB: 1UKF, purple sticks). **f)** Activity of wildtype DrT1-VTD2 (WT) against K48-linked di-ubiquitin, compared to the catalytically inactive mutants C322A, D451A and H453A. **g)** Activity of 5 nM wildtype DrT1-VTD2 (WT, black) against Ubiquitin-AMC, compared to 5 nM of the catalytically inactive mutants C322A (red), D451A (blue) and H453A (yellow). The RFU values are the means of triplicates. **h)** Activity of 5 nM wildtype DrT1-VTD2 (WT, black) against Ubiquitin-AMC compared to 5 nM of the binding mutants D376A (green), E377A (red) and Δhairpin (blue). The RFU values are the means of triplicates. **i)** Activity of wildtype DrT1-VTD2 (WT) against K48-linked di-ubiquitin compared to the binding mutants Δhairpin, F398A, P405A, A406G, L407A, D376A and E377A. **j)** Reaction of DrT1-VTD2 with the wildtype Ub (Ub-PA) and Ub^{I44A} activity-based probe after overnight incubation (18h). Asterisks mark the shifted bands after reaction.




Figure 4

Activity of bacterial VTDs. **a-b)** Reaction of Wc-VTD1 (a) and Wc-VTD2 (b) with Ub and Ubl activity-based probes. Asterisks mark the shifted bands after reaction. **c-d)** Activity of Wc-VTD1 and Wc-VTD2 against Ubiquitin-AMC (c) and NEDD8-AMC (d). The RFU values are the means of triplicates. **e-f)** Linkage specificity analysis of Wc-VTDs. A panel of homotypic di-ubiquitin chains was incubated with Wc-VTD1 (e) or Wc-VTD2 (f) for the indicated time points.




Figure 5

Crystal structure of *Waddlia VTD1*. **a)** Crystal structure of Wc-VTD1 (teal) in covalent complex with Ub-PA (orange) in cartoon representation. The $\alpha 9$ helix is colored blue, $\beta 2$ is pink and the P383F384 loop is depicted in yellow. **b)** Structural superposition of apo Wc-VTD1 (yellow) and ubiquitin-bound Wc-VTD1 (teal). The RMSD is 0.54 Å over 205 residues. **c)** Magnification of $\alpha 9$ interaction with S1 ubiquitin. Residues of $\alpha 9$ in Wc-VTD1 are shown as blue sticks and the hydrophobic interaction interface of

ubiquitin as orange sticks. **d)** Putative interactions between ubiquitin's Ile44 patch (residues shown as orange sticks) with Wc-VTD1 through its α 9 (blue), β 2 (pink) and P383F384 loop (yellow) Mutated residues are shown as sticks. **e)** Activity of wildtype Wc-VTD2 (WT) against K6-linked di-ubiquitin compared to the binding mutants F384A, A392G, L481A. **f)** Reaction of Wc-VTD1 with the wildtype Ub (Ub-PA) and Ub^{I44A} activity-based probe after overnight incubation (18h). Asterisk marks the shifted bands after reaction. **g)** Interaction of Wc-VTD1 (teal) with ubiquitin's C-terminus (orange). The C-terminus of ubiquitin is shown as orange sticks and the propargylamide bond is marked with an asterisk. The catalytic cysteine of Wc-VTD1 is colored yellow and E363 and N364 are shown as grey sticks. The electron density map highlights the flexible R74. **h)** Activity of wildtype Wc-VTD1 (WT) against K6-linked di-ubiquitin compared to the catalytically inactive mutants C313A, D440A and H442A, and against binding mutants E363A and N364A.




Figure 6

Activity of fungal and protozoan VTDs. a-c) Reaction of Nf-VTD (a), Si-VTD (b) or SI-VTD with Ub and Ub1 activity-based probes. Asterisks mark the shifted bands after reaction. **d-e)** Activity of Nf-VTD against Ubiquitin-AMC (d) and NEDD8-AMC (e). The RFU values are the means of triplicates. **f-g)** Activity of Si-VTD against Ubiquitin-AMC (f) and NEDD8-AMC (g). The RFU values are the means of triplicates. **h-i)** Activity of SI-VTD against Ubiquitin-AMC (d) and NEDD8-AMC (e). The RFU values are the means of

triplicates. **j-l)** Linkage specificity analysis of eukaryotic VTDs. A panel of homotypic di-ubiquitin chains was incubated with Nf-VTD (j), Si-VTD (k) or SI-VTD (l) for the indicated time points.




Figure 7

Activity of invertebrate VTDs. a) Reaction of Dm-VTD with Ub and Ubl activity-based probes. Asterisks mark the shifted bands after reaction. **b-c)** Activity of Dm-VTD against Ubiquitin-AMC (b) and NEDD8-AMC (c). The RFU values are the means of triplicates. **d)** Linkage specificity analysis. A panel of homotypic di-ubiquitin chains was incubated with Dm-VTD for the indicated time points.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [D1292124205valreportfullP1.pdf](#)
- [D1292124207valreportfullP1.pdf](#)
- [D1292124209valreportfullP1.pdf](#)
- [SuppFigure1.pdf](#)
- [SuppFigure2.pdf](#)
- [SuppFigure3.pdf](#)
- [SuppLegends.docx](#)