

Interventions in the first 1000 days to prevent childhood obesity: a systematic review and quantitative content analysis

Celia Pérez-Muñoz (✉ celia.perez@uca.es)

University of Cádiz

Jesús Carretero-Bravo

University of Cádiz

Esther Ortega-Martín

University of Cádiz

Begoña Ramos-Fiol

University of Cádiz

Bernardo Ferriz-Mas

Andalusian Health System

Mercedes Díaz-Rodríguez

University of Cádiz

Research Article

Keywords: childhood obesity, first 1000 days, intervention

Posted Date: July 27th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1846058/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Childhood obesity poses a global health challenge. Despite efforts to reverse this situation in recent years, the figures remain high. In recent years, there has been an increase in interventions that begin in pregnancy, putting the concept of early programming into practice. The present study aims to update the findings regarding interventions in the first 1000 days of life.

Methods: A systematic review of the literature was carried out in PubMed, WoS, Scopus and CINAHL to obtain the articles to be analysed. We included those studies published between 2016 and 2021. Human interventions that started within the first 1000 days of life and acted on at least one programming factor were included. Once selected, coding and quantitative content analysis was carried out to obtain a profile of the interventions during the first 1000 days.

Results: From all screened articles, 51 unique interventions, which met the selection criteria, were included. The majority of interventions (81%) take place in developed areas. Almost all (86%) are targeted at the general population. The majority (54%) start in the second trimester of pregnancy. A clear majority (61%) end at the time of birth. 44% of the interventions include all pregnant women. Only 48% of these interventions are focused on improving the nutritional status of the offspring in the short term. Most interventions collect the baby's weight at birth (68%).

Conclusions: It can be concluded that current interventions are not covering as many aspects as they should. Future research should be conducted more frequently in developing countries and target disadvantaged groups. These interventions should include all pregnant women, aiming to cover as many programming factors as possible and extending through the first 1000 days of life, with BMI or skinfolds as measures of effectiveness during this period.

Introduction

Childhood obesity poses a global health, policy and research challenge [1]. In 1975, 32 million children under five years of age worldwide had overweight or obesity, growing to 42 million by 2020 [1, 2]. Of more significant concern is the data for children as young as two years old, with this group already accounting for 10% of obesity in the US [1].

In the short and medium-term, childhood obesity is associated with multiple "adult" diseases such as type 2 diabetes mellitus, hypertension, non-alcoholic fatty liver disease and dyslipidaemia [3, 4]. In the long term, children suffering from obesity have an increased risk of developing type II diabetes, hypertension, dyslipidaemia and atherosclerosis in adulthood [3]. According to a systematic review, most children with obesity become adults with obesity [5]. This, coupled with associated comorbidities, is why the increasing prevalence of obesity in childhood poses a challenge to health systems [6].

Although it may seem to be a problem in developed countries, according to WHO estimates, of the 42 million children under five years of age who are overweight globally, 35 million belong to developing countries [1]. On the other hand, studies reveal inequalities between different social classes, with children from the most disadvantaged social groups being more affected by being overweight [7, 8].

Despite efforts to reverse this situation in recent years, the figures remain high, suggesting that the approach to interventions is inadequate. Research into the effect of a range of environmental factors in the first 1,000 days of life, including pregnancy and the first two years of life, on developing this type of disease has led to changing strategies [9]. In recent years, there has been an increase in interventions that begin in pregnancy, putting the concept of early programming into practice [10] and are leading the way to prevent childhood obesity. Early programming is considered an adaptive response to the environment, based on the

information received during fetal life and the first years of postnatal life [10].

Many risk factors at this stage of development have been shown to affect future childhood obesity: maternal weight at the start of pregnancy, gestational weight gain (GWG), gestational diabetes mellitus (GDM), smoking, maternal undernutrition, caesarean section, high protein intake, no or short breastfeeding, vitamin D deficiency, excessive weight gain during the first year of life, and others such as sleep quality, screen viewing, attachment, appetite control, physical activity or complementary feeding [11–15]. In addition, it is common for the same child to have more than one of these factors, which has been shown to have a cumulative effect, increasing the risk of obesity [16, 17]. However, current interventions generally do not cover all of these factors but focus on some of them, especially through nutrition and physical activity [18], leaving aside others as important as rest or the use of screens, among others.

It is essential to analyse the implemented interventions to prevent childhood obesity in the first 1000 days of life. Previous reviews up to 2016 have done a similar analysis of these interventions [19–22]. However, these reviews cover interventions targeting a broader age range, including birth to 18 years. The present study aims to update the findings regarding interventions in the first 1000 days of life by reviewing the scientific literature in the main databases and conducting a quantitative analysis of the content of these databases [22–25]. With this, we will be able to outline the main characteristics of the interventions and identify gaps and shortcomings to target future interventions to address these possible deficiencies.

Methodology

A systematic review of the literature based on the guidelines for Systematic Reviews and Meta-Analyses (PRISMA) was carried out to obtain the articles to be analysed [26], proceed to their screening, select against the inclusion criteria, and select those finally included for analysis. The protocol had previously been defined, detailing the different phases of the process and the inclusion and exclusion criteria, data extraction and analysis methods. The protocol was registered in the PROSPERO platform on 02 November 2021 (CRD42021282951).

We included those studies published between 01 January 2016 (the period covered by the previous reviews) [19, 23] and 31 December 2021. Studies published in English and Spanish were included, languages that the authors were fluent in and did not require external translation. Human interventions that started

within the first 1000 days of life and acted on at least one programming factor were included [11–15]. Articles were excluded if they were not interventions, were not original articles or were only the publication of a protocol, with no publications having been carried out at the review date. Following Cochrane guidelines for systematic reviews, only one article per study was included. [28]. If more than one article per study was found, the one with the most recent data on offspring follow-up was included.

Once selected, coding and quantitative content analysis was carried out to obtain a profile of the interventions during the first 1000 days.

Search strategy and initial screening

The search was carried out in four databases: PubMed, WoS, Scopus and CINAHL. These were the databases selected as they are the most commonly used in the most recent systematic reviews and to not miss relevant information regarding interventions for the prevention of childhood obesity during the first 1000 days. For this purpose, the conjunction of different terms was adapted to each database to obtain the best results. The complete search strategy could be seen on additional file 1. The terms used were derived from the research question and were related to obesity, the first 1000 days and interventions. The filters offered in the databases regarding publication date, language and human research were used.

The searches were conducted during September 2021, and a final search was conducted in January 2022 to cover articles published between September and December 2021. This last search found only 1,903 new articles, of which, after the screening, none were included in the final analysis as they did not meet the inclusion criteria.

Figure 1 summarises the process from search to final inclusion of the selected articles. A total of 45,675 articles were identified, of which 19,789 were eliminated as duplicates using a reference manager. The remaining 25,886 articles were screened by title, eliminating all those not written in the target language, did not involve humans, had other objectives or were not original research articles. Finally, 2,631 studies were selected in this phase to continue with the process.

Application of eligibility criteria

The articles selected so far were reviewed by abstracts. Eligibility criteria were applied, which were guidelines for screening based on study abstracts. Interventions were included if they had their start in the first 1000 days of life, collected some offspring variable, at least birth weight, and aimed, directly or indirectly, to prevent childhood obesity in the short or long term.

In this step, 2,119 articles were eliminated, resulting in 512 studies being selected for the next stage. In the full-text reading, the same eligibility criteria were applied, checking for any remaining doubts by simply reading the abstract and confirming that all inclusion criteria were met in detail. Finally, 456 articles were excluded at this stage, so 56 studies were included in the review. The reasons for exclusion at this stage were: being only the protocol of intervention, not being an intervention, not being original research articles, not collecting any offspring variables, or being an intervention prior to the defined date. The latter case refers to interventions that have been conducted well before the stated publication date and whose results were previously published and included in previous reviews. The reviewers reached a 92% level of agreement, while discrepancies were resolved by discussion.

The 56 selected articles were reviewed according to the first author's surname, name of the intervention and country where it was developed to identify duplicate studies from the publication of several articles of the same intervention. Finally, five articles were eliminated, resulting in a sample of 51 articles belonging to 51 unique interventions included for analysis. The complete list of included articles can be found in additional file 2.

Data extraction

To standardise the coding process, the different categories from which information has been extracted were coded. Content analysis methodology was used in order to extract and analyse the characteristics of each intervention [22–25]. Three articles, identified in the initial search and excluded during the screening, were used to test the coding that had been established. Finally, four articles included in the final sample (approximately 10%) were used to confirm the codes used.

Having carried out this preliminary step and the assessment of inter-rater reliability, the reviewers carried out the coding of a part of the final sample of 51 articles, each one separately.

General characteristics of the interventions

The general characteristics of the articles and the interventions they showed were coded. For the characteristics of the articles themselves, the year of publication and the main subject of the journal in which they were published were coded. Six categories were established for the year of publication (2016, 2017, 2018, 2019, 2020 and 2021), in the same way as for the main topic of the journal of publication (Nutrition, Obesity, Childhood obesity, Gynaecology and obstetrics, Medicine and Paediatrics).

Concerning the general characteristics of the interventions, we coded the geographical region in which it took place, the type of population targeted, the setting in which it took place and how it was developed.

Six categories were defined in terms of geographic region, including Europe/UK, USA, Australia/New Zealand, South America, Asia and Africa, thus covering all possible geographic areas. The target population was coded into general, ethnic/racial, and low-income groups.

Because the interventions being evaluated in this article started in the first 1000 days of life, the setting in which they are developed was coded into four categories (community-based, health care, home-based, multi-setting) since it is not possible to develop them in other settings such as schools. Finally, all interventions were concentrated in two established categories (community-based and health care).

The mode in which the intervention took place was also coded. Three categories were established: in person, with technology, both.

Specific characteristics of the interventions

All the characteristics of the interventions of interest for profiling and gap identification, in order to guide future research, have been coded. In this way, the time at which the intervention begins was coded classified into three categories: first trimester, second trimester and third trimester, given that all the interventions began during the period of pregnancy. As for the end of the intervention, six categories were established according to whether the interventions ended during pregnancy, at birth, during the first year, at the first year, at the second year or after the second year.

Five categories were coded for the mothers included in terms of their nutritional status at the start of pregnancy: all pregnant women, overweight pregnant women, pregnant women with obesity, only normal-weight pregnant women, or pregnant women with a clinical history of GWG, GDM or macrosomic offspring. About the aspect that the intervention addresses intending to improve it, four categories were coded: excessive weight gain, gestational diabetes mellitus, pregnancy outcomes in general and nutritional status of the offspring. On the other hand, how the intervention addresses these aspects is categorised, resulting in six categories: nutrition, physical activity, nutrition and physical activity, general lifestyle including breastfeeding promotion, general lifestyle including breastfeeding and sleep, and supplement use.

Five categories were coded for the main variables collected: GWG, DMG, measures of children, breastfeeding and pregnancy outcomes. The measures collected in the offspring were classified into birth weight, body mass index (BMI) at different times or skinfolds. Finally, the time these measures were taken was coded into four categories: at birth, during the first year, from birth to two years and after two years.

Inter-rater reliability

Two observers were responsible for data extraction. To test for inter-rater reliability, 26 articles were randomly selected from those that would ultimately form part of the review (50%), and inter-rater agreement was tested using Cohen's kappa coefficient [29]. Two rounds of coding training were conducted to achieve sufficient reliability. Reliability was assessed for each variable. The mean kappa score for all variables was 0.80, while the percentage of simple agreement reached 94.4%. Variables with a minimum kappa of 0.7 [30]. Two variables had a kappa value below this score: What does it aim to address? (0.66) and Variables Collected Baby (0.61), but as they had a high total agreement and were variables with more than four categories they were considered for analysis, as the kappa coefficient is difficult to interpret as the number of categories increases [30, 31].

Data synthesis and analysis

First, the data was cleaned. In cases where data were missing or in doubt, the entire article was reviewed to confirm the information. The unit of analysis is the intervention, so a denominator of 51 interventions was used to assess the characteristics of the interventions. The tables presented show the number of interventions, accompanied by the percentage of the sample in brackets, that meet a given characteristic. Inter-rater reliability, as well as all other analyses, were performed with SPSS (version 24).

Results

From all screened articles, 51 unique interventions, which met the selection criteria, were included. The general characteristics of these interventions are summarised in Table 1.

As for the year of publication, we observe a peak of interventions published in 2016 (24%) that declines in the following years, although it seems to increase again in the last years, 2019 and 2021. Suppose we refer to the main topic of the journals in which the selected interventions are published. In that case, we find that most interventions are published in gynaecology (25%) and nutrition journals (28%), with only 8% being published in specific journals of childhood obesity or 10% in obesity journals.

Focusing on the geographical region in which the intervention takes place, it stands out that the majority of interventions (81%) take place in developed areas, such as the USA (33%), Australia/New Zealand (24%) and Europe/UK (24%). There are very few interventions in developing areas such as Africa (2%), South America (5%) or Central America (0%) or Asia (12%).

Table 1

General characteristics of childhood obesity prevention interventions published from 2016 to 2021 (n = 51)

	n (%)
Publication Year	
2016	12 (24)
2017	5 (10)
2018	8 (16)
2019	9 (18)
2020	7 (13)
2021	10 (19)
Topic of the Journal of publication	
Nutrition	14 (28)
Obesity	5 (10)
Childhood obesity	4 (8)
Gynaecology	13 (25)
Medicine	10 (19)
Paediatrics	5 (10)
Geographic region	
Europe/United Kingdom	12 (24)
United States	17 (33)
Australia/New Zealand	12 (24)
South America	3 (5)
Asia	6 (12)
Africa	1 (2)
Targeted population	
General population	44 (86)
Racial/ethnic groups	6 (12)
Low-income groups	1 (2)
Setting	
Community-based	16 (31)
Health care	35 (69)
Mode of delivery	
In-person	31 (61)
Using technology	2 (4)
Both	18 (35)

In terms of the target population of these interventions, almost all (86%) are targeted at the general population, with very few interventions specifically targeting ethnic or racial groups (12%) or the lower-income population (2%).

It is only in the USA that interventions targeting racial or ethnic groups are developed, specifically 33% of all interventions developed in this country. On the other hand, in South America, the only intervention aimed at the lower-income population has been developed.

In terms of the setting of the interventions included in this review, we found that most interventions (69%) have been developed in health care settings.

Finally, most interventions continue to be carried out face-to-face (61%). However, there is a growing tendency to use technology in combination with face-to-face development of interventions (35%).

The characteristics of the interventions are summarised in Table 2.

Table 2

Intervention characteristics of childhood obesity prevention interventions published from 2016 to 2021 (n = 51)

	n (%)
Start of the intervention	
First trimester	14 (28)
Second trimester	28 (54)
Third trimester	4 (8)
Postpartum	5 (10)
End of the intervention	
During pregnancy	5 (10)
At birth	31 (61)
During the first year	6 (12)
At first year	4 (8)
At second year	2 (4)
> 2 years	3 (5)
Included mothers	
All pregnant	22 (44)
Normal weight	5 (10)
With obesity and overweight	15 (29)
With obesity	6 (12)
With clinic antecedents	3 (5)
Targeted issue	
Gestational weight gain	13 (25)
Gestational Diabetes Mellitus	3 (5)
Pregnancy outcomes	11 (22)
Children nutritional status	24 (48)
Type of intervention ^a	
Nutrition	8 (16)
Physical Activity	11 (22)
Nutrition and Physical Activity	17 (33)
Lifestyle (including breastfeeding)	13 (25)
Lifestyle (including sleep)	3 (5)
Use of supplements	2 (4)
Main outcomes	
GWG	18 (36)
GDM	10 (19)
Breastfeeding	6 (12)
Measures of children	14 (28)
Pregnancy outcomes	3 (5)
Measures of children	
Birthweight	35 (68)
Body mass index (BMI)	8 (16)
Skinfolds	8 (16)
Moment of children measures	

Birth	30 (59)
During the first year	10 (19)
Birth to 2 years	4 (8)
> 2 years	7 (14)

a Groups are not mutually exclusive thus, totals may exceed 100%.

In terms of the time these interventions begin, all of the interventions included in the review begin during pregnancy. Specifically, the majority (54%) start in the second trimester of pregnancy, i.e. from the 14th week. Only 28% of the interventions start before this time, during the first trimester.

There are more differences in the timing of the end of the interventions. A clear majority (61%) end at the time of birth. Of the remainder, 10% end before pregnancy comes to an end, while 12% end at different times during the first year of life. 8% of interventions continue into the second year of life and end at some point during the second year of life, while only 4% end at the age of two years. On the other hand, only 5% of the interventions continue after the second year of life.

Regarding the group of women targeted by the intervention, in terms of their nutritional status at the start of pregnancy, 44% of the interventions include all pregnant women. In comparison, 29% include only those with overweight, i.e. with overweight or obesity. 12% of the interventions target only women with obesity, and 10% target only women with average weight. Finally, 5% of the interventions target mothers with a medical history of gestational diabetes, a newborn with macrosomia or some other complication of pregnancy.

We find different categories if we analyse the issue on which the interventions are focused. Only 48% of these interventions are focused on improving the nutritional status of the offspring in the short term. 25% focus on controlling the weight gain of pregnant women during this period in order to avoid excessive weight gain. 22% of interventions aim to improve pregnancy outcomes, while 5% focus on preventing the development of gestational diabetes mellitus.

Regarding the type of intervention, most interventions are nutritional and physical activity at the same time (33%). While 25% address lifestyle to improve health, including breastfeeding, only 5% also include rest as a determining factor.

In terms of the main variables collected, there are many interventions whose main outcome measure is excessive weight gain (36%). The rest focus on collecting the development of gestational diabetes mellitus (19%), initiation of breastfeeding (12%), anthropometric measures in offspring (28%) or pregnancy outcomes in general (5%).

Most interventions collect the baby's weight at birth (68%). Only 16% take BMI measurements, which is valid for skinfolds. As for the timing of these measurements, 19% take them at different times during the first year of life, only 8% of the interventions take them at two years of age, while 14% take them after this age.

Table 3 represents a cross-tabulation of the mothers included, the type of intervention, the main variables, the variables collected from the baby and the time of collection of these variables according to the aspect addressed.

Table 3

Included mothers, type of intervention, main outcomes, children's measures and moment of collection depending on the targeted issue (n=51)

n (%)	Targeted issue		
	GWG + GDM	Pregnancy outcomes	Children nutritional status
Included mothers			
All pregnant	8 (50)	5 (45)	10 (42)
Normal weight	1 (6)	3 (28)	0 (0)
With obesity and overweight	5 (32)	1 (9)	9 (38)
With obesity	1 (6)	2 (18)	3 (12)
With clinic antecedents	1 (6)	0 (0)	2 (8)
Type of intervention			
Nutrition	3 (19)	0 (0)	5 (21)
Physical Activity	5 (32)	3 (28)	3 (12)
Nutrition and Physical Activity	7 (43)	6 (54)	4 (17)
Lifestyle (including breastfeeding/sleep)	1 (6)	2 (18)	10 (42)
Use of supplements	0 (0)	0 (0)	2 (8)
Main outcomes ^a			
GWG	16 (100)	5 (45)	4 (17)
GDM	9 (56)	7 (63)	0 (0)
Breastfeeding	0 (0)	2 (18)	4 (17)
Measures of children	0 (0)	5 (46)	13 (54)
Pregnancy outcomes	0 (0)	3 (28)	3 (12)
Measures of children			
Birthweight	14 (88)	6 (55)	8 (33)
Body mass index (BMI)	2 (12)	1 (9)	11 (46)
Skinfolds	0 (0)	4 (36)	5 (21)
Moment of children measures			
Birth	14 (88)	7 (63)	9 (38)
During the first year	1 (6)	3 (28)	6 (25)
Birth to 2 years	0 (0)	1 (9)	3 (12)
> 2 years	1 (6)	0 (0)	6 (25)

^a Groups are not mutually exclusive thus, totals may exceed 100%.

It is observed that those interventions that focus on controlling GWG or GDM include mostly (50%) all types of women in terms of nutritional status. However, a high percentage (32%) include only those women with excess weight. They are mainly interventions that combine nutrition and physical activity as tools (43%). Finally, they are interventions that mainly collect the GWG of mothers (100%) and the birth weight of offspring (88%).

Interventions that aim to improve pregnancy outcomes generally are more diverse in terms of the nutritional status of the mothers included, although the majority (45%) include all types of women. 54% of these interventions address nutrition and physical activity together. As for the collection of variables, there is diversity among the interventions, finding those that collect GWG, GDM, different measures in the offspring or at the end of pregnancy.

Finally, interventions focused on improving the nutritional status of offspring mainly include all mothers (42%), but remains high the interventions that only include those mothers who have obesity or overweight (38%). They improve overall lifestyle, including breastfeeding or sleep, to achieve their goals (42%). Most of these interventions collect measures of offspring at birth (38%). These measures include skinfolds and BMI, as well as birth weight.

Figure 2 shows the type of developed interventions, depending on the targeted issue. It highlights how most interventions focus on nutrition or physical activity while only those that focus on improving the nutritional status of offspring cover lifestyle in general.

Discussion

As early as the first half of the 20th century, thanks to Kermack's epidemiological studies, it became known that early exposure to environmental factors could have a long-term effect on the health of offspring [32, 33]. Subsequently, it was thanks to Barker and his formulation of the hypothesis of the foetal origin of adult diseases that a direct link was established between the incidence of certain adult diseases and an environment of malnutrition during pregnancy [34].

Early programming is now considered an adaptive response to the environment, which develops as a function of information received through the womb, during gestation and, after birth, up to two years of age [35]. This concept has taken centre stage in recommendations for preventing childhood obesity [36, 37].

Following a systematic review process, this study shows the results of a quantitative content analysis of interventions to prevent childhood obesity, directly or indirectly, published between 2016 and 2021 to obtain a profile and identify gaps to guide future research towards effectiveness.

Gaps have been identified concerning the geographical region in which the interventions are developed and the social groups targeted. On the other hand, gaps have been identified in terms of the start and end time of the interventions, the nutritional status of the mothers included, the type of intervention and the factors it addresses, and the measures collected in the offspring.

In terms of geographical region, it has been found that most interventions in recent years have been developed in North America, Oceania and Europe. Previous studies have highlighted the need for such interventions in developing countries where the prevalence of childhood obesity remains very high [1, 38].

Similarly, it has been noted that most interventions are targeted at the general population, which is necessary given the data on childhood obesity worldwide. However, there is a recognised need to develop interventions targeted at the most disadvantaged groups, be they racial/ethnic or economically disadvantaged groups, as this is where inequalities originate, increasing the prevalence of childhood obesity [39–43].

When it comes to the timing of interventions, the answer is clear: the earlier, the better [44]. However, it has been observed that most interventions start in the second trimester of pregnancy, some even after birth. Interventions should start from the first trimester, spanning the entire pregnancy, a period that has been linked to future obesity in offspring [45, 46]. Similarly, the time of termination of interventions has been positioned, in most of the interventions analysed, at the time of birth, with few interventions continuing into the postnatal period and only five, of the 51 included, ending at two years of age or more, covering the period of the first thousand days of life. It has been shown that this is the period of greatest plasticity and where the actions carried out will have the greatest effect on the future since once this period is over, plasticity decreases so that the actions carried out will not be as successful [45]. Moreover, the cumulative effect of many exposures over this period results in poor future health [35]. Therefore, the entire period of the first 1000 days should be covered to minimise exposures.

On the other hand, interventions should be targeted at all pregnant women, regardless of their weight at the start of pregnancy, as weight gain during this period is an early programming factor that will affect offspring [48–51]. This review found that many of the interventions include all pregnant women (44%), but there are still many interventions that focus only on those who are overweight (29%) or only on those who have obesity (12%) at the start of pregnancy.

In terms of the factors covered by current interventions, it has been found that almost all interventions target a single factor (GWG, GDM) or generally improve the Pregnancy outcomes or nutritional status of future offspring, but only cover nutrition, physical activity or a combination of both, while very few include factors such as breastfeeding promotion or sleep hygiene. Interventions should be comprehensive, targeting all possible programming factors, as there is often more than one factor in the same individual, which has been shown to have a cumulative effect on future obesity risk [44, 52, 53].

Finally, concerning measures collected in offspring, as a measure of the effectiveness of interventions, most of the studies analysed collect weight at birth, which is the end of the intervention. As mentioned above, interventions should be continued until two years, so the most effective measures would be BMI or skinfolds at this age. BMI is the most widely used. Although it has some limitations, skinfold measurement could overcome these limitations by allowing estimation of body fat percentage [54–56].

Compared previous systematic reviews of interventions in the first 1000 days of life [20, 21], they found efficacy for interventions that began during pregnancy and continued postnatally. Of these, those that only intervened in one aspect were effective. Those based on home visits and group sessions focusing on diet, infant feeding practices and physical activity were effective. On the other hand, reviews focusing on the content analysis [18, 22] found that most interventions were developed in the general population, without taking into account ethnic or racial minorities and focusing on a single factor, mainly diet and physical activity.

Limitations

The present study has some limitations that should be taken into account. Firstly, the quality of the included studies was not assessed, as the aim of the review was not to evaluate their effectiveness but to profile the interventions. Secondly, we did not include studies that only presented an intervention protocol, without any related publication on outcomes, to avoid biasing the profile obtained with protocols that were not carried out. This allows those interventions that are already being carried out do not appear in this review and should appear in future reviews.

Abbreviations

US: United States; WHO: World Health Organisation; GWG: Gestational weight gain; GDM: Gestational Diabetes mellitus; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analysis; UK: United Kingdom; BMI: Body mass index; PA: Physical Activity.

Conclusions

From the analyses presented here, it can be concluded that current interventions are not covering as many aspects as they should. Future research should be conducted more frequently in developing countries and target disadvantaged groups. In addition, these interventions should include all pregnant women, aiming to cover as many programming factors as possible and extending through the first 1000 days of life, with BMI or skinfolds as measures of effectiveness during this period. Ensuring that the interventions that are carried out cover all the aforementioned aspects will lead to better prevention of childhood obesity and an improvement in the epidemic that adult obesity entails.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

The authors received no funding for this study and have no relevant financial relationships to disclose.

Authors' contributions

CPM, JCB, BFM and MDR developed the search strategy. EOM and BRF performed the literature search, the screening and the data extraction. CPM and JCB ran the analyses and generated the Tables. CPM, BFM and MDR drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgments

Not applicable.

References

1. OMS. Obesity and Overweight. What are obesity and overweight? [Internet]. Vol. 2007, WHO Fact Sheet. 2020 [cited 2020 Dec 1]. p. 2. Available from: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
2. NCD-RisC NRFC. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet* [Internet]. 2017 Dec 16 [cited 2021 Apr 15];390(10113):2627–42. Available from: <https://pubmed.ncbi.nlm.nih.gov/29029897/>
3. Kumar S, Kelly AS. Review of Childhood Obesity: From Epidemiology, Etiology, and Comorbidities to Clinical Assessment and Treatment [Internet]. Vol. 92, *Mayo Clinic Proceedings*. Elsevier Ltd; 2017 [cited 2021 May 5]. p. 251–65. Available from: <http://dx.doi.org/10.1016/j.mayocp.2016.09.017>
4. Morales Camacho WJ, Molina Díaz JM, Plata Ortiz S, Plata Ortiz JE, Morales Camacho MA, Calderón BP. Childhood obesity: Aetiology, comorbidities, and treatment. *Diabetes/Metabolism Research and Reviews*. 2019 Nov 1;35(8).
5. Simmonds M, Burch J, Llewellyn A, Griffiths C, Yang H, Owen C, et al. The use of measures of obesity in childhood for predicting obesity and the development of obesity-related diseases in adulthood: a systematic review and meta-analysis. *Health Technol Assess (Rockv)* [Internet]. 2015 Jun [cited 2019 Feb 15];19(43):1–336. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26108433>
6. Yusefzadeh H, Rashidi A, Rahimi B. Economic burden of obesity: A systematic review. *Soc Heal Behav* [Internet]. 2019 [cited 2022 Feb 14];2(1):7. Available from: <https://www.shbonweb.com/article.asp?issn=2589-9767;year=2019;volume=2;issue=1;spage=7;epage=12;aulast=Yusefzadeh>
7. Stamatakis E, Wardle J, Cole TJ. Childhood obesity and overweight prevalence trends in England: Evidence for growing socioeconomic disparities. *Int J Obes* [Internet]. 2010 Jan [cited 2020 Sep 8];34(1):41–7. Available from: <https://pubmed.ncbi.nlm.nih.gov/19884892/>
8. Wang YC, Gortmaker SL, Taveras EM. Trends and racial/ethnic disparities in severe obesity among US children and adolescents, 1976-2006. *Int J Pediatr Obes* [Internet]. 2011 Feb [cited 2020 Sep 8];6(1):12–20. Available from: <https://pubmed.ncbi.nlm.nih.gov/20233157/>
9. González-Muniesa P, Martínez-González MA, Hu FB, Després JP, Matsuzawa Y, Loos RJJ, et al. Obesity. *Nat Rev Dis Prim* 2017 31 [Internet]. 2017 Jun 15 [cited 2022 Feb 14];3(1):1–18. Available from: <https://www.nature.com/articles/nrdp201734>
10. Hanson MA, Gluckman PD. Early developmental conditioning of later health and disease: physiology or pathophysiology? [Internet]. Vol. 94, *Physiological reviews*. *Physiol Rev*; 2014 [cited 2020 Oct 7]. p. 1027–76. Available from: <https://pubmed.ncbi.nlm.nih.gov/25287859/>
11. Moreno Villares JM. Los mil primeros días de vida y la prevención de la enfermedad en el adulto. *Nutr Hosp* [Internet]. 2016 Jul 12 [cited 2019 Feb 15];33(Suppl 4):337. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27571856>

37. UNICEF. MALNUTRICIÓN, OBESIDAD INFANTIL Y DERECHOS DE LA INFANCIA EN ESPAÑA [Internet]. 2019 Oct [cited 2021 Apr 15]. Available from: www.unicef.es
38. White J, Rehkopf D, Mortensen LH. Trends in Socioeconomic Inequalities in Body Mass Index, Underweight and Obesity among English Children, 2007–2008 to 2011–2012. Baker JL, editor. PLoS One [Internet]. 2016 Jan 26 [cited 2019 Jun 18];11(1):e0147614. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26812152>
39. Fundación Gasol. Estudio PASOS resultados finales [Internet]. 2019 [cited 2020 Dec 1]. Available from: <https://www.gasolfoundation.org/es/estudio-pasos/>
40. AESAN. ESTRATEGIA NAOS - Estudio ALADINO 2019. 2020.
41. El-Sayed AM, Scarborough P, Galea S. Socioeconomic inequalities in childhood obesity in the United Kingdom: A systematic review of the literature. *Obes Facts* [Internet]. 2012 Nov [cited 2020 Sep 27];5(5):671–92. Available from: <https://pubmed.ncbi.nlm.nih.gov/23108336/>
42. Gillman MW, Ludwig DS. How Early Should Obesity Prevention Start? *N Engl J Med* [Internet]. 2013 Dec 5 [cited 2020 Mar 17];369(23):2173–5. Available from: <http://www.nejm.org/doi/10.1056/NEJMp1310577>
43. Ravelli GP, Stein ZA, Susser MW. Obesity in Young Men after Famine Exposure in Utero and Early Infancy. *N Engl J Med* [Internet]. 1976 Aug 12 [cited 2020 Oct 5];295(7):349–53. Available from: <https://pubmed.ncbi.nlm.nih.gov/934222/>
44. Roseboom TJ, Painter RC, Van Abeelen AFM, Veenendaal MVE, De Rooij SR. Hungry in the womb: What are the consequences? Lessons from the Dutch famine [Internet]. Vol. 70, *Maturitas*. Maturitas; 2011 [cited 2020 Oct 5]. p. 141–5. Available from: <https://pubmed.ncbi.nlm.nih.gov/21802226/>
45. Gluckman PD, Hanson MA. Living with the past: evolution, development, and patterns of disease. *Science* [Internet]. 2004 Sep 17 [cited 2022 Feb 25];305(5691):1733–6. Available from: <https://pubmed.ncbi.nlm.nih.gov/15375258/>
46. Wrotniak BH, Shults J, Butts S, Stettler N. Gestational weight gain and risk of overweight in the offspring at age 7 y in a multicenter, multiethnic cohort study. *Am J Clin Nutr* [Internet]. 2008 Jun 1 [cited 2020 Oct 2];87(6):1818–24. Available from: <https://pubmed.ncbi.nlm.nih.gov/18541573/>
47. Samuelsson AM, Matthews PA, Argenton M, Christie MR, McConnell JM, Jansen EHJM, et al. Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity, hypertension, and insulin resistance: A novel murine model of developmental programming. *Hypertension* [Internet]. 2008 Feb [cited 2020 Sep 27];51(2):383–92. Available from: <https://pubmed.ncbi.nlm.nih.gov/18086952/>
48. Oken E, Taveras EM, Kleinman KP, Rich-Edwards JW, Gillman MW. Gestational weight gain and child adiposity at age 3 years. *Am J Obstet Gynecol* [Internet]. 2007 [cited 2021 Apr 13];196(4):322.e1-322.e8. Available from: <https://pubmed.ncbi.nlm.nih.gov/17403405/>
49. Oken E, Rifas-Shiman SL, Field AE, Frazier AL, Gillman MW. Maternal gestational weight gain and offspring weight in adolescence. *Obstet Gynecol* [Internet]. 2008 Nov [cited 2020 Oct 2];112(5):999–1006. Available from: <https://pubmed.ncbi.nlm.nih.gov/18978098/>
50. Iguacel I, Escartín L, Fernández-Alvira JM, Iglesia I, Labayen I, Moreno LA, et al. Early life risk factors and their cumulative effects as predictors of overweight in Spanish children. *Int J Public Health*. 2018;63(4):501–12.
51. Robinson SM, Crozier SR, Harvey NC, Barton BD, Law CM, Godfrey KM, et al. Modifiable early-life risk factors for childhood adiposity and overweight: An analysis of their combined impact and potential for prevention. *Am J Clin Nutr*. 2015;101(2):368–75.
52. Chung S. Body mass index and body composition scaling to height in children and adolescent. *Ann Pediatr Endocrinol Metab* [Internet]. 2015 [cited 2022 Feb 25];20(3):125. Available from: [/pmc/articles/PMC4623339/](http://pmc/articles/PMC4623339/)
53. Ros Arnal I, Herrero Álvarez M, Castell Miñana M, López Ruzafa E, Galera Martínez R, Moráis López A, et al. Valoración sistematizada del estado nutricional. *Acta Pediátrica Española* [Internet]. 2011 [cited 2022 Feb 25];69(4):165–72. Available from: <https://www.actapediatrica.com/index.php/secciones/nutricion-infantil/19-valoracion-sistematizada-del-estado-nutricional#.YhjGa-jP3IU>
54. Díaz-Rodríguez M, Pérez-Muñoz C, Carretero-Bravo J, Ruíz-Ruíz C, Serrano-Santamaría M, Ferriz-Mas BC. Early Risk Factors for Obesity in the First 1000 Days-Relationship with Body Fat and BMI at 2 Years. *Int J Environ Res Public Health* [Internet]. 2021 Aug 1 [cited 2022 Apr 22];18(15). Available from: <https://pubmed.ncbi.nlm.nih.gov/34360471/>

Figures




Figure 1

PRISMA flow diagram for identifying and screening eligible childhood obesity prevention interventions.




Figure 2

Distribution of the type of intervention according to the targeted issue of the interventions developed.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.docx](#)
- [Additionalfile2.docx](#)