

Magnitude of metabolic syndrome and its associated factors among patients with type 2 diabetes mellitus in Ayder comprehensive specialized hospital, Tigray, Ethiopia, 2019: a cross sectional study

Gebreamlak Gebremedhn Gebremeskel (✉ gmost1221@gmail.com)

Kalayou Kidanu Berhe

Mekele University

Desta Siyoum Belay

Mekele University

Berihu Hailu Kidanu

Aksum University

Assefa Iyasu Negash

Aksum University

Kfle Tekulu Gebreslasse

Aksum University

Degenah Bahrey Tadesse

Aksum University

Mulugeta Molla Birhanu

St. paulos medical college

Research note

Keywords: Ethiopia, Associated Factors, Magnitude, Metabolic syndrome

Posted Date: September 19th, 2019

DOI: <https://doi.org/10.21203/rs.2.10927/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 18th, 2019. See the published version at <https://doi.org/10.1186/s13104-019-4609-1>.

Abstract

Abstract Objective : The objective of this study was to assess magnitude of metabolic syndrome and its associated factors among type 2 diabetes mellitus patients in Ayder comprehensive specialized hospital. A hospital based cross sectional study design was used. Binary logistic regression model was used.

Result : A total of 419 respondents (208 males and 211 females) were enrolled; the mean age was 56.39 (SD 10.18), 51.1% of the respondents had metabolic syndrome according to international diabetes federation. Sex and age were statistically associated with metabolic syndrome with [AOR (95% CI) = 1.93(1.057, 3.533) and 1.04 (1.012, 1.072)] respectively. Regular physical exercise, overweight and obesity were statistically associated with metabolic syndrome with [AOR (95% CI) =1.84 (1.002, 3.362), 2.68 (1.518, 4.747) and 3.55 (1.254, 10.074)] respectively. To conclude, Magnitude of metabolic syndrome was high. The associated factors for metabolic syndrome are physical inactivity, inadequate intake of fruits, family history, overweight, and obesity. **Key words :** Ethiopia, Associated Factors, Magnitude, Metabolic syndrome

Introduction

Metabolic syndrome was defined by, WHO, NCEP ATP III and IDF etc. National cholesterol education program's adult treatment panel III (NCEPATP III) defines metabolic syndrome as having three or more of the five risk factors: raised fasting plasma glucose, high blood pressure, abdominal obesity, low HDL, and/or raised plasma triglyceride (1). The world health organization (WHO) defines metabolic syndrome as insulin resistance [impaired glucose tolerance (IGT) or diabetes] to be present for the diagnosis to be made. In addition to insulin resistance, at least two other components must also be present: obesity, hypertension, dyslipidemia and/or micro albuminuria (2).

Because of the study patients are type 2 DM, we select IDF criteria to define MetS. This is particularly evident for the risk of type 2 DM, which is apparent at much lower levels of obesity in Asians compared to Europeans. The IDF, having recognized the difficulties in identifying unified criteria for MetS that were applicable across all ethnicities, has proposed a new set of criteria with ethnic/racial specific cut-offs and IDF states that because of the limited data in Sub-Saharan Africa including Ethiopia, they should use European criteria (3).

According to the new international diabetes federation (IDF) definition, metabolic syndrome is defined as central obesity (defined by waist circumference) **plus** any two of the four risk factors: raised triglyceride, low high density lipoprotein (HDL) cholesterol, raised blood pressure and/or raised fasting plasma glucose level. Metabolic syndrome is a complex disorder which leads type 2 diabetes patients and cardiovascular disease (CVD) to be a twin global epidemic problem (4).

About 70-80% of diabetes mellitus (DM)(6)and 20-25% of adult population in the world is estimated to have metabolic syndrome and they are two times as likely to die from and three-fold as likely to have a heart attack or stroke as compared to without the syndrome. Besides, metabolic syndrome has five times

increased risk of acquiring type 2 diabetes mellitus(5). Consumption of calorie-dense foods, sedentary lifestyle, and tobacco consumption are considered as potential risk factors for MetS(7). In Ethiopia, the prevalence of metabolic syndrome among type 2 diabetes mellitus according to NCEP on (ATP) III was 45.9% (10).

There is limited data on metabolic syndrome in Ethiopia and in the horn of Africa at large. Therefore, this study is helpful to the development of life style modification, control and management of the metabolic syndrome. Finally, this will bring non-communicable disease in general and metabolic syndrome in particular in to the clinical area, and research agenda.

Methods And Patients

A hospital based cross section study design was conducted in Ayder comprehensive specialized hospital (ACSH) from February to June 2018. ACSH is found in Mekelle the capital city of [Tigray region](#) northern of [Ethiopia](#). Mekelle is located around 780 kilometers north of the Ethiopian capital city [Addis Ababa](#). The study participants were previously diagnosed/known type 2 DM patients. DM clinic is served for about 500 type 2 DM patients per month that is estimated from the average of the last six months report.

Study design: A hospital based cross section study design.

Source patients: All type 2 diabetes patients attending DM clinic at ACSH.

Study patients: All volunteer type 2 DM patients attending DM clinic and available at ACSH during the data collection period.

Study subjects: All sampled subjects (419) participated in this study

Inclusion criteria

- All type 2 diabetes mellitus patients

Exclusion criteria

- Critically ill patient
- Pregnant mother (difficult to measure BMI and waist circumference)

Sample size determination

The sample size was calculated by using single proportion formula: (See Formula in Supplemental Files)

Considering 10% non-response rate, because of some respondents may withdraw in between the interview, the total estimated study subjects were 419 and selected by systematic random sampling technique with k^{th} interval of 2 and every second (2) was participated by random selection from 1- 2(k^{th}).

Data collection procedure

Data collection was carried out from February to March 2018. To collect data a standardized, interviewer administered questionnaire and document review checklist of physical measurements, residence (urban or rural) for the last six month, co-morbid disease like hypertension, CHF, stroke were used and contains three parts. Thus are socio-demographic and medical history, the life style and related information.

Measurements and tools

Anthropometric measurements: Anthropometric measurements of weight and height was measured using Seca weighing scale and stadiometer respectively and participants were wearing light clothing (single and thin) and without shoes to the nearest Kg and CM respectively. A simple flexible steel metric tape calibrated in meters was used for measuring waist circumference. Waist circumference was measured midway between the iliac crest and the lower rib margin in the horizontal plane while the participant is standing to the nearest 0.5 centimeter.

Blood pressure: Two blood pressure measurements taken after five minutes apart were determined for each participant using standard adult digital blood pressure apparatus with the correct size arm cuff. Participants was measured after 5 minutes of rest in sitting position, arm should be rest on table at heart level, back supported, on the same arm and legs rest on ground (no crossed). And the average readings of the two measurements was recorded in questionnaire (11).

Data quality assurance

Questionnaire was prepared in English and translated in to local language Tigrigna by expert in language. Pretest was done on 5% of the subjects in Mekelle hospital two weeks before the actual data collection. Because Mekelle hospital is near to ACSH and have similar socio-demographic characteristics with patients in ACSH. Two BSC nurses and one supervisor were recruited for data collection and training was given. The collected data was checked by supervisor and principal investigator daily.

Data processing and analysis

After the data collection, the data was entered in to Epi-info version 7 and exported to SPSS version 23 statistical program. Descriptive characteristics were presented in text, frequency percentage tables, and graphs. Binary logistic regression model was used to see the association between the outcome variable with each independent variables. Then those variables with P-value <0.25 at bivariate analysis was included in the multivariable analysis. Finally P-value < 0.05 is describe as a statistically associated with 95% confidence level. Model fitness was checked by using Hosmer and Lemeshow goodness fit model.

Operational definition

Metabolic syndrome: As per the definition of international diabetes federation it is defined as having central obesity (defined by waist circumference \geq 94 cm for male and \geq 80 cm for female) plus any two

of the following four factors: raised triglycerides, reduced HDL Cholesterol, raised blood Pressure, and/or raised fasting plasma glucose.

Central obesity: defined by waist circumference ≥ 94 cm for male and ≥ 80 cm for female putting to the nearest centimeter.

Raised triglycerides: ≥ 150 mg/dL (1.7 mmol/L) or specific treatment for this lipid abnormality.

Reduced HDL Cholesterol: < 40 mg/dL (1.03 mmol/L) in male < 50 mg/dL (1.29 mmol/L) in female or specific treatment for this lipid abnormality.

Raised blood Pressure: Is a systolic BP ≥ 130 mm Hg or diastolic BP ≥ 85 mm Hg, or any patient on treatment of previously diagnosed hypertension..

Reduce salt intake: about 1 tea spoon of table salt per meal

Results

Socio-demographic characteristics

A total of 419 type 2 DM patients were enrolled in this study. Binary logistic regression was used. Of these respondents 211 (50.4%) were females. The mean age of the respondents was 56.39 SD ± 10.18 with minimum of 35 and maximum of 90 years old. Of the total respondents 79.7% were married, 83.1% were orthodox Christian followers (**Table 1**).

Metabolic syndrome and its component

According to IDF criteria 51.1% of the respondents had metabolic syndrome, with higher prevalence in females when compared to males (57.5% and 42.5% respectively). Central obesity (59.7%) was highly prevalent component of metabolic syndrome followed by elevated triglyceride (45.1%) and then hypertension and low HDL-c 41.3% and 34.4% respectively. Females were also having higher proportion of central obesity and reduced HDL than males (**figure 1**).

Factors associated to metabolic syndrome

In the multivariable analysis, age ([AOR (95% CI) = 1.04 (1.01, 1.07)], sex ([AOR (95% CI) = 1.94 (1.06, 3.54)], eating fruits once, twice, and four and above times in a typical week ([AOR (95% CI) = 0.41 (0.21, 0.78), 0.35 (0.16, 0.76), and 0.24 (0.07, 0.75)], regular physical exercise (AOR 1.83 (1.05, 3.36)], family history of DM ([AOR (95% CI) = 0.54 (0.29, 0.99)], chronic disease co-morbid (AOR 0.42 (0.25, 0.70)) Overweight ([AOR (95% CI) = 2.66 (1.50, 4.69)) and obesity (AOR 3.50 (1.23, 9.91) shows statistical association with metabolic syndrome (**Table 2**)

Discussion

This study was aimed to assess magnitude of metabolic syndrome and its associated factors among type 2 DM patients who have follow up in ACSH.

The magnitude of metabolic syndrome among type 2 DM patients in this study was 51.1%; which lies within a range of 12% - 86% from a review study done in sub Saharan Africa(12). But, this study is higher than from a study conducted in Ethiopia (45.9%), in 10 European countries (24%) and UK (32%)(10, 13). This difference is due to difference in study setting, sample size, and the criteria used to define metabolic syndrome. The magnitude of MetS in this study is lower than from studies done in Gahanna (68.6%), Nigeria (68.7%) and Iran (64.9%)(14-16). This variation could be due to difference in socio-cultural, study setting, study design and life style.

This study reveals that, age was significantly associated with metabolic syndrome, and this is in line with a study conducted in Iran(14). This association might be due to older age participants were physically inactive and adoption of unhealthy life styles. Sex was another variable that shows statistical association with metabolic syndrome. This is consistent with studies in Hawassa Ethiopia, Addis Ababa Ethiopia, Nigeria, and Iran (10, 14, 15, 17).

Respondents who eat fruit twice, four and above times in a typical week were 65% and 76% protected from metabolic syndrome respectively as compared to those who never eat fruit in a typical week. This is supported by EPHA (Ethiopia) and in line with study conducted in brazil(18, 19).

Respondents who did not do regular physical exercise were 1.83 odds to have MetS as compared to those who did regular physical exercise. This is supported by EPHA report (Ethiopia) and in line with studies done in US, Canada, and Sub-Saharan Africa (18, 20-22).

In addition, the odds of MetS were 2.66 and 3.5 times higher among overweight and obesity compared to those normal weigh respondents respectively among type 2 DM patients. This is in line with studies done in Ethiopia, Gahanna, Iran and Nigeria (10, 14, 15, 23). But lower in AOR from a study in Ethiopia. This may be due to sample size and difference in reference category.

Conclusion And Recomendations

Conclusion

The magnitude of metabolic syndrome among type 2 diabetes mellitus is 51.5%. The associated factors for metabolic syndrome are physical inactivity, advanced age, inadequate intake of fruits, family history, overweight and obesity.

Recommendations

To Ayder comprehensive specialized hospital

- Regular screening of type 2 DM patients in ACSH for components of MetS is vital in order to avert/limit the risks before developing cardiovascular related morbidity and mortality.

To type 2 DM in ACSH

- Should do regular physical exercise as recommended
- Should reduce and manage their weight if they are told to do so by health care professionals

To further study

- Further cohort or interventional studies should be done to address other predictors of MetS.

Limitations

Limitation of the study

- The cross sectional nature of the study; temporal relationship between exposure and disease cannot be clearly determined or not powerful
- The study was conducted only in a single public hospital

Declarations

Ethical approval

Ethical approval was obtained from the institutional review board of Mekelle University College of health science on 13 February 2018. Then letter of permission was received from medical director of ACSH and given to head department in the DM clinic. All of the study participants was informed about the purpose of the study, about their right to participate or to with draw at any time if they don't want and their confidentiality. Verbal consent to participate was taken because there are participants who cannot read and approved by the IRB. This research is original and not considered in another journal for publication.

Consent to publish

Not applicable

Availability of data and materials

I the undersigned, declare that this is our original work and has not been submitted and considered for publish in any journal and all sources of materials and data used for this research have been secured and acknowledged.

Competing interests

Authors declared that, they have no conflict of interest and approved for publication.

Funding

No funding was received for this research study. Because of we are from low income countries, we politely asked you to waive us.

Authors' contributions

All authors' listed in this research article have been involved and contribute for this thesis. **'GG'** contributes in conception, design, analysis, data interpretation, data acquisition, drafting and writing the final manuscript **'MM'** contributes in conception, design, analysis, data acquisition, drafting and revising the manuscript, **'KK'** contributes in design, analysis, data interpretation, drafting and revising, **'DS', 'BH', 'AI'** and **'KT'** contributes in design, analysis, drafting and revising, and **'DB'** contributes in design, analysis, data acquisition, drafting and revising it critically for important intellectual content and all authors read and given final approval of the manuscript to be published.

Acknowledgment

We would like to express our thanks to Mekelle University Department of Nursing and Aksum University Department of Nursing. Then, we would like to extend our appreciation to supervisor, data collectors and study participants for their cooperation, participation and willingness.

Abbreviations

ACSH: Ayder Comprehensive Specialized Hospital, **DM:** Diabetes Mellitus, **EPHA:** Ethiopian Public Health Association, **HDL:** High Density Lipoprotein, **IDF:** International Diabetes Federation, **MetS:** Metabolic Syndrome, **NCEP/ATP:** National Cholesterol Education Adult Treatment Panel, **US:** United States

References

1. Expert Panel on Detection E. Executive summary of the Third Report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). *Jama*. 2001;285(19):2486.
2. WHO. Definition, diagnosis and classification of diabetes mellitus and its complications: report of a WHO consultation. Part 1. 1999.
3. Alberti K, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA. Harmonizing the metabolic syndrome. *Circulation*. 2009;120(16):1640-5.
4. Alberti KG, Zimmet P, Shaw J, Grundy SM. The IDF consensus worldwide denition of the metabolic syndrome. 2006:4-10.
5. Alberti, George K, Mayer M, Paul Z, Jonathan S. Metabolic syndrome—a new world-wide definition. A consensus statement from the international diabetes federation. *Diabetic medicine*. 2006;23(5):469-80.

6. Marchesini G, Forlani G, Cerrelli F, Manini R, Natale S, Etal. WHO and ATPIII proposals for the definition of the metabolic syndrome in patients with type 2 diabetes. *Diabet Med.* 2004;21:383-7.
7. Young F, Critchley JA, Johnstone LK, Unwin NC. "A review of co-morbidity between infectious and chronic disease in Sub Saharan Africa: TB and diabetes mellitus, HIV and metabolic syndrome, and the impact of globalization." *Global Health.* 2009;5(9).
8. WHO. Global report on diabetes geneva,2016.
9. Jeon CY, Murray MB. Diabetes mellitus increases the risk of active tuberculosis: a systematic review of 13 observational studies. *PLoS medicine.* 2008;5(7):e152.
10. Agete Tadewos H, Demissie Assegu. Pattern of Metabolic Syndrome in Relation to Gender among type-II DM Patients in Hawassa University Comprehensive Specialized Hospital, Hawassa, Southern Ethiopia. . *Health Sci J.* 2017;11.
11. JNC. PL Detail-Document, Treatment of Hypertension: JNC 8 and More. 2014.
12. Okafor CI. The metabolic syndrome in Africa: Current trends. *Indian Journal of Endocrinology and Metabolism.* 2012;16(1):56-66.
13. Scuteri A, Laurent, S., Cucca, F., Cockcroft, J., Cunha, P. G., Mañas, L. R., Nilsson, P. M. Metabolic syndrome across Europe: different clusters of risk factors. *European Journal of Preventive Cardiology.* 2015;22(4):486-1.
14. Foroozanfar Z, Najafipour H, Khanjani N, Bahrampour A, Ebrahimi H. The prevalence of metabolic syndrome according to different criteria and its associated factors in type 2 diabetic patients in Kerman, Iran. *Iranian journal of medical sciences.* 2015;40(6):522.
15. Soyoye D, Adebayo O, Kolawole B, Ikem R. Prevalence and associated risks for metabolic syndrome in Nigerians with type 2 diabetes mellitus. 2013.
16. Abagre TA. Metabolic Syndrome among People with Type 2 Diabetes Mellitus in Two Selected Hospitals in the Brong Ahafo Region. 2016.
17. Tran A, Gelaye B, Girma B, Lemma S, Berhane Y, Bekele T, et al. Prevalence of metabolic syndrome among working adults in Ethiopia. *International journal of hypertension.* 2011;2011.
18. MOH. The Federal Democratic Republic of Ethiopia Ministry of Health: health sector transformation plan. 2015/2016.
19. Oliviera EPd, McLellan kCP, Silveira IVdA, Burini RC. Dietary factors associated with metabolic syndrome in Brazilian adults. *Nutrition.* 2012;11(3).
20. Ford ES, Kohl HW, Mokdad AH, Ajani UA. Sedentary behavior, physical activity, and the metabolic syndrome among US adults. *Obesity.* 2005;13(3):608-14.
21. Ross R, Dagnone D, Jones PJ, Smith H, Paddags A, Hudson R, et al. Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in menA randomized, controlled trial. *Annals of internal medicine.* 2000;133(2):92-103.
22. Assah FK, Ekelund U, Brage S, Mbanya JC, Wareham NJ. Urbanization, physical activity, and metabolic health in sub-Saharan Africa. *Diabetes Care.* 2011;34(2):491-6.

23. Nsiah K, Shang V, Boateng K, Mensah F. "Prevalence of metabolic syndrome in type 2 diabetes mellitus patients,". International Journal of Applied and Basic Medical Research. 2015;5:133.

Tables

Table 1; Frequency distribution of socio-demographic characteristics of respondents among type 2 diabetes mellitus patients in ACSH, Tigray, Ethiopia (n = 419)

Variable	Category	No.	Percent
Sex	Female	211	50.4%
	Male	208	49.6%
Ethnicity of participants	Tigray	395	94.3%
	Amara	24	5.7%
Marital status	Single	19	4.5%
	Married	334	79.7%
	Divorced	24	5.7%
	Widowed	42	10%
Religion	Orthodox	348	83.1%
	Muslim	65	15.5%
	Protestant	6	1.4%
Occupation of participant	Farmer	50	11.9%
	Merchant	110	26.3%
	G. Employed	115	27.4%
	Daily laborer	25	6%
	House wife	78	18.6%
	P. Employed	41	9.8%
Educational status	Unable to read and write	131	31.3%
	Primary school	118	28.2%
	High school	69	16.5%
	College and above	101	24.1%
Place of residence	Urban	361	86.2%
	Rural	58	13.8%

N.B: **G. employed:** - government employed, **P. employed:** - private employed

Table 2: Multivariable analysis of the independent variables with metabolic syndrome among patients with type 2 diabetes mellitus in ACSH, Mekelle, Tigray, Ethiopia (n= 419)

Variables	Metabolic syndrome		Crude OR		Adjusted OR	
	No	Yes	P-value	95% CI	P-value	95% CI
Sex						
Male	117	91		1		1
Female	88	123	0.003	1.79 (1.22, 2.64)	0.031	1.94(1.06, 3.54)*
Age			0.00	1.05 (1.03, 1.08)	0.005	1.04 (1.01, 1.07)*
Ethnicity of participant						
Tigray (Ref)	187	208		1		1
Amara	18	6	0.012	0.3 (0.11, 0.77)	0.1357	0.40 (0.11, 1.42)
Marital status			0.068		0.549	
Single	14	5		1		1
Married	164	170	0.045	2.9 (1.02, 8.24)	0.287	2.01 (0.55, 7.32)
Divorced	12	12	0.12	2.8 (0.76, 10.24)	0.509	1.73 (0.34, 8.79)
Widowed	15	17	0.008	5.04 (1.52, 16.7)	0.157	2.97 (0.65, 13.50)
Occupation			0.002		0.322	
Farmer	37	13		1		1
Merchant	52	58	0.02	3.17 (1.52, 6.62)	0.062	3.36 (0.94, 12.01)
Government employee	57	58	0.04	2.89 (1.39, 6.01)	0.039	3.88 (1.07, 14.06)
Daily laborer	14	11	0.119	2.24 (0.82, 6.15)	0.294	2.25 (0.49, 10.24)
House wife	33	45	0.001	3.88 (1.78, 8.43)	0.073	3.3 (0.89, 12.22)
Private employee	12	29	0.00	6.87 (2.73, 17.31)	0.027	4.78 (1.19, 19.11)
Residence of participant						
Urban	163	198		1		1
Rural	42	16	0.00	0.31 (0.17, 0.57)	0.817	1.14 (0.37, 3.45)

Variables	Metabolic syndrome		Crude OR		Adjusted OR	
	No	Yes	P-value	95% CI	P-value	95% CI
Physical exercise						
Yes	60	34		1		1
No	145	180	0.001	2.19 (1.36, 3.52)	0.048	1.83 (1.05, 3.36)*
Eat fruits in a typical week			0.221		0.027	
None per week	54	78		1		1
One times a week	83	77	0.062	0.64 (0.4, 1.02)	0.007	0.41 (0.21, 0.78)*
Twice a week	42	36	0.07	0.59 (0.34, 1.04)	0.008	0.35 (0.16, 0.76)*
Three times a week	11	7	0.111	0.44 (0.16, 1.20)	0.088	0.31(0.08, 1.18)
Four and above a week	15	16	0.449	0.74 (0.36, 1.62)	0.014	0.24 (0.07, 0.75)*
Eat vegetables in a typical week			0.028		0.068	
None per week						
One times a week	17	22		1		1
Twice a week	37	21	0.051	0.43 (0.19, 1.05)	0.122	0.42 (0.14, 1.25)
Three times a week	69	58	0.242	0.65 (0.31, 1.33)	0.345	0.62 (0.23, 1.66)
Four and above a week	28	43	0.672	1.18 (0.53, 2.62)	0.650	1.27 (0.44, 3.65)
	54	70	0.996	1.02 (0.48, 2.07)	0.718	1.21 (0.43, 3.40)
Any alcoholic drinks						
Yes	31	23		1		1
No	174	191	0.183	1.48 (0.83, 2.63)	0.611	1.23 (0.54, 2.78)
Ever smoke any tobacco product ever						
Yes						
No	2	6		1		1
	203	208	0.191	0.34 (0.68, 1.72)	0.092	0.17 (0.02, 1.33)
Low salt intake						
Yes	148	174		1		1
No	57	40	0.028	0.59 (0.37, 0.94)	0.344	.074 (0.41, 1.36)

Variables	Metabolic syndrome		Crude OR		Adjusted OR	
	No	Yes	P-value	95% CI	P-value	95% CI
Body mass index			0.000		0.001	
Normal weight	147	108		1		1
underweight	15	1	0.021	0.91 (0.01, 0.69)	0.089	0.14 (0.01, 1.34)
Overweight	37	76	0.000	2.79 (1.76, 4.45)	0.001	2.66 (1.50, 4.69)*
Obese	6	29	0.000	6.58 (2.64, 16.4)	0.018	3.50 (1.23, 9.91)*
Duration of DM			0.000		0.038	
< 5 years duration	134	84		1		1
6-10 years duration	46	63	0.001	2.18 (1.37, 3.48)	0.334	1.35 (0.73, 2.50)
>10 years duration	25	67	0.000	4.27 (2.5, 7.29)	0.010	2.49 (1.23, 5.01)*
Treatment			0.012		0.073	
Metformin	70	57		1		1
Glibenclamide	13	4	0.104	0.38 (0.12, 1.22)	0.156	0.35 (0.08, 1.47)
Metformin plus Glibenclamide	67	70	0.313	1.28 (0.79, 2.08)	0.604	0.84 (0.45, 1.58)
Insulin plus oral hypoglycemic agents	55	83	0.013	1.85 (1.14, 3.02)	0.148	1.62 (0.84, 3.12)
Family hx of DM						
Yes	33	74		1		1
No	172	140	0.000	.36 (0.228, 0.58)	0.047	0.54 (0.29, 0.99)*
Chronic disease						
Yes	70	123		1		1
No	135	91	0.000	0.38 (0.26, 0.57)	0.001	0.42 (0.25, 0.70)*

N.B; * states significant association, and 1; reference category

Figures




Figure 1

Frequency of component of MetS in relation to sex of respondents among type 2 patients with DM in ACSH, Tigray, Ethiopia, 2019 (n=419)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Formula.jpg](#)