

Association between war-related traumatic events and blood pressure trajectory: A population-based study among Palestinian adults living in Gaza

Majed Jebril

Global Health Institute, School of Public Health, Xi'an Jiaotong University Health Science Center, Xi'an, Shaanxi, China

Mohsen Mazidi

Department of Twin Research and Genetic Epidemiology, King's College London, London, UK and Medical Research Council Population Health Research Unit, University of Oxford, Oxford, UK

Xin Liu

Global Health Institute, School of Public Health, Xi'an Jiaotong University Health Science Center, Xi'an, Shaanxi, China

Mi Baibing

Global Health Institute, School of Public Health, Xi'an Jiaotong University Health Science Center, Xi'an, Shaanxi, China

Heba Arafat

Department of Laboratory Medicine, Faculty of Applied Medical Science, Al Azhar University

Zumin Shi

Department of Human Nutrition, College of Health Sciences, QU Health, Qatar University, Doha, Qatar

Youfa Wang (✉ youfawang@gmail.com)

Global Health Institute, School of Public Health, Xi'an Jiaotong University Health Science Center, Xi'an, Shaanxi, China

Research Article

Keywords: Blood pressure, Palestine, Traumatic event, War event

Posted Date: July 21st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1865021/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Little is known regarding health status in environment characterized by instability and ongoing war risks. This study investigated hypertension disease burden and associations of war-related traumatic events with blood pressure (BP) trajectory over time amongst Palestinian adults in Gaza Strip.

Methods: From nine primary healthcare centers, medical records between 2013 and 2019 were collected for 1000 Palestinian adults living in Gaza. Multinomial logistic regression analysis examined associations between war-related traumatic events and BP trajectories derived using latent class trajectory analysis (LCTA).

Results: The prevalence of self-reported injury (of participants or their family members), death of a family member, and violence due to house bombing was 51.4%, 54.1%, and 66.5%, respectively. In total, 22.4% and 21.4% participants had constant-very-high (CVH) SBP (>160 mmHg) and DBP (>95 mmHg), and normal-stable SBP and DBP was found only 54.9% and 52.6%, respectively. Injury (participants or family members), death of a family member, and violence due to house bombing during wars were associated with CVH SBP with odds ratios (95%CI) of 1.79 (1.28-2.48), 1.90 (1.36-2.65), and 1.44 (1.01-2.05), respectively. The corresponding figures were 1.92 (1.36-2.71), 1.90 (1.35-2.68), and 1.62 (1.13-2.38) for CVH DBP. Living in debt was positively associated with CVH SBP, OR (95%CI) of 2.49 (1.73-3.60) and CVH DBP, OR (95%CI) of 2.37 (1.63-3.45).

Conclusions: The disease burden related to war-related traumatic events is high, and positively related to adverse BP trajectory among Palestinians living in Gaza. Intervention programs are needed to manage and prevent chronic diseases for this vulnerable population.

Funding: This research was funded in part by the US-based China Medical Board (CMB, grant number 16-262); National Natural Science Foundation of China, grant number 82173504, 82011530197; and the Chinese National Key Research and Development Program (grant number 2017YFC0907200, 2017YFC0907201).

Introduction

Many people are exposed to political violence, conflict, war and war-related events worldwide (1). Gaza, Palestine is a unique place in the world, where people have been exposed to four fierce wars over the past thirteen years (2). Those wars left 4160 martyrs and more than 18,300 injured, in which most of the victims were civilians, and many of their camps and buildings were destroyed (3). Often these populations also suffer from high prevalence of non-communicable diseases (NCDs) and lack of related medical services (4).

Exposure to war-related traumatic events may increase the risk for developing hypertension (5), which became one of the significant health concerns in Gaza (6). Those stressful experiences in war life may increase circulating catecholamine and cortisol levels and blood pressure (BP) over time (7-9).

Several studies have revealed that frequent exposure to stressful life events is a risk factor for developing post-traumatic stress disorder (PTSD) and hypertension (10-12). A meta-analysis of six cohort studies suggested that chronic exposure to stress may influence increased BP; where individuals who had stronger responses to stressor events were 21% more likely to develop BP increase than those with less intense responses (13).

Besides the actual BP level, the overtime changes of BP (i.e., BP trajectory) could be used to assess the risk of cardiovascular diseases (CVDs) (14). In Palestine, no previous study has examined the effect of traumatic life events on BP trajectories, which must be explored in the light of experienced living circumstances for a long time. Therefore, monitoring BP trajectories is indispensable for CVDs prevention (15, 16).

This study aimed to investigate the burden of hypertension (as an indicator of NCDs) and the association between war-related traumatic events (injury, death, serious illness, loss of job, business bankruptcy, exposure to violence due to house bombing, living in debt, marital separation, or divorce; and exposure to major disasters during wars) with BP trajectory over time in Palestinian adults living in Gaza Strip. We hypothesized that these war-related life events affect their health and BP trajectory.

Methods And Materials

Study design and participants

This study used a mixed-method study design and was based on a survey conducted in 2019 and historical longitudinal data extracted (for 2013-2019) among residents regularly visiting nine primary healthcare centers (PHCs) of the Ministry of Health (MoH) in Gaza Strip, Palestine. The participants who met the following criteria were enrolled: (1) Palestinian adult males and females (refugees or citizens), who have lived in Gaza Strip for ≥ 14 years; and (2) Registered at PHCs and have performed multiple anthropometric and biochemical measurement data during 2013 to 2019. Pregnant women and those having missing data were excluded from the study. A total of 1120 potential residents were invited to participate in our survey. We excluded those without anthropometric data ($n=83$) and pregnant ($n=37$). Ultimately, 1000 participants were included in the analysis (17).

The Xi'an Jiaotong University Health Science Center Ethical Committee approved this research. It was also approved by the Helsinki Committee affiliated to the Palestinian Health Research Council (PHRC) in Gaza and the Research Department at the Department of Human Resources Development, Ministry of Health, Gaza (PHRC/HC/576/19). All participants provided written informed consent.

Data collection

We selected the largest nine clinics out of 51 PHCs located in Gaza governorates to enhance the representativeness. One to three PHCs were chosen from each governorate, matching each governorate's population size. The data were collected in two phases: 1) The sociodemographic and war-related traumatic

events data were collected by face-to-face questionnaire survey administered by trained health workers. 2) The anthropometric measurements, including weight, height, waist circumference, systolic and diastolic BP, in addition to biochemical parameters, were extracted (from 2013 to 2019) from the electronic health record (E-Health) system by matching the participant's registration ID (17).

Questions on the war-related traumatic events were adopted from the Traumatic Life Events Questionnaire (TLEQ) (18). The questions were chosen based on possible war-traumatic life events to be exposed to and are also common among the Palestinian people. All of these events were caused by the three recent wars in Gaza between 2008 and 2020, including nine binary items (yes, or no): (1) Had injury of any family members; (2) Death of any family member; (3) Had serious illness of any family member based on the affliction with any critical diagnosed diseases including cancer, heart attack, stroke, kidney failure, and liver cirrhosis; (4) Had a loss of a job; (5) Had business bankrupt; (6) Exposure to violence due to house bombing; (7) Living in debt; (8) Had marital separation or divorce; and (9) Had major disasters that families have been exposed to during wars and caused a real tragedy for them to lose their livelihood by bombing their private properties, i.e., destroying factories, crops, shops, and vehicles.

BP was measured through a calibrated sphygmomanometer after a quiet rest time of 3-5 minutes, an average measurement of 2-3 BPs, use of an adequately sized cuff placed on a bare arm, and proper patient positioning, including back and heart-level arm support. Each year's average BP measurements were extracted for each participant.

Definitions of hypertension and trajectories of BP

Hypertension was defined as having a documented diagnosis by general practitioners in the PHCs, with measurements of SBP ≥ 140 mmHg and/or DBP ≥ 90 mmHg, based on WHO criteria (19).

This study identified BP trajectories using latent class trajectory analysis (LCTA), a method used to identify unobserved trajectory classes in epidemiological data (20). LCTA has the advantage of identifying distinct groups with similar underlying trajectories (21-23). These trajectories can vary in functional form across different order polynomials, allowing the best-fitting polynomial form to be specified for each trajectory separately, including the coefficients' polynomial order. The Bayesian Information Criterion (BIC) was calculated to evaluate the number of distinct trajectories and choose the best fit model (24).

Statistical analysis

We conducted descriptive- and in-depth modeling analyses. Continuous variables were summarized as mean values and standard deviations, while categorical variables were described as frequency and percentage. To compare categorical data and continuous variables, respectively, chi-square and t-tests were used. One-way variance analysis (ANOVA) was used to compare the demographic characteristics between groups with three or more groups.

The dynamic patterns (trajectories) in SBP and DBP between 2013 and 2019 were derived using LCTA, and their associations with war-related traumatic events were analyzed using multinomial logistic regression analysis. The model was adjusted for potential covariates. Principal components analysis (PCA) was used to develop a composite score based on nine war-related traumatic events.

All analyses were conducted using STATA software (version 14.0) and SAS (version 9.4). $P < 0.05$ was considered statistically significant.

Results

1|BP trajectories

Three distinct trajectory changes of SBP, and five DBP were identified, with 13,978 measurements of SBP and DBP (from 2013 to 2019) for our 1000 participants (Figure 1). The three SBP trajectory groups were: Group 1 (Normal-stable, 54.9%), Group 2 (High-stable, 22.7%), and Group 3 (Constant-very-high, CVH, 22.4%). In most years, SBP was above 160 mmHg in the CVH group.

Among the five DBP trajectory groups (normal-stable, high-stable, moderate to high, elevated to decreasing, and CVH), the prevalence of CVH DBP was 21.4%. Over 2013-2019, DBP was above 95 mmHg in the CVH DBP group.

Figure 1 displays the definition of each BP trajectory group based on their follow-up data: The 'Normal-stable' had a stable SBP at 120 mmHg or stable DBP at 80 mmHg, 'High-stable' had a stable SBP at 150 mmHg or a stable DBP at 86 mmHg, 'Moderate to high' had a rapid gain of nearly 10 mmHg from DBP at 85 mmHg and then decreased by almost 5 mmHg but within a high level of DBP, and 'Elevated to decreasing' had a rapid decrease of nearly 15 mmHg from DBP at 100 mmHg.

2|Characteristics across BP trajectory groups

Across the BP trajectory groups, more than 41% of those with CVH SBP and CVH DBP were refugees ($P = 0.851$ and 0.165 , respectively, Table 1). Additionally, 88.8% and 92.1% of those with CVH SBP and CVH DBP were unemployed. Besides, more than 86% of both have no stable monthly income among those clinically diagnosed with hypertension, 24.4% were non-compliant with antihypertensive medication (51.3% and 46.3% in CVH SBP and CVH DBP, respectively). Individuals with hypertension were more likely to use diuretic drugs than angiotensin-converting enzyme (ACE) inhibitors drugs (10.9% vs. 2.3%), whereas 7.5% take ACE inhibitors or diuretic drugs. There was no difference in all the sociodemographic factors except gender among the different BP trajectory groups.

3. Distribution of war-related traumatic events by trajectories of DBP and SBP

The overall prevalence of self-reported injury (or family member), death of a family member, violence due to house bombing, and living in debt was 51.4%, 54.1%, 66.5%, and 65.1%, respectively (Table 2).

The prevalence of self-reported injury or their family member, death of a family member during wars was 63.8% and 65.6% for CVH SBP. The corresponding figures were 66.4% and 66.4% for CVH DBP.

Moreover, the composite score based on the overall events indicated that CVH for SBP and DBP trajectory changes have the highest adverse event than other trajectory groups. The mean score for CVH SBP and DBP trajectories was 0.44 ± 0.10 and 0.49 ± 0.11 , respectively ($P < 0.001$) (Table 2).

4. Association between war traumatic events and BP trajectories

Multinomial logistic regression analysis revealed that war-related traumatic events were significantly associated with the trajectory groups of CVH SBP and DBP (Table 3). After adjusting for covariates, self-reported injury or their family member, death of a family member during wars were positively associated with CVH SBP with an odds ratio (95%CI) of 1.79 (1.28-2.48) and 1.90 (1.36-2.65), respectively. The corresponding figures were 1.92 (1.36-2.71) and 1.90 (1.35-2.68) for CVH DBP (Table 3).

Moreover, living in debt was positively associated with CVH SBP with an odds ratio (95%CI) of 2.49 (1.73-3.60) and CVH DBP with an odds ratio (95%CI) of 2.37 (1.63-3.45) (Table 3).

Discussion

This study investigated the trajectory of BP using longitudinal data collected from a uniquely vulnerable population, the Palestinian adults living in Gaza Strip, 64% of them are refugees. They have faced adverse and challenging living conditions, including war trauma. Our findings show that exposure to frequent traumatic events is linked to chronic disease risks, as indicated by elevated BP over time. In our trajectory analyses of SBP and DBP measurements (2013-2019), 22.4% and 21.4% of participants had CVH SBP (>160 mmHg) and DBP (>95 mmHg). The war-related traumatic events were positively associated with adverse BP trajectories.

This study provides evidence of the ongoing hypertension burden and the particular difficulties this vulnerable population faces. Because of exposure to war-related traumatic events might cause a double or triple burden in many families in Gaza, or perhaps more. One may find people from the same family have been injured or killed or even exposed to violence due to the bombing of their houses and others. The burden of the presence of all these events caused by wars might increase the frequency of developing hypertension or lead to worsening the levels of systolic and diastolic BP more and more. The Palestinians in Gaza are greatly affected by events surrounding oscillating stability and cumulative traumatic life (25). We believe that living in a situation of ongoing trauma may lead to further suffering in an agitated environment, leaving them vulnerable to developing hypertension.

The high affliction of war-related traumatic events indicates a significant association of war-related traumatic events, which are: injury of any family member, exposure to violence due to house bombing, and living in debt, with the highest trajectory change 'CVH' of SBP and DBP, where this trajectory characterized with a rapid gain of BP values, throughout the follow-up (2013-2019), within a constant high level of SBP and DBP. The relatively high proportion of 'CVH' trajectory change (22.4%) in SBP and (21.4%) in DBP confirm that large numbers of hypertension patients could be affected by the war-related traumatic events, which is a significant concern and should be taken into consideration in the prevention/management of NCDs.

The Palestinian people have been facing four wars in the past 14 years, besides the post-war repercussions of low living standards, poverty, and livelihood in debt. In 2019, Gaza Strip residents' poverty rate was 86% (33% severe poverty) (25, 26). Perhaps living at this poverty level among Gazans would make them live their daily lives dependent on others' debts. Meanwhile, life for the average Palestinian in Gaza is getting more and more wretched (27).

We found injury due to wars was positively associated with adverse BP trajectories. The finding is in line with another study, which reported that initial injury severity was independently associated with hypertension (28). Another study conducted in Iraq and Afghanistan veterans reported that the severity of combat injury was associated with the subsequent development of hypertension, coronary artery disease, diabetes mellitus, and chronic kidney disease (29).

Moreover, Gaza Strip is a complicated environment with structural challenges such as the lack of medicines, low salaries, and specialized training abroad (30). It challenges the health system's capacity to respond to the populations' basic needs (31). There is a noticeable shortage of medicines for cancer and chronic diseases due to those wars' repercussions. According to the Ministry of Health in Gaza, over 50% of the essential medications for chronic diseases were unavailable (32). In light of those mentioned earlier, the presence of 24.4% of the diagnosed hypertension to be non-compliant to antihypertensive medication in our study is reasonable; these findings support the figures of a survey that was undertaken at the outpatient clinics of the Ministry of Health in West Bank, Palestine ($n=450$), which indicated that more than half of the hypertension patients (54.2%) had poor adherence with medications, where one of the reasons for the low-adherence to medications was cost and unavailability of these medications at the healthcare centers (33).

Various potential risk factors, including psychosocial factors and stressful experiences in war life in Gaza, may play a role in developing hypertension (34). Frequent exposure to stressful events is thought to be one of the most common environmental causes of hypertension on a physiological basis (35). Studies have shown that those stressful events induce various neurochemical, neurotransmitter, and hormonal changes, predominantly by triggering the sympathetic nervous system (SNS) and hypothalamic-pituitary-adrenal (HPA) axes (36). SNS and HPA axes are woken up to release chemical mediators to protect the body from stress (37). This is in line with a cohort study of 122,816 adults aged ≥ 30 years, with a different cultural and socioeconomic setting population in France, where perceived stress was significantly associated with high BP (38).

More future research will help explain the biological mechanisms for the effect of war events on BP. There is research suggest that the body creates a surge of hormones in a stressful environment, which then causes the heart to beat faster and blood vessels to narrow (39). Stability and living in peace without wars can help people with hypertension trigger the relaxation response and reduce stress. Thus, reducing the risk of CVDs.

The major strengths of this study are targeting a uniquely vulnerable population, the Palestinian adults, who face adverse and challenging living conditions and war trauma. It is the first study to investigate the longitudinal trajectory of SBP and DBP in Gaza, a very special place in the world. This study involved both cross-sectional and historical longitudinal designs with repeated SBP and DBP measurements for a representative population, enabling us to perform the trajectory analyses.

This study has limitations. First, the study design cannot prove causal relationships. Second, the sociodemographic and the war-related traumatic events were collected in 2019 using a survey questionnaire. Third, we cannot rule out selection bias as only those who regularly visited the primary healthcare centers in Gaza Strip were enrolled in the study. In addition, the sample size and study duration are limited.

Future efforts are needed to enhance implementing care models to improve NCD management, including hypertension, health education, and medication services in Gaza Strip. In addition, an in-depth trauma care system evaluation is needed in Gaza's health system. Therefore, a comprehensive intervention towards war-related traumatic events should be provided to the people in Gaza Strip.

In conclusion, this study provides evidence that the burden of war-related traumatic events among Palestinians living in Gaza Strip is high. Such war-related traumatic events are positively associated with adverse BP trajectory. Efforts and sustainable programs are needed to enhance health care services to improve health education and medication services and for NCD prevention and management, including hypertension, mental health, and other prevalent health conditions in Gaza Strip.

Declarations

Conflicts of interest

The authors declare no conflict of interest. The funders had no role in the study's design, collection, analyses, or interpretation of the data, writing of the manuscript, or deciding to publish the results.

Author contributions

MJ, XL and YW designed the study. MJ collected the data, made data entries with Gaza's teamwork, analysed the data, and drafted the manuscript. MM assisted data analysis, interpretation of results, and manuscript drafting. XL, ZS and YW designed the research hypothesis and guided data analysis. MB analysed the trajectory data and assisted in the data interpretation and editing. HA helped in data collection and entry. All authors revised the manuscript and approved the final version to be submitted. YW provided administration support for the study and is the guarantor of this work. This work is part of MJ's PhD dissertation research in Xi'an Jiaotong University.

Acknowledgments

This research was funded in part by the US-based China Medical Board (CMB, grant number 16-262); National Natural Science Foundation of China, grant number 82173504, 82011530197; and the Chinese National Key Research and Development Program (grant number 2017YFC0907200, 2017YFC0907201). We would like to thank all the study participants and our collaborators in Gaza Strip and China for their support. Special thanks to the Ministry of Health of Gaza.

Availability of data and materials

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

References

1. Pedersen D. Political violence, ethnic conflict, and contemporary wars: broad implications for health and social well-being. *Social science & medicine*. 2002;55(2):175–90.
2. Filii J-P. The twelve wars on Gaza. *Journal of Palestine Studies*. 2014;44(1):52–60.
3. Bregman A. *Israel's wars: a history since 1947*: Routledge; 2016.
4. Baidoun SD, Salem MZ, Omran OA. Assessment of TQM implementation level in Palestinian healthcare organizations: The case of Gaza Strip hospitals. *The TQM Journal*. 2018.
5. Munakata M. Clinical significance of stress-related increase in blood pressure: current evidence in office and out-of-office settings. *Hypertension Research*. 2018;41(8):553–69.
6. Sa'adeh HH, Darwazeh RN, Khalil AA, Sa'ed HZ. Knowledge, attitudes and practices of hypertensive patients towards prevention and early detection of chronic kidney disease: a cross sectional study from Palestine. *Clinical hypertension*. 2018;24(1):6.
7. Feldman I. The humanitarian condition: Palestinian refugees and the politics of living. *Humanity: an international journal of human rights, humanitarianism, and development*. 2012;3(2):155–72.
8. Cuevas AG, Williams DR, Albert MA. Psychosocial factors and hypertension: a review of the literature. *Cardiology clinics*. 2017;35(2):223–30.
9. Leng B, Jin Y, Li G, Chen L, Jin N. Socioeconomic status and hypertension: a meta-analysis. *Journal of hypertension*. 2015;33(2):221–9.
10. Johnson H, Thompson A. The development and maintenance of post-traumatic stress disorder (PTSD) in civilian adult survivors of war trauma and torture: A review. *Clinical psychology review*. 2008;28(1):36–47.

11. Friedman MJ, Schnurr PP, McDonagh-Coyle A. Post-traumatic stress disorder in the military veteran. *Psychiatric Clinics*. 1994;17(2):265–77.
12. Mendlowicz V, Garcia-Rosa ML, Gekker M, Wermelinger L, Berger W, de Luz MP, et al. Post-traumatic stress disorder as a predictor for incident hypertension: a 3-year retrospective cohort study. *Psychological Medicine*. 2021:1–8.
13. Gasperin D, Netuveli G, Dias-da-Costa JS, Pattussi MP. Effect of psychological stress on blood pressure increase: a meta-analysis of cohort studies. *Cadernos de saude publica*. 2009;25:715–26.
14. Mosleh M. Non-communicable diseases and war injuries in Palestine: burden, incidence and management in the health system: Mid Sweden University/Mittuniversitetet; 2020.
15. Allen N, Berry JD, Ning H, Van Horn L, Dyer A, Lloyd-Jones DM. Impact of blood pressure and blood pressure change during middle age on the remaining lifetime risk for cardiovascular disease: the cardiovascular lifetime risk pooling project. *Circulation*. 2012;125(1):37–44.
16. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic review and meta-regression analysis. *Circulation*. 2008;117(25):3171.
17. Jebri M, Liu X, Shi Z, Mazidi M, Altaher A, Wang Y. Prevalence of Type 2 Diabetes and Its Association with Added Sugar Intake in Citizens and Refugees Aged 40 or Older in the Gaza Strip, Palestine. *International journal of environmental research and public health*. 2020;17(22):8594.
18. Kubany ES, Leisen MB, Kaplan AS, Watson SB, Haynes SN, Owens JA, et al. Development and preliminary validation of a brief broad-spectrum measure of trauma exposure: the Traumatic Life Events Questionnaire. *Psychological assessment*. 2000;12(2):210.
19. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Himmelfarb CD, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Journal of the American College of Cardiology*. 2018;71(19):e127–e248.
20. Croudace TJ, Jarvelin M-R, Wadsworth ME, Jones PB. Developmental typology of trajectories to nighttime bladder control: epidemiologic application of longitudinal latent class analysis. *American Journal of Epidemiology*. 2003;157(9):834–42.
21. Nagin DS, NAGIN D. *Group-based modeling of development*: Harvard University Press; 2005.
22. Andruff H, Carraro N, Thompson A, Gaudreau P, Louvet B. Latent class growth modelling: a tutorial. *Tutorials in Quantitative Methods for Psychology*. 2009;5(1):11–24.
23. Nagin DS. Analyzing developmental trajectories: a semiparametric, group-based approach. *Psychological methods*. 1999;4(2):139.
24. Nylund KL, Asparouhov T, Muthén BO. "Deciding on the number of classes in latent class analysis and growth mixture modeling: A Monte Carlo simulation study": Erratum. 2008.
25. Hammad J, Tribe R. Social suffering and the psychological impact of structural violence and economic oppression in an ongoing conflict setting: The Gaza Strip. *Journal of community psychology*. 2020;48(6):1791–810.
26. Mosleh M, Dalal K, Aljeesh Y, Svanström L. The burden of war-injury in the Palestinian health care sector in Gaza Strip. *BMC international health and human rights*. 2018;18(1):1–9.
27. Al-Salehi J, Twam F, Atamanov A, Palaniswamy N. *Measuring Poverty in West Bank and Gaza*. 2018.
28. Howard JT, Sosnov JA, Janak JC, Gundlapalli AV, Pettey WB, Walker LE, et al. Associations of initial injury severity and posttraumatic stress disorder diagnoses with long-term hypertension risk after combat injury. *Hypertension*. 2018;71(5):824–32.
29. Stewart IJ, Sosnov JA, Howard JT, Orman JA, Fang R, Morrow BD, et al. Retrospective analysis of long-term outcomes after combat injury: a hidden cost of war. *Circulation*. 2015;132(22):2126–33.
30. Mosleh M, Al Jeesh Y, Dalal K, Eriksson C, Carlerby H, Viitasara E. Barriers to managing and delivery of care to war-injured survivors or patients with non-communicable disease: a qualitative study of Palestinian patients' and policy-makers' perspectives. *BMC health services research*. 2020;20:1–11.
31. Shabab QS. The Gaza health system. *AMSA Journal of Global Health*. 2018;12(1):6–9.
32. Mosleh M, Dalal K, Aljeesh Y, Svanström L. The burden of war-injury in the Palestinian health care sector in Gaza Strip. *BMC international health and human rights*. 2018;18(1):28.
33. Al-Ramahi R. Adherence to medications and associated factors: A cross-sectional study among Palestinian hypertensive patients. *Journal of epidemiology and global health*. 2014;5(2):125–32.
34. Schoenthaler AM, Rosenthal DM. *Stress and hypertension. Disorders of Blood Pressure Regulation*: Springer; 2018. p. 289–305.
35. Pickering TG, Devereux RB, James GD, Gerin W, Landsbergis P, Schnall PL, et al. Environmental influences on blood pressure and the role of job strain. *Journal of hypertension Supplement: official journal of the International Society of Hypertension*. 1996;14(5):S179-85.
36. Sugama S, Kakinuma Y. Stress and Brain Immunity: Microglial Homeostasis through Hypothalamus-Pituitary-Adrenal Gland Axis and Sympathetic Nervous System. *Brain, Behavior, & Immunity-Health*. 2020:100111.
37. Liu Y-Z, Wang Y-X, Jiang C-L. Inflammation: the common pathway of stress-related diseases. *Frontiers in human neuroscience*. 2017;11:316.
38. Wiernik E, Pannier B, Czernichow S, Nabi H, Hanon O, Simon T, et al. Occupational status moderates the association between current perceived stress and high blood pressure: evidence from the IPC cohort study. *Hypertension*. 2013;61(3):571–7.
39. Rahma VA, Rau M, Krisnasari S. Effects of Psychological Distress on Hypertension among Liquefaction Refugees in Petobo Shelter. *European Journal of Molecular & Clinical Medicine*. 2020;7(7):89–96.

Tables

Table 1. Characteristics of the study participants of Palestinian adults living in Gaza

A. Trajectory group of SBP	Total	Normal-stable	High-stable	Constant very-high	P-value
	N=1000	n=549	n=227	n=224	
Age, years (mean±SD)	59.2±7.5	58.9±7.7	59.7±7.2	59.3±7.6	0.333
Gender					0.039
Male	532 (53.2%)	296 (53.9%)	132 (58.2%)	104 (46.4%)	
Female	468 (46.8%)	253 (46.1%)	95 (41.8%)	120 (53.6%)	
Study region					0.897
North	220 (22.0%)	127 (23.1%)	48 (21.1%)	45 (20.1%)	
Gaza	122 (12.2%)	59 (10.8%)	33 (14.5%)	30 (13.4%)	
Midzone	214 (21.4%)	117 (21.3%)	47 (20.7%)	50 (22.3%)	
Khan Younis	223 (22.3%)	123 (22.4%)	48 (21.2%)	52 (23.2%)	
Rafah	221 (22.1%)	123 (22.4%)	51 (22.5%)	47 (21.0%)	
Refugees	422 (42.2%)	236 (43.0%)	93 (41.0%)	93 (41.5%)	0.851
Household size, person (mean±SD)	6.9±1.5	6.9±1.5	6.9±1.5	6.8±1.5	0.623
Marital status					0.473
Married	844 (84.4%)	456 (83.1%)	191 (84.1%)	197 (88.0%)	
Non-married	156 (15.6%)	93 (16.9%)	36 (15.9%)	27 (12.0%)	
Education*					0.614
Low	451 (45.1%)	257 (46.8%)	99 (43.6%)	95 (42.4%)	
Moderate	295 (29.5%)	154 (28.1%)	74 (32.6%)	67 (29.9%)	
High	254 (25.4%)	138 (25.1%)	54 (23.8%)	62 (27.7%)	
Employment					0.936
Employed	117 (11.7%)	66 (12.0%)	26 (11.4%)	25 (11.2%)	
Non-employed					
Household income					0.858
<500 NIS	51 (5.1%)	25 (4.5%)	15 (6.6%)	11 (4.9%)	
500-1000 NIS	41 (4.1%)	25 (4.5%)	9 (4.0%)	7 (3.1%)	
>1000 NIS	54 (5.4%)	34 (6.2%)	11 (4.8%)	9 (4.0%)	
No stable income	854 (85.4%)	465 (84.8%)	192 (84.6%)	197 (88.0%)	
Physical activity					0.382
Low	654 (65.4%)	355 (64.6%)	144 (63.4%)	155 (69.2%)	

Moderate	247 (24.7%)	133 (24.3%)	60 (26.5%)	54 (24.1%)	
High	99 (9.9%)	61 (11.1%)	23 (10.1%)	15 (6.7%)	
Regular cigarette smokers	447 (44.7%)	250 (55.9%)	101 (22.6%)	96 (21.5%)	0.792
Antihypertensive drugs					0.485
ACE inhibitors	23 (2.3%)	-	12 (5.3%)	11 (4.9%)	
Diuretics	109 (10.9%)	-	54 (23.7%)	55 (24.5%)	
Either	75 (7.5%)	-	32 (14.1%)	43 (19.3%)	
Neither	549 (54.9%)	549 (54.9%)	-	-	
Non-compliant	244 (24.4%)	-	129 (56.9%)	115 (51.3%)	

B. Trajectory group of DBP	Total	Normal-stable	High-stable	Moderate to high	Elevated to decreasing	Constant-very-high	P-value
	<i>N</i> =1000	<i>n</i> =526	<i>n</i> =99	<i>n</i> =64	<i>n</i> =97	<i>n</i> =214	
Age, years (mean±SD)	59.2±7.5	58.8±7.7	59.0±6.8	60.1±6.8	59.6±7.2	59.4±7.8	0.686
Gender							0.055
Male	532 (53.2%)	287 (54.6%)	58 (58.6%)	36 (56.3%)	56 (57.7%)	95 (44.4%)	
Female	468 (46.8%)	239 (45.4%)	41 (41.4%)	28 (43.7%)	41 (42.3%)	119 (55.6%)	
Study region							0.567
North	220 (22.0%)	124 (23.6%)	15 (15.1%)	11 (17.2%)	25 (25.8%)	45 (21.0%)	
Gaza	122 (12.2%)	57 (10.8%)	11 (11.1%)	13(20.3%)	9 (9.3%)	32 (15.0%)	
Midzone	214 (21.4%)	114 (21.7%)	23 (23.2%)	13 (20.3%)	16 (16.5%)	48 (22.4%)	
Khan Younis	223 (22.3%)	112 (21.3%)	26 (26.3%)	15 (23.4%)	23 (23.7%)	47 (22.0%)	
Rafah	221 (22.1%)	119 (22.6%)	24 (24.3%)	12 (18.8%)	24 (24.7%)	42 (19.6%)	
Refugees	422 (42.2%)	227 (43.2%)	31 (31.3%)	25 (39.1%)	46 (47.4%)	93 (43.5%)	0.165
Household size, person (mean±SD)	6.9±1.5	6.9±1.5	7.2±1.6	6.9±1.4	6.8±1.5	6.8±1.5	0.336
Marital status							0.765
Married	844 (84.4%)	437 (83.1%)	82 (82.8%)	53 (82.8%)	81 (83.5%)	191 (89.3%)	
Non-married	156 (15.6%)	89 (16.9%)	17 (17.2%)	11 (17.2%)	16 (16.5%)	23 (10.7%)	
Education*							0.658
Low	451 (45.1%)	245 (46.6%)	46 (46.5%)	30 (46.9%)	41 (42.3%)	89 (41.6%)	
Moderate	295 (29.5%)	149 (28.3%)	33 (33.3%)	16 (25.0%)	34 (35.0%)	63 (29.4%)	
High	254 (25.4%)	132 (25.1%)	20 (20.2%)	18 (28.1%)	22 (22.7%)	62 (29.0%)	
Employment							0.361
Employed	117 (11.7%)	64 (12.2%)	14 (14.1%)	8 (12.5%)	14 (14.4%)	17 (7.9%)	
Non-employed		462 (87.8%)	126 (85.9%)	56 (87.5%)	83 (85.6%)	197 (92.1%)	
Household income							0.977
<500 NIS	51 (5.1%)	25 (4.7%)	4 (4.0%)	5 (7.8%)	5 (5.2%)	12 (5.6%)	
500-1000 NIS	41 (4.1%)	22 (4.2%)	5 (5.1%)	1 (1.6%)	6 (6.2%)	7 (3.3%)	
>1000 NIS	54 (5.4%)	32 (6.1%)	6 (6.1%)	2 (3.1%)	4 (4.1%)	10 (4.6%)	
No stable income	854 (85.4%)	447 (85.0%)	84 (84.8%)	56 (87.5%)	82 (84.5%)	185 (86.5%)	
Physical activity							0.689

Low	654 (65.4%)	64 (64.6%)	39 (60.9%)	67 (69.1%)	145 (67.8%)	145 (67.8%)	
Moderate	247 (24.7%)	28 (28.3%)	16 (25.0%)	23 (23.7%)	52 (24.3%)	52 (24.3%)	
High	99 (9.9%)	7 (7.1%)	9 (14.1%)	7 (7.2%)	17 (7.9%)	17 (7.9%)	
Regular cigarette smokers	447 (44.7%)	40 (40.4%)	31 (48.4%)	36 (37.1%)	98 (45.8%)	98 (45.8%)	0.430
Antihypertensive drugs							0.106
ACE inhibitors	23 (2.3%)	8 (8.1%)	3 (4.7%)	4 (4.1%)	10 (4.6%)	10 (4.6%)	
Diuretics	109 (10.9%)	18 (18.2%)	12 (18.7%)	29 (30.0%)	58 (27.1%)	58 (27.1%)	
Either	75 (7.5%)	9 (9.1%)	8 (12.5%)	17 (17.5%)	47 (22.0%)	47 (22.0%)	
Neither	549 (54.9%)	-	-	-	-	-	
Non-compliant	244 (24.4%)	64 (64.6%)	41 (64.1%)	47 (48.4%)	99 (46.3%)	99 (46.3%)	

-The associations were studied using ANOVA and X^2 test.

Abbreviations: SBP, Systolic blood pressure; DBP, Diastolic blood pressure; NIS, New Israeli Shekel; ACE inhibitors, Angiotensin-converting enzyme.

*Education level (low: secondary or more subordinate, moderate: diploma or two years after high school, and high: bachelor's and above).

Trajectory groups of SBP: 1) Normal-stable (54.9 %): had a stable SBP at 120 mmHg. 2) High-stable (22.7 %): had a stable SBP at 150 mmHg. 3) Constant-very-high (22.4%): SBP was above 160 mmHg.

Trajectory groups of DBP: 1) Normal-stable (52.6 %): had a stable DBP at 80 mmHg. 2) High-stable (9.9%): had a stable DBP at 86 mmHg. 3) Moderate to high (6.4%): had a rapid gain of nearly 10 mmHg from DBP at 85 mmHg and then decreased by almost 5 mmHg but within the high level of DBP. 4) Elevated to decreasing (9.7%): had a rapid decrease of nearly 15 mmHg from DBP at 100 mmHg. 5) Constant-very-high (21.4%): DBP was above 95 mmHg.

Table 2. Distribution of experiencing war-related traumatic events on the trajectory of systolic and diastolic BP of Palestinian adults living in Gaza, from 2013 to 2019 (N=1000)

War-related traumatic events¶	Total N=1000	Trajectory group of SBP			P-value	Trajectory group of DBP					P-value
		Normal-stable n=549	High-stable n=227	Constant-very-high n=224		Normal-stable n=526	High-Stable n=99	Moderate to high n=64	Elevated to decreasing n=97	Constant-very-high n=214	
1) Injury of any family member	514 (51.4%)	278 (50.6%)	93 (41.0%)	143 (63.8%)	<0.001	269 (51.1%)	48 (48.5%)	18 (28.1%)	37 (38.1%)	142 (66.4%)	<0.001
2) Death of any family member	541 (54.1%)	279 (50.8%)	115 (50.6%)	147 (65.6%)	<0.001	269 (51.1%)	48 (48.5%)	32 (50.0%)	50 (51.5%)	142 (66.4%)	0.001
3) Serious illness of any family member	303 (30.3%)	154 (28.1%)	76 (33.5%)	73 (32.6%)	0.228	146 (27.7%)	33 (33.3%)	23 (35.9%)	34 (35.1%)	67 (31.3%)	0.001
4) Loss of job	137 (13.7%)	79 (14.4%)	31 (13.7%)	27 (12.1%)	0.693	74 (14.1%)	14 (14.1%)	12 (18.8%)	14 (14.4%)	23 (10.8%)	0.001
5) Business Bankrupt	73 (7.3%)	35 (6.4%)	19 (8.4%)	19 (8.5%)	0.463	32 (6.1%)	6 (6.1%)	10 (15.6%)	10 (10.3%)	15 (7.0%)	0.001
6) Exposure to violence due to house bombing	665 (66.5%)	362 (65.9%)	138 (60.8%)	165 (73.6%)	0.014	347 (66.0%)	70 (70.7%)	40 (62.5%)	46 (47.4%)	162 (75.7%)	<0.001
7) Living in debt	651 (65.1%)	326 (59.4%)	151 (66.5%)	174 (77.7%)	<0.001	312 (59.3%)	69 (69.7%)	37 (57.8%)	68 (70.1%)	165 (77.1%)	<0.001
8) Marital separation or divorce	59 (5.9%)	26 (4.7%)	21 (9.3%)	12 (5.4%)	0.049	27 (5.1%)	6 (6.1%)	4 (6.3%)	9 (9.3%)	13 (6.1%)	0.001
9) Major disasters of any family member	149 (14.9%)	79 (14.4%)	38 (16.7%)	32 (14.3%)	0.675	74 (14.1%)	15 (15.2%)	13 (20.3%)	18 (18.6%)	29 (13.5%)	0.001
Composite score* (Mean±SD)	-0.16±0.09	-0.10±0.07	-0.19±0.11	0.44±0.10	<0.001	-0.10±0.08	-0.00±0.17	-0.39±0.19	-0.31±0.15	0.49±0.11	<0.001

¶Variables were recoded binary (yes or no).

*Principal component analysis (PCA) was used to develop the composite score based on nine war-related traumatic events.

Table 3. Association between war-related traumatic events and trajectory of systolic BP of Palestinian adults living in Gaza, from 2013 to 2019 (N=1000)

War-related traumatic events¶	Trajectory group of SBP			
	High-stable		Constant-very-high	
	Crude OR (95% CI)	Adjusted OR (95% CI)	Crude OR (95% CI)	Adjusted OR (95% CI)
1. Injury of any family member	0.68 (0.49-1.03)	0.73 (0.51-1.08)	1.72 (1.25-2.37)*	1.79 (1.28-2.48)*
1. Death of any family member	0.99 (0.73-1.35)	1.03 (0.75-1.42)	1.85 (1.34-2.55)*	1.90 (1.36-2.65)*
1. Serious illness of any family member	1.29 (0.93-1.80)	1.31 (0.93-1.85)	1.24 (0.89-1.73)	1.23 (0.87-1.74)
1. Loss of job	0.94 (0.60-1.47)	0.96 (0.61-1.52)	0.82 (0.51-1.30)	0.86 (0.53-1.35)
1. Business bankrupt	1.34 (0.75-2.40)	1.35 (0.75-2.45)	1.36 (0.76-2.43)	1.36 (0.75-2.49)
1. Exposure to violence due to house bombing	0.80 (0.58-1.10)	0.81 (0.58-1.12)	1.44 (1.02-2.04)*	1.44 (1.01-2.05)*
1. Living in debt	1.36 (0.98-1.88)	1.37 (0.98-1.91)	2.38 (1.66-3.40)*	2.49 (1.73-3.60)*
1. Marital separation or divorce	0.92 (0.66-1.27)	1.21 (0.86-1.64)	0.88 (0.44-1.77)	0.83 (0.40-1.73)
1. Major disasters of any family member	1.19 (0.77-1.83)	1.20 (0.78-1.85)	0.99 (0.64-1.54)	1.01 (0.66-1.59)

War-related traumatic events¶	Trajectory group of DBP							
	High-stable		Moderate to high		Elevated to decreasing		Constant-very-high	
	Crude OR (95% CI)	Adjusted OR (95% CI)	Crude OR (95% CI)	Adjusted OR (95% CI)	Crude OR (95% CI)	Adjusted OR (95% CI)	Crude OR (95% CI)	Adjusted OR (95% CI)
1) Injury of any family member	0.89 (0.59-1.38)	0.96 (0.62-1.50)	0.78 (0.59-1.09)	0.81 (0.63-1.16)	0.95 (0.61-1.49)	0.99 (0.62-1.55)	1.88 (1.35-2.62)*	1.92 (1.36-2.71)*
2) Death of any family member	0.90 (0.59-1.38)	0.95 (0.61-1.49)	0.95 (0.57-1.61)	1.02 (0.60-1.76)	1.02 (0.66-1.56)	1.02 (0.65-1.61)	1.80 (1.33-2.58)*	1.90 (1.35-2.68)*
3) Serious illness of any family member	1.30 (0.82-2.06)	1.33 (0.83-2.14)	1.46 (0.85-2.52)	1.49 (0.85-2.61)	1.40 (0.89-2.22)	1.40 (0.88-2.24)	1.19 (0.84-1.68)	1.20 (0.84-1.71)
4) Loss of job	1.01 (0.54-1.86)	1.01 (0.54-1.88)	1.41 (0.72-2.77)	1.44 (0.71-2.88)	1.03 (0.55-1.91)	1.04 (0.57-1.94)	0.74 (0.45-1.21)	0.75 (0.44-1.23)
5) Business bankrupt	1.00 (0.41-2.45)	1.02 (0.41-2.49)	2.86 (1.33-4.16)*	2.87 (1.26-4.20)*	1.77 (0.84-3.74)	1.85 (0.86-3.96)	1.16 (0.61-2.21)	1.17 (0.62-2.24)
6) Exposure to violence due to house bombing	1.24 (0.78-1.99)	1.29 (0.80-2.08)	0.85 (0.50-1.47)	0.85 (0.49-1.49)	0.94 (0.60-1.47)	0.96 (0.60-1.49)	1.59 (1.11-2.30)*	1.62 (1.13-2.38)*
7) Living in debt	1.58 (0.99-2.51)	1.65 (1.03-2.66)*	0.94 (0.55-1.59)	0.95 (0.56-1.61)	1.61 (1.01-2.57)*	1.69 (1.05-2.73)*	2.35 (1.63-3.41)*	2.37 (1.63-3.45)*
8) Marital separation or divorce	0.84 (0.34-2.09)	0.94 (0.34-2.57)	0.81 (0.27-2.39)	0.82 (0.27-2.52)	0.53 (0.24-1.16)	0.48 (0.22-1.13)	0.84 (0.42-1.65)	0.82 (0.40-1.66)
9) Major disasters of any family member	1.09 (0.59-1.99)	1.13 (0.61-2.09)	1.55 (0.81-3.00)	1.58 (0.80-3.14)	1.39 (0.78-2.45)	1.45 (0.80-2.63)	0.96 (0.60-1.52)	0.94 (0.57-1.50)

Multinomial logistic regression analysis was conducted. The reference group was normal-stable SBP. **The models** adjusted for age, gender, region, family size, citizenship, marital status, income, education, employment, smoking, physical activity, body mass index, fruits and vegetable intake, type 2 diabetes, and lipid profile (total cholesterol, triglycerides, HDL-C, and LDL-C).

- ¶Variables were recoded binary (yes or no).

*P-value < 0.05, statistically significant association.

Figures




Figure 1

Trajectory modeling of BP of Palestinian adults living in Gaza, from 2013 to 2019 (N=1000)

Trajectory groups of SBP: 1) Normal-stable (54.9 %): stable SBP at 120 mmHg. 2) High-stable (22.7 %): had a stable SBP at 150 mmHg.

3) Constant very high (22.4%): had a rapid gain of nearly 10 mmHg from SBP at 165 mmHg and then decreased but within the high level of SBP.

Trajectory groups of DBP: 1) Normal-stable (52.6 %): had a stable DBP at 80 mmHg. 2) High-stable (9.9%): had a stable DBP at 86 mmHg.

3) Moderate to high (6.4%): had a rapid gain of nearly 10 mmHg from DBP at 85 mmHg and then decreased by almost 5 mmHg but within the constant high level of DBP. 4) Elevated to decreasing (9.7%): had a rapid decrease of nearly 15 mmHg from DBP at 100 mmHg.

5) Constant very high (21.4%): had a gain of nearly 3 mmHg from DBP at 95 mmHg and decreased but within the constant high level of DBP.