

Associated factors with awareness, treatment and control of hypertension among 3,579 hypertensive adults in China: data from the China Health and Nutrition Survey

junxiang Wei (✉ junxiang.wei@xjtu.edu.cn)

Xi'an Jiaotong University School of Medicine

Bo Xin

Shanxi University of Traditional Chinese Medicine

Yan Li

Icahn School of Medicine at Mount Sinai Department of Medicine

Youfa Wang

Xi'an Jiaotong University

Research article

Keywords: Hypertension, awareness, treatment, control, China

Posted Date: April 1st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-18826/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on March 1st, 2021. See the published version at <https://doi.org/10.1186/s12889-021-10417-4>.

Abstract

Background: Hypertension prevalence is high and rising in China, but it is inadequately controlled. This study investigated hypertension awareness, treatment, and control and their associated factors among Chinese adults.

Methods: Data collected from the 2011 China Health and Nutrition Survey (CHNS) from 12,991 Chinese adults were used. Hypertension was defined as systolic blood pressure ≥ 140 mm Hg, diastolic blood pressure ≥ 90 mm Hg, self-reported prior diagnosed hypertension, or taking antihypertensive medications. Hypertension awareness, treatment, and control were defined as a self-reported diagnosis of hypertension, current use of antihypertensive medication, and blood pressure $< 140/90$ mm Hg, respectively. Multivariate logistic regression was performed to examine factors associated with hypertension awareness, treatment, and control.

Results: Overall, 3,579 (27.6%) of the CHNS adult participants had hypertension, of whom 55.7% were aware of their diagnosis, 46.5% were treated with antihypertensive medications, but only 20.3% had their blood pressure controlled. Rates of hypertension awareness and treatment varied across population subgroups; higher likelihood was associated with being female (OR = 1.37; 95%CI, 1.12-1.66), older age (1.57; 1.65-4.02), urban residence (1.50; 1.14-1.97), living in the Eastern region (1.52; 1.14-2.01), having overweight/obesity (1.99; 1.39-2.84), and engaging in less healthy lifestyles. Lower control rate was associated with obesity (0.56; 0.42-0.76) and minority ethnicity (0.52; 0.31-0.86).

Conclusion: Hypertension is a major public health burden in China. Rates of hypertension awareness, treatment, and control are still low despite efforts that have been made. More intensive screening and treatment intervention programs are needed in the future.

Introduction

Hypertension is one of the most prevalent chronic diseases in many countries and is the most common modifiable risk factor for other health conditions such as heart disease, stroke, kidney disease, and subsequent morbidity and mortality.⁽¹⁾ It leads to 9.4 million deaths each year worldwide, with the condition particularly affecting low- and middle-income countries.^(2, 3)

China has the largest population with hypertension in the world. Recent population-based surveys estimated that almost one third of Chinese adults have hypertension.⁽⁴⁻⁶⁾ As the society ages and continues an uncurbed trend of unhealthy lifestyles such as high sodium consumption, hypertension has become a major threat to the improvement of population health. To address this problem, effective prevention programs and timely treatment and control are critical.

Some recent studies showed that the awareness, treatment, and control rates of hypertension in China have increased significantly over the past two decades.^(7, 8) This is mainly attributed to the increases in health insurance coverage and utilization of healthcare resources as well as a wider implementation of

evidence-based blood pressure management programs.(9, 10) However, the awareness, treatment, and control rates of hypertension in China remain low compared to most developed countries. There is a need to study these issues further to alleviate the burden of hypertension and improve public health.

This study aimed to: (a) examine the status of hypertension awareness, treatment, and control among Chinese adults and by age, sex, region, and socioeconomic status; and (b) assess the factors that might affect hypertension awareness, treatment, and control to help shed light on the implementation of policies and interventions in the future.

Methods

METHODS

1. Data source

We used data from the 2011 China Health and Nutrition Survey (CHNS), which is an ongoing, large-scale, population-based survey initiated in 1989. CHNS used a stratified multistage, random cluster sampling strategy to select participants from 288 communities across a large number of primary provinces/autonomous cities.(11) The CHNS 2011 was conducted across more than 10 provinces and autonomous cities, including Beijing, Chongqing, Guangxi, Guizhou, Heilongjiang, Henan, Hubei, Hunan, Jiangsu, Liaoning, Shaanxi, Shandong, Shanghai, Yunnan, and Zhejiang. Our sample from the CHNS 2011 includes 5,884 households and 12,991 adults, which provides a large nationwide sample of the Chinese population in terms of major behavioral health and disease burden.

We restricted our analysis to those aged ≥ 18 years ($n = 12,991$) and focused on those with hypertension ($n = 3,645$) in 2011. Excluding missing values, 3,579 adults were included in the final dataset for analysis. Written informed consent was obtained from each participant before any data were collected. Note that newer CHNS data have not yet been released for use.

2. Measurement and definitions

We selected variables regarding knowledge about and medical history of hypertension, as well as demographics, socioeconomic status, and lifestyle behaviors from the adult household questionnaire in CHNS. We categorized geographic regions into three groups: Western China (Guangxi, Guizhou, and Chongqing), Eastern China (Shanghai, Beijing, Jiangsu, Liaoning, and Shandong) and Central China (Henan, Hubei, and Heilongjiang). Marital status was classified into currently married and single (i.e., never married, divorced, widowed, or separated). There were six a priori characteristics selected: residency (urban vs. rural), ethnicity (Han vs. minority), occupation (employed vs. unemployed), educational attainment (elementary school, middle school, high school or technical school, and college or above), smoking status (non-smoker, ex-smoker, and current smoker), alcohol consumption status (current drinker vs. non-drinker), and medical insurance status (insured vs. not insured). The physical examination, including height and weight, were measured by health care professionals. Weight was measured in light

closing to the nearest 0.1kg on a calibrated beam balance, while height was measured to the nearest 0.1cm using a portable stadiometer. Body mass index (BMI) was calculated as weight (kg) divided by the squared height (m²). Overweight and obesity were defined as a BMI of at least 24kg/m² and 28kg/m², respectively, based on the recommendations of the Working Group on Obesity in China.(12)

3. Assessment of hypertension and hypertension awareness, treatment, and control

The measurement and definition of hypertension was reported according to the 7th Chinese Joint National Commission guidelines.(13) A standard mercury sphygmomanometer was used by well-trained physicians to measure SBP and DBP on the right arm in triplicate after a 10-minute seated rest. The mean of the three readings was calculated and used in all analysis. Hypertension was defined as having an average SBP \geq 140 mm Hg and/or an average DBP \geq 90 mm Hg, and/or a self-reported previous diagnosis of hypertension by a health care provider, and/or taking antihypertensive drugs currently. Awareness of hypertension was defined as a self-report of any previous diagnosis of hypertension by a health care provider. Treatment of hypertension was defined as a self-reported use of antihypertensive medications at present. Control of hypertension was defined as having an average SBP < 140 mmHg and an average DBP < 90 mmHg while under pharmacological treatment for hypertension.

4. Statistical analysis

Data were presented as meanSD or portions/percentage for continuous and categorical variables according to gender, respectively. Differences between groups were tested using two-sample student t-tests for continuous variables and the Chi-square test for categorical variables. Multivariable logistic regression models were fit to explore the associations between relevant risk factors and hypertension awareness and treatment.

We also investigated the adjusted associations between independent variables and taking antihypertensive drugs among subjects who were aware of hypertension. Finally, characteristics and proportions of subjects by age groups were analyzed to identify the subpopulations that were more likely to take antihypertensive medications (“adherence”) in the subsample of participants who took treatments for hypertension. All the analysis was done using Stata 15.0 (StataCorp., 2017). P values were 2-tailed and $p < 0.05$ was considered to be statistically significant.

Results

1. General characteristics

The mean age of the patients was 60 years. About half of them were men (50.4%) and 58.5% were from rural areas. **Table 1** presents their demographic characteristics, behavioral risk factors, and weight status.

Rates of smoking and alcohol consumption were significantly higher in men than in women ($p < 0.001$). Women had a higher obesity (BMI ≥ 28) prevalence (22.6% vs. 18.4%) than men ($p = 0.01$).

// Table 1 here //

2. Awareness, treatment, and control of hypertension

Overall, the estimated awareness, treatment, and control rates were 55.7%, 46.5%, and 20.3% among Chinese adults with hypertension, respectively. Of those who were aware of their hypertension, 83.0% ($n = 1,664$) received antihypertensive treatment. In addition, 43.8% ($n = 728$) of those who received treatment had their hypertension controlled (SBP/DBP $< 140/90$ mmHg). The awareness and treatment rates were higher in women compared to men, but the blood pressure control rate was higher in men than in women though not significant (**Figure 1**). **Table 2** presents the proportions of hypertension outcomes in various characteristic groups. Older adults (≥ 65 yrs) had higher hypertension awareness and treatment rates compared to young adults (18-49 yrs). Both treatment and control rates were higher in urban areas compared to rural areas and were higher in Eastern China compared to Central China. Participants who had a higher educational attainment were more likely to have controlled hypertension. In contrast, participants who were ethnic minorities and obese were less likely to have controlled hypertension.

// Figure 1 here //

// Table 2 here //

3. Factors associated with hypertension awareness and treatment

Table 3 presents findings from multivariate logistic regression that identified factors associated with hypertension awareness, treatment, and control. Women were more likely to be aware of their hypertension than men, with an OR of 1.39 (95% CI, 1.14-1.70). Rural residency, minority ethnicity, and unemployment were negatively associated with hypertension awareness. Adults who were overweight and obese were more likely to be aware of their hypertension compared to those with normal weights. Factors associated with treatment of hypertension were similar to those related to hypertension awareness except for smoking status and drinking. Smoking status was not associated with treatment of hypertension. Current drinkers were less likely to receive antihypertensive medication (OR, 0.53; 95% CI, 0.39-0.72).

// Table 3 here //

4. Factors associated with hypertension control

The odds ratio (and 95% CI) of hypertension control was 1.32 (1.04-1.68) for women versus men and 1.84 (1.34-2.52) for older adults (≥ 65 years old) compared to young or mid-aged adults. People in urban areas (2.16; 1.79-2.60) and those from Eastern China (2.12; 1.72-2.63) were more likely to have controlled hypertension. In contrast, minority ethnicity (0.47; 0.31-0.73) and obese (0.77; 0.60-0.98) participants

were less likely to have their hypertension controlled. Among participants taking medication for hypertension, those in an urban area, located in Eastern China, and with higher educational attainment had higher odds of hypertension control, whereas minorities (0.52; 0.31-0.86) and obesity people (0.56; 0.42-0.76) were less likely to have their blood pressure controlled.

Discussion

This study provides important evidence about the burden of hypertension in China. Over a quarter (27.6%) of Chinese adults had hypertension, but only 55.7% of the hypertensive patients were aware of their disease, and 83% of those who were aware of their hypertension took antihypertensive medication treatment. Among those who were treated, only 43.8% had their hypertension controlled. Overall, only 20.3% of participants with hypertension achieved controlled blood pressure. Being female, of older age, a high socioeconomic status, and a healthy lifestyle were associated with a higher rate of hypertension awareness, treatment, and control. Low hypertension treatment and control rates and large variations of these rates across different subgroups highlight the urgent need for more effective and affordable strategies for hypertension control in China.

Our findings on hypertension awareness, treatment, and control are consistent with similar studies.(14) As shown in **Supplemental table 1**, we compared the prevalence, awareness, treatment, and control of hypertension among Chinese adults from two waves of CHNS (2001 and 2011) and from the International Collaborative Study of Cardiovascular Disease in ASIA (InterASIA 2000-2001). We would like to evidence what are the improvements in hypertension prevalence, awareness, treatment and control over ten years interval in the same survey and another coetaneous study. This comparison reveals that progress has been made in hypertension awareness, treatment, and control. For example, the hypertension awareness and treatment rates in the present study were more than two times greater than those seen a decade earlier(15) and the control rate was substantially higher than that reported in the InterASIA study.(16)

Despite the progress, awareness, treatment, and control of hypertension in China were still lower than the global average level,(17, 18) and they were much worse than those in developed countries.(19, 20) According to a systematic review based on data from 35 countries, the hypertension awareness and control rates in China were even lower than the average level of many developing countries.(21) The poor performance in hypertension control in China may be caused by, among others, unequal socioeconomic status, which limits access to basic health care facilities and primary health care practitioners, and the prevalence of related risk factors in the general population.(22, 23)

The low hypertension control rate may be impacted by two factors. First, BP screening was not included in many primary health care service practices yet, even though it has been required by government health programs. Second, the lack of appropriate antihypertensive medication may partly explain the low control rate. Therefore, the direct and intuitive implication would be that a more intensive education program and

interventions should be launched to strengthen outpatient and community-based screening for hypertension. In addition, more standardized and comprehensive hypertension treatment guidelines must be released to improve the knowledge of primary care physicians in China who prescribe evidence-based medications to obtain the optimal control of hypertension management.

Our multivariate logistic regression analysis identified several factors being associated with awareness, treatment, and control of hypertension in China. Consistent with other studies,(24-26) we found that women were more likely to be aware of their hypertension and receiving treatment than men. The control rate did not differ by sex among those who received medication treatment. In accordance with evidence from other studies,(8, 27, 28) urban residence and older age were associated with higher awareness, treatment, and control rates. A potential reason is that urban residents are more wary of their health, and so does the elderly population. Participants in Eastern China had higher rates of hypertension awareness, treatment, and control compared to those in Central China, which is likely due to the fact that Eastern China has an advantage in economic development and, thus, has more health care resources.(29, 30)

Being an ethnic minority was shown to be associated with lower rates of hypertension awareness, treatment, and control. This difference may be attributed to variations in genetics, culture, and lifestyles between ethnic minorities and the majority Han ethnicity. A higher education level was found to be associated with a higher hypertension control rate. This result was consistent with a study based on data collected from adults in Southern China,(31) but was different from what was reported by the CHPSNE study.(32) The disparity may reflect differences in the study sample and treatment of hypertension in these studies.

Our results showed that having overweight and obesity was positively associated with higher rates of hypertension awareness and treatment, which was consistent with several other studies.(33, 34) However, obesity was associated with a lower control rate among patients receiving medication treatment. This could be explained by poor adherence to medical treatment among subjects with a suboptimal body weight.(35, 36) In addition, our study found that several lifestyle factors were associated with hypertension outcomes. The awareness and control rates were greater among former smokers and lower among current smokers and drinkers. The reasons behind these associations require further investigation that is beyond the scope of this study.

Our study has several limitations. First, cross-sectional data analysis does not support causal inferences between risk factors and hypertension outcomes. Second, blood pressure was not measured on separate occasions. Since the definition of hypertension was partly based on the blood pressure measurement, the absolute hypertension burden might be overestimated. Third, the CHNS did not capture non-pharmacological treatment strategies such as lifestyle modifications.

In conclusion, this study adds to the growing evidence that China suffers from a serious hypertension burden. Hypertension awareness, treatment, and control rates are still low compared to global averages and even compared to the average rates in many developing countries. The situation is worse among certain population groups, such as those who live in rural areas and underdeveloped regions. Substantial

effort and resources—such as increasing physician-training programs and expanding screening services—are needed to alleviate the burden and hypertension and, eventually, improve population health in China.

Abbreviations

OR, odds ratio;

95% CI, confidence interval;

SD, standard deviation;

BMI, body mass index;

BP, blood pressure;

SBP, systolic blood pressure;

DBP, diastolic blood pressure;

HTN, hypertension;

CHNS, China Health and Nutrition Survey;

Declarations

Ethics approval and consent to participate

Written informed consent for CHNS was approved by the institutional review committees of the University of North Carolina at Chapel Hill and the national institute of Nutritional and Food safety, China Centre for Disease Control and Prevention.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests:

The authors declared that they have no competing interests.

Funding:

Part of Drs. Junxiang Wei and Youfa Wang's effort was supported by a research grant from the China Medical Board (Grant No. 16-262).

Author contributions

JW designed the overall study with contributions from YL. JW collected, analyzed data and drafted the manuscript. BX critically revised the manuscript. YL and YW interpreted the results and critically revised the manuscript. All authors approved the final manuscript.

Acknowledgments

We thank the National Institute of Nutrition and Food Safety, China Center of Disease Control and Prevention, and the Carolina Population Center at University of North Carolina at Chapel Hill for making the data available. The CHNS data collection was supported in part by research grants from the National Institute for Health (NIH) (R01-HD30880, R24 HD050924, R01-HD38700 and DK050924) and the NIH Fogarty International Center (D43 TW009077, D43 TW007709). Part of Drs. Junxiang Wei and Youfa Wang's effort in this project was supported by a research grant from the China Medical Board (Grant No. 16-262).

References

1. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet*. 2005;365(9455):217-23.
2. Dhillon RS, Clair K, Fraden M, Abdalla M. Hypertension in populations of different ethnic origins. *Lancet*. 2014;384(9939):234.
3. Hypertension: an urgent need for global control and prevention. *Lancet*. 2014;383(9932):1861.
4. Lewington S, Lacey B, Clarke R, Guo Y, Kong XL, Yang L, et al. The Burden of Hypertension and Associated Risk for Cardiovascular Mortality in China. *JAMA Intern Med*. 2016;176(4):524-32.
5. Gao Y, Chen G, Tian H, Lin L, Lu J, Weng J, et al. Prevalence of hypertension in china: a cross-sectional study. *PLoS One*. 2013;8(6):e65938.
6. Lu J, Lu Y, Wang X, Li X, Linderman GC, Wu C, et al. Prevalence, awareness, treatment, and control of hypertension in China: data from 1.7 million adults in a population-based screening study (China PEACE Million Persons Project). *Lancet*. 2017;390(10112):2549-58.
7. Hesketh T, Zhou X. Hypertension in China: the gap between policy and practice. *Lancet*. 2017;390(10112):2529-30.

8. Wang J, Zhang L, Wang F, Liu L, Wang H, China National Survey of Chronic Kidney Disease Working G. Prevalence, awareness, treatment, and control of hypertension in China: results from a national survey. *Am J Hypertens*. 2014;27(11):1355-61.
9. Meng Q, Fang H, Liu X, Yuan B, Xu J. Consolidating the social health insurance schemes in China: towards an equitable and efficient health system. *Lancet*. 2015;386(10002):1484-92.
10. Su M, Zhang Q, Bai X, Wu C, Li Y, Mossialos E, et al. Availability, cost, and prescription patterns of antihypertensive medications in primary health care in China: a nationwide cross-sectional survey. *Lancet*. 2017;390(10112):2559-68.
11. Zhang B, Zhai FY, Du SF, Popkin BM. The China Health and Nutrition Survey, 1989-2011. *Obes Rev*. 2014;15 Suppl 1:2-7.
12. Zhou BF, Cooperative Meta-Analysis Group of the Working Group on Obesity in C. Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults—study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Biomed Environ Sci*. 2002;15(1):83-96.
13. Du S, Batis C, Wang H, Zhang B, Zhang J, Popkin BM. Understanding the patterns and trends of sodium intake, potassium intake, and sodium to potassium ratio and their effect on hypertension in China. *The American journal of clinical nutrition*. 2014;99(2):334-43.
14. Li Y, Yang L, Wang L, Zhang M, Huang Z, Deng Q, et al. Burden of hypertension in China: A nationally representative survey of 174,621 adults. *Int J Cardiol*. 2017;227:516-23.
15. Wu Y, Huxley R, Li L, Anna V, Xie G, Yao C, et al. Prevalence, awareness, treatment, and control of hypertension in China: data from the China National Nutrition and Health Survey 2002. *Circulation*. 2008;118(25):2679-86.
16. Muntner P, Gu D, Wu X, Duan X, Wenqi G, Whelton PK, et al. Factors associated with hypertension awareness, treatment, and control in a representative sample of the chinese population. *Hypertension*. 2004;43(3):578-85.
17. Chow CK, Teo KK, Rangarajan S, Islam S, Gupta R, Avezum A, et al. Prevalence, awareness, treatment, and control of hypertension in rural and urban communities in high-, middle-, and low-income countries. *JAMA*. 2013;310(9):959-68.
18. Pereira M, Lunet N, Azevedo A, Barros H. Differences in prevalence, awareness, treatment and control of hypertension between developing and developed countries. *J Hypertens*. 2009;27(5):963-75.
19. Egan BM, Zhao Y, Axon RN. US trends in prevalence, awareness, treatment, and control of hypertension, 1988-2008. *JAMA*. 2010;303(20):2043-50.
20. Diederichs C, Neuhauser H. Regional variations in hypertension prevalence and management in Germany: results from the German Health Interview and Examination Survey (DEGS1). *J Hypertens*. 2014;32(7):1405-13; discussion 14.
21. Mohsen Ibrahim M. Hypertension in Developing Countries: A Major Challenge for the Future. *Curr Hypertens Rep*. 2018;20(5):38.

22. Grotto I, Huerta M, Sharabi Y. Hypertension and socioeconomic status. *Curr Opin Cardiol.* 2008;23(4):335-9.
23. Wang Y, Wang L, Qu W. New national data show alarming increase in obesity and noncommunicable chronic diseases in China. *Eur J Clin Nutr.* 2017;71(1):149-50.
24. Yang L, Yan J, Tang X, Xu X, Yu W, Wu H. Prevalence, Awareness, Treatment, Control and Risk Factors Associated with Hypertension among Adults in Southern China, 2013. *PloS one.* 2016;11(1):e0146181.
25. Meisinger C, Heier M, Volzke H, Lowel H, Mitusch R, Hense HW, et al. Regional disparities of hypertension prevalence and management within Germany. *J Hypertens.* 2006;24(2):293-9.
26. Wyatt SB, Akylbekova EL, Wofford MR, Coady SA, Walker ER, Andrew ME, et al. Prevalence, awareness, treatment, and control of hypertension in the Jackson Heart Study. *Hypertension.* 2008;51(3):650-6.
27. Liu X, Gu W, Li Z, Lei H, Li G, Huang W. Hypertension prevalence, awareness, treatment, control, and associated factors in Southwest China: an update. *J Hypertens.* 2017;35(3):637-44.
28. Lao XQ, Xu YJ, Wong MC, Zhang YH, Ma WJ, Xu XJ, et al. Hypertension prevalence, awareness, treatment, control and associated factors in a developing southern Chinese population: analysis of serial cross-sectional health survey data 2002-2010. *Am J Hypertens.* 2013;26(11):1335-45.
29. Li Y, Wang L, Feng X, Zhang M, Huang Z, Deng Q, et al. Geographical variations in hypertension prevalence, awareness, treatment and control in China: findings from a nationwide and provincially representative survey. *J Hypertens.* 2018;36(1):178-87.
30. Cai L, Liu A, Zhang L, Li S, Wang P. Prevalence, awareness, treatment, and control of hypertension among adults in Beijing, China. *Clin Exp Hypertens.* 2012;34(1):45-52.
31. Wang H, Zhang X, Zhang J, He Q, Hu R, Wang L, et al. Factors associated with prevalence, awareness, treatment and control of hypertension among adults in Southern China: a community-based, cross-sectional survey. *PloS one.* 2013;8(5):e62469.
32. Tian S, Dong GH, Wang D, Liu MM, Lin Q, Meng XJ, et al. Factors associated with prevalence, awareness, treatment and control of hypertension in urban adults from 33 communities in China: the CHPSNE Study. *Hypertens Res.* 2011;34(10):1087-92.
33. Ahn S, Zhao H, Smith ML, Ory MG, Phillips CD. BMI and lifestyle changes as correlates to changes in self-reported diagnosis of hypertension among older Chinese adults. *J Am Soc Hypertens.* 2011;5(1):21-30.
34. Dallongeville J, Bhatt DL, Steg PH, Ravnaud P, Wilson PW, Eagle KA, et al. Relation between body mass index, waist circumference, and cardiovascular outcomes in 19,579 diabetic patients with established vascular disease: the REACH Registry. *Eur J Prev Cardiol.* 2012;19(2):241-9.
35. Farah R, Zeidan RK, Chahine MN, Asmar R, Chahine R, Salameh P, et al. Predictors of Uncontrolled Blood Pressure in Treated Hypertensive Individuals: First Population-Based Study in Lebanon. *J Clin Hypertens (Greenwich).* 2016;18(9):871-7.

36. Bailey JE, Hajjar M, Shoib B, Tang J, Ray MM, Wan JY. Risk factors associated with antihypertensive medication nonadherence in a statewide Medicaid population. *Am J Med Sci.* 2014;348(5):410-5.

Tables

Table 1. Demographic characteristics of Chinese adults with hypertension based on China Health and Nutrition Survey 2011

	All	Women	Men	P-value**
Sample size (N)	3,579	1,775	1,804	
Age (year, mean±SD)	60.0±12.4	61.6±11.7	58.5±12.9	< 0.001
BMI (kg/m ²)	25.3±5.4	25.4±5.2	25.2±5.6	0.13
Residence (%)				
Urban	41.5	41.5	41.5	1.0
Rural	58.5	58.5	58.5	
Geographic region [†] (%)				0.33
Western China	20.1	21.1	19.1	
Eastern China	49.4	48.6	50.3	
Central China	30.5	30.3	30.6	
<u>Marital_status</u> * (%)				< 0.001
Married	83.2	76.0	90.3	
Single	16.8	24.0	9.7	
Ethnicity				1.0
Han	92.0	92.0	90.0	
Minority	8.0	8.0	8.0	
Occupational <u>status</u> (%)				< 0.001
Employed	44.1	32.1	56.0	
Unemployed	55.9	67.9	44.0	
Health insurance (%)				
Insured	95.8	95.9	95.7	0.82
Not insured	4.2	4.1	4.3	
Education (%)				< 0.001
≤ Elementary school	48.2	60.8	35.7	
Middle school	27.4	22.4	32.4	
High school	17.1	13.0	21.1	
≥ College	7.3	3.8	10.8	
Smoking (%)				< 0.001
Non-smoker	65.9	94.3	38.0	

Ex-smoker	6.4	0.8	11.8	
Smoker	27.7	4.9	50.2	
Alcohol consumption (%)				< 0.001
Drinking	34.5	8.7	59.9	
Not drinking	65.5	91.3	40.1	
Weight status (%)				0.01
Normal weight (18.5≤BMI<24)	36.4	36.0	36.8	
Underweight (BMI<18.5)	2.2	2.3	2.0	
Overweight (24≤BMI<28)	41.0	39.1	42.8	
Obese (BMI≥28)	20.5	22.6	18.4	

Hypertension was defined as having an average systolic BP \geq 140 mmHg, diastolic BP \geq 90mmHg, self-reported being previously diagnosed as hypertension by a physician or taking antihypertension drugs currently;

*Single includes never married, divorced, widowed and separated;

†Western China includes Guangxi, Guizhou and Chongqing; Eastern China includes Shanghai, Beijing, Jiangsu, Liaoning and Shandong; Central China includes Henan, Hubei, Henan and Heilongjiang;

***p*-value was calculated from t-test for continuous variables and *chi*-square test for categorical variables.

Table 2. Proportion[¶] (%) of hypertension awareness, treatment, and control among Chinese adults with hypertension (HTN) based on CHNS 2011

	Awareness	Treatment		Control	
	(1,993)	(1,664)		(728)	
		HTN	Awareness	HTN	Treated
Age group (years)					
18-49	36.2	25.9	71.7	12.7	49.0
50-64	58.1**	48.6**	83.6**	22.2**	45.7
65+	64.8**	56.8**	87.6**	22.8**	32.9
Residence					
Rural	48.0	38.4	80.0	13.5	35.2
Urban	66.6**	58.0**	87.0**	30.0**	51.7**
Geographic region [†]					
Central China	50.1	39.6	79.0	13.3	33.6
Western China	47.2	39.1	82.9	15.7	40.2
Eastern China	62.6**	53.8**	85.1**	26.6**	49.4**
Ethnicity					
Han	56.7	47.5	83.7	21.3	44.9
Minority	44.1**	35.0**	79.4	9.1**	26.0 [‡]
Marital status*					
Unmarried	58.9	49.8	84.5	19.6	39.5
Single	55.0	45.8	83.3	20.5	44.7
Occupational status					
Unemployed	64.5	56.4	87.4	24.4	43.3
Employed	44.6**	34.0**	76.3**	15.2**	44.7
Health insurance					
Not insured	51.3	42.0	81.8	19.3	46.0
Insured	55.9	46.7	83.6	20.4	43.7
Education					
≤ Elementary school	54.2	45.0	82.9	16.0	35.6
Middle school	52.9	43.8	82.9	21.2 [‡]	48.4**

High school	60.0 [§]	51.6 [‡]	86.1	27.1 ^{**}	52.5 ^{**}
≥ Collage	65.9 ^{**}	54.8 [‡]	83.1	29.9 ^{**}	54.5 ^{**}
Smoking					
Non-smoker	58.4	50.1	85.7	21.7	43.3
Ex-smoker	68.4 [‡]	55.3	80.8	28.5 [§]	51.6
Smoker	46.3 ^{**}	36.0 ^{**}	77.8 ^{**}	15.3 ^{**}	42.6
Alcohol consumption					
Not drinking	58.7	50.9	86.8	21.5	42.1
Drinking	50.0 ^{**}	38.1 ^{**}	76.0 ^{**}	18.2 [§]	47.9
Weight status					
Normal weight (BMI<24)	50.7	40.5	79.8	20.2	50.0
Overweight (24≤BMI<28)	58.6 ^{**}	49.6 ^{**}	84.8 [§]	21.8	43.8
Obese (BMI≥28)	59.4 ^{**}	51.6 ^{**}	83.5 [‡]	17.7	34.4 ^{**}

[‡]3,579 adults with hypertension;

Hypertension was defined as having an average systolic BP ≥140 mmHg, diastolic BP ≥ 90 mmHg, self-reported being previously diagnosed as hypertension by a physician or taking antihypertension drugs currently;

Awareness of hypertension was defined as a self-report of hypertension diagnosed by a doctor before, or taking any antihypertensive drugs;

Control of hypertension was defined as blood pressure (SBP/DBP) < 140/90 mmHg;

*Single includes never married, divorced, widowed and separated;

[†]Western China includes Guangxi, Guizhou and Chongqing; Eastern China includes Shanghai, Beijing, Jiangsu, Liaoning and Shandong; Central China includes Henan, Hubei, Henan and Heilongjiang;

[§]p<0.05, [‡]p<0.01, ^{**}p<0.001.

Table 3. Multivariable logistic regression analysis of factors associated with hypertension awareness, treatment and control among Chinese adults with hypertension (n=3,579) in CHNS 2011

	Awareness	Treatment [¶]	Control	Control [‡]
Women (vs. men)	1.37 (1.12 - 1.66)	0.94 (0.65 - 1.34)	1.32 (1.04 - 1.68)	1.17 (0.88 - 1.57)
Age group (years)				
18-49 (ref)				
50-64	2.34 (1.92 - 2.85)	1.87 (1.32 - 2.65)	1.87 (1.43 - 2.44)	1.33 (0.92 - 1.92)
65+	3.13 (2.45 - 3.99)	2.57 (1.65 - 4.02)	1.84 (1.34 - 2.52)	1.06 (0.70 - 1.62)
Residence				
Urban (vs. rural)	1.75 (1.50 - 2.04)	1.50 (1.14 - 1.97)	2.16 (1.79 - 2.60)	1.79 (1.43 - 2.24)
Geographic region [†]				
Central (ref)				
Western	0.92 (0.75 - 1.12)	1.35 (0.93 - 1.96)	1.33 (1.01 - 1.75)	1.33 (0.95 - 1.88)
Eastern	1.50 (1.27 - 1.77)	1.52 (1.14 - 2.01)	2.12 (1.72 - 2.63)	1.78 (1.37 - 2.31)
Ethnicity (vs. Han)				
Minority	0.74 (0.57 - 0.96)	0.78 (0.49 - 1.26)	0.47 (0.31 - 0.73)	0.52 (0.31 - 0.86)
Occupational status (vs. unemployed)				
Employed	0.78 (0.66 - 0.93)	0.74 (0.54 - 1.00)	0.92 (0.74 - 1.15)	0.96 (0.72 - 1.26)
Marital status* (vs. single)				
Married	1.09 (0.89 - 1.34)	1.27 (0.89 - 1.83)	1.11 (0.87 - 1.43)	1.16 (0.86 - 1.57)
Health insurance (vs. not insured)				
Insured	1.32 (0.93 - 1.87)	1.44 (0.77 - 2.68)	1.08 (0.70 - 1.66)	0.90 (0.52 - 1.55)
Weight status				
Normal weight ((BMI<24) (ref)				
Overweight (24≤BMI<28)	1.41 (1.20 - 1.66)	1.52 (1.15 - 2.01)	0.96 (0.79 - 1.16)	0.83 (0.65 - 1.05)
Obese (BMI≥28)	1.57 (1.29 - 1.92)	1.99 (1.39 - 2.84)	0.77 (0.60 - 0.98)	0.56 (0.42 - 0.76)
Education				
≤ Elementary school (ref)				
Middle school	1.11 (0.92 - 1.33)	1.08 (0.78 - 1.50)	1.39 (1.11 - 1.74)	1.57 (1.20 - 2.06)
High school	1.34 (1.07 - 1.67)	1.42 (0.97 - 2.08)	1.65 (1.28 - 2.12)	1.53 (1.13 - 2.07)
≥ College	1.69 (1.23 - 2.32)	1.04 (0.64 - 1.71)	1.77 (1.26 - 2.48)	1.80 (1.20 - 2.70)

Smoking status				
Non-smoker (ref)				
Ex-smoker	1.62 (1.16 - 2.24)	0.75 (0.46 - 1.23)	1.47 (1.04 - 2.09)	1.23 (0.80 - 1.88)
Smoker	0.86 (0.70 - 1.04)	0.83 (0.59 - 1.18)	0.81 (0.63 - 1.05)	0.83 (0.61 - 1.14)
Alcohol consumption				
Drinking	0.95 (0.79 - 1.13)	0.53 (0.39 - 0.72)	0.96 (0.77 - 1.19)	1.03 (0.78 - 1.35)
(vs. not drinking)				

[†]Treatment among subjects who were aware of their hypertension;

[‡]control among participant who took antihypertension medications;

Hypertension was defined as having an average systolic BP \geq 140 mmHg, diastolic BP \geq 90mmHg, self-reported being previously diagnosed as hypertension by a physician or taking antihypertension drugs currently;

Awareness of hypertension was defined as a self-report of hypertension diagnosed by a Physician before, or taking any antihypertensive drugs;

Control of hypertension was defined as blood pressure (SBP/DBP) < 140/90 mmHg;

*Single includes never married, divorced, widowed and separated;

[†]Western China includes Guangxi, Guizhou and Chongqing; Eastern China includes Shanghai, Beijing, Jiangsu, Liaoning and Shandong; Central China includes Henan, Hubei, Henan and Heilongjiang.

Figures




Figure 1

Rates of hypertension awareness, treatment, and control among Chinese adults with hypertension (n=3,579) by sex based on CHNS 2011

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaltable1.docx](#)