

## Lactoferrin feeding deficiency during lactation increases the risk of depression in adult mice by affecting microbiota-gut-brain axis development

Wenli Wang China Agricultural University Zhimei Cheng China Agricultural University Xiong Wang China Agricultural University An Oin China Agricultural University **Kunlun Huang** China Agricultural University Yunping Dai China Agricultural University Qingyong Meng China Agricultural University yali zhang (Zzhangyali@cau.edu.cn) China Agricultural University https://orcid.org/0000-0002-5354-1973

#### **Research Article**

Keywords: lactation, lactoferrin, depression, microbial-intestinal-brain, innate immunity, inflammation

Posted Date: December 13th, 2022

DOI: https://doi.org/10.21203/rs.3.rs-1883790/v2

**License:** (a) This work is licensed under a Creative Commons Attribution 4.0 International License. Read Full License

## Abstract Background

Lactoferrin is an active protein in breast milk that plays an important role in the growth and development of infants and is implicated as a neuroprotective agent. The incidence of depression is currently increasing, and it is unclear whether the lack of lactoferrin during lactation affects the incidence of depression in adulthood.

## Objectives

The purpose of our work was to investigate the effect of lactoferrin feeding deficiency during lactation on adult depression and related mechanisms.

## Methods

Lactoferrin knockout mice provided lactoferrin-free milk to establish a lactoferrin feeding deficiency model during lactation. The chronic unpredictable mild stress (CUMS) depression model was established when the mice reached 9 weeks of age. Behavioural performance, serum indicators, and the microbiota, intestine, and hippocampus injury of the CUMS mice were investigated; RNA-seq of the small intestine and hippocampus, and gut microbiota compositions in 18-day-old lactoferrin-free mice were analysed; the development of the intestine and hippocampus and microbiota compositions in adult lactoferrin-free mice were also examined.

## Results

Lack of lactoferrin feeding during lactation affected the barrier and innate immune functions of the intestine, disrupted the intestinal microflora, and led to neuroimmune dysfunction and neurodevelopmental delay in the hippocampus. When exposed to external stimulation, adult lactoferrin feeding-deficient mice presented with worse depression-like symptoms; the mechanisms involved were activation of the LPS-TLR4 signalling pathway in the intestine and hippocampus, reduced BDNF-CREB signalling pathway in hippocampus, increased abundance of depression-related bacteria, and decreased abundance of beneficial bacteria.

# Conclusions

Overall, our findings reveal that feeding lactoferrin during lactation can reduce the risk of depression in adults. The mechanism is related to the regulatory effect of lactoferrin on the development of the "microbial-intestinal-brain" axis.

### Introduction

Depressive disorders are a global public health concern and a leading cause of disease burden, substantially impairing patients' quality of life and increasing mortality risk. According to the World Health Organization (WHO), the number of people affected by depression increased by 18% worldwide in the decade from 2005 to 2015. In the first year of the COVID-19 pandemic, the global prevalence of anxiety and depression increased by a massive 25%, according to a scientific brief released by the WHO. Depression is caused by complex interactions of multiple factors. The pathogenesis is not fully understood, although many hypotheses have been proposed, such as the inflammatory hypothesis [1], "leaky gut" hypothesis [2], gut microbiota hypothesis [3], and monoamine hypothesis [4]. Biochemical signals that occur between the gastrointestinal tract and central nervous system are known as the gut brain axis. The bidirectional communication between the brain and gut occurs through the autonomic, enteric, neuroendocrine, and immune systems [3]. Studies of the microbiota–gut–brain axis may lead to novel approaches for preventing and treating mental illnesses, including anxiety and depression.

Immune system disorders caused by impaired gastrointestinal barrier integrity are considered as important mechanisms of depression [5]. The intestinal epithelial barrier serves as the first boundary of defence between the organism and luminal environment [6]. Increased intestinal permeability caused by intestinal barrier injury leads to systemic chronic inflammation, whereby nervous system inflammation is an important pathological feature of depression. Lipopolysaccharide (LPS) from bacteria in the intestinal lumen, mainly gram-negative bacteria, enters the systemic circulation through the increased intercellular space and cross-cellular pathways [7] and, LPS binds to TLR-4 in the central nervous system, thus triggering neuroinflammation [8]. Depression is a common comorbidity associated with gastrointestinal diseases, such as irritable bowel syndrome and inflammatory bowel diseases (IBDs) [9]. A previous study showed that chronic gastrointestinal inflammation induces anxiety-like behaviour and alters the central nervous system biochemistry [10]. A population- based study revealed that individuals with IBD had a higher prevalence of depression than matched controls [11]. Gampierakis et al. showed that acute and chronic experimental colitis affects adult hippocampal neurogenesis and innate immune cell responses, which may be mechanisms of cognitive and mood dysfunction in patients with IBD [12].

The gut microbiota is a key regulator of the gut-brain axis and influences brain function and behaviour through the microbiota-gut-brain axis. Many antibodies against intestinal symbiotic bacteria LPS have been detected in the serum of patients with depression, supporting the notion that inflammation is related to intestinal symbiotic bacteria [2]. Animal experiments add further support to this idea; interestingly, faecal microbiota transplantation from depressed patients to microbiota-depleted rats induced anxiety and depressive behaviours and altered tryptophan metabolism in recipient animals [13]. Furthermore, bacterial species regulate the production of neurotransmitters and their precursors, such as serotonin, γ-aminobutyric acid type A (GABA), and tryptophan [14]. Gut microbes also promote the release of short-chain fatty acids (SCFAs) and brain-derived nutritional factors (BDNFs), which have been shown to ameliorate depression [10].

Lactation is a critical period for the development of the intestinal tract and nervous system and the establishment of the gut microbiota; breast milk plays important roles in these processes. However, many newborns cannot be breastfed due to the nutritional and health status of their mothers, social factors, and uncertain circumstances. In low- and middle-income countries, only 37% of children younger than six months of age are exclusively breastfed; this rate is even lower in high-income countries [15]. Lactoferrin (LF) is an important active whey protein in breast milk present at concentrations of 1-7 and 0.2-1.5 g/L in humans and bovines, respectively [16]. LF plays an important role in neonatal development by promoting maturation of the neonatal small intestine and increasing the integrity of the intestinal barrier [17]. Several studies have shown that LF can improve neural development, neuroprotection, and cognitive abilities [18]. The early gut microbiota has been predicted to contribute to disease progression later in life, and the foundation for a stable adult gut microbiota has already been established in infancy [19]. According to a previous study, early life LF intervention can modulate the microbial community of the cecum [20]; however, whether the effects of LF on the microbiota of newborns affect adult depression remains unclear. Formula is an important alternative to breast milk, but the addition of LF to formula is not widespread. Thus, considering that exclusive breastfeeding rates are < 50% in most countries, LF deficiency during lactation may be prevalent worldwide. The effect of LF feeding deficiency during lactation on adult health, particularly depression, remains unknown.

In this study, we generated an LF-knockout (KO) mouse model as a mother mouse that provided LF-free milk. Two groups of mice were evaluated: wild-type (WT) neonates breast-fed by WT or KO female mice (WT-WT or KO-WT). The chronic unpredictable mild stress (CUMS) depression model was established when the mice reached 9 weeks of age, and differences in depressive phenotypes were compared among groups. To investigate the mechanism by which LF feeding deficiency during lactation increases the risk of depression in adult mice, the intestinal and brain development and intestinal microbe composition at different growth stages of mice were investigated.

### Material And Methods

# Animal

The heterozygous LF gene knockout mice were customized from Biocytogen Co., Ltd. (Beijing, China). Heterozygous breeding produced *Ltf* gene knockout (KO) and wild-type littermate (WT) C57BL/6N mice, which were used for homozygous breeding to produce a larger number of age-matched WT and KO mice for the current experiments. Before the experiment, KO and WT homozygous breeding produced mice at the same time. On the second day after birth, all litters born to KO dams were replaced with pups of the same age born to WT dams (KO-WT group). Control litters suckling WT dams were also standardized to pups on the second day after birth (WT-WT group). The mice in KO-WT group drunk LF-free milk during sucking time, and these two group mice would be used in subsequent experiments.

Mice were housed in a controlled environment (12:12 h light: dark cycle; temperature: 22 ± 2°C; humidity: 60%; 15 – 20 fresh air changes per hour) with free access to food and water. All mice were treated in

accordance with the guidelines of the Institutional Animal Care and Use Committee (SYXK 2020-0052). All experiments were approved by the Animal Experimentation Committee of the China Agricultural University (Beijing, China).

# Procedure of CUMS

16 male 9-week-old mice (8 from WT-WT group and 8 from KO-WT group) and 16 female 9-week-old mice (8 from WT-WT group and 8 from KO-WT group), were divided into 4 groups: wt-wtM, ko-wtM, wt-wtF, ko-wtF. The CUMS procedure was a variation of methods described by Agnieszka [21]. The list of stressors, their details, and the day they were applied was given in Table. S1.

## **Behavioral evaluations**

# Open field test (OFT)

The test consisted of one 5 min trial in a white opaque 50 cm × 50 cm × 50 cm arena. The center zone was defined as a 25 cm × 25 cm central square. To begin each test, a mouse was introduced to the center of the square and its behavior was captured on video (Samsung, South Korea) over the course of 5min, the area was cleaned with 50% ethanol and allowed to dry completely between each test. The distance that each mouse walked in either the peripheral or central regions was quantified using KEmaze software. **Sucrose preference test (SPT)** 

The sucrose preference test was performed according to a previous report [22] with minor modifications. Before the test, mice were acclimatized to sucrose solution. At first, two bottles of 1% sucrose solution (w/v) were placed in each cage for 24 h to avoid sucrose neophobia. After adaptation, mice were deprived of liquid and food for 24 h and then given the sucrose preference test. Each mouse had free access to 2 bottles for 24 h: one bottle with 1% sucrose solution (w/v) and another bottle with tap water. To avoid position effects, the positions of the 2 bottles were reversed at intervals of 12 h. Sucrose consumption and water consumption were measured by comparing the weights of the bottles before and after the test. The sucrose preference was calculated as sucrose preference (%) = (sucrose solution intake)/ (sucrose solution intake + tap water intake) \*100%.

# Forced swimming test (FST)

Each mouse was placed in an open cylindrical container (diameter 10 cm, height 30 cm) with 23 ± 1°C water to a depth of 20 cm and allowed to swim for 6 min. Immobility times were measured during the last 4 min of the test. During the test, the behavior of each mouse was recorded using a video camera (Samsung, South Korea), and immobility time was measured using SuperFst software (KEWBIO Co., Ltd. Nanjing, China).

# Tail suspension test (TST)

Mice were habituated to the testing room for 30 min before the experiments. Animals were attached by their tails to a shelf with medical adhesive tape (placed approximately 1 cm from the distal end of the

tail). The last 4 min was recorded for immobility time during a 6-min test. During the test, the behavior of each mouse was recorded using a video camera (Samsung, South Korea), and immobility time was measured using SuperTst software (KEWBIO Co., Ltd. Nanjing, China).

# Dextran Sodium Sulfate (DSS)-induced colitis

Colitis was induced by the administration of 2.5% w/v DSS (MW = 36,000–50,000 kDa; MP Biomedicals) in the drinking water for 7 days [12]. The 8-week-old female mice (fDWT-WT, fDKO-WT) and male mice (mDWT-WT, mDKO-WT) were weighed daily and the disease activity index (DAI) score can be assessed. The DAI is the combined score of weight loss compared to initial weight, stool consistency and bleeding [23].

## Sample Collection and Measurement

After 4 weeks of CUMS, all animals were sacrificed under anesthesia with 10% chloral hydrate and the colon tissue, hippocampus, serum and cecal content of all animals were collected and stored at – 80°C for the following experiments. Blood was collected from the retro-orbital sinus under 10% chloral hydrate general anesthesia. Plasma was obtained from the whole blood by centrifugation at 5000 rpm for 20 min. Collected colon samples were fixed in 4% paraformaldehyde and embedded in paraffin. Sections of the colon were stained with H.E.

18-day-old mice (wt-wt, ko-wt) were sacrificed under anesthesia with isoflurane without fasting. The body weight, whole brain, small intestinal and colon were weighted, length of small intestinal and colon were measured. The small intestinal, colon, hippocampus and cecal content were collected and immediately snapfrozen in liquid nitrogen. Collected colon and hippocampus samples were fixed in 4% paraformaldehyde and embedded in paraffin. Sections of the colon were stained with H.E and alcian blue (AB) staining by using commercial kits (Beijing Solarbio Technology Co., Ltd. Beijing, China) according to the manufacturer's instructions. Sections of the hippocampus were stained with H.E.

9 weeks female mice (fWT-WT, fKO-WT) and male mice (mWT-WT, mKO-WT) were sacrificed under anesthesia with 10% chloral hydrate. The small intestine and colon were removed, and total length and weight were measured. Intestinal density is indices of intestinal mucosal growth. Intestinal density = weight of small intestine or colon/length of small intestine or colon. The small intestinal, half of colon, hippocampus and cecal content were collected and immediately snap-frozen in liquid nitrogen. The other half of colon was fixed in 4% buffered formaldehyde for the histological observation. The activities of lactase and maltase in each section of the small intestine were measured using commercial kits (Nanjing Jiancheng Institute of Bioengineering, Jiangsu, China) according to manufacturer's instructions. Sections of the intestinal were stained with H.E.

The DSS induced colitis mice (mDWT-WT, mDKO-WT, fDWT-WT, fDKO-WT) were sacrificed on day 7 and colon length was measured. Half of colon and cecal content of all animals were collected and stored at – 80°C for following experiments. The other half of colon was fixed in 4% buffered formaldehyde for the histological observation. Cytokine levels in colonic tissues were detected by enzyme-linked

# immunosorbent assay (ELISA) kit (Cloud-Clone, Wuhan, China) according to the manufacturer's instructions.

# **RNA sequencing**

The small intestinal and hippocampus of 18-day-old mice (wt-wt, ko-wt) were collected and the total RNA was extracted by using TRIZOL reagent (Invitrogen, USA). The concentration and purity of the RNA were detected by Nanodrop2000, RNA integrity was detected by agarose gel electrophoresis, the RIN value was measured by Agilent2100. The construction of a single database requires that the total amount of RNA  $\geq$  1µg, concentration  $\geq$  35 ng/µl, OD260/280 = 1.8 ~ 2.2, OD260/230  $\geq$  2.0.RIN  $\geq$  6.5, 28S:18S  $\geq$  1.0, Libraries were constructed using the Illumina TruseqTM RNA sample prep Kit (Illumina, San Diego, CA, USA) according to the manufacturer's instructions.

Sequencing of the libraries was performed on an Illumina HiSeq2000 instrument by Shanghai Majorbio Biopharm Biotechnology (Shanghai, China), and individually assessed for quality using FastQC. To identify DEGs between two different samples, the expression level of each transcript was calculated according to the transcripts per million reads (TPM) method. RSEM

(http://deweylab.biostat.wisc.edu/rsem/) was used to quantify gene abundances. Analysis of differential expression was carried out using edgeR. Statistical significance was assessed using a negative binomial Wald test, then corrected for multiple hypothesis testing with the Benjamini-Hochberg method.

## Quantitative real-time PCR

Total RNA was extracted from the tissue using an RNA extraction kit (Magen, Guangzhou, China) according to the manufacturer's instructions. The purity and concentration of RNA were determined using a NanoDrop spectrophotometer (Thermo Fisher, Waltham, MA). One microgram of total RNA was reverse transcribed into cDNA using a HiFiScript cDNA synthesis kit (NUOWEIZAN, Nanjing, China). RT-PCR was performed using Rotor-Gene Q (Qiagen, Hilden, Germany) with an Taq Pro Universal SYBR qPCR Master Mix Kit (NUOWEIZAN, Nanjing, China). RT-PCR amplification conditions included denaturation at 95°C for 10 s, annealing at 55°C for 30 s, and extension for 32 s at 72°C. Geometric mean glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used for normalization. All primer pairs were synthesized by Synbio Technologies (Jiangsu, China), the sequence was showed in Table 1.

Gene name	F-primer (5'-3')	R-primer (5'-3')
Gapdh	TCTCCTGCGACTTCAACA	TGTAGCCGTATT CATTGTCA
ZO-1	GCATCATTCGCCTTCATAC	GACACAACCTCATCCTCAT
occludin	GGCTGCTGCTGATGAATA	ATCCTCTTGATGTGCGATAA
TLR4	CACTGTTCTTCTCCTGCCTGA	GGAATGTCATCAGGGACTTTGC
LBP	GTCGTGGGCAGTACGAGTTT	CCTTCCATTTGCCTCGGACA
MyD88	CGCATGGTGGTGGTTGTTTC	AGTCGCTTCTGTTGGACACC
CD14	ACAATTCACTGCGGGATGCT	AGCTCATCTGGGCTAGGGTT
JNK	CCACCAAAGATCCCGGACAA	GGCTGCCCTCTTATGACTCC
ΝϜκΒ	CCCTACGGAACTGGGCAAAT	GCAAATTTTGACCTGTGGGT
ІкВа	CCTGACCTGGTTTCGCTCTT	CTGTATCCGGGTACTTGGGC
P38MAPK	TTCTACCGGCAGGAGCTGAA	ATCAAAAGCAGCACACACCG
IL-1β	CTTCAGGCAGGCAGTATC	CAGCAGGTTATCATCATCATC
TNF-a	ACTGAACTTCGGGGTGATCG	CCACTTGGTGGTTTGTGAGTG
BDNF	TAAACGTCCACGGACAAGGC	AGTGTCAGCCAGTGATGTCG
GRB2	AACATCCGTGTCCAGGAACC	AAGTCTCCTCTGCGAAAGCC
CaMK1	TGATCCTGGCAGAGGACAAGA	GCTACAATGTTGGGGTGCTTG
CREB	AACCAGCAGAGTGGAGATGC	GATGTTGCATGAGCTGCTGG
Lrg1	TGTCCAGCCTCAAGGAATGC	TTCCACCGACAGATGGACAG
Gbp4	TGAACCAGGAAGCCATAGAGA	GGGAAACCTTTGGCTGGTAGG
Stx11	ATGACTTTGACGCTCCTCGG	GCGCCTCACGTCTATCAGAA
Pigr	GACTCTCGCTGGAGAACCAC	ACGGATAGTGGCAGGAAACG
Ltf	AGCTGAAGTCTACGGGACCA	CTCAGGCCTTGGAGTTGGTT
Lcn2	ACAACCAGTTCGCCATGGTAT	AAGCGGGTGAAACGTTCCTT
S100a9	ACCACCATCATCGACACCTTC	AAAGGTTGCCAACTGTGCTTC
Ccl28	CTCACACTCATGGCTGTGGCT	AGTACGATTGTGCGGGCTGA
IL-25	GACCTGTACCACGCTCGATG	AGAAGACCGTCTGGTTGTGG
Ccn5	TGTGTGACCAGGCAGTGATG	GTGCTCCAGTTTGGACAGGG

Table 1 Sequences of primers for real-time PCR

Gene name	F-primer (5'-3')	R-primer (5'-3')
Cdh3	GCATGCACCATGCAGACAAT	GTGGCATCACCCACTCTCTC
Prdm12	TCAAGTGTGCCCGGAATGAG	TCCTGGTCTGGAGGGATCAT
Pcdha8	TCAAAGGGTTTTGGGGACCT	GAGCCTTTTGTGTCGCTGGA
Trpv4	TGGAACCAGAACTTGGGCAT	GGACCAACGATCCCTACGAA
Gabra6	AGGAGTCAGTCCCAGCAAGA	ATGAACCAATCCATGGCGGT
Kcne2	CCAGAGTGGATGCCGAGAAC	CTTCGACTTCACCGTGCTCA
Kcnj13	TTTGTGGCGAAGATTGCACG	ACACGAACGTTGGTCAGAGG

## 16S rRNA gene sequencing and analysis

Microbial community genomic DNA was extracted from cecal content samples using the E.Z.N.A.® soil DNA Kit (Omega Bio-Tek, Norcross, GA, USA) according to the manufacturer's instructions. The DNA extract was checked on 1% agarose gel, and DNA concentration and purity were determined with NanoDrop 2000 UV-vis spectrophotometer (Thermo Scientific, Wilmington, USA). The hypervariable region V3-V4 of the bacterial 16S rRNA gene were amplified. The PCR product was extracted from 2% agarose gel and purified using the AxyPrep DNA Gel Extraction Kit (Axygen Biosciences, Union City, CA, USA) according to the manufacturer's instructions and quantified using Quantus<sup>™</sup> Fluorometer (Promega, USA).

Purified amplicons were pooled in equimolar and paired-end sequenced on an Illumina MiSeq PE300 platform/NovaSeq PE250 platform (Illumina, San Diego, USA) according to the standard protocols by Majorbio Bio-Pharm Technology Co. Ltd. (Shanghai, China).

The raw 16S rRNA gene sequencing reads were demultiplexed, quality-filtered by fast version 0.20.0 and merged by FLASH version 1.2.7. Operational taxonomic units (OTUs) with a 97% similarity cutoff were clustered using UPARSE version 7.1, and chimeric sequences were identified and removed. The taxonomy of each OTU representative sequence was analyzed by RDP Classifier version 2.2 against the 16S rRNA database using a confidence threshold of 0.7.

## Statistical analysis

The results are expressed as mean ± SEM. Statistical analyses were performed using Statistical analysis was performed using Prism 8.0 (GraphPad Software, CA). Student's t-test was used to compare two groups, and an ANOVA and post hoc Duncan test were used to compare more than two groups. Statistical significance was set at P < 0.05.

### Results

# LF feeding deficiency during suckling period increased depression risk in adult mice

Male and female mice that consumed normal milk and LF-free milk during the suckling period were fed a normal diet until 9 weeks of age, after which, the CUMS model was established for 4 weeks (Fig. 1A). As shown in Fig. 1B and 1C, wt-wt and ko-wt mice did not differ before CUMS; however, after 4 weeks of CUMS treatment, mice without LF intake during the suckling period (ko-wtM and ko-wtF) showed a lower sucrose preference in the SPT. In the TST (Fig. 1D), there was no significant difference between the wtwtM and ko-wtM groups during the first week of CUMS. At week 4, the immobility time of the wt-wtM and ko-wtM groups was higher than that of week 1, and the immobility time of the ko-wtM group was longer than that of the wt-wtM group. Although a similar trend was observed in the female group (Fig. 1E), no significant difference was observed between the wt-wtF and ko-wtF groups at weeks 1 and 4. In the FST (Fig. 1F), there was no difference between wt-wtM and ko-wtM mice at week 1; at week 4, the immobility time of the wt-wtM and ko-wtM groups increased compared with that of week 1, and the immobility time of the ko-wtM mice was longer than that of the wt-wtM mice. Figure 1G shows that the immobility time in the FST of the ko-wt female mice (ko-wtF) was significantly longer than that of the wt-wtF group at week 1, and increased compared with that of the wt-wtF mice at week 4, but the difference was not significant. In the OFT, the distance in the central area for the ko-wtM group was significantly smaller than that for the wt-wtM group, with no difference between the ko-wtF and wt-wtF groups (Fig. 1H). Male mice with LF feeding deficiency during lactation showed more severe anxiety-like behaviours in the OFT compared to male mice that drank normal milk. Figure 1H also shows that the distance in the central area for the wtwtF group was significantly lower than that for the wt-wtM group, indicating that female mice exhibited more severe anxiety-like behaviours compared to male mice.

Serum indicators associated with depression were measured. BDNF is a well-known growth factor in the brain, and the level of BDNF in the ko-wt group was significantly lower than that in the wt-wt group, for both male and female mice (ko-wtM vs. wt-wtM; ko-wtF vs. wt-wtF). The levels of BDNF in female mice were significantly lower than those in male mice (wt-wtF vs. wt-wtM; ko-wtF vs. ko-wtM) (Fig. 1). Corticosterone (CORT) and adrenocorticotropic hormone (ACTH) are hormones related to the hypothalamic–pituitary–adrenal axis and are associated with depression. Our results revealed a significantly increased production of CORT and ACTH in the ko-wt mice compared to that in the wt-wt mice (ko-wtM vs. wt-wtM; ko-wtF vs. wt-wtF) (Fig. 1J, 1K); the wt-wtF group showed significantly higher CORT levels than the wt-wtM group. The levels of pro-inflammatory cytokines, such as tumour necrosis factor (TNF)- $\alpha$  (Fig. 1L) and interleukin (IL)-1 $\beta$  (Fig. 1M), in the ko-wt group were significantly higher than those in the wt-wtM group. Additionally, LPS production was significantly increased in the ko-wt mice serum (ko-wtM vs. wt-wtM; ko-wtF vs. wt-wtF) (Fig. 1N), which can induce inflammation. Furthermore, the LPS levels in the ko-wtF group were significantly higher than those in the ko-wtM group.

# CUMS mice with lactation LF feeding deficiency showed intestinal, brain and microbial flora disorders

Dysfunction of the microbiota-gut-brain axis is thought to be the main pathological basis of depression. We compared the histological changes in the colon, inflammation of the colon and hippocampus and composition of intestinal microorganisms in depressed mice with LF feeding deficiency during lactation with those in depressed mice that consumed normal mouse milk during lactation. As shown in Fig. 2A, the crypt depth of ko-wtM mice was significantly lower than that of wt-wtM mice, there was no significant difference in colonic crypt depth between the wt-wtF and ko-wtF groups. Damage to the intestinal epithelial barrier increased the translocation of this luminal LPS to systemic circulation, and expression of the zonula occludens gene (*zo-1*) in ko-wt mice was significantly lower than that in wt-wt mice (ko-wtM vs wt-wtM, analysis of variance, P < 0.05; ko-wtF vs wt-wtF, t test, P < 0.05). *Occludin* expression in ko-wtM mice was lower than that in wt-wtM mice but the difference was not significant (Fig. 2B). *Occludin* expression in wt-wtF was significantly lower than that in wt-wtM. The lactating LF feeding deficient mice showed more severe damage to their intestinal barrier integrity after CUMS, which may explain the increased LPS concentration in the serum of the ko-wtF and ko-wtM groups. Moreover, the intestinal barrier function was more severely damaged in female mice than in male mice.

As shown in Fig. 2C, the colons of the ko-wtM mice showed more severe inflammatory infiltration than that of wt-wtM mice; however, among females, inflammatory infiltration was more severe in wt-wtF mice than that in ko-wtF mice. We further examined the activation of the LPS-TLR4 signaling pathway in the colon and hippocampus. In the colon (Fig. 2D) of male mice, the expression of *TLR-4, Myd88, CD14*, nuclear factor (*NF*)-*κB, TNF-a* and *IL-1β* was significantly higher in ko-wtM group than in the wt-wtM group and expression of *P38MAPK* was higher in the ko-wtM group than in wt-wtM group, but the difference was not significant (P = 0.07). *JNK* expression in the wt-wtM group was significantly higher than that in the ko-wtM group; other genes showed no significant differences between these two groups. In female mice, the expression of *TLR-4, CD14, p38MAPK, JNK, IL-1β* and *TNF-a* was significantly lower in ko-wtF group than in the wt-wtF group. In the hippocampus (Fig. 2E), the expression of *Myd88, CD14, NF-κB*, and *p38MAPK* tended to be higher in the ko-wtM group than in the wt-wtM group, but the difference was not significant. The expression of *TLR-4, LBP, Myd88, NFĸB, IL-1β*, and *TNF-a* was significantly higher in ko-wtF mice. The expression of *ILR-4, LBP, Myd88, NFĸB, IL-1β*, and *TNF-a* was significantly higher in ko-wtF mice.

Thus, after 4 weeks of CUMS, the colon of LF-free male mice showed more severe inflammation which occurred through activation of the *TLR4-Myd88-NF-kB* signaling pathway. However, female mice that consumed normal mouse milk showed more severe intestinal inflammation via activation of the *MAPK* and *JNK* signals in the TLR4 signaling pathway. The difference in colonic inflammation between female and male mice may be related to different levels of oestrogen [24]. Based on the significantly different

levels of BDNF in the serum, we measured the expression of genes related to the *BDNF* signalling pathway in the hippocampus of depressed mice. None of the genes differed between the ko-wtM and wt-wtM group (Fig. 2F); however, in the female groups, the expression of *GRB2, CaMK*, and *CREB* in the ko-wtF group was significantly lower than that in the wt-wtF group, and the expression of *BDNF* in the ko-wtF mice was lower than that in the wt-wtF group but not significantly (P = 0.09).

Next, we examined the composition of the gut flora. Alpha diversity represents the richness and diversity of the microbial community. The Shannon formula was used to estimate diversity, and the richness of the microbial community was estimated using the Chao index. Mice without LF intake during the suckling period (ko-wtM and ko-wtF) showed a lower Shannon index of the intestinal microbiota but the difference was not significant (Fig. 2G). The Chao index was significantly lower in the ko-wtF than in the wt-wtF mice, but there was no significant difference between the ko-wtM and wt-wtM groups (Fig. 2G). Taken together, LF-free feeding during the suckling period decreased the microbial community richness compared with LF feeding after CUMS. Principal coordinate analysis (PCoA) revealed major differences between the ko-wtM and wt-wtM mice as well as differences in the microbial composition between the kowtF and wt-wtF mice (Fig. 2H, P = 0.097). There was a significant difference in the pattern of intestinal flora between the ko-wt-depressed and wt-wt-depressed mice. The predominant phyla (Fig. 2I) were Firmicutes, Bacteroidetes, Desulfobacterota, Patescibacteria, and Actinobacteria in the faecal samples of the four groups, and there was no difference between ko-wtM and wt-wtM mice. In contrast, the abundances of Desulfobacterota, Actinobacteria, and Chloroflexi significantly increased in the ko-wtF group (Fig. 2J). At the genus level, we used linear discriminant analysis effect size (LEfSe) to identify bacteria whose relative abundance significantly differed between ko-wt and wt-wt mice. Figure 2K shows that there was a greater abundance of *Eubacterium\_xylanophilum\_group*, and a lower abundance of Lactobacillus and SCFA-producing bacteria (Alistipes and Dubosiella) in the ko-wtM group than in the wtwtM group. SCFAs exert anti-inflammatory functions and are beneficial to the composition of the gut microbiota and intestinal barrier [25]. Figure 2K shows that Lactobacillus, Desulfovibrio, Arenimonas, Enterorhabdus, and Bifidobacterium were significantly enriched in the ko-wtF group, and Ileibacterium was significantly enriched in the wt-wtF group.

To further explore the correlation between the gut microbiota (significantly different microbiota at the genus level) and depression-related behavioural indices (OFT, SPT, TST, and FST), depression-related hormones (ACTH and CORT), serum inflammatory factors (IL-1 $\beta$  and TNF- $\alpha$ ), and BDNF, a heatmap of Spearman's correlation analysis was generated. As shown in Fig. 2L, *Desulfovibrio* was significantly negatively correlated with SPT and positively correlated with FST (P = 0.06); *Bifidobacterium* was negatively correlated with TST (P = 0.05); and *Alistipes* was significantly positively correlated with OFT. Furthermore, *Lactobacillus* and *Enterorhabdus* were positively correlated with TST, ACTH, CORT, IL-1 $\beta$ , TNF- $\alpha$ , and LPS, and negatively correlated with OFT, SPT, and BDNF. *Blautia* showed a positive correlation with ACTH, FST, TST, CORT, IL-1 $\beta$ , TNF- $\alpha$ , LPS, and depression.

Intestinal growth retardation and intestinal microbiological disorder in 18-day-old suckling mice with LF feeding deficiency during lactation

Our experiments showed that LF intake during lactation significantly affected depression in adult mice, which was induced by differences in body development caused by LF intake during lactation. Therefore, we used 18-day-old mice with almost no feed intake for analysis. The body weight of 18-day-old suckling mice without LF feeding during lactation did not differ from that of control mice (Fig. 3A) but the intestinal index and density of the small intestine and colon decreased significantly (Fig. 3B, 3C). Crypt depth did not differ significantly between the two groups (Fig. 3D). The number of colonic goblet cells in the ko-wt group was significantly lower than that in the control group (Fig. 3E).

To further explore the effect of LF feeding deficiency on intestinal development during lactation, we performed genome-wide transcriptional profiling of the small intestine of the two groups of mice for RNA sequencing. PCoA showed that the transcriptomes of ko-wt and wt-wt mice were clearly separated (Fig. 3F). Transcriptome sequencing showed that 143 genes were upregulated and 72 genes downregulated in the small intestine of the ko-wt mice compared to those of the wt-wt mice (fold-change > 1, P adjust < 0.05, Fig. 3G). These DEGs were subjected to pathway analysis using the Kyoto Encyclopedia of Genes and Genomes (KEGG) database, which showed that upregulated mRNAs were mainly involved in infectious, immune, and neurodegenerative diseases, and were enriched in the nervous, immune, and endocrine systems (Fig. 3H). We explored immune-related genes in the small intestine using an RNA-seq transcriptomic profiling approach, as shown in Fig. 3I; genes encoding  $\alpha$ -defensin, such as *Defa23, Defa36, Defa3, Defa17, Defa38*, and *Defa39*, were significantly decreased in the ko-wt group. The RNA-Seq results showed no difference in the expression of the tight junction proteins *zo-1* and *occludin* between the two groups. However, the expression of the gene encoding the actin protein (*Gm12715*) was significantly decreased (Fig. 3J); this gene is related to the assembly of tight junctions.

To determine the effect of LF deficiency during lactation on intestinal microorganisms, we analysed the gut microbiota composition by 16S rRNA gene amplicon sequencing of the caecal contents of 18-day-old mice. Alpha diversity measures (Shannon and Chao diversity indices), calculated for each group, showed that the Chao diversity index of the ko-wt group did not differ from that of the wt-wt group; however, the Shannon index was significantly higher than that of the control group (Fig. 3K). These results indicate that the intestinal microbial alpha diversity was increased in 18-day-old mice not fed LF. PCoA of the beta-diversity comparison revealed significant separation between the two groups of microbial communities (Fig. 3L). At the phylum level, Firmicutes, Bacteroidetes, and Desulfobacterota accounted for the majority of the bacteria (Fig. 3M), with no differences between the two groups (Fig. 3N). At the genus level, the LEfSe results revealed a greater abundance of *Muribaculum, Oscillibacter, Bilophila, Colidextribacter, Harryflintia, Bifidobacterium, Odoribacter, UCG-009, Lachnospiraceae\_UCG-006, Eubacterium\_xylanophilum\_group, Ruminococcus, and some unclassified microbiotas; there was a lower abundance of <i>Blautia, Desulfovibrio, Parabacteroides, Romboutsia, Eubacterium\_brachy\_group*, and *Turicibacter* in the ko-wt group (Fig. 30).

# LF feeding deficiency during lactation induced hippocampal growth retardation

Considering that depression is a mental illness, it is necessary to explore the effects of LF deficiency during lactation on the hippocampus. Figure 4A shows that there was no significant difference in the brain index between ko-wt and wt-wt 18-day-old mice. The CA1 region of the hippocampus is involved in cognitive processes, learning, and memory[26]. The hippocampal dentate gyrus (DG) is a key target region for both antidepressant effects and stress-related processes [27]. In the wt-wt group, pyramidal cells in the CA1 region were arranged neatly in dense layers and had regular and clear structures. No significant difference was observed between the ko-wt and wt-wt groups in cell morphology in the DG region (Fig. 4B).

RNA-seq of the hippocampi of 18-day-old mice was performed. PCoA showed that the microbiota profiles of the two groups were clearly distinct (Fig. 4C). Transcriptome sequencing results showed that 199 genes were upregulated and 297 genes downregulated in the hippocampi of the ko-wt mice compared to those of the wt-wt mice (fold-change > 1, P < 0.05, Fig. 4D). The KEGG pathway annotation results showed that the downregulated mRNAs mainly participated in signalling molecules and interactions, and signal transduction, and were enriched in the immune system (Fig. 4E). Figure 4F shows that the expression of genes involved in innate immunity, such as Lrg1, Gbp4 [28], Gbp10 [28], and Stx11 [29], was decreased. Genes involved in adaptive immunity, such as Ccl21a [30], Cd247 [31], Cd28 [32], and lL25 [33], showed increased expression. The gene expression of Pigr [34], Ltf, Lcn2 [35], S100a9 [36], and Ccl28 [37], which are involved in both innate and adaptive immune responses, decreased. Some genes involved in the development of the nervous system in the ko-wt group were significantly different from those of the wt-wt group. The gene expression of BDNF, Cdh3, Ccn5, and Folr1 decreased, and that of Prdm12 and Pcdha8 increased (Fig. 4G). These differences may lead to the delayed development of the nervous system. Figure 4H shows that the expression of genes related to neuronal signal transduction (Trpv4, Gabra6, Kcne2, and Kcnj13) decreased in the ko-wt group.

To explore whether the influence of LF-deficient feeding during lactation on the hippocampus extends into adulthood, we assayed differential gene expression in the hippocampus of adult mice. Figure 4I shows that there was no significant difference between KO-WT mice (mKO-WT, fKO-WT) and WT-WT mice (mWT-WT, fWT-WT) in immune-related differential expression genes (DEGs), but gene expression in female mice was significantly higher than that in male mice. Figure 4J shows that the expression of *BDNF, Ccn5,* and *Cdn3* in KO-WT mice (mKO-WT, fKO-WT) did not differ from that in WT-WT mice (mWT-WT, fWT-WT). Similar to in 18-day-old mice, the expression of Prdm12 and Pcdha8 in adult female KO-WT mice (fKO-WT) was significantly higher than that in WT-WT mice (fWT-WT); however, this difference was not observed in adult male mice. Figure 4K shows that the expression of *Trpv4, Gabra6, Kcne2,* and *Kcnj13* in KO-WT mice (mKO-WT, fKO-WT) did not differ from that in WT-WT mice (mWT-WT), and the expression of *Gabra6, Kcne2,* and *Kcnj13* was significantly higher in female mice than in male mice. When the mice reached adulthood, the difference induced by LF feeding deficiency in the hippocampus was decreased, but some genetic differences related to the nervous system development process persisted.

# Lactation LF feeding-deficient adult mice still showed some differences in intestinal and intestinal microorganisms compared with the control group

To explore whether the effects of LF feeding deficiency during lactation on intestinal and intestinal microorganisms continue into adulthood, mice were fed normally until adulthood after weaning.

Intestinal development was evaluated using four groups (male: mWT-WT, mKO-WT; female: fWT-WT, fKO-WT) of mice drinking normal water which was different from that of 18-day-old mice. There were no significant differences in the body weight (Fig. 5A), small intestinal index (Fig. 5B), small intestinal density (Fig. 5C) and colon length (Fig. 5D) between WT-WT and KO-WT mice either male or female. The maltase/lactase values of the ileum in the mKO-WT and fKO-WT mice were significantly lower than those in the mWT-WT and fWT-WT mice, respectively (Fig. 5E, 5F, respectively). As shown in Fig. 5G, there was no difference between KO-WT and WT-WT mice in the crypt depth (mKO-WT vs. mWT-WT; fKO-WT vs. fWT-WT). Figure 5H shows that the expression of *zo-1* and *occludin* in KO-WT mice was lower than that in WT-WT (mKO-WT vs. mWT-WT; fKO-WT vs. fWT-WT), but not significantly. *Occludin* expression was significantly lower in female mice than in male mice (fWT-WT vs. mWT-WT; fKO-WT vs. mKO-WT). Thus, the influence of LF feeding deficiency on the intestines of normal adult mice was weakened; however, female mice that consumed LF-free milk during lactation may have a greater risk of the intestinal barrier dysfunction in adulthood. In addition, the ileal maturity of LF-deficient mice was lower than that of LF-drinking mice, and we have previously shown that the maturity of the three segments of the small intestine of 18-day old mice drinking LF-free milk was lower than that of normal mice [17].

To examine the effects of different feeding methods during lactation on the composition of intestinal microflora in healthy adult mice (male: mWT-WT, mKO-WT; female: fWT-WT, fKO-WT), the cecal content was investigated using 16S rRNA gene sequencing. Alpha-diversity analysis revealed no difference in the Shannon and Chao indices between WT-WT and KO-WT mice (mKO-WT vs mWT-WT; fKO-WT vs fWT-WT, Fig. 5I). PCoA of the beta-diversity comparison revealed separation between the mKO-WT and mWT-WT groups, but the difference was not significant (P = 0.053). PCoA showed no separation among fKO-WT and fWT-WT groups on OUT level (Fig. 5J). These results suggest that there was no difference in alpha and beta diversity between lactating LF-deficient adult mice and normal mice. The phylum level was dominated by Firmicutes, Bacteroidetes, Desulfobacterota, Actinobacteria, and Patescibacteria (Fig. 5K). Patescibacteria was significantly higher in the mWT-WT group, and there was no difference between the fKO-WT and fWT-WT groups at the phylum level (Fig. 5L). At the genus level, the LEfSe results demonstrated a greater abundance of Lactobacillus, Lysinibacillus, Ileibacterium, Bifidobacterium, and Lachnospiraceae\_UCG-001, whereas there was a lower abundance of Kurthia, Enterorhabdus, Candidatus, Saccharimonas, Eubacteriumxylanophilum\_group, Monoglobus, and ASF356 in the mKO-WT group. Dubosiella, Lactococcus, Sporosarcina, Pseudogracilibacillus, and Eubacteriumbrachygroup were significantly higher in the fKO-WT group; Staphylococcus and Eubacterium\_nodatum\_group were significantly higher in the fWT-WT group (Fig. 5M).

# Lactation LF feeding-deficient adult mice showed severe intestinal injury and intestinal microbial disorder based on DSS-induced colitis

Following LF deficiency during lactation, the intestinal development and composition of intestinal microorganisms in 18-day-old mice significantly differed from those of normal mice; however, these differences were covered and became not obvious after 9 weeks of normal feeding under physiological conditions. To examine whether the two types of mice exhibited the same pathological reaction, a DSS model was established in adult mice.

In the DSS colitis model mice, the body weight of mDKO-WT mice was lower than that of mDWT-WT mice at d 1-5, but this difference disappeared at d 6-7 (body weight decreased to 73.4% and 73.1% in mDWT-WT and mDKO-WT mice, respectively), possibly because of the longer treatment time with DSS (Fig. 6A). In female mice, the body weights of fDKO-WT mice were consistently lower than those of fDWT-WT mice from d 3 to 7, but the difference was not significant, and the body weights of fDWT-WT and fDKO-WT mice decreased to 77.3% and 74.6%, respectively (Fig. 6B). The disease activity index (DAI) score is an effective indicator of colon inflammation and damage. The DAI score of the mDKO-WT group was higher than that of the mDWT-WT group but not significantly (Fig. 6C). The DAI score of the fDKO-WT group was also higher than that of the fDWT-WT group, and the difference was significant at d 6 and 7 (Fig. 6D). The survival rate of the mDKO-WT group was lower than that of the mDWT-WT group, and there were no deaths in the two female mouse groups (Fig. 6E). There was no difference in colon length between the two groups of male mice but the colon length of fDKO-WT mice was significantly shorter than that of fDWT-WT mice (Fig. 6F). Hematoxylin and eosin (H.E) staining of the colonic tissue sections showed that DSS-treated colons had great histological damage (cellular infiltration, goblet cell depletion, and damage to the crypt architecture), but the difference between KO-WT and WT-WT mice was unclear. The crypt depth of fDKO-WT mice was lower than that of fDWT-WT mice, but the difference was not significant (Fig. 6G). Figures 6H–K shows that the expression of IL-1 $\beta$  and IL-10 in the colon in the fDKO-WT group was significantly higher than that in the fDWT-WT group. Thus, LF-deficient feeding of mice during lactation led to more severe DSS-induced weight loss (female), an increased DAI score (male and female), decreased survival (male mice), and colon shortening (female mice).

Ulcerative colitis is typically associated with dysbiosis of the gut microbiota. Alpha-diversity analysis showed that the Shannon index of the fDKO-WT group was significantly lower than that of the fDWT-WT group. The Chao index at the OTU level of fDKO-WT mice was significantly lower than that of fDWT-WT mice (Fig. 6L). These results suggest that alpha diversity was lower in LF-deficient female mice compared to those with normal milk; no difference was observed in male mice after acute DSS injury. PCoA revealed significant separation between the mDKO-WT and mDWT-WT groups at the OTU level, and the microflora of female LF-deficient DSS mice (fDKO-WT) differed significantly from that of the fDWT-WT group (Fig. 6M). At the phylum level, Bacteroidetes, Firmicutes, Proteobacteria, Campilobacterota, Desulfobacterota, Actinobacteriota, and Deferribacterota accounted for the majority of the bacteria (Fig. 6N). The relative abundance of Firmicutes was significantly higher in the mDKO-WT group, the relative

abundance of Campilobacterota was significantly higher, whereas that of Desulfobacterota and Cyanobacteria was significantly lower (Fig. 60). At the genus level, the relative abundance of Lachnospiraceae\_NK4A136 group, Ruminococcus\_torques\_group, Odoribacter, Romboutsia, Marvinbryantia, and Family\_XIII\_AD3011\_group was higher in the mDKO-WT group than in the mDWT-WT group (Fig. 6P). As shown in Fig. S2B, the abundance of Ruminococcus\_torgues \_group, Odoribacter, *Romboutsia*, and *Family\_XIII\_AD3011\_group* was higher in the mDKO-WT group than in the mKO-WT group, indicating that mDKO-WT mice experienced a more substantial increase in these genera. The relative abundances of Leucobacter and Parvibacter were lower in the mDKO-WT group than in the mDWT-WT group (Fig. 6P). In female mice, the relative abundances of pathogenic bacteria such as Escherichia-Shigella, Enterococcus, Clostridium\_sensu\_stricto\_1, Erysipelatoclostridium, Colidextribacter, and Akkermansia were higher in the fDKO-WT group than in the fDWT-WT group (Fig. 6P). The relative abundances of beneficial genera, such as Alloprevotella, Desulfovibrio, and many unclassified bacteria on genus level were lower in the fDKO-WT group than in the fDWT-WT group (Fig. 6P). Similarly, the abundance of Escherichia-Shigella, Enterococcus, Erysipelatoclostridium, Colidextribacter and Akkermansia were increased and Desulfovibrio was decreased in the DSS model (Fig. S2 D). fDKO-WT mice showed a more significant increase in Escherichia-Shigella, Enterococcus, Erysipelatoclostridium, Colidextribacter and Akkermansia and a serious decline in Desulfovibrio. LF feeding-deficient mice clearly showed more severe microbial disorders after DSS processing.

### Discussion

LF is an important functional protein in breast milk that has profound effects on the growth and development of infants and young children. Formulas are the main breast milk substitute for infants who cannot breastfeed. However, LF has not been widely added to formulas, commonly leading to LF feeding deficiency during lactation. LF has a protective effect on newborns, particularly premature infants. However, the effects of lactation LF feeding deficiency on adult health remain unclear. Depression is the most common mental health problem in adults and a leading risk factor for physical disability. Here, we explored the influence of LF feeding deficiency during lactation on the incidence of depression in adult mice and related mechanisms. Unexpectedly, we found that lactating LF feeding-deficient mice showed more severe depressive behaviour after induction of CUMS depression in adulthood (Fig. 1).

# LF feeding deficiency during lactation impairs intestinal barrier function and immunomodulatory function

Dysfunction of the microbiota-gut-brain axis is the main pathological basis of depression and may directly influence and cause psychiatric disorders [38]. Our results showed that lactating LF feeding-deficient mice had more severe damage of intestinal barrier function after depression in adulthood, which led to more severe intestinal microbial translocation, higher concentrations of LPS in systemic circulation and more severe inflammation in the colon. Based on these observations in the intestines, we further explored the effects of LF deficiency during lactation on intestinal development in mice in different

growth periods. The small intestine of 18-day-old LF-deficient mice showed decreased intestinal barrier function and innate immunity. ZO family proteins bind directly to F-actin and barrier-forming claudins are necessary for the assembly of tight junctions [39]. Actin cytoskeletal can also regulate epithelial permeability, and knockdown of myosin IIA decreases epithelial permeability in several intestinal epithelial cell lines [40]. Therefore, the low expression of actin protein in the ko-wt group (18-day-old) may have weakened intestinal barrier function. When the mice reached 9 weeks of age, LF feeding-deficient mice still showed lower mRNA expression of tight junctions than LF-fed mice. The DEGs related to immunity were mainly enriched in the NOD-like receptor signaling pathway, which plays a key role in pathogen identification and innate immune response (Fig. S1 A). Paneth cells in the crypts of the small intestine secrete microbicidal α-defensins as components of the enteric innate immunity. Moreover, dysregulation of α-defensins has been observed under pathogenic conditions, such as in IBD [41]. The lack of LF during lactation reduces the secretion of  $\alpha$ -defensin in the small intestine of suckling mice, which may reduce innate immunity and increase the incidence of inflammatory intestinal disease. When the mice reached adulthood, the difference between the KO-WT and WT-WT groups (mKO-WT vs. mWT-WT, fKO-WT vs. fWT-WT) in the intestinal physiological structure of healthy mice was decreased. However, after DSS enteritis treatment, KO-WT mice showed more serious damage and inflammation, particularly female mice. Thus, loss of LF feeding during lactation hinders the development of intestinal physiological structure, barrier function, and innate immune function in sucking mice, and the gap in intestinal physiological structure gradually narrows with increasing age but a functional gap remains, particularly during negative stress. Therefore, LF feeding deficiency during lactation impairs intestinal barrier and immunomodulatory functions, causing serious damage to the intestinal barrier and inducing systemic inflammation, under the stimulation of depression modelling, which in turn leads to neuroinflammation and depression.

# Effect of LF feeding deficiency during lactation on gut microbiota from infancy to adulthood

The gut microbiota plays a prominent role in gut-brain interactions. The results of gut microbiota are summarized in Table. S2. LF feeding deficiency during lactation affects the composition of intestinal microorganisms in mice, and this effect continues into adulthood. The overall microbial composition of 18-day-old ko-wt mice deviated from that of the control group. This separation disappeared in 9-week-old healthy mice, possibly because they had the same dietary conditions. However, this separation reappeared under stimulation of depression and DSS. The effects of LF deficiency during lactation on gut microbes are persistent. Alpha diversity is often used as a proxy for community stability and functioning. The effect of depression on intestinal microbial alpha-diversity is inconclusive [14]. In our study, the microbiota of LF feeding-deficient mice showed lower alpha diversity after CUMS. Similarly, LF feeding-deficient mice showed lower alpha diversity after Se netritis. Alpha diversity was increased in 18-day-old ko-wt mice, possibly because of the antibacterial activity of LF. In contrast, another study showed that LF intervention in early life increased the diversity of the caecal microbiota [20].

The effect of LF deficiency during lactation at the genus level of intestinal microorganisms was complex and diverse. In the intestine of 18-day-old suckling mice, many microorganisms enriched in the ko-wt group were related to inflammation. For example, many studies have reported that Oscillibacter is increased in depression [42] and is positively correlated with obesity and increased permeability of the mouse colon [43]. Similarly, the increasing relative abundance of clusters in the ko-wt group, including Bilophila, Colidextribacter, Harryflintia [43], Odoribacter [44], Lachnospiraceae\_UCG-006 [45], and Ruminococcus [46], has also been reported as positively associated with inflammation due to progressive obesity or type 2 diabetes mellitus. In our study, Blautia, Desulfovibrio, Parabacteroides, Romboutsia, Eubacterium\_brachy\_group, and Turicibacter were enriched in wt-wt group. Blautia may function as a probiotic in the host and has gained attention because of its ability to alleviate inflammatory diseases and metabolic diseases, as well as because of its antibacterial activity against specific microorganisms [47]. However, we found that Blautia was positively related to depression. Parabacteroides and Turicibacter resulted in increased formation of the faecal SCFAs propionate and butyrate [48]. In our RNAseq results, LF feeding deficiency during lactation decreased the expression of a-defensin in the small intestine of 18-day-old sucking mice (Fig. 3I), which may contribute to host-microbe dysbiosis and enhance inflammatory responsiveness associated with the pathogenesis of IBD [49]. Therefore, the disorder of intestinal microorganisms in 18-day-old ko-wt mice may be directly related to the antibacterial effect of LF. Additionally, it may be indirectly related to the effect of LF on the development and gene expression of the intestine.

When the mice reached adulthood, the abundance of beneficial bacteria in LF-feeding deficient mice was lower than in LF-fed mice. Among the intestinal microorganisms of 9-week-old healthy mice, the abundance of Kurthia, Enterorhabdus, Candidatus\_Saccharimonas, Monoglobus, ASF356, and Eubacterium\_xylanophilum\_group was lower in the mKO-WT group and decreased in the DSS model compared to in healthy mice (Fig. S2 B). A previous study showed that the abundance of Enterorhabdus spp. decreased after chronic DSS induction [50]. According to previous reports, ASF356 exerts probiotic effects [51]. Candidatus\_Saccharimonas can suppress the production of TNF-α in macrophages, suggesting its potential for immune suppression [52]. Eubacterium\_nodatum\_group enriched in fWT-WT is SCFAs-producing bacteria [53]. These results suggest that the adverse effect of LF feeding deficiency during lactation on the composition of intestinal microorganisms continued into adulthood. This effect was also observed in the DSS model. Among intestinal microorganisms in the DSS model, more harmful bacteria were enriched in LF-deficient mice (Fig. 6P). Ruminococcus\_torques\_group and *Family\_XIII\_AD3011\_group* were enriched in the mDKO-WT group; these genera are significantly positively corelated with inflammation [25, 54]. Escherichia-Shigella, Enterococcus, Clostridium\_sensu\_stricto\_1, Erysipelatoclostridium, Colidextribacter and Akkermansia were increased in the fDKO-WT group compared to in the fDWT-WT group. Escherichia-Shigella and Enterococcus are potential pathogens [55, 56], and studies found that Clostridium\_sensu\_stricto\_1, Erysipelatoclostridium and Colidextribacter were increased in high-fat diet-fed mice [43, 44, 57]. Similar to our results, Xu et al. found that the relative abundance of Akkermansia was markedly increased after chronic DSS induction [50]. Considering the close relationship between the intestinal microbial composition and depression, LF deficiency during

lactation is likely to affect intestinal microorganisms in the CUMS model. In the CUMS model, *Eubacterium\_xylanophilum\_group* was enriched in the ko-wtM group and was positively correlated with FST (Fig. 2L), which was positively correlated with the severity of depression. Study found that *Eubacterium\_xylanophilum\_group* was the characteristic genera of valproic acid-induced rat autism model [58]. To the best of our knowledge, our results are the first to report that Eubacterium\_xylanophilum\_group might be associated with depression. We also found that Alistipes was positively correlated with the OFT, indicating that Alistipes is negatively correlated with the severity of depressive symptoms. The abundance of *Alistipes* is decreased in patients with IBD, which is characterised by overexpression of indoleamine 2,3-dioxygenase (IDO) in the colon [59]. IDO can reduce 5-hydroxytryptamine (5-HT) levels by promoting the metabolism of tryptophan, a precursor of 5-HT synthesis, leading to depression. In our study, the abundance of Lactobacillus was decreased in the kowtM group but increased in the ko-wtF group (Fig. 2K). Studies of the role of Lactobacillus in depression have reported conflicting results. Mice with chronic restraint stress-induced depression showed a significant increase in Lactobacillus [60], whereas a low abundance of Lactobacillus was found in people with depression [61]. In addition, this difference may be sex-dependent, and Lactobacillus was shown to be inversely associated with depression scores among males but was positively associated with anxiety scores among females [62], which is similar to our results. Bifidobacterium was enriched in the ko-wtF group, and is present at high levels in major depressive disorder/depression groups in many studies [14]. Interestingly, specific strains of *Bifidobacterium* are commonly considered as anti-inflammatory and used as probiotics. However, a higher abundance of *Bifidobacterium* has been associated with IBD, indicating that specific strains have inflammatory potential [63]. However, we found that Bifidobacterium was negatively corelated with depression (Fig. 2L), based on the negative correlation between *Bifidobacterium* and depression in male mice (Fig. S1 B), and the correlation was greater than that in female mice, thus affecting the overall correlation. *Bifidobacterium* was also enriched in the 18-day-old ko-wt (Fig. 30) group and 9-week mKO-WT (Fig. 5M) group, indicating that colonization of the neonatal intestine by Bifidobacterium can be reduced if milk is concurrently supplemented with bovine LF [64]. Therefore, a lack of LF during lactation may be beneficial for colonization of *Bifidobacterium*. Many studies revealed higher levels of *Desulfovibrio* in patients with major depressive disorder than in controls [14, 61]. Similarly, Desulfovibrio was enriched in ko-wtF group (Fig. 2K) and was positively related to the severity of depressive symptoms in our study (Fig. 2L).

The lack of LF during lactation increased the pro-inflammatory microflora in the intestinal microflora of 18-day-old mice and decreases the number of beneficial bacteria in adult mice, which is disadvantageous for establishing a healthy intestinal ecosystem. LF feeding deficiency during lactation may decreased the stability of intestinal microorganisms, rendering them more vulnerable to dysregulation during adverse stimuli.

# LF feeding deficiency during lactation inhibits hippocampal development from infancy to adulthood

The hippocampus is a key brain region involved in cognitive processes, mood regulation, and the pathophysiology of depression [12]. In our study, the ko-wtF group exhibited more severe hippocampal inflammation (Fig. 2E) and lower BDNF levels (Fig. 1I, 2F). In the hippocampus of 18-day-old mice, genes involved in innate immunity were downregulated in the ko-wt group (Fig. 4F). Family-wide loss-of-function analysis showed that *Gbp10* conferred cell-autonomous immunity to listerial or mycobacterial infection within macrophages and gene-deficient animals [28]. Another study revealed that *stx11* knockdown reduced phagocytosis of *Escherichia coli* in interferon-y-activated macrophages [29]. The innate immune response can eliminate most invading pathogenic microorganisms and participate in immune selfstabilization of the body. Low expression of innate immune related genes is unfavorable for pathogen clearance in time, thus aggravating infection. Furthermore, activation of innate immunity is indispensable for inducing adaptive immune responses. Downregulation of innate immune function may affect the intensity and type of adaptive immunity. In our study, the genes involved in adaptive immunity were upregulated in the hippocampus of LF-deficient mice (Fig. 4F). Ccl21a promotes the establishment of self-tolerance in T cells in the thymic medulla [30]. CD28 is the second signal required for T-cell activation during the immune response [32]. CD247 is part of the T-cell antigen receptor complex, which plays a key role in receptor expression and signaling leading to optimal effector T-cell functions [31]. IL-25 can induce and enhance Th2 type immune response [33]. Some genes participate not only in the innate immunity of the mucous membrane but also in acquired immunity. Pigr mainly functions to cooperate with IgA transport, which plays an important role in mucosal immunity [34]. CCL28 is an anchoring point that bridge innate and adaptive immunities [37]. The Ltf gene encodes LF, which has a defensive role in the body as its levels were found to be profoundly elevated under pathological conditions such as neurodegeneration and inflammatory disease [65]. Lcn2 can control neuroinflammation by modulating cytokine production and exerts neuroprotective functions in the brain by suppressing the secretion of proinflammatory cytokines [35]. S100a9 encodes a pro-inflammatory protein (calgranulin) that has been implicated in multiple diseases. A previous study has shown that S100a9 prevents hyperinflammatory responses without impairing pathogen defence [36]. The Ltf, Lcn2, and S100a9 genes play important roles in controlling the balance of the immune response, and low expression of these genes may induce immune response disorders. Although our results showed that the disorder of hippocampal immunomodulatory function caused by LF feeding deficiency in 18-day-old mice disappeared in adult mice (Fig. 4I), another study showed that early life inflammation causes depressive symptoms in adolescence [66], suggesting that disorder of hippocampal immunomodulatory function likely led to neuroinflammation in 18-day-old ko-wt mice, which may have been related to depression in adult mice.

Mice fed an LF-deficient diet during lactation may be at risk of neurodevelopmental delay. *Prdm12* regulates the proliferation and differentiation of neural stem cells. Overproduction of *prdm12* impaired cell proliferation and increased the G1 population [67], and high expression of *Prdm12* in the ko-wt group (18-day-old mice (Fig. 4G)) and fKO-WT group (adult mice (Fig. 4J)) may inhibit neurogenesis. *Pcdha8* is a member of the protocadherin cluster, which regulates the ability of neurons to migrate and of dendrites to avoid each other. Many brain diseases, such as autism spectrum disorders and depression, are caused by abnormal neuronal migration and connections [68]. Low *Pcdha8* expression increases the risk of

depression [68], but the influence of high *Pcdha8* expression on depression is unclear. In our study, the expression of *Pcdha8* was higher in the ko-wt group (18-day-old mice (Fig. 4G)) and fKO-WT group (adult mice (Fig. 4J)). The *Cdh3* gene encodes cadherin, a cell adhesion molecule that is important in a variety of morphogenetic events during neural development including cell migration, segmentation of the neural tube, neurite outgrowth, axon targeting, and synapse formation [69]. A previous study showed that the overexpression of *Wisp2/Ccn5* potently enhanced neurite outgrowth. The low expression of *Cdh3* and *Ccn5* in 18-day-old ko-wt mice (Fig. 4G) may negatively affect neural development. We found that *BDNF* expression was significantly lower in the 18-day-old ko-wt mice (Fig. 4G); similarly, the ko-wtM and ko-wtF mice showed lower BDNF serum levels after CUMS (Fig. 11). LF has been suggested to promote early neurodevelopment and cognition in postnatal piglets by upregulating the *BDNF* signalling pathway [70]. Therefore, a lack of LF during lactation reduces the expression of *BDNF*, which may increase the risk of depression in adult mice. In the hippocampi of 18-day old mice, the expression of *Folr1* was lower in the ko-wt group (Fig. 4G). *Folr1* is essential for neural tube closure and can induce central nervous system axon regeneration mediated by folate [71].

In our study, the expression of genes related to neuronal signal transduction in the 18-day-old ko-wt mice significantly decreased (Fig. 4H), but did not differ from that in adult mice (Fig. 4K). Gabra6 is a GABA receptor. GABA is a neurotransmitter involved in the mechanism of anxiety and depression. A rat experiment suggested that depressive behaviour caused by maternal-foetal separation is related to decreased expression of Gabra6 [72], and low GABA transmission has been reported in victims of suicide [73]. TRPV4 is a Ca2<sup>+</sup> nonselective cation channel involved in modulating membrane potentials in neurons. A previous study showed that downregulation of hippocampal TRPV4 affects depression-like behavior in mice by regulating neurogenesis [74]. Kcne2 and Kcnj13 regulate the return of the membrane potential to resting potential, and loss of *Kcne2* can increase excitability in neurones [75]. Decreased expression of *Kcne2* and *Kcnj13* may lead to continuous excitation of neurons, resulting in neurotoxicity. In summary, LF feeding deficiency during lactation decreased the expression of Gabra6, TRPV4, Kcne2, and Kcnj13 in 18-day-old mice, which may increase the depression rate of in mice; however, as the mice aged, this difference disappeared. The choroid plexus is the site of cerebrospinal fluid (CSF) production and of the blood-CSF barrier, and claudin2 is present in the epithelial tight junctions of the choroid plexus forming the blood-CSF barrier [76]. Our results showed that *Cldn2* expression was decreased in 18-day old ko-wt mice (Fig. S1C), which may have led to increase in permeability of the blood-CSF barrier and thus more severe neuroinflammation and an increased risk of depression.

## Sex differences

We evaluated mice of both sexes and observed differences in some results between male and female mice. Epidemiological studies have shown that women have much higher rates of major depression and higher levels of inflammatory markers than men [77]. Similarly, female mice showed a more severely depressed phenotype in our study (Fig. 1B, 1I–N). Furthermore, female mice had lower intestinal tight junction expression and higher level of inflammatory cytokines, which induced more severe hippocampal

inflammation compared to in male mice in our CUMS model. LF-feeding deficient male mice showed more severe intestinal inflammatory damage after CUMS (Fig. 2). In contrast, the LF feeding-deficient female mice showed mild intestinal inflammatory damage after CUMS. This difference may be related to the anti-inflammatory effect of oestrogen, study found that oestrogen decreased the production of macrophage migration inhibitory factor, which affected the susceptibility to inflammation in the colon by reducing the TNF-α and IL-1β production in colitis [24, 78]. However, why did the anti-inflammatory effects of oestrogen manifest only in the ko-wtF and not in wt-wtF mice? We speculated that the LF feeding deficiency during lactation may influenced the signal transduction of estrogen. In 18-day-old suckling mice, the genes encoding FKBP52 and Hsp70 in the oestrogen signalling pathway were up regulated in the ko-wt group (Fig. S1 D, E). As receptor associated proteins, FKBP52 and Hsp70 affect oestrogen receptor nucleocytoplasmic shuttling [79]. Oestrogen receptors are abundantly expressed in the intestine and cells of the immune system [80], and binding of these receptors to oestrogen induces gene regulation. This phenomenon did not occur in the DSS enteritis model, possibly because of the different pathogeneses of chronic enteritis caused by depression and acute enteritis established by DSS. Further studies are needed to investigate the underlying reasons for this result. In our DSS model, in agreement with the results of previous studies [78], male mice showed more severe weight loss and higher mortality than female mice, which was also related to the anti-inflammatory function of oestrogen. Sex-specific relationships between gastrointestinal microbiota and depression were also observed in this study. The abundance of Lactobacillus and Bifidobacterium was negatively related to depression among male mice and positively related to depression among female mice. Currently, the relationship between Lactobacillus and Bifidobacterium and depression in individuals of different sex is still inconclusive, which is also an important reason to prevent these two bacteria from being widely used in the clinical treatment of depression [62].

### Conclusions

Our study demonstrated for the first time that LF feeding deficiency during lactation increases the risk of depression in adults. During negative stress, LF deficiency causes more serious intestinal barrier damage and inflammatory responses, thus increasing systemic inflammation and neuroinflammation, leading to depression. Lack of LF during lactation disrupts the homeostasis of intestinal microorganisms in mice from infancy to adulthood and causes more serious flora disorders after negative stress stimulation. Additionally, lack of LF during lactation leads to neuroimmune dysfunction and neurodevelopmental retardation in the hippocampus, which in turn leads to increased susceptibility to depression. In short, LF feeding deficiency during lactation increases the risk of depression in adult mice by inducing microbiota–gut–brain axis disorders.

### List Of Abbreviations

5-HT Hydroxytryptamine

AB Alcian blue ACTH Adrenocorticotropic hormone BDNFs Brain-derived nutritional factors CORT Corticosterone CSF Cerebrospinal fluid CUMS Chronic unpredictable mild stress DAI Disease activity index DEGs Differentially expressed genes DG dentate gyrus DSS Dextran sodium sulfate **ELISA** Enzyme-linked immunosorbent assay FST Forced swimming test GABA γ-Aminobutyric acid type A GAPDH glyceraldehyde-3-phosphate dehydrogenase H.E Haematoxylin and eosin IBD Inflammatory bowel disease IDO Indoleamine 2,3-dioxygenase IL. Interleukin KEGG Kyoto Encyclopedia of Genes and Genomes KO Knockout

LF Lactoferrin LPS Lipopolysaccharide MIF Macrophage migration inhibitory factor NF-κB Nuclear factor-kB OFT Open field test OTU Operational taxonomic unit PCoA Principal coordinate analysis **RT-PCR Ouantitative real-time PCR** SCFAs Short-chain fatty acids SPT Sucrose preference test TLR Toll-like-receptor TNF-α Tumour necrosis factor-a TPM Transcripts per million reads TST Tail suspension test WHO World Health Organization WT Wild-type zo-1 zonula occludens

### Declarations

### Ethical Approval and Consent to participate

All mice were treated in accordance with the guidelines of the Institutional Animal Care and Use Committee (SYXK 2020-0052). All experiments were approved by the Animal Experimentation Committee of the China Agricultural University (Beijing, China).

### **Consent for publication**

All the co-authors and participants have given their consent for publication in Journal of neuroinflammation.

### Availability of data and materials

All data are available in the main text or the supplementary materials.

### Competing interests

The authors declare no competing financial interests or potential conflicts of interest.

### Funding

We gratefully acknowledge the financial support provided by National Key R&D Program of China (2021YFD1200903), the China Agriculture Research System of MOF and MARA, Ministry of Agriculture and Villages Transgenic Major Project of China (2018ZX08007001).

### Authors' contributions

YZ, YD, WW and ZC conceived and designed the study. WW and ZC conducted the experiments and performed the analyses. WW wrote the manuscript. XW, QA helped collect and analyze the data. KH, QM checked the manuscript. All authors read and approved the final manuscript.

### Acknowledgments

We would like to thank Editage (www.editage.cn) for English language editing.

### References

- 1. Krishnadas R, Cavanagh J (2012) Depression: an inflammatory illness? J NEUROL NEUROSUR PS 83(5):495–502
- Maes M, Kubera M, Leunis J, Berk M (2012) Increased IgA and IgM responses against gut commensals in chronic depression: Further evidence for increased bacterial translocation or leaky gut. J AFFECT DISORDERS 141(1):55–62
- 3. Foster JA, Neufeld KM (2013) Gut-brain: how the microbiome influences anxiety and depression. TRENDS NEUROSCI 36(5):305–312
- 4. Lin Y, Zhang R, Lan M, Zhang T, Yan Z (2013) Relationship between BDNF-TrkB signaling pathway and the generation of anxiety disorders. Chem Life 33(6):668–672

- 5. Liu Y, Xu F, Liu S, Liu G, Yang X, Gao W et al (2020). Significance of gastrointestinal tract in the therapeutic mechanisms of exercise in depression: Synchronism between brain and intestine through GBA. PROG NEURO-PSYCHOPH. ; 103(109971)
- 6. Yu Y, Li Y (2014) Enteric glial cells and their role in the intestinal epithelial barrier. WORLD J GASTROENTERO 20(32):11273–11280
- 7. Schachter J, Martel J, Lin C, Chang C, Wu T, Lu C et al (2018) Effects of obesity on depression: A role for inflammation and the gut microbiota. BRAIN BEHAV IMMUN 69:1–8
- 8. Garate I, Garcia-Bueno B, Madrigal JLM, Bravo L, Berrocoso E, Caso JR et al (2011). Origin and consequences of brain Toll-like receptor 4 pathway stimulation in an experimental model of depression. J NEUROINFLAMM. ; 8(151)
- 9. Gonzalez Olmo BM, Butler MJ, Barrientos RM (2021) Evolution of the Human Diet and Its Impact on Gut Microbiota, Immune Responses, and Brain Health.NUTRIENTS. ; 13(1)
- Bercik P, Verdu EF, Foster JA, Macri J, Potter M, Huang X et al (2010) Chronic Gastrointestinal Inflammation Induces Anxiety-Like Behavior and Alters Central Nervous System Biochemistry in Mice. Gastroenterology 139(6):2102–2409
- 11. Blackwell J, Saxena S, Petersen I, Hotopf M, Creese H, Bottle A et al (2020) Depression in individuals who subsequently develop inflammatory bowel disease: a population-based nested case-control study. Gut 0:1–7
- Gampierakis I, Koutmani Y, Semitekolou M, Morianos I, Charalampopoulos I, Xanthou G et al (2020). Hippocampal neural stem cells and microglia response to experimental inflammatory bowel disease (IBD).MOL PSYCHIATR.
- 13. Kelly JR, Borre Y, Brien CO, Patterson E, El Aidy S, Deane J et al (2016) Transferring the blues: Depression-associated gut microbiota induces neurobehavioural changes in the rat. J PSYCHIATR RES 82:109–118
- 14. Simpson CA, Diaz-Arteche C, Eliby D, Schwartz OS, Simmons JG, Cowan CSM (2021) The gut microbiota in anxiety and depression A systematic review. CLIN PSYCHOL REV 83:101943
- 15. Victora CG, Bahl R, Barros AJ, França GV, Horton S, Krasevec J et al (2016) Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. Lancet 387(10017):475–490
- 16. Nguyen DN, Jiang P, Stensballe A, Bendixen E, Sangild PT, Chatterton DEW (2016) Bovine lactoferrin regulates cell survival, apoptosis and inflammation in intestinal epithelial cells and preterm pig intestine. J Proteom 139:95–102
- Wang W, Cheng Z, Wang X, An Q, Huang K, Dai Y et al (2021) Lactoferrin, a Critical Player in Neonate Intestinal Development: RHLF may be a Good Choice in Formula. J AGR FOOD CHEM 69(31):8726– 8736
- 18. Wang B (2016) Molecular Determinants of Milk Lactoferrin as a Bioactive Compound in Early Neurodevelopment and Cognition. J PEDIATR-US 173S:S29–S36
- 19. Milani C, Duranti S, Bottacini F, Casey E, Turroni F, Mahony J et al (2017) The First Microbial Colonizers of the Human Gut: Composition, Activities, and Health Implications of the Infant Gut

Microbiota. MICROBIOL MOL BIOL R 81(4):e17-e36

- 20. Zhao F, Hu P, Wang J, Zhu W (2021) Effects of early life lactoferrin intervention on cecal microbiota, the concentration of short chain fatty acids and the level of mucosal cytokines in suckling piglets. J Nanjing Agricultural Univ 44(1):151–159
- 21. Palucha-Poniewiera A, Podkowa K, Rafalo-Ulinska A, Branski P, Burnat G (2020) The influence of the duration of chronic unpredictable mild stress on the behavioural responses of C57BL/6J mice. BEHAV PHARMACOL 31(6):574–582
- 22. Shen Z, Xu Y, Jiang X, Wang Z, Guo Y, Pan W et al (2019) Avicularin Relieves Depressive-Like Behaviors Induced by Chronic Unpredictable Mild Stress in Mice. MED SCI MONITOR 25:2777–2784
- 23. Kim JJ, Shajib MS, Manocha MM, Khan WI (2012) Investigating Intestinal Inflammation in DSSinduced Model of IBD.Journal of Visualized Experiments. ;(60):e3678
- 24. Houdeau E, Moriez R, Leveque M, Salvador-Cartier C, Waget A, Leng L et al (2007) Sex steroid regulation of macrophage migration inhibitory factor in normal and inflamed colon in the female rat. Gastroenterology 132(3):982–993
- 25. Shi J, Zhao D, Song S, Zhang M, Zamaratskaia G, Xu X et al (2020) High-Meat-Protein High-Fat Diet Induced Dysbiosis of Gut Microbiota and Tryptophan Metabolism in Wistar Rats. J Agric Food Chem 68(23):6333–6346
- 26. Chandrasekaran K, Choi J, Arvas MI, Salimian M, Singh S, Xu S et al (2020) Nicotinamide Mononucleotide Administration Prevents Experimental Diabetes-Induced Cognitive Impairment and Loss of Hippocampal Neurons. INT J MOL SCI 21(11):3756
- 27. Herzog DP, Pascual Cuadrado D, Treccani G, Jene T, Opitz V, Hasch A et al (2021). A distinct transcriptional signature of antidepressant response in hippocampal dentate gyrus granule cells.TRANSL PSYCHIAT. ; 11(1)
- 28. Kim BH, Shenoy AR, Kumar P, Das R, Tiwari S, MacMicking JD (2011) A family of IFN-γ-inducible 65kD GTPases protects against bacterial infection. Science 332(6030):717–721
- 29. Kinoshita D, Sakurai C, Morita M, Tsunematsu M, Hori N, Hatsuzawa K et al (2019) Syntaxin 11 regulates the stimulus-dependent transport of Toll-like receptor 4 to the plasma membrane by cooperating with SNAP-23 in macrophages. MOL BIOL CELL 30(9):1085–1097
- 30. Kozai M, Kubo Y, Katakai T, Kondo H, Kiyonari H, Schaeuble K et al (2017) Essential role of CCL21 in establishment of central self-tolerance in T cells. J EXP MED 214(7):1925–1935
- 31. Eldor R, Klieger Y, Sade-Feldman M, Vaknin I, Varfolomeev I, Fuchs C et al (2015) CD247, a Novel T Cell-Derived Diagnostic and Prognostic Biomarker for Detecting Disease Progression and Severity in Patients With Type 2 Diabetes. Diabetes Care 38(1):113–118
- 32. Hao XL, Zhang FC (2004) Research progress of CD28 molecules.J. Beihua Uni; 5(2)
- 33. Wang YH, Angkasekwinai P, Lu N, Voo KS, Arima K, Hanabuchi S et al (2007) IL-25 augments type 2 immune responses by enhancing the expansion and functions of TSLP-DC-activated Th2 memory cells. J EXP MED 204(8):1837–1847

- 34. Shan DF, Wang HC, Dong Y, Yu Z (2021) The function and regulation of plgR and its relationship with diseases. Chin J Immunol 37(22):2738–2744
- 35. Lim D, Jeong J, Song J (2021) Lipocalin 2 regulates iron homeostasis, neuroinflammation, and insulin resistance in the brains of patients with dementia: Evidence from the current literature. CNS NEUROSCI THER 27(8):883–894
- 36. Ulas T, Pirr S, Fehlhaber B, Bickes MS, Loof TG, Vogl T et al (2017) S100-alarmin-induced innate immune programming protects newborn infants from sepsis. NAT IMMUNOL 18(6):622
- 37. Mohan T, Deng L, Wang BZ (2017) CCL28 chemokine: An anchoring point bridging innate and adaptive immunity. INT IMMUNOPHARMACOL 51:165–170
- 38. Sun Q, Cheng L, Zhang X, Wu Z, Weng P (2021) The interaction between tea polyphenols and host intestinal microorganisms: an effective way to prevent psychiatric disorders.FOOD FUNCT.
- 39. Rodgers LS, Fanning AS (2011) Regulation of epithelial permeability by the actin cytoskeleton. CYTOSKELETON 68(12):653–660
- 40. Ivanov Al, Bachar M, Babbin BA, Adelstein RS, Nusrat A, Parkos CA (2007) A Unique Role for Nonmuscle Myosin Heavy Chain IIA in Regulation of Epithelial Apical Junctions.PLOS ONE. ;2(e6588)
- 41. Ou J, Liang S, Guo X, Hu X (2020) alpha-Defensins Promote Bacteroides Colonization on Mucosal Reservoir to Prevent Antibiotic-Induced Dysbiosis. FRONT IMMUNOL. ; 11(2065)
- 42. Rong H, Xie X, Zhao J, Lai W, Wang M, Xu D et al (2019) Similarly in depression, nuances of gut microbiota: Evidences from a shotgun metagenomics sequencing study on major depressive disorder versus bipolar disorder with current major depressive episode patients. J PSYCHIATR RES 113:90–99
- 43. Jo J, Seo S, Park S, Kim H, Kim E, Kim J et al (2021) Gut Microbiome and Metabolome Profiles Associated with High-Fat Diet in Mice. Metabolites 11(8):482
- 44. Wu T, Zhang Y, Li W, Zhao Y, Long H, Muhindo EM et al (2021) Lactobacillus rhamnosus LRa05 Ameliorate Hyperglycemia through a Regulating Glucagon-Mediated Signaling Pathway and Gut Microbiota in Type 2 Diabetic Mice. J Agric Food Chem 69(31):8797–8806
- 45. Li X, Wang Y, Xing Y, Xing R, Liu Y, Xu Y (2020) Changes of gut microbiota during silybin-mediated treatment of high-fat diet-induced non-alcoholic fatty liver disease in mice. HEPATOL RES 50(1):5–14
- 46. Yu M, Yue J, Hui N, Zhi Y, Hayat K, Yang X et al (2021) Anti-Hyperlipidemia and Gut Microbiota Community Regulation Effects of Selenium-Rich Cordyceps militaris Polysaccharides on the High-Fat Diet-Fed Mice Model. Foods 10(10):2252
- 47. Liu X, Mao B, Gu J, Wu J, Cui S, Wang G et al (2021) Blautia-a new functional genus with potential probiotic properties? Gut Microbes 13(1):1–21
- 48. Henning SM, Yang J, Hsu M, Lee RP, Grojean EM, Ly A et al (2018) Decaffeinated green and black tea polyphenols decrease weight gain and alter microbiome populations and function in diet-induced obese mice. EUR J NUTR 57(8):2759–2769

- 49. Salzman NH, Hung K, Haribhai D, Chu H, Karlsson-Sjoeberg J, Amir E et al (2010) Enteric defensins are essential regulators of intestinal microbial ecology. NAT IMMUNOL 11(1):71–76
- 50. Xu HM, Huang HL, Liu YD, Zhu JQ, Zhou YL, Chen HT et al (2021) Selection strategy of dextran sulfate sodium-induced acute or chronic colitis mouse models based on gut microbial profile. BMC MICROBIOL 21(1):279
- 51. Zhang J, Zhou HC, He SB, Zhang XF, Ling YH, Li XY et al (2021) The immunoenhancement effects of sea buckthorn pulp oil in cyclophosphamide-induced immunosuppressed mice. FOOD FUNCT 12(17):7954–7963
- 52. Dos Santos Cruz BC, Da Conceicao LL, de Oliveira Mendes TA, de Luces Fortes Ferreira CL, Goncalves RV, Gouveia Peluzio MDC (2020) Use of the synbiotic VSL#3 and yacon-based concentrate attenuates intestinal damage and reduces the abundance of Candidatus Saccharimonas in a colitis-associated carcinogenesis model. FOOD RES INT. ; 137(109721)
- 53. Xiao S, Liu C, Chen M, Zou J, Zhang Z, Cui X et al (2020) Scutellariae radix and coptidis rhizoma ameliorate glycolipid metabolism of type diabetic rats by modulating gut microbiota and its metabolites. Appl Microbiol Biotechnol 104(1):303–317
- 54. Yue C, Li M, Li J, Han X, Zhu H, Yu G et al (2020) Medium-, long- and medium-chain-type structured lipids ameliorate high-fat diet-induced atherosclerosis by regulating inflammation, adipogenesis, and gut microbiota in ApoE(-/-) mice. FOOD FUNCT 11(6):5142–5155
- 55. Tang S, Zhang S, Zhong R, Su D, Xia B, Liu L et al (2021) Time-course alterations of gut microbiota and short-chain fatty acids after short-term lincomycin exposure in young swine. Appl Microbiol Biotechnol 105(21–22):8441–8456
- 56. Sun L, Zhang H, Cao Y, Wang C, Zhao C, Wang H et al (2019) Fluoxetine ameliorates dysbiosis in a depression model induced by chronic unpredicted mild stress in mice. INT J MED SCI 16(9):1260– 1270
- 57. Yang B, Zheng F, Stanton C, Ross RP, Zhao J, Zhang H et al (2021). Lactobacillus reuteri FYNLJ109L1 Attenuating Metabolic Syndrome in Mice via Gut Microbiota Modulation and Alleviating Inflammation. Foods. ; 10(9): 2081
- 58. Gu Y, Han Y, Ren S, Zhang B, Zhao Y, Wang X et al (2022). Correlation among gut microbiota, fecal metabolites and autism-like behavior in an adolescent valproic acid-induced rat autism model. BEHAV BRAIN RES. ; 417(113580)
- 59. Frank DN, St AA, Feldman RA, Boedeker EC, Harpaz N, Pace NR (2007) Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. Proc Natl Acad Sci U S A 104(34):13780–13785
- 60. Yang HL, Li MM, Zhou MF, Xu HS, Huan F, Liu N et al (2021) Links Between Gut Dysbiosis and Neurotransmitter Disturbance in Chronic Restraint Stress-Induced Depressive Behaviours: the Role of Inflammation. INFLAMMATION 44(6):2448–2462
- 61. Barandouzi ZA, Starkweather AR, Henderson WA, Gyamfi A, Cong XS (2020) Altered Composition of Gut Microbiota in Depression: A Systematic Review. FRONT PSYCHIATRY 11:541

- 62. Taylor AM, Thompson SV, Edwards CG, Musaad SMA, Khan NA, Holscher HD (2020) Associations among diet, the gastrointestinal microbiota, and negative emotional states in adults.NUTR NEUROSCI. ; 23(12)
- 63. Wang W, Chen L, Zhou R, Wang X, Song L, Huang S et al (2014) Increased Proportions of Bifidobacterium and the Lactobacillus Group and Loss of Butyrate-Producing Bacteria in Inflammatory Bowel Disease. J CLIN MICROBIOL 52(2):398–406
- 64. Woodman T, Strunk T, Patole S, Hartmann B, Simmer K, Currie A (2018) Effects of lactoferrin on neonatal pathogens and Bifidobacterium breve in human breast milk.PLOS ONE. ;13(e02018198)
- 65. Sriramoju B, Kanwar RK, Kanwar JR (2015) Lactoferrin induced neuronal differentiation: A boon for brain tumours. INT J DEV NEUROSCI 41:28–36
- 66. Cao P, Changmao, Liu CA, Jin Y, Xu L, Zhang Z (2021) Early-life inflammation promotes depressive symptoms in adolescence via microglial engulfment of dendritic spines.NEURON. ;(109):2573–2589
- 67. Yang CM, Shinkai Y (2013) Prdm12 is induced by retinoic acid and exhibits anti-proliferative properties through the cell cycle modulation of P19 embryonic carcinoma cells. CELL STRUCT FUNCT 38(2):197–206
- 68. Fan L, Lu Y, Shen X, Shao H, Suo L, Wu Q (2018) Alpha protocadherins and Pyk2 kinase regulate cortical neuron migration and cytoskeletal dynamics via Rac1 GTPase and WAVE complex in mice.ELIFE. ;7
- 69. Polanco J, Reyes-Vigil F, Weisberg SD, Dhimitruka I, Bruses JL (2021) Differential Spatiotemporal Expression of Type I and Type II Cadherins Associated With the Segmentation of the Central Nervous System and Formation of Brain Nuclei in the Developing Mouse. FRONT MOL NEUROSCI. ; 14(633719)
- 70. Chen Y, Zheng Z, Zhu X, Shi Y, Tian D, Zhao F et al (2015) Lactoferrin Promotes Early Neurodevelopment and Cognition in Postnatal Piglets by Upregulating the BDNF Signaling Pathway and Polysialylation. MOL NEUROBIOL 52(1):256–269
- 71. Iskandar BJ, Rizk E, Meier B, Hariharan N, Bottiglieri T, Finnell RH et al (2010) Folate regulation of axonal regeneration in the rodent central nervous system through DNA methylation. J CLIN INVEST 120(5):1603–1616
- 72. Zheng Y, Zhou LJ, Wei ZS, Dong JY, Yang LJ, Cai X (2018) Effects of maternal-infant separation on depressive behavior and expression of Gabra6 protein in hippocampus of adult rats. Guangdong Med J 39(23):3469–3472
- 73. Gonda X, Sarginson J, Eszlari N, Petschner P, Toth ZG, Baksa D et al (2017) A new stress sensor and risk factor for suicide: the T allele of the functional genetic variant in the GABRA6 gene. Sci Rep 7(1):12887
- 74. Lou D (2020) The role of hippoocampal TRPV4 in diabetic cognitive impairment. master thesis, Nanhua university,
- 75. Ying SW, Kanda VA, Hu Z, Purtell K, King EC, Abbott GW et al (2012) Targeted deletion of Kcne2 impairs HCN channel function in mouse thalamocortical circuits. PLoS ONE 7(8):e42756

- 76. Wolburg H, Wolburg-Buchholz K, Liebner S, Engelhardt B (2001) Claudin-1, claudin-2 and claudin-11 are present in tight junctions of choroid plexus epithelium of the mouse. NEUROSCI LETT 307(2):77–80
- 77. Angst J, Gamma A, Gastpar M, Lépine JP, Mendlewicz J, Tylee A (2002) Gender differences in depression. Epidemiological findings from the European DEPRES I and II studies. Eur Arch Psychiatry Clin Neurosci 252(5):201–209
- 78. Nie X, Xie R, Tuo B (2018) Effects of Estrogen on the Gastrointestinal Tract. DIGEST DIS SCI 63(3):583-596
- 79. DeFranco DB, Ramakrishnan C, Tang YT (1998) Molecular chaperones and subcellular trafficking of steroid receptors. J STEROID BIOCHEM 65(1–6):51–58
- 80. Tran A, Scholtes C, Songane M, Champagne C, Galarneau L, Levasseur M et al (2021) Estrogenrelated receptor alpha (ERR alpha) is a key regulator of intestinal homeostasis and protects against colitis. SCI REP-UK. ; 11(150731)

### **Figures**



### Figure 1

**LF feeding deficiency during suckling period increased depressive-like behaviors of CUMS mice in adult.** (A) Study design of mice experiment. (B) Percent of sucrose consumption in SPT of male mice. (C) Percent of sucrose consumption in SPT of female mice. (D) Immobility time in the TST of male mice. (E) Immobility time in the TST of female mice. (F) Immobility time in the FST of male mice. (G) Immobility time in the FST of female mice. (H) Trajectory map and distance in central area in OFT of male and female mice. (I) Serum levels of brain-derived neurotrophic factor (BDNF), (J) corticosterone (CORT), (K) adrenocorticotropic hormone (ACTH), (L) tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ). (M) interleukin-1 $\beta$  (IL-1 $\beta$ ), (N) lipopolysaccharide (LPS). Data are presented as mean ± standard error (n=8). Student's t-test was employed for comparisons between two groups; \*P < 0.05; \*\*P < 0.01; \*\*\*P < 0.001. Comparisons between two groups were analyzed using one-way ANOVA analysis followed by the Duncan test; different letters represent significant differences, P < 0.05.



### Figure 2

The damage of colon, hippocampus and microorganism composition in lactation LF feeding deficient mice after CUMS model. (A) Crypt depth in the colon of mice and representative images of H.E-stained colonic sections, 200×, scale bar = 100  $\mu$ m. (B) Zo-1, occludin mRNA expression in the colon. (C) Infiltration in the submucosa, 200×, scale bar = 100  $\mu$ m. (D) Activation of LPS-TLR4 signaling pathway in the colon. (E) Activation of LPS-TLR4 signaling pathway and microglia in the hippocampus. (F) The mRNA expression of BDNF signaling pathway in the hippocampus; (G) The alpha diversity analysis, including Shannon diversity and Chao diversity, Wilcoxon rank-sum test. (H) PCoA analysis results of male and female CUMS mice. (I) Analysis on the composition of microbial community at phylum level. (J) Student's t-test on Phylum level of male and female depression mice. (K) Linear discriminant analysis (LDA >3) scores derived from LEfSe analysis at genus level of male and female depression mice. (L Heatmap of Spearman's correlation between gut microbiota (at the genus level) and depression related indices. Data are presented as mean  $\pm$  standard error (n=8). Student's t-test was employed for comparisons between two groups; \*P < 0.05; \*\*P < 0.01; \*\*\*P < 0.001. Comparisons between more than two groups were analyzed using one-way ANOVA analysis followed by the Duncan test; different letters represent significant differences, P < 0.05.



### Figure 3

### Effects of LF feeding deficiency during lactation on intestinal development and intestinal

**microorganisms in 18-day-old suckling mice.** (A) Body weight of 18-day-old mice. (B)Small intestine and colon index of 18-day-old mice. (C) Intestinal density of small intestinal and colon of 18-day-old mice. (D) Crypt depth in the colon of 18-day-old mice and representative images of H.E-stained colonic sections, 200×, scale bar = 100 µm. (E) Counts of the goblet cells in the colon of 18-day-old mice and

representative images of AB-stained colonic sections, 200×, scale bar = 100 µm. (A-E, data are presented as mean ± standard error, n=8. Student's t-test was employed for comparisons between two groups; \*P < 0.05; \*\*P < 0.01; \*\*\*P < 0.001). (F) Principal components analysis (PCA) of small intestinal RNAseq. (G) Differential expression genes (DEGs) between wt-wt group and ko-wt group mice. (P adjust < 0.05, fold change  $\geq$  2). (H) KEGG functional annotation analysis of differential genes of wt-wt and ko-wt group mice. (I) The clustering heatmap of immune related DEGs between wt-wt group and ko-wt group. (J) Tight junction genes expression in RNAseq (F-J, n=3). (K) Alpha diversity index difference analysis of the wt-wt and ko-wt group: Shannon index, Chao index. Wilcoxon rank-sum test. (L) Principal coordinates analysis (PCoA) of bacterial beta-diversity based on the Bray-Curtis dissimilarity index. Each symbol represents a single sample of feces. (M) Composition of microbial community at phylum level of 18-day-old sucking mice. (N) Student's t-test on Phylum level of 18-day-old sucking mice. (O) Linear discriminant analysis (LDA >3) scores derived from LEfSe analysis at genus level of 18-day-old sucking mice, (K-O, n=8).



### Figure 4

LF feeding deficiency during lactation affects the development of hippocampus in 18-day old and adult mice. (A) The brain index of 18-day-old mice in wt-wt and ko-wt group. (B) The representative H.E staining of hippocampus ( $50\times$ , scale bar =  $500 \mu$ m), hippocampus CA1 and DG region ( $200\times$ , scale bar =  $100 \mu$ m) in 18-day-old mice. (A-B), n=8-9. (C)Principal components analysis (PCA) of hippocampus RNAseq in 18-day-old mice. (D)Differential gene expression between wt-wt group and ko-wt group mice (18-day-old), (P

value <0.05, fold change $\geq$ 2). (E) KEGG functional annotation analysis of differential genes in 18-day-old mice. (F, G) The clustering heatmap of immune, nervous development response-related DEGs between wt-wt group and ko-wt group (18-day-old). (H) Expression of nervous signal transduction related genes in 18-day old mice RNAseq. (I) Expression of immune related genes in the hippocampus of adult mice. (J) Expression of nervous development related genes in the hippocampus of adult mice. (K) Expression of nervous signal transduction related genes in the hippocampus of adult mice. (K) Expression of nervous signal transduction related genes in the hippocampus of adult mice. (K) Expression of nervous signal transduction related genes in the hippocampus of adult mice. (C-K), n=3.



### Figure 5

### Effects of LF feeding deficiency during lactation on intestinal development and intestinal

**microorganisms in adult mice.** (A) Body weight of four groups of adult mice. (B) Small intestinal index of four groups of adult mice. (C) Small intestinal density of four groups of adult mice. (D) Colon length of four groups of adult mice. (E) The ratio of maltase/ lactase in the duodenum, jejunum and ileum of male mice. (F) The ratio of maltase/ lactase in the duodenum, jejunum and ileum of female mice. (G) H.E stain colonic tissue sections and crypt depth of four groups of adult mice, 200×, scale bar = 100 µm. (H) The alpha diversity analysis of four groups of adult mice, including Shannon diversity and Chao diversity, Wilcoxon rank-sum test. (I) PCoA analysis results of male and female adult mice. (J) The composition of microbial community at phylum level of four groups of adult mice. (K) Student's t-test on Phylum level between WT-WT and KO-WT of male and female mice. (L) Linear discriminant analysis (LDA >3) scores derived from LEfSe analysis at genus level of male and female mice. n=7-8.

Fig. 6



#### Figure 6

Lactation LF feeding deficiency mice show more severe intestinal damage and gut microbial disorders in DSS model. (A) Changes in body weight of male mice in DSS model within 7 days. (B) Changes in body weight of female mice in DSS model within 7 days. (C) DAI scores of male mice during 7 days of DSS-induced colitis experiments. (D) DAI scores of female mice during 7 days of DSS-induced colitis experiments. (E) Survival rate of mice in DSS model. (F) Colon length of mice in DSS model. (G) H.E stain

colonic tissue sections and crypt depth of DSS model mice, 200×, scale bar = 100  $\mu$ m. (H-K) The level of TNF-α, IL-1β, IL-6, IL-10 in colon tissue of DSS model mice. Data are presented as mean ± standard error (n=8). Student's t-test was employed for comparisons between two groups. (L)Alpha diversity index difference analysis in DSS model: Shannon index, Chao index. Wilcoxon rank-sum test. (M) Principal coordinates analysis (PCoA) of bacterial β-diversity based on the Bray-Curtis dissimilarity index in DSS model. (N) Composition of microbial community at phylum level in DSS model. (O) Student's t-test on Phylum level in DSS model. (P) Linear discriminant analysis (LDA >3) scores derived from LEfSe analysis at genus level. n=6-8, \*P < 0.05; \*\*P < 0.01; \*\*\*P < 0.001.

### **Supplementary Files**

This is a list of supplementary files associated with this preprint. Click to download.

• Supplementalfiles.docx