Compound Heterozygous Mutations in *TGFBI* Cause a Severe Phenotype of Granular Corneal Dystrophy Type 2

Ikhyun Jun

Yonsei University

Yong Woo Ji

Yonsei University

Seung-il Choi

Yonsei University

Bo Ram Lee

Yonsei University

Ji Sang Min

Yonsei University

Eung Kweon Kim (✉ eungkim@yuhs.ac)

Yonsei University

Research Article

Keywords: Compound heterozygosity, granular corneal dystrophy, transforming growth factor- β -induced (TGFB1) gene

Posted Date: February 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-189647/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on March 26th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-86414-9>.

Abstract

We investigated the clinical and genetic features of patients with severe phenotype of granular corneal dystrophy type 2 (GCD2) associated with compound heterozygosity in the transforming growth factor- β -induced (*TGFB1*) gene. Patients with severe GCD2 underwent ophthalmic examination (best-corrected visual acuity test, intraocular pressure measurement, slit-lamp examination, and slit-lamp photograph analysis) and direct Sanger sequencing of whole-*TGFB1*. The patient's family was tested to determine the pedigrees. Five novel mutations (p.His174Asp, p.Ile247Asn, p.Tyr88Cys, p.Arg257Pro, and p.Tyr468*) and two known mutations (p.Asn544Ser and p.Arg179*) in *TGFB1* were identified, along with p.Arg124His, in the patients. Trans-phase of *TGFB1* second mutations was confirmed by pedigree analysis. Multiple, extensive discoid granular, and increased linear deposits were observed in the probands carrying p.Arg124His and other nonsense mutations. Some patients who had undergone phototherapeutic keratectomy experienced rapid recurrence (p.Ile247Asn and p.Asn544Ser); however, the cornea was well-maintained in a patient who underwent deep anterior lamellar keratoplasty (p.Ile247Asn). Thus, compound heterozygosity of *TGFB1* is associated with the phenotypic variability of *TGFB1* corneal dystrophies, suggesting that identifying *TGFB1* second mutations may be vital in patients with extraordinarily severe phenotypes. Our findings indicate the necessity for a more precise observation of genotype-phenotype correlation and additional care when treating *TGFB1* corneal dystrophies.

Introduction

Several transforming growth factor beta-induced gene (*TGFB1*) mutations, with corneal deposit formation and specific clinical phenotypes, have been reported to date^{1,2}. Components of *TGFB1* product, *i.e.*, transforming growth factor beta-induced protein (TGFB1p), have been detected in the deposits^{3,4}, indicating a relation between *TGFB1* mutation and deposit formation. Granular corneal dystrophy type 2 (GCD2) is an autosomal dominant disorder caused by p.Arg124His mutation of *TGFB1*⁵. While a homozygote may show severe corneal deposits from the age of 3 years⁶, slow and progressive accumulation of granular deposits, linear deposits, and diffuse haze in the corneal stroma with aging is a characteristic pathologic feature of heterozygous GCD2^{7,8}. The phenotypic variability of heterozygous GCD2 (p.Arg124His mutation) in the cornea is considerable, from few subtle white dots to multiple severe opacities throughout the stroma, even at relatively young ages^{9,10}. However, the possible cause or mechanism of this variability has not yet been elucidated.

Severe phenotypic features due to double mutations in *TGFB1* have been reported^{11–23}. In the current study, we detected five novel and two known mutations, occurring as a compound heterozygote in *TGFB1* along with p.Arg124His, which result in a very severe phenotypic variant of GCD2. In addition, we have presented each proband's family pedigree. Interestingly, individuals carrying a nonsense mutation in an allele and no p.Arg124His mutation in the opposite allele did not show an abnormal phenotype. We further performed *in vitro* experiments to investigate the effect of the identified mutations as well as p.Arg124His on TGFB1p aggregation.

Results

The compound heterozygous mutations p.Arg124His and p.His174Asp in *TGFB1*

A 35-year-old woman (Proband 1; Family 1-Patient II-3) with confluent granular deposits, dense linear deposits, and extremely severe diffuse haze on her cornea visited our clinic. Her best-corrected visual acuity (BCVA) was 20/40 in both eyes (Fig. 1a). Genetic analysis revealed p.Arg124His and p.His174Asp missense mutations in *TGFB1* (Table 1). Her 38-year-old brother showed more diffuse haze in both eyes than in any usual GCD2 heterozygote of his age (Fig. 1a). The cornea of the proband's father, who only harboured a heterozygous p.His174Asp mutation, was clear, whereas the mother, who was a GCD2 heterozygote, showed typical heterozygotic GCD2 phenotype, *i.e.*, annular granular deposits and deep linear deposits in the cornea. The pedigree analysis showed both the proband and her brother to have inherited the mutations in trans-phase (*i.e.*, p.His174Asp from the father and p.Arg124His from the mother) (Fig. 1b).

Table 1

Detected mutations increasing severity of granular corneal dystrophy 2 as a compound heterozygote with p.Arg124His in this study

Family	Nucleotide change ^a	Amino acid change	Amino acid sequence conservation ^b	Frequencies in the dbSNP database ^c	Frequencies in the gnomAD database ^d	PP2 ^e	MT ^f	PROVEAN ^g	SIFT ^h
YUGCDF-1 YUGCDF-2 YUGCDF-3	c.520C > G	p.His174Asp	<i>D. rerio</i>	NA	0.000012	Dam (0.999)	DC (0.999)	Del (-7.81)	Dam (0.000)
YUGCDF-4	c.740T > A	p.Ile247Asn	<i>D. rerio</i>	rs376340498 C = 0.000008/1 (ExAC) A = NA	C = 0.00001444 A = NA	Dam (0.804)	DC (0.999)	Del (-5.14)	Dam (0.000)
YUGCDF-5	c.263A > G	p.Tyr88Cys	<i>D. rerio</i>	NA	NA	Dam (0.863)	DC (0.999)	Del (-6.93)	Dam (0.000)
YUGCDF-6	c.770G > C	p.Arg257Pro	<i>D. rerio</i>	rs377709778 A = 0.00003/4 (ExAC) C = NA	A = 0.00002890 C = NA	Dam (0.977)	DC (0.999)	Del (-2.86)	Dam (0.010)
YUGCDF-7	c.1631A > G	p.Asn544Ser	<i>D. rerio</i>	rs777288957 G = 0.00008/10 (ExAC)	0.000053	Benign (0.281)	DC (0.999)	Del (-2.97)	Dam (0.003)
YUGCDF-8	c.535C > T	p.Arg179*	<i>D. rerio</i>	rs886059924 (MAF = NA)	A = 0.000008133 T = NA	NA	DC (1.000)	NA	NA
YUGCDF-9	c.1404T > A	p.Tyr468*	<i>X. tropicalis</i>	rs781643226 C = 0.000008/1 (ExAC) A = NA	C = 0.000004064 A = NA	NA	DC (1.000)	NA	NA

Abbreviations: Dam, damaging; DC, disease causing; Del, deleterious; MAF, minor allele frequency; MT, mutation taster; NA, not available; PP2, PolyPhen-2 prediction score Humvar; PROVEAN, Protein Variation Effect Analyzer; SIFT, Sorting Intolerant from Tolerant; SNP, single nucleotide polymorphism

^acDNA mutations are numbered according to the human cDNA reference sequence NM_004183; +1 corresponds to the A of ATG translation initiation codon.

^bAmino acid residue is continually conserved throughout evolution including the species as indicated.

^cdbSNP database (<http://www.ncbi.nlm.nih.gov/SNP>)

^dgnomAD browser (<http://gnomad.broadinstitute.org/>)

^ePolyPhen-2 prediction score HumVar ranges from 0 to 1.0; 0 = benign, 1.0 = probably damaging (<http://genetics.bwh.harvard.edu/pph2/>)

^fMutation taster (<http://www.mutationtaster.org/>). ^gPROVEAN (<http://provean.jcvi.org/index.php>). ^hSIFT (<http://sift.jcvi.org/>)

A 49-year-old man (Proband 2; Family 2-Patient II-2), not related to Family 1, visited our clinic with visual disturbances complaint; he presented with very severe diffuse haze and stromal opacities compared to other patients (Fig. 1c). Genetic and segregation analyses of the patient's family showed compound heterozygous p.Arg124His and p.His174Asp mutations in *TGFBI* (Supplementary Fig. S1a).

A 41-year-old woman (Proband 3; Family 3), not related to either Family 1 or 2, presented with severe phenotypic variation (Fig. 1d) and showed both p.His174Asp and p.Arg124His mutations in *TGFBI* upon genetic analysis. We, however, could not trace her pedigree, since the patient refused clinical evaluation and genetic analysis of her family members.

To investigate the molecular mechanism of why the phenotype of GCD2 aggravate when p.His174Asp mutation is accompanied, the degree of aggregation of p.His174Asp, and p.Arg124His mutant proteins was examined. Mixture of the cultured media of p.His174Asp expressing cells and p.Arg124His corneal fibroblasts, showed increased TGFBIp oligomer and aggregate formation during *in vitro* experiments (Supplementary Fig. S2, see also Supplementary Note).

Compound heterozygous mutations, p.Arg124His and p.Ile247Asn, in TGFBI

A 35-year-old man (Proband 4; Family 4-Patient III-1) was referred to our hospital with severe diffuse corneal haze causing visual disturbance, which recurred after phototherapeutic keratectomy (PTK) at another clinic. We obtained photographs of the cornea, acquired at the other clinic at the age of 25 years; he had undergone PTK of the right cornea at the age of 24 years, while the left cornea had not been treated. The photograph of his left untreated cornea acquired at the age of 25 years (Fig. 2a left first) showed extremely severe diffuse haze and granular deposits. Both p.Ile247Asn and p.Arg124His missense mutations were detected in *TGFBI* (Table 1), and pedigree analysis confirmed the mutations to be located in different alleles (Fig. 2b). The proband's father, aged 77 years, showed clear corneas (Fig. 2c) with only p.Ile247Asn mutation, whereas the proband's mother harboured p.Arg124His mutation and her cornea showed typical GCD2 features. Moreover, the 62-year-old paternal uncle of the proband harboured only p.Ile247Asn mutation with a clear cornea. Since the proband experienced decreased visual acuity, additional PTK for each eye was performed separately in our clinic. Corneal opacity of the proband, however, recurred rapidly, becoming diffuse and dense even after several PTK ablations. After deep anterior lamellar keratoplasty (DALK) of his left eye, the cornea has remained clear since the past 3 years (Fig. 2a).

Similar to p.His174Asp, the p.Ile247Asn mutation was also examined for the aggregation of TGFBIp. Mixture of the cultured media of p.Ile247Asn expressing cells and p.Arg124His corneal fibroblasts, showed increased TGFBIp oligomer and aggregate formation during *in vitro* experiments (Supplementary Fig. S2, see also Supplementary Note).

Heterozygous patients with GCD2 having additional p.Tyr88Cys, p.Arg257Pro, or p.Asn544Ser mutation in a different TGFBI allele

A 17-year-old woman (Proband 5; Family 5-Patient II-2) with numerous large granular deposits in the cornea visited our clinic. She experienced intermittent eye pain due to recurrent corneal erosion, although her BCVA was 20/25 in both eyes (Fig. 3a). Whole-*TGFBI* sequencing and pedigree analysis of the patient revealed that she harboured compound heterozygous p.Arg124His and p.Tyr88Cys mutations (Table 1, Supplementary Fig. S1b). No corneal deposit was detected in her 46-year-old father who harboured p.Tyr88Cys heterozygous mutation alone, whereas her 45-year-old mother harbouring p.Arg124His heterozygous mutation showed typical GCD2 heterozygous phenotype with discoid granular and star-shaped corneal deposits in both eyes (Fig. 3a). The proband's sibling, with no mutation, showed no corneal opacity.

A 28-year-old man (Proband 6; Family 6-Patient III-1) was referred to our clinic owing to visual disturbance due to corneal opacities in both eyes. BCVA of the patient was 20/40 for the right eye and 20/35 for the left eye, with both eyes showing confluent granular deposits and dense lattice deposits (Fig. 3b). Genetic analysis of the proband showed p.Arg124His and p.Arg257Pro missense mutations in *TGFBI* (Table 1, Supplementary Fig. S1c). His 54-year-old father, who only harboured heterozygous p.Arg257Pro mutation, did not show any deposit (Fig. 3b), whereas his 51-year-old mother and 49-year-old maternal uncle, both GCD2 heterozygotes, showed typical annular granular deposits with deep linear deposits (Fig. 3b), confirming trans-phase mutations in the proband.

A 32-year-old woman (Proband 7; Family 7-Patient III-11), who had undergone laser-assisted sub-epithelial keratomileusis (LASEK) in both her eyes at another clinic, visited our clinic with complains of visual disturbance, with severe diffuse stromal opacities. She underwent PTK in our clinic to remove the opacity in her right eye; however, her corneal opacity recurred rapidly after PTK (Fig. 3c). Genetic and pedigree analyses revealed that she carried both p.Arg124His and p.Asn544Ser mutations in a different allele of *TGFBI*.

(Table 1, Supplementary Fig. S1d). We could not find older living family members with p.Asn544Ser mutation alone in her family, and the younger members (aged 15 and 4 years) who harboured p.Asn544Ser mutation alone did not show any opacity.

We happened to identify a 22-year-old woman with p.Asn544Ser mutation alone in a different family during routine genetic screening before refractive surgery, following which we analysed the whole family of the patient for p.Asn544Ser mutation (Supplementary Fig. S1g). Two family members, one 82-year-old woman and the other 74-year-old woman, who were p.Asn544Ser heterozygotes, did not show any corneal opacity. A 56-year-old woman with p.Asn544Ser mutation alone (Family 10-Patient III-5), who had undergone LASEK 4 years ago on her left cornea only, to correct myopia of 2.5 D, showed single, small, white opacity, 1.4 mm away from the pupil centre (Supplementary Fig. S1h).

Heterozygous patients with GCD2 and additional nonsense mutations (p.Arg179* or p.Tyr468*) in the opposite allele of *TGFBI*

A 35-year-old woman (Proband 8; Family 8-Patient III-2) visited our clinic for treating extensive severe multiple discoid granular corneal deposits, resembling those observed in homozygous GCD2, and diffuse haze in both eyes. She had BCVA of 20/50 in both eyes and no other ocular history (Fig. 4a). After whole-*TGFBI* sequencing, she was found to carry compound heterozygous p.Arg124His and p.Arg179* [CGA (Arg) → TGA (stop codon)] mutations (Table 1). The proband's 76-year-old father and 36-year-old elder sister were found to be p.Arg124His heterozygotes with their corneas expressing the typical GCD2 phenotypes (father: superficial breadcrumb-like deposits with deep spiny deposits) (Fig. 4a). The younger sister, without mutation, had no corneal abnormality. The proband's mother and two maternal aunts, who were 70-, 64-, and 56-year-old, respectively, showed clear corneas despite having a missense mutation (p.Arg179*) in *TGFBI* (Fig. 4a, and Supplementary Fig. S1e). The findings for this family have been previously reported,²¹ except for the results of the genetic tests of the proband's mother and maternal aunts.

A 21-year-old man (Proband 9; Family 9-Patient II-1) with numerous large granular deposits in both eyes visited our clinic (Fig. 4b). Both p.Arg124His and p.Tyr468* [TAT (Tyr) → TAA (stop codon)] mutations were detected in *TGFBI* (Table 1). The proband's 44-year-old mother harbouring p.Arg124His mutation showed only mild typical corneal opacities compared to the proband (Fig. 4b). Although we could not test whether the his father harboured p.Tyr468* mutation, the proband was likely compound heterozygous for these mutations, assuming that p.Tyr468* mutation did not occur *de novo* (Supplementary Fig. S1f).

Discussion

The current study demonstrated that simultaneous presence of mutations, such as p.His174Asp, p.Ile247Asn, p.Tyr88Cys, p.Arg257Pro, p.Asn544Ser, p.Arg179*, and p.Tyr468*, along with p.Arg124His, in *TGFBI* in a compound heterozygous pattern could result in the severe phenotype of GCD2. p.Tyr88Cys, p.Arg179*, or p.Tyr468* mutation showed increased number or size of granules, while the other mutations showed increased amounts of linear deposits. Compound mutations of p.Arg124His with p.Asn544Ser or p.Arg179* have been reported previously, without identifying the phenotype of family members having p.Asn544Ser or p.Arg179* mutation alone^{16,21}; such aspects have been presented herein. The remaining five variants are newly detected mutations.

Double mutations with p.Arg124His in *TGFBI* have been identified in only 3 cases to date^{16,20,21}, out of 12 *TGFBI* double mutations reported in 13 previous studies (Supplementary Table S2)^{11–23}. Whether the previously reported p.Ser104Lysfs*27 and p.Arg179* mutations carrying p.Arg124His^{20,21} were compound heterozygous, could not be confirmed because of the refusal of family members for evaluation. We found five novel and two known *TGFBI* variants in compound heterozygotes with p.Arg124His and verified that each of the second mutation identified caused no corneal abnormality when present by itself. These results had significant clinical implications, since unexpected severe p.Arg124His variant can arise in families when p.Arg124His mutation is combined with the silent mutations mentioned above.

The cause of the presentation of the severe phenotype of GCD2 in individuals with compound heterozygous mutations of *TGFBI* is not yet clear. Mixture of TGFBIps obtained from the culture media of cells expressing p.His174Asp- or p.Ile247Asn, and p.Arg124His-mutant corneal fibroblasts, showed increased oligomer and aggregate formation in *in vitro* experiments. Several studies have reported that homozygous GCD2 corneal fibroblasts showed more oxidative stress and impaired autophagy functions, resulting in a greater accumulation of TGFBIp in the corneal fibroblasts^{24,25}. Further, intracellular TGFBIp is reportedly cleared out via lysosomes, and this function is impaired in cultured GCD2 corneal fibroblasts²⁶. We suspected the double production of mutated TGFBIps from p.Arg124His allele and heterozygous mutation in opposite alleles to result in an intracellular TGFBIp status closer to that seen in GCD2 homozygotes

than in heterozygotes. Elucidating the mechanism underlying the formation of differential phenotypes for each different second mutation requires further studies.

Interestingly, all the heterozygotes of p.His174Asp (70-year-old), p.Ile247Asn (77- and 62-year-old), p.Tyr88Cys (46-year-old), p.Arg257Pro (64- and 54-year-old), and p.Arg179* (70-, 64-, 61-, and 56-year-old) mutations showed clear corneas even at advanced age. Moreover, p.Asn544Ser heterozygotes, both 82- and 74-year-old Korean women with the mutation, showed clear cornea (Supplementary Fig. S1h). The 56-year-old woman with p.Asn544Ser mutation alone, who had undergone LASEK 4 years ago, showed only a small opacity. We could not determine whether this opacity was a result of scarring after LASEK or the exacerbation of silent cornea having p.Asn544Ser-only mutation. A study conducted in Japan reported lattice corneal dystrophy in a 68-year-old patient with a p.Asn544Ser mutation only²⁷, although the underlying cause could not be precisely explained. However, the data suggested that more reports of cases with silent second mutations would be required to verify the safety of refractive surgery. There are abundant reports of exacerbation of *TGFBI*-related corneal dystrophies, including GCD2 or GCD1 with deposits, following corneal trauma such as laser-assisted *in situ* keratomileusis^{28,29}, LASEK, photorefractive keratectomy, and refractive keratotomy^{4,30,31}.

Two types of mutations with stop codons, p.Arg179* or p.Tyr468*, combined with p.Arg124His mutation in *TGFBI* showed similar phenotypes in our study, with extensive discoid granular deposits resembling those in homozygous GCD2. Yam *et al.* reported a 31-year-old patient carrying compound heterozygous p.Arg124His and p.Leu103Leufs*28 mutations (c.307_308delCT induced frame shift and finally resulted in premature stop at position 130) in *TGFBI*, showing a pattern of corneal opacities very similar to that in our cases²⁰. In the present study, p.Arg179* or p.Tyr468* stop codon was generated at codon 179 or 468, while in the study conducted by Yam *et al.*, the stop codon occurred with a frame shift; however, the phenotype remained very similar. These data collectively showed that the presence of compound heterozygous mutation with the stop codon in one allele and p.Arg124His mutation in another could result in extensive granular deposits. Poulsen *et al.*³² reported that the macroscopic, microscopic, and ultrastructural appearance of *TGFBI*-null mouse cornea remain unaffected, suggesting that partial or complete knockdown of *TGFBI* could be a potential therapy against *TGFBI*-linked corneal dystrophies. Currently, correction of p.Arg124His mutation in cornea by knocking out the mutated allele is being attempted for the treatment of GCD2^{33,34}. Our present data indicate that the knockout of the allele should be very precise during gene therapy, in case of the heterozygote, such that only the allele containing p.Arg124His mutation is treated, leaving the normal-sequence allele intact. If the wild-type allele is altered to stop codon while the p.Arg124His-mutant allele is kept unchanged, it may result in the final corneal lesion being similar to that observed in compound heterozygotes with stop codon in the present study.

Family 1 in this study was previously reported to have extremely varied GCD2 heterozygote phenotypes¹⁰. However, the exact reason for severe phenotypic variability in the family was not known then; here, we showed the compound heterozygous mutation to be a causative factor among multiple genetic and environmental factors. Moreover, we showed that the patients with compound heterozygous *TGFBI* mutations experienced rapid recurrence after PTK. Identifying the reason underlying why the cornea remaining clear for an extended period after DALK in Proband 4 of this study would require further investigation.

In conclusion, this study reported five novel and two known genetic mutations that can predict the occurrence of the severe phenotype of heterozygous GCD2. Since phenotypic variations may indicate the presence of a new genetic change, whole-*TGFBI* sequencing would be necessary when a heterozygous patient with severe GCD2 is identified. Identification of cases with second mutations could more thoroughly explain the pathophysiology of GCD2.

Methods

Clinical investigation

This study was approved by the Institutional Review Board of Yonsei University College of Medicine (IRB No. 4-2012-0209) and followed the tenets of the Declaration of Helsinki. Written informed consent was obtained from all participants. All participants agreed to have their photos published. The patients who visited Severance Hospital from January 2007 to October 2020 underwent a detailed ophthalmological examination, including BCVA test, intraocular pressure measurement, slit-lamp examination, and slit-lamp photograph analysis. Among the patients with GCD2, the ones who presented severe phenotypes and had no other history of ocular or systemic diseases were selected as candidates for whole-*TGFBI* sequencing. After identifying other mutations, in addition to p.Arg124His, family members were enrolled in the study to perform segregation analysis.

Genomic DNA preparation and mutation analysis

TGFBI was analysed as described previously, after receiving informed consent from the participants²⁹. Briefly, 2 ml of blood was drawn from each subject and genomic DNA was extracted from their peripheral leukocytes using a QIAamp DNA Blood Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. Primers were designed to amplify all 17 exons of *TGFBI* (Supplementary Table S1). Each polymerase chain reaction (PCR) was performed using a 20- μ l reaction mixture (Maxime PCR Premix kit; iNtRON Biotechnology, Seongnam-si, Gyeonggi-do, Korea) containing 100 ng of genomic DNA, 10 pmol each of forward and reverse primers, and distilled water. Samples were amplified through 35 PCR cycles; each cycle consisted of denaturation for 20 s at 94°C, annealing for 15 s at 58°C, and extension for 50 s at 72°C. PCR was performed using a 96-well thermal cycler (Applied Biosystems, Foster City, CA, USA). Subsequently, Sanger sequencing was performed for all exons; to screen for mutations, we compared the DNA sequence of patients with the complementary DNA sequence of *TGFBI* obtained from GenBank (NC_000005.10).

TGFB1p oligomer formation, aggregate analysis, and immunoblotting

To determine oligomer formation and aggregation ability, both wild-type and p.Arg124His-mutant TGFB1p were incubated with either p.His174Asp- or p.Ile247Asn-mutant TGFB1p for 1 h at 37°C, following which immunoblotting was performed.³⁵ Details are available in the Supplementary Methods.

Declarations

Author Contributions

IJ, YWJ, and EKK conceived and designed the study. SIC and BRL conducted the experiments. IJ and JSM analysed the results. IJ, YWJ, and EKK wrote the main manuscript. SIC, BRL, and JSM performed the critical revision of the manuscript for important intellectual content. All authors contributed to the review of the literature and drafting of the manuscript, and approved the final manuscript.

Competing Interests

EKK is a medical advisory board member of Avellino Lab, USA. All other authors declare no competing interests.

References

1. Runager, K., Klintworth, G. K., Karring, H. & Enghild, J. J. The insoluble TGFB1p fraction of the cornea is covalently linked via a disulfide bond to type XII collagen. *Biochemistry* **52**, 2821-2827 (2013).
2. Lakshminarayanan, R. *et al.* Biochemical properties and aggregation propensity of transforming growth factor-induced protein (TGFB1p) and the amyloid forming mutants. *Ocul Surf* **13**, 9-25 (2015).
3. Courtney, D. G. *et al.* Protein Composition of TGFB1-R124C- and TGFB1-R555W-Associated Aggregates Suggests Multiple Mechanisms Leading to Lattice and Granular Corneal Dystrophy. *Investigative ophthalmology & visual science* **56**, 4653-4661 (2015).
4. Kim, T. I., Kim, T., Kim, S. W. & Kim, E. K. Comparison of corneal deposits after LASIK and PRK in eyes with granular corneal dystrophy type II. *Journal of refractive surgery* **24**, 392-395 (2008).
5. Han, K. E. *et al.* Pathogenesis and treatments of TGFB1 corneal dystrophies. *Prog Retin Eye Res* **50**, 67-88 (2016).
6. Moon, J. W. *et al.* Homozygous granular corneal dystrophy type II (Avellino corneal dystrophy): natural history and progression after treatment. *Cornea* **26**, 1095-1100 (2007).
7. Kim, S. W. *et al.* Characteristic features of granular deposit formation in granular corneal dystrophy type 2. *Cornea* **30**, 848-854 (2011).
8. Folberg, R. *et al.* Clinically atypical granular corneal dystrophy with pathologic features of lattice-like amyloid deposits. A study of these families. *Ophthalmology* **95**, 46-51 (1988).
9. Abazi, Z., Magarasevic, L., Grubisa, I. & Risovic, D. Individual phenotypic variances in a family with Avellino corneal dystrophy. *BMC ophthalmology* **13**, 30 (2013).
10. Han, K. E. *et al.* Extremely varied phenotypes in granular corneal dystrophy type 2 heterozygotes. *Molecular vision* **18**, 1755-1762 (2012).

11. Dighiero, P. *et al.* A novel variant of granular corneal dystrophy caused by association of 2 mutations in the TGFBI gene-R124L and DeltaT125-DeltaE126. *Archives of ophthalmology* **118**, 814-818 (2000).
12. Sakimoto, T. *et al.* A novel nonsense mutation with a compound heterozygous mutation in TGFBI gene in lattice corneal dystrophy type I. *Japanese journal of ophthalmology* **47**, 13-17 (2003).
13. Klintworth, G. K., Bao, W. & Afshari, N. A. Two mutations in the TGFBI (BIGH3) gene associated with lattice corneal dystrophy in an extensively studied family. *Investigative ophthalmology & visual science* **45**, 1382-1388 (2004).
14. Aldave, A. J. *et al.* Lattice corneal dystrophy associated with the Ala546Asp and Pro551Gln missense changes in the TGFBI gene. *American journal of ophthalmology* **138**, 772-781 (2004).
15. Frising, M., Wildhardt, G., Frisch, L. & Pitz, S. Recurrent granular dystrophy of the cornea: an unusual case. *Cornea* **25**, 614-617 (2006).
16. Yamada, N. *et al.* Double mutation (R124H, N544S) of TGFBI in two sisters with combined expression of Avellino and lattice corneal dystrophies. *Molecular vision* **15**, 974-979 (2009).
17. Zenteno, J. C., Correa-Gomez, V., Santacruz-Valdez, C., Suarez-Sanchez, R. & Villanueva-Mendoza, C. Clinical and genetic features of TGFBI-linked corneal dystrophies in Mexican population: description of novel mutations and novel genotype-phenotype correlations. *Experimental eye research* **89**, 172-177 (2009).
18. Zhong, X. *et al.* Novel and known mutations of TGFBI, their genotype-phenotype correlation and structural modeling in 3 Chinese families with lattice corneal dystrophy. *Molecular vision* **16**, 224-230 (2010).
19. Niel-Butschi, F. *et al.* Genotype-phenotype correlations of TGFBI p.Leu509Pro, p.Leu509Arg, p.Val613Gly, and the allelic association of p.Met502Val-p.Arg555Gln mutations. *Molecular vision* **17**, 1192-1202 (2011).
20. Yam, G. H. *et al.* In vitro amyloid aggregate forming ability of TGFBI mutants that cause corneal dystrophies. *Investigative ophthalmology & visual science* **53**, 5890-5898 (2012).
21. Song, J. S., Lim, D. H., Chung, E. S., Chung, T. Y. & Ki, C. S. Mutation Analysis of the TGFBI Gene in Consecutive Korean Patients With Corneal Dystrophies. *Annals of laboratory medicine* **35**, 336-340 (2015).
22. Chae, H. *et al.* Mutational spectrum of Korean patients with corneal dystrophy. *Clinical genetics* **89**, 678-689 (2016).
23. Ann, L. B. *et al.* Variant lattice corneal dystrophy associated with compound heterozygous mutations in the TGFBI gene. *Br J Ophthalmol* **101**, 509-513 (2017).
24. Kim, T. I. *et al.* Altered mitochondrial function in type 2 granular corneal dystrophy. *The American journal of pathology* **179**, 684-692 (2011).
25. Choi, S. I. *et al.* Impaired autophagy and delayed autophagic clearance of transforming growth factor β -induced protein (TGFBI) in granular corneal dystrophy type 2. *Autophagy* **8**, 1782-1797 (2012).
26. Choi, S. I., Woo, J. H. & Kim, E. K. Lysosomal dysfunction of corneal fibroblasts underlies the pathogenesis of Granular Corneal Dystrophy Type 2 and can be rescued by TFEB. *J Cell Mol Med* (2020).
27. Kawashima, M., Yamada, M., Funayama, T. & Mashima, Y. [Six cases of late-onset lattice corneal dystrophy associated with gene mutations induced by the transforming growth factor- β]. *Nippon Ganka Gakkai zasshi* **109**, 93-100 (2005).
28. Roh, M. I. *et al.* Avellino corneal dystrophy exacerbated after LASIK: scanning electron microscopic findings. *Cornea* **25**, 306-311 (2006).
29. Jun, R. M. *et al.* Avellino corneal dystrophy after LASIK. *Ophthalmology* **111**, 463-468 (2004).
30. Feizi, S., Pakravan, M., Baradaran-Rafiee, A. R. & Yazdani, S. Granular corneal dystrophy manifesting after radial keratotomy. *Cornea* **26**, 1267-1269 (2007).
31. Lee, J. H. *et al.* Exacerbation of granular corneal dystrophy type II (Avellino corneal dystrophy) after LASEK. *Journal of refractive surgery* **24**, 39-45 (2008).
32. Poulsen, E. T. *et al.* Proteomic profiling of TGFBI-null mouse corneas reveals only minor changes in matrix composition supportive of TGFBI knockdown as therapy against TGFBI-linked corneal dystrophies. *The FEBS journal* **285**, 101-114 (2018).
33. Christie, K. A. *et al.* Towards personalised allele-specific CRISPR gene editing to treat autosomal dominant disorders. *Scientific reports* **7**, 16174 (2017).
34. Taketani, Y. *et al.* Repair of the TGFBI gene in human corneal keratocytes derived from a granular corneal dystrophy patient via CRISPR/Cas9-induced homology-directed repair. *Scientific reports* **7**, 16713 (2017).

35. Choi, S. I. *et al.* Decreased catalase expression and increased susceptibility to oxidative stress in primary cultured corneal fibroblasts from patients with granular corneal dystrophy type II. *The American journal of pathology* **175**, 248-261 (2009).

Figures

Figure 1

TGFBI p.His174Asp variant aggravates granular corneal dystrophy 2 caused by p.Arg124His mutation in a compound heterozygote. (a) Slit-lamp photographs of the members of Family 1. Severe confluent granular deposits with lattice deposits were observed in Proband 1. Her brother's eyes also showed severe snowflake-like corneal deposits. (b) Pedigree of Family 1 showed both proband and her brother to have inherited the mutations in trans-phase. (c) Slit-lamp photographs of Family 2. The cornea of Proband 2 showed intensive opacities, whereas that of his mother showed mild phenotype. (d) Slit-lamp photograph of Proband 3 showed coarse granular and lattice deposits.

Family 4 – p.Arg124His + p.Ile247Asn

Figure 2

Compound heterozygous mutations, p.Ile247Asn and p.Arg124His, in TGFBI. (a) Slit-lamp photographs of Proband 4 (Family 4). Intensive snowflake-like corneal deposits were noted at the age of 25 years, and rapid recurrence occurred after phototherapeutic keratectomy (PTK). Following deep anterior lamellar keratoplasty (DALK), the cornea has been well-maintained since the past 3 years. (b) Pedigree of Family 4 confirmed compound heterozygosity. (c) Slit-lamp photographs of the parents of Proband 4. The mother, who only had heterozygous p.Arg124His mutation in TGFBI, showed milder phenotypes than the son, and the father, who had heterozygous p.Ile247Asn mutation in TGFBI, showed a clear cornea. Although TGFBI p.Ile247Asn mutation itself was silent, when accompanying p.Arg124His mutation, phenotypes of granular corneal dystrophy 2 became extensive and recurrence occurred rapidly after PTK.

Figure 3

TGFBI p.Tyr88Cys, p.Arg257Pro, and p.Asn544Ser mutations were accompanied by p.Arg124His mutation in patients with severe phenotypes. (a) Slit-lamp photographs of Proband 5 and her parents (Family 5). The proband showed numerous large discoid granular deposits. However, her mother carrying heterozygous p.Arg124His variant showed age-appropriate disease pattern, while the father carrying heterozygous p.Tyr88Cys variant showed no disease phenotype. (b) Slit-lamp photographs of the members of Family 6. Compound heterozygosity of TGFBI p.Arg124His and p.Arg257Pro mutations caused severe GCD2 phenotypes compared to the heterozygosity of TGFBI p.Arg124His mutation alone, although heterozygosity of TGFBI p.Arg257Pro mutation did not cause disease. (c) Slit-lamp photographs of Proband 7 carrying compound heterozygous mutations of TGFBI (p.Arg124His and p.Asn544Ser). The recurrence was very rapid after phototherapeutic keratectomy (PTK).

Figure 4

Nonsense mutations of TGFBI (p.Arg179* and p.Tyr468*) manifested severe granular corneal dystrophy 2 (GCD2) phenotypes. (a) Slit-lamp photographs of members of Family 8. Extensive multiple discoid granular deposits, with some annular deposits, were detected, resembling the features of homozygous GCD2 in Proband 8 carrying TGFBI p.Arg179* and p.Arg124His mutations. Her father carrying heterozygous TGFBI p.Arg124His mutation showed milder phenotypes than the proband, and the mother carrying heterozygous TGFBI p.Arg179* showed no corneal opacity. (b) Slit-lamp photographs of Proband 9 and his mother. Proband 9 harboured compound heterozygous TGFBI p.Tyr468* and p.Arg124His mutations and showed more severe discoid granular deposits than his mother who harboured heterozygous TGFBI p.Arg124His mutation.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryInformationsver210130bylJun.pdf](#)