

# Evaluating the Efficacy of Seattle-PAP for the Respiratory Support of Premature Neonates: Study Protocol for a Randomized Controlled Trial

**Carl H. Backes, MD**

Center for Perinatal Research, The Research Institute at Nationwide Children's Hospital, Department of Pediatrics, The Ohio State University Wexner Medical Center, and The Heart Center, Nationwide Children's Hospital

**Apurwa Naik, MD**

OhioHealth, Grant Medical Center

**Thomas N. Hansen, MD**

Seattle Children's Research Institute, Seattle Children's Hospital

**Larissa I. Stanberry, PhD**

Minneapolis Heart Institute Foundation

**John A. F. Zupancic, MD**

Division of Newborn Medicine, Harvard Medical School and Department of Neonatology, Beth Israel Deaconess Medical Center

**Noah H. Hillman, MD**

SSM Health, Cardinal Glennon Children's Hospital, St. Louis University

**Stephen E. Welty, MD**

Seattle Children's Research Institute, Seattle Children's Hospital and Seattle Children's Neonatology Program, CHI Franciscan Health

**Christopher R. Howard, MBA**

Seattle Children's Research Institute, Seattle Children's Hospital

**Jennifer N. Cooper, PhD**

Department of Pediatrics, The Ohio State University Wexner Medical Center, Center for Surgical Outcomes, The Research at Nationwide Children's Hospital, and Division of Epidemiology, The Ohio State University College of Public Health

**Randy R. Miller, MD**

Pediatrix Medical Group of Ohio and Mt. Carmel St. Ann's Hospital

**Jennifer L. Notestine, BSN, RNC-NIC**

Center for Perinatal Research, The Research Institute at Nationwide Children's Hospital

**Mohannad Moallem, MD**

Department of Pediatrics, The Ohio State University Wexner Medical Center

**Brian K. Rivera, MS**

Center for Perinatal Research, The Research Institute at Nationwide Children's Hospital

**Michael R. Stenger, MD**

Department of Pediatrics, The Ohio State University Wexner Medical Center

**Juli M. Kern, MD**

Pediatrix Medical Group of Ohio

**Crystal M. Alfred, MS, RRT**

Department of Pediatrics, The Ohio State University Wexner Medical Center

**Allison M. Notestine**

Center for Perinatal Research, The Research Institute at Nationwide Children's Hospital

**Jeanne C. Balough, MSN, RN, FACCE, IBCLC**

Center for Perinatal Research, The Research Institute at Nationwide Children's Hospital

**Jane M. Lamp, MS, RN-BC**

OhioHealth Research Institute, Riverside Methodist Hospital

**Charles V. Smith, PhD**

Seattle Children's Research Institute, Seattle Children's Hospital

---

**Method Article**

**Keywords:** Neonate, Continuous positive airway pressure (CPAP), Preterm Infant, Infant, Costs, Costs analysis, Randomized controlled trial

**Posted Date:** December 17th, 2018

**DOI:** <https://doi.org/10.21203/rs.1.18/v2>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on January 18th, 2019. See the published version at <https://trialsjournal.biomedcentral.com/articles/10.1186/s13063-018-3166-6>.

# Abstract

*Background:* At birth, the majority of neonates born at <30 weeks of gestation require respiratory support to facilitate transition and ensure adequate gas exchange. Although the optimal approach to the initial respiratory management is uncertain, the American Academy of Pediatrics endorses noninvasive respiratory support with nasal continuous positive airway pressure (nCPAP) for premature neonates with respiratory insufficiency. Despite evidence for its use, nCPAP failure, requiring intubation and mechanical ventilation is common. Recently, investigators have described a novel method to deliver bubble nCPAP, termed Seattle-PAP. While preclinical and pilot studies are encouraging with regard to the potential value of Seattle-PAP, a large trial is needed to compare Seattle-PAP directly with the current standard of care for bubble nCPAP (Fisher Paykel-CPAP; FP-CPAP).

*Methods:* We designed a multicenter, non-blinded, randomized controlled trial that will enroll 230 premature infants (220/7 to 296/7 weeks of gestation). Infants will be randomized to receive Seattle-PAP or FP-CPAP. The primary outcome is respiratory failure requiring intubation and mechanical ventilation. Secondary outcomes include measures of short and long-term respiratory morbidity and cost effectiveness.

*Discussion:* This trial will assess whether Seattle-PAP is more efficacious and cost effective than FP-CPAP in real-world practice among premature neonates.

*Trial Registration:* This trial has been registered with the United States National Library of Medicine ([www.clinicaltrials.gov](http://www.clinicaltrials.gov), Trial Identifier #NCT03085329). Registered on 21 March 2017.

## Background

Neonates face significant challenges for successful transition from fetal life.[1, 2] Aeration of liquid-filled lungs and the establishment of functional residual capacity are critical tasks in newborn adaptation to the extrauterine environment. Beginning with their initial breath after birth, term infants clear fluid from their lungs and establish functional residual capacity to facilitate gas exchange.[2] In contrast, among many preterm neonates, surfactant deficiency, immature epithelial sodium channels, instability of the thorax, respiratory fatigue, and apnea contribute to respiratory insufficiency, often resulting in respiratory failure and the need for respiratory support.[3-5] Although the best approach to the initial respiratory management of these infants is uncertain, evidence that even brief exposures to tracheal intubation and mechanical ventilation cause lung injury have led to interest among health care providers in the use of noninvasive respiratory support devices, primarily nasal continuous positive airway pressure (nCPAP).[6]

The American Academy of Pediatrics endorsed the use of nCPAP among premature neonates with respiratory distress.[7] This recommendation is based on data indicating lower rates of bronchopulmonary dysplasia (BPD) in infants treated with nCPAP than are observed with more invasive modes of ventilatory support.[8-10] However, nCPAP failure, requiring intubation and mechanical ventilation in preterm infants, is common, with failure rates exceeding 50% in large clinical trials.[8-10] In

developed countries, CPAP failure is associated with greater morbidity; in developing countries, CPAP failure is associated with greater mortality.[11, 12]

More recently, bubble nCPAP has re-emerged as a potential strategy to address high nCPAP failure rates. [13] Although data are limited, neonates on bubble nCPAP had lower incidence of respiratory failure (tracheal intubation and mechanical ventilation) than did infants supported on ventilator-derived CPAP. [12, 14] Despite evidence that bubble nCPAP may be advantageous for preterm infants, optimal delivery for newborns with respiratory distress is unknown. To address high failure rates associated with CPAP among preterm neonates, investigations of novel strategies to deliver more effective bubble nCPAP are warranted.[15] One potential strategy is with use of Seattle-PAP.[12, 13, 16, 17]

In general, bubble nCPAP delivery systems consist of six primary components: sources of air, oxygen, and other breathing gases, a blender to mix these gases together, a gas heater and humidifier, inspiratory and expiratory tubing (breathing circuit), a patient interface (e.g., bi-nasal prongs or nasal mask), and a pressure generator (e.g., bubbler device). The design of a bubbler device can take many forms, but is generally comprised of a tube with its distal end submerged in a body of water. The airway pressures provided by the system are determined largely by the depth of the distal end of the bubbler tube below the surface of the water (5 cm below surface = 5 cm H<sub>2</sub>O). In conventional bubble nCPAP systems, the bubbler tube points straight down (perpendicular to the surface of the water), which we define as 0°. With Seattle-PAP, the bubbler tube points initially straight down with a transition to a horizontal, then upward sloping section, which we optimized at 135° (**Figure 1**).

Preclinical evidence demonstrated that this 135° modification provides fluctuations in airway pressures, including lower frequencies pressure oscillations than those created by conventional systems.[17, 18] Moreover, a recent study showed that prematurely-born neonates (mean gestational age of 29 weeks) with minimal parenchymal lung disease exhibited lower effort of spontaneous breathing using Seattle-PAP than with conventional bubble nCPAP (Fisher Paykel; FP-CPAP).[13] While the results of preliminary studies are encouraging, safety and efficacy data among smaller, more immature preterm infants are lacking.

If the Seattle-PAP arm proves superior to conventional bubble nCPAP, the likely benefits would be large, with clear applications in both the developed world and in low and middle income countries (LMIC). However, to recommend Seattle-PAP over the current standard of care for bubble nCPAP (FP-CPAP), a large comparative trial is needed. Additionally, since respiratory failure is associated with higher daily costs related to more intensive monitoring and personnel requirements,[19] Seattle-PAP may result in lower overall treatment costs if it prevents respiratory failure. Similarly, since infants failing CPAP have longer lengths of hospital stay than those successfully supported on CPAP, total costs should be lower if use of Seattle-PAP is associated with lower incidence of CPAP failure than is use of FP-CPAP.[20] To estimate the cost effectiveness of Seattle-PAP, a formal economic evaluation ancillary to the proposed randomized controlled trial (RCT) is necessary.[21]

The present report describes the rationale and design for an ongoing, randomized controlled trial (RCT) that aims to compare the effectiveness of Seattle-PAP versus conventional FP-CPAP in the prevention of respiratory failure of premature neonates born at 220/7 - 296/7 weeks' gestational age (GA).

**Hypothesis:** We are testing the hypothesis that premature neonates supported by Seattle-PAP will have lower rates of respiratory failure (need for tracheal intubation and mechanical ventilation) than will neonates supported with conventional bubble CPAP (FP-CPAP).

**Aims:** The primary objective of this study is to compare the rates of respiratory failure from 72 hours post-delivery to 32 weeks' postmenstrual age () in neonates born at 220/7 to 296/7 weeks of gestational age (GA) who receive either Seattle-PAP or conventional bubble nCPAP (FP-CPAP).

## Methods/design

### Study Settings:

This is a multicenter, non-blinded, RCT in premature neonates born at 220/7 to 296/7 weeks' GA. The trial is ongoing at 5 hospitals within the Nationwide Children's Hospital Neonatal Research Network (NCH-NRN): Nationwide Children's Hospital (NCH) Main Campus, The Ohio State University Wexner Medical Center, OhioHealth Riverside Methodist Hospital, Mt. Carmel St. Ann's Hospital, and OhioHealth Grant Medical Center. NCH is a Level IV referral facility covering Central Ohio and parts of West Virginia and Kentucky. All neonatal intensive care units (NICUs) in the NCH-NRN routinely care for premature neonates with respiratory insufficiency, using FP-CPAP as a non-invasive respiratory support modality. Participating sites did not use Seattle-PAP prior to study commencement.

All participating sites administer exogenous surfactant if the infant requires tracheal intubation for respiratory distress. One NICU (Mount Carmel St. Ann's Hospital) has some experience with less invasive surfactant administration, in which infants on bubble nCPAP are provided surfactant via thin catheters directly placed into infants' tracheas while spontaneously breathing, thereby avoiding the need for intubation for surfactant administration.[22] All participating centers have 24 hour on-site neonatal staff.

The Nationwide Children's Hospital (NCH) Institutional Review Board (IRB) approved the trial (IRB #16-00678). The IRB at Mt. Carmel St. Ann's Hospital granted separate approval for the trial (IRB #170817-2). Of note, prior to study commencement, the Seattle-PAP device design received US FDA clearance via the 510(k) pathway [K131502] (October 11, 2013).

### Inclusion/exclusion criteria:

*Inclusion criteria are as follows:*

1. Parental consent and/or legal guardian consent given to participate in this research study.
2. Preterm infants delivered at 22 - <30 weeks of completed gestation by best obstetrical estimate.

3. Weight of <1500 grams at birth.

4. Able to be weaned from mechanical ventilator support within 72 hours of birth.

*Exclusion criteria are as follows:*

1. Infants with known major congenital cardiac (e.g., transposition of the great arteries), pulmonary (e.g., pulmonary and/or tracheal hypoplasia), or physiological (e.g., anencephaly, omphalocele, congenital diaphragmatic hernia) anomalies. For the purposes of this definition, common preterm cardiac issues such as patent ductus arteriosus and patent *foramen ovale* / atrial septal defect will not be grounds for exclusion.

2. Infants with known genetic anomalies (e.g., Trisomy 21).

3. Infants born to mothers who are unable to give informed consent and/or who do not have a legal, surrogate guardian who can provide consent.

**Recruitment:** A member of the study team will approach parents of potentially eligible infants with threatened preterm delivery between 220/7 and 296/7 weeks' GA to offer study participation and to obtain written informed consent. Consent will be obtained by team members who have been trained in obtaining consent for clinical trials and who are familiar with trial protocol. Whenever possible, consent will be obtained by someone not directly involved in the clinical care of the infant. If obtaining consent is not possible during the antenatal period, as soon as possible following birth, a member of the study team will approach the infant's parents or legal guardians to obtain consent (up to 72 hours post-delivery). Parents or guardians of infants who are not yet eligible but are likely to become eligible (e.g., infants requiring mechanical ventilation who are likely to transition from mechanical ventilation to bubble nCPAP prior to 72 hours) will also be approached. A schedule of enrollment, interventions, and assessments is shown in **Table 1**. Recruitment began in March, 2017.

**Randomization:** Eligible infants enrolled are randomly assigned to one of two treatment arms (Seattle-PAP vs. FP-CPAP) (**Figure 2**). The allocation sequence is generated using an online, computer-generated randomizer (<https://sealedenvelope.com>), with a block size of 6, stratified by gestational age (27 – 296/7 and 22 – <27 weeks' gestation). We did not stratify by site, as the treatment of premature neonates is based on shared guidelines throughout the NCH-NRN.[23] Multiple gestations (twins, triplets) are assigned to the same treatment arm.

**Intervention:** Allocated treatment is applied immediately after randomization. Infants whose condition cannot be maintained with the assigned method of noninvasive respiratory support will be intubated and the originally assigned intervention resumed after extubation. In the NCH-NRN, bubble nCPAP is given through nasal prongs/mask with an initial pressure of 5 to 6 cmH<sub>2</sub>O. As described above, the primary difference between the groups is the "bubbler" generating the pressure for the circuit (refer to **Figure 1**).

**Blinding:** Blinding of the allocated treatment is not feasible, as the mode of respiratory support is apparent to health care professionals and families. However, statisticians performing the data analyses will be blinded to group assignment. All enrolled neonates were provided clinical care according to established guidelines, including weaning, extubation, and reintubation.[23] Moreover, the primary outcome (respiratory failure, defined below) is based on an objective set of criteria applied to both groups.

**Clinical and respiratory guidelines:** Because the duration and type of respiratory support are critical end points, we have taken a number of steps to ensure that respiratory support is applied similarly to both groups: 1) NCH-NRN follows a respiratory algorithm for care (see **Supplemental Figure**); 2) thresholds for achieving the primary endpoint of respiratory failure are clearly defined (see below). In addition to standardized respiratory support, all aspects of neonatal management and treatment will be in accordance with local guidelines.[23] Internal audits to investigate maintenance and adherence to the guidelines are conducted on a routine basis.

*Treatment Failure* is defined as any of the following:

1. Tracheal intubation within 72 hours for surfactant administration after initiation of bubble nCPAP and then not extubated by 72 hours.
2. Tracheal intubation of the infant or support with biphasic CPAP (SiPAP) in the neonatal intensive care unit (NICU) after 72 hours and up to 32 weeks GA. As adjudicated by an independent party, tracheal intubation for non-respiratory issues after 72 hours and up to 32 weeks GA (e.g., surgery for retinopathy of prematurity) will not be considered a treatment failure.
3. The infant cannot sustain SpO<sub>2</sub> of at least 90%, despite noninvasive respiratory support of up to and including 8 cm H<sub>2</sub>O bubble nCPAP, and FiO<sub>2</sub> greater than 0.40 for more than 1 hour.

**Study conclusion:** Patient participation in the study will conclude at 36 weeks' corrected gestational age or at discharge, whichever comes first.

**Primary Outcome measure:** The primary outcome is treatment failure (as defined above).

**Secondary Outcomes:** We will capture important secondary outcomes, including short and longer-term respiratory morbidities (**Additional file 1: Table S1**).

**Economic Outcomes:** With a birth to hospital discharge time horizon, cost-effectiveness analysis from the perspective of the health-care system perspectives was planned *a priori* and will be conducted alongside the RCT. From a health care sector perspective, cost-effectiveness analyses include formal health sector costs (medical) costs, such as those paid by a third-party such as government or a private insurer.[24]

**Sample Size:** Using data from other recent, large neonatal randomized trials with similar populations, including the SUPPORT trial and the COIN trial,[9, 10] as well as contemporary data within the NCH-NRN,

we estimate that the proportion of patients experiencing respiratory failure in the control arm (FP-CPAP) will be 50%. Based on preclinical work, we estimate the failure rate in the Seattle PAP arm will be 30%. [13, 17] Accounting for two interim analyses, 200 neonates are needed with a two-tailed type I error rate of 0.05 and a power of 80%. [25] Multiples will be randomized as a set to the same study arm, requiring an inflation of the estimate by 1.12 to allow for the design effect due to clustering. [26, 27] Thus, the calculated sample size is 115 infants per treatment arm.

### **Data collection:**

*Clinical:* With the exception of the data related to the screening log, all remaining clinical data will be obtained from the electronic medical records (**Additional file 2: Table S2**).

*Economic:* Routinely available costs of inpatient stay will be sourced from the hospital costing units. Direct medical resource utilization will be ascertained through collection of itemized billing records and UB-04 forms, a uniform billing statement recommended by the National Uniform Billing Committee and utilized for reporting of hospital expenditures by third party payers including the Centers for Medicare and Medicaid Services (CMS). We will convert hospital-reported charges to costs by applying the appropriate CMS cost-center specific ratio of costs to charges. Total hospital costs will be the sum of the product of the number of days in each cost category and the calculated per diem costs. Physician professional fees for the initial hospitalization will be based on CMS reimbursement levels for each day of stay and non-bundled procedures.

### **Safety**

**Data and safety monitoring board (DSMB):** We have assembled a data and safety monitoring board (DSMB) to protect study subjects and monitor the overall conduct of the trial. Prior to study commencement, the 4-person DSMB agreed to the following activities: 1) review the protocol and amendments; 2) participate in the development, finalization, and approval of the DSMB Charter; 3) recommend discontinuation of the trial for safety reasons; 4) recommend changes to the study protocol (amendments) for safety reasons; 5) evaluate emergent safety information, evaluate any risk, and identify any potential safety concerns for study patients; 6) request additional data not included in the report, if deemed necessary for effective safety monitoring; 7) communicate DSMB findings and recommendations (to stop, continue, or modify the study) to the site-PI (C.B.); 8) provide written minutes on an ongoing basis following scheduled and ad hoc meetings.

Adverse events and their relationship to study, severity, time of experience, expectation, actions taken to resolve the event, and final outcome will be recorded. All serious adverse events (SAEs) will be sent within 48 hours to the DSMB and local IRB. A SAE for this study is any untoward medical occurrence that is believed by the investigators to be causally related to the study intervention and results in any of the following: life-threatening condition (that is, immediate risk of death), severe or permanent disability, and prolonged hospitalization. SAEs occurring after a subject is discontinued from the study will not be reported, unless the investigators feel that the event may have been caused by the study device. All SAEs

will be followed until satisfactory resolution is achieved or until the health care provider responsible for the care of the participant deems the event to be chronic or the patient to be stable. All expected and unexpected SAEs, whether or not they are attributable to the study intervention, will be reviewed by the site-PI (C.B.) to determine if there is a reasonable suspected causal relationship to the intervention. If the relationship is reasonable, SAEs will be reported to DSMB and local IRB for consideration.

**Interim analysis:** Interim analyses were conducted by the DSMB following enrollment of 25 and 110 infants, respectively. The analyses compared the two groups with respect to efficacy, safety, and futility. A Haybittle-Peto stopping guideline was set at  $P < 0.001$  for each interim analysis. The interim analyses were completed in June 2017 and January 2018. On the basis of these analyses, and on safety reviews conducted to date, the DSMB recommended that the trial continue without modification.

**Duration of study:** The projected study duration is 3 years, including 2½ years for subject recruitment.

**Training of clinical staff:** Comprehensive education and training was undertaken to ensure technical proficiency and compliance with the intervention and study protocol at all sites. The site-PI (C.B.), study coordinator (J.N.), and nursing leadership on the study (J.L. and J.B.), ran a 'boot-camp' at the start of the study for all participating sites, with mandatory attendance for physicians, respiratory therapists, and nursing staff to accomplish the following: 1) review differences between Seattle-PAP and the conventional FP-CPAP device; 2) identify areas for device troubleshooting; 3) demarcate study goals and objectives; 4) review safety protocols, procedures, and guidelines for clinical and respiratory care. This effort was intended to optimize reproducibility and consistency. All centers received detailed written instruction on study protocol. The PI and study coordinator were available 24 hours/day to answer any questions or concerns.

**Data management, processing, monitoring, and security:** Data generated in this study will be appropriately documented and checked for validity and accuracy. Data will be entered into the database, then the data will be matched and checked for validity and accuracy by a second person before being endorsed for analysis. A record of all discrepancies and resolutions will be kept by the study coordinator (J.N.). Outlier data will be investigated. Data will be primarily managed using REDCap (Research Electronic Data Capture).[28] REDCap is a secure, web-based application designed to support data capture for clinical studies. All data with identifiers will be stored on firewall-protected secure servers. Study monitoring visits are performed by the sponsor and their representatives after the enrollment of 50, 100, 150, and 200 infants, and upon the closure of the study.

## **Statistical analysis**

*Clinical outcomes:* Analyses will be performed using Statistical Analysis Software (SAS) Enterprise Guide version 7.15 (SAS Institute Inc., Cary, NC) using an 'as randomized' ('intention-to-treat') principle to compare the primary outcome between treatment arms. Continuous data will be expressed as means with standard deviations or as medians with ranges, whereas categorical variables will be expressed with frequencies and proportions. The primary outcome, and other binary outcomes occurring in at least 5% of

patients, will be analyzed using Pearson chi-square tests. Less common binary outcomes will be compared using Fisher exact tests. Risk differences and the number needed to treat (NNT), along with their 95% Newcombe hybrid score confidence intervals (CI) will be calculated, and risk ratios, and their 95% Koopman asymptotic score CIs, will also be calculated. Normally distributed, continuous outcomes will be compared using Student's t test, whereas the Wilcoxon rank-sum test will be used to compare continuous outcomes with skewed distributions.

In additional analyses, log binomial regression models will be used to evaluate heterogeneity in the effect of treatment with Seattle PAP on the primary outcome. Treatment effect heterogeneity will be explored across several clinical characteristics that are known to be associated with CPAP treatment failure: gestational age, birth weight, and exposure to antenatal corticosteroids.[15] Treatment effect heterogeneity will be tested by evaluating the significance of interactions between the factors of interest and treatment arm in log binomial regression models that include the factor, treatment arm, and their interaction. Effect estimates within subgroups defined by these factors will be expressed as risk ratios (RRs) with maximum likelihood-based 95 % confidence limits. Regression models will be estimated using generalized estimating equations in order to account for the inherent correlation expected with multiples. [27].

*Economic outcomes:* To inform whether it is cost-effective to incorporate FP-CPAP or Seattle-PAP into the existing health system, decision analysis will be constructed based on the primary outcome and associated hospital costs. Univariate and probabilistic sensitivity analyses will be conducted to test the impact of uncertainty in data. We will first compare the mean patient-level costs for each treatment arm, without consideration of effectiveness. Because of the anticipated skewed nature of cost data, we will model the logarithm of the mean costs directly, using a generalized linear model with a logarithmic link function and gamma distribution.[24, 29] In addition to treatment assignment, covariates will be entered into this model to account for any differences in baseline prognostic indicators (e.g., exposure to antenatal corticosteroids) that are evident despite randomization. The model will take into account clustering among twins.

We will then determine the simultaneous outcome of cost and effect, or value for money, expressed as the incremental cost effectiveness ratio, which is calculated as the difference in mean cost per patient in the Seattle-PAP and FP-CPAP groups divided by the difference in the mean effect between the 2 groups. We will determine the statistical uncertainty in the joint distribution of costs and effects using nonparametric bootstrapping, in which we will draw 1000 samples of 220 infants, with replacement, from the study data set.[30, 31] For each of the 1000 samples, we will calculate the mean cost, mean effect, and incremental cost effectiveness ratio. Results of this analysis will be presented using incremental cost-effectiveness plot and cost-effectiveness acceptability curves.[32] Parameter uncertainty, for variables such as price weights, will be estimated using deterministic sensitivity analysis, in which the results are generated again after varying the parameter through its plausible range.[30]

**Dissemination of results:** The results of the trial will be published in a peer-reviewed journal with open access. To facilitate clinical research data sharing, data from the present study will be made available at the global data sharing enterprise, Vivli ( <https://vivli.org/resources>). The Vivli platform provides an independent data repository, search engine, and a cloud-based, secure analytics platform.

**Study Progress:** The trial began recruiting in March 2017. It is expected that recruitment for the study will be completed in 2019.

## Discussion

Previous investigators have shown that tracheal intubation and mechanical ventilation are associated with ventilator-induced lung injury and airway inflammation, which contribute to long-term adverse respiratory consequences.[33, 34] Walsh *et al.* reported that duration of tracheal intubation and mechanical ventilation among premature neonates is associated with an increased risk of death or survival with neurocognitive impairment.[35] These observations have led health care providers to pursue less invasive forms of respiratory support for premature neonates, including nCPAP.[9, 10]

Nasal-CPAP (nCPAP) is an established alternative to intubation and mechanical ventilation among premature infants with respiratory distress.[7] Although meta-analytic data show that early adoption of nCPAP was associated with lower risk of BPD than was observed with intubation and ventilation, a high proportion of neonates are not adequately supported on nCPAP and require tracheal intubation and mechanical ventilation (CPAP failure).[8-10] In fact, data suggest that 34 to 83% of premature neonates maintained on nCPAP in the delivery room require tracheal intubation and mechanical ventilation.[8-10]

More recently, bubble nCPAP has re-emerged as a potential strategy to address high nCPAP failure rates.[13] Although neonates on bubble nCPAP had lower incidence of respiratory failure (tracheal intubation and mechanical ventilation) than did infants supported on ventilator-derived CPAP, optimal delivery of nCPAP to premature neonates remains unknown. [12, 14, 15]. Recognizing the need for a more efficient bubble nCPAP system among preterm infants,[15] investigators designed and developed an alternative device, termed Seattle-PAP.[12, 13, 17, 18] In preclinical studies with juvenile rabbits lavaged to produce pulmonary surfactant deficiency,[12, 13, 17, 18] Seattle-PAP provides oscillations in airway pressures at lower frequencies than are observed with other devices.[17, 18] The frequencies of airway pressure oscillations generated by Seattle-PAP are lower than are those generated by conventional bubble nCPAP. [17, 18] While conventional bubble CPAP provides stabilization and distension of small airways and alveolar spaces, the range of fluctuations in airway pressure provided by Seattle-PAP may improve recruitment of and ventilation to the low ventilation-perfusion compartments of the lung that contribute to hypoxemia.[12]

These observations led to a recent study that compared Seattle-PAP and conventional bubble nCPAP (FP-CPAP) among premature infants with an average gestational age of 29 weeks. The authors observed, among 40 neonates, lower effort of breathing with Seattle-PAP than with FP-CPAP; however, the study was not designed to detect important clinical differences between the two devices.[13] To recommend

Seattle-PAP over the current standard of care in bubble nCPAP, a larger comparative trial is needed. Thus, the present study is designed to compare the incidence of respiratory failure over the first weeks of life between the two groups. While the present study is not designed to determine differences in longer-term clinical outcomes (BPD), pre-specified secondary analyses are planned.

We acknowledge a number of challenges in the design of the trial. First, the potential benefits and risks of Seattle-PAP versus conventional bubble nCPAP (FP-CPAP) may not be expressed uniformly across infants born at all gestational ages. To that end, we determined *a priori* to perform subgroup analysis on infants born <27 weeks versus those born 27- <30 weeks of gestation. Second, our recruitment window provides enrollment up to 72 hours of life, irrespective of previous respiratory support (mechanical ventilation, CPAP) and likely over a wide range of acutely impaired lung function. We recognize that this potential heterogeneity increases the risk of a Type II error.[36] Third, some infants allocated to Seattle-PAP may receive a brief period of FP-CPAP prior to randomization, which conceivably could affect interpretation of the results. We attempted to restrict the impact of this by excluding from the trial infants who have received more than 72 hours of FP-CPAP, which we felt to be the shortest window in which seeking parental consent would be feasible. Finally, blinding of the interventions, which would have been preferable from a study design perspective, was not feasible, due to the complexity of building and operating a system that disguised which device was providing respiratory support to the infant, yet allowed the clinical care team to monitor effective engagement of the infant with the pressure generating system.

The results of the present study will inform the design of larger, multicenter investigations to understand more fully the benefits and risks of bubble nCPAP in low and middle income countries. Since CPAP failure in developing countries may result in neonatal death, potential use in low and middle income countries would be even more attractive.[11, 14] While previous investigators have described barriers to successful use of bubble CPAP in such settings (fixation devices are bulky and cover much of the infant's face; interference with parental interaction and feeding; trauma to the nasal skin or septum; need for nursing vigilance to ensure that an adequate seal), others have shown that bubble nCPAP can be applied effectively in resource-limited situations.[14, 37, 38]

The airway pressure oscillations generated with Seattle-PAP, including lower frequencies of pressure oscillations, are modestly different than are those generated with conventional bubble nCPAP.[17, 18] Although preclinical and small clinical studies conducted to date have not indicated greater risk of adverse events associated with Seattle-PAP than with conventional bubble nCPAP[13, 17, 18] the present trial will provide an opportunity to observe any unanticipated severe adverse events.

Use of CPAP among preterm infants is associated with reduced hospital stay[16], with savings exceeding \$10,000 for every six neonates treated with CPAP.[39] Given the interrelatedness of health, availability of resources, and the constraints on healthcare budgets, as well as the substantial resource utilization by preterm neonates, improving health care status is likely to have important cost implications; thus, we will

perform a formal economic evaluation ancillary to the proposed RCT with the goal of informing the practice of effective and efficient health care.

## Potential Impact

Seattle-PAP is a promising new bubble CPAP delivery system that may reduce the need for tracheal intubation and mechanical ventilation among preterm neonates. The Seattle-PAP Trial will provide relevant and timely evidence on the efficacy and safety of the Seattle PAP device compared to conventional bubble CPAP (FP-CPAP) in this vulnerable subgroup of patients.

## Declarations

*Ethics approval and consent to participate:* The Nationwide Children's Hospital (NCH) Institutional Review Board (IRB) approved the trial (IRB #16-00678). The IRB at Mt. Carmel St. Ann's Hospital granted separate approval for the trial (IRB #170817-2).

*Consent for publication:* Not applicable.

*Availability of Data and Material:* The results of the trial will be published in a journal with open-access policy to allow for data dissemination. Data will be made available at the global data sharing enterprise, Vivli (<https://vivli.org/resources>). The Vivli platform provides an independent data repository, an in-depth search engine, and a cloud-based, secure analytics platform.

*Competing interests:* T.N.H. and C.V.S. are included among the holders of patents on the Seattle-PAP device, specifically United States Patent #8499759, under the title "Broad-band, low frequency, high-amplitude, long time duration, oscillating airway pressure breathing apparatus and method utilizing bubbles". This listing is readily available on-line. The patent is also registered with the World Intellectual Property Organization (#PCT/US2009/039957), with patents/letters granted in several regions or countries, including Europe (#2303373), Korea (#10-1540948), New Zealand (#588682), China (#201310269076.X), Canada (#2720976), Vietnam (#13756), and India (#7898/DELNP/2010). A patent has also been issued in Australia (#2010241205), under a modified name. The patent holders are committed to the Global Access Policy (<http://globalaccess.gatesfoundation.org>) of the Bill & Melinda Gates Foundation. The rights to the Seattle-PAP device have been licensed to Draeger, which will have no role in the study. The remaining authors have no potentially competing interests.

*Study Sponsor:* The study sponsor is the Seattle Children's Research Institute, Seattle, Washington.

*Funding Source:* Trial funding was provided by The Bill & Melinda Gates Foundation ([www.gatesfoundation.org](http://www.gatesfoundation.org)). The Foundation played no role in the design of the study or the preparation of the manuscript. Further, there is no intent to play any role in the collection, analysis, and interpretation of the data when the study is complete.

### *Author's Contributions:*

All authors agree to be held accountable for all aspects of the work reported herein and ensure that questions related to accuracy and integrity of the work are appropriately investigated and resolved. Specific contributions are as follows:

- 1) Design and/or conception of the study and protocol: CHB, NHH, JAZ, LIS, TNH, CVS
- 2) Drafting, revision, and preparation of the manuscript: All authors
- 3) Acquisition, analysis, and/or interpretation of data: CHB, JLN, JL, JCB, AMN, CMA, BKR, JNC, LIS
- 4) Approval of the final, submitted manuscript: All authors

*Acknowledgements:* The authors wish to acknowledge the doctors, nurses, respiratory therapists, and other hospital staff in the cooperating NCH-NRN hospitals for their efforts.

The authors wish to acknowledge Edward Shepherd MD, Thomas Bartman MD, and the Cure-Me BPD team at Nationwide Children's Hospital who designed and implemented the respiratory treatment algorithm for premature neonates.

## References


1. Malkar MB, Gardner WP, Mandy GT, Stenger MR, Nelin LD, Shepherd EG, Welty SE: **Respiratory severity score on day of life 30 is predictive of mortality and the length of mechanical ventilation in premature infants with protracted ventilation.** *Pediatr Pulmonol* 2015, **50**:363-9.
2. Vyas H, Field D, Milner AD, Hopkin IE: **Determinants of the first inspiratory volume and functional residual capacity at birth.** *Pediatr Pulmonol* 1986, **2**:189-93.
3. Heldt GP, McIlroy MB: **Distortion of chest wall and work of diaphragm in preterm infants.** *J Appl Physiol (1985)* 1987, **62**:164-9.
4. Heldt GP, McIlroy MB: **Dynamics of chest wall in preterm infants.** *J Appl Physiol (1985)* 1987, **62**:170-4.
5. Hillman NH, Kemp MW, Miura Y, Kallapur SG, Jobe AH: **Sustained inflation at birth did not alter lung injury from mechanical ventilation in surfactant-treated fetal lambs.** *PLoS One* 2014, **9**:e113473.
6. Subramaniam P, Ho JJ, Davis PG: **Prophylactic nasal continuous positive airway pressure for preventing morbidity and mortality in very preterm infants.** *Cochrane Database Syst Rev* 2016:CD001243.
7. American Academy of Pediatrics: Committee on Fetus and Newborn: **Respiratory support in preterm infants at birth.** *Pediatrics* 2014, **133**:171-4.

8. Dunn MS, Kaempf J, de Klerk A, de Klerk R, Reilly M, Howard D, Ferrelli K, O'Connor J, Soll RF, Vermont Oxford Network DRMSG: **Randomized trial comparing 3 approaches to the initial respiratory management of preterm neonates.** *Pediatrics* 2011, **128**:e1069-76.
9. Morley CJ, Davis PG, Doyle LW, Brion LP, Hascoet JM, Carlin JB, Investigators CT: **Nasal CPAP or intubation at birth for very preterm infants.** *N Engl J Med* 2008, **358**:700-8.
10. Support Study Group of the Eunice Kennedy Shriver NICHD Neonatal Research Network, Finer NN, Carlo WA, Walsh MC, Rich W, Gantz MG, Laptook AR, Yoder BA, Faix RG, Das A, *et al.*: **Early CPAP versus surfactant in extremely preterm infants.** *N Engl J Med* 2010, **362**:1970-9.
11. Field S, Kelly SM, Macklem PT: **The oxygen cost of breathing in patients with cardiorespiratory disease.** *Am Rev Respir Dis* 1982, **126**:9-13.
12. Welty SE: **Continuous positive airway pressure strategies with bubble nasal continuous positive airway pressure: Not all bubbling is the same: The Seattle positive airway pressure system.** *Clin Perinatol* 2016, **43**:661-71.
13. Welty SE, Rusin CG, Stanberry LI, Mandy GT, Gest AL, Ford JM, Backes CH, Jr., Richardson CP, Howard CR, Hansen TN, *et al.*: **Short term evaluation of respiratory effort by premature infants supported with bubble nasal continuous airway pressure using Seattle-PAP and a standard bubble device.** *PLoS One* 2018, **13**:e0193807.
14. Martin S, Duke T, Davis P: **Efficacy and safety of bubble CPAP in neonatal care in low and middle income countries: a systematic review.** *Arch Dis Child Fetal Neonatal Ed* 2014, **99**:F495-504.
15. Dargaville PA, Gerber A, Johansson S, De Paoli AG, Kamlin CO, Orsini F, Davis PG, Australian, New Zealand Neonatal N: **Incidence and Outcome of CPAP Failure in Preterm Infants.** *Pediatrics* 2016, **138**.
16. Diblasi RM: **Nasal continuous positive airway pressure (CPAP) for the respiratory care of the newborn infant.** *Respir Care* 2009, **54**:1209-35.
17. Diblasi RM, Zignego JC, Smith CV, Hansen TN, Richardson CP: **Effective gas exchange in paralyzed juvenile rabbits using simple, inexpensive respiratory support devices.** *Pediatr Res* 2010, **68**:526-30.
18. Diblasi RM, Zignego JC, Tang DM, Hildebrandt J, Smith CV, Hansen TN, Richardson CP: **Noninvasive respiratory support of juvenile rabbits by high-amplitude bubble continuous positive airway pressure.** *Pediatr Res* 2010, **67**:624-9.
19. Levesque BM, Kalish LA, LaPierre J, Welch M, Porter V: **Impact of implementing 5 potentially better respiratory practices on neonatal outcomes and costs.** *Pediatrics* 2011, **128**:e218-26.
20. Merritt TA, Pillers D, Prows SL: **Early NICU discharge of very low birth weight infants: a critical review and analysis.** *Semin Neonatol* 2003, **8**:95-115.

21. Mowitz ME, Zupancic JA, Millar D, Kirpalani H, Gaulton JS, Roberts RS, Mao W, Dukhovny D: **Prospective economic evaluation alongside the non-invasive ventilation trial.** *J Perinatol* 2017, **37**:61-6.
22. Isayama T, Iwami H, McDonald S, Beyene J: **Association of noninvasive ventilation strategies with mortality and bronchopulmonary dysplasia among preterm infants: A systematic review and meta-analysis.** *JAMA* 2016, **316**:611-24.
23. Nankervis CA, Martin EM, Crane ML, Samson KS, Welty SE, Nelin LD: **Implementation of a multidisciplinary guideline-driven approach to the care of the extremely premature infant improved hospital outcomes.** *Acta Paediatr* 2010, **99**:188-93.
24. Glick H, Doshi JA, Sonnad SS, Polsky D: **Economic evaluation in clinical trials.** Second edition. ed. Oxford: Oxford University Press; 2015. x, 252 pages p.
25. Rosner B: **Fundamentals of biostatistics.** 8th edition. ed. Boston, MA: Cengage Learning; 2016. xix, 927 pages p.
26. Donner A, Birkett N, Buck C: **Randomization by cluster. Sample size requirements and analysis.** *Am J Epidemiol* 1981, **114**:906-14.
27. Hibbs AM, Black D, Palermo L, Cnaan A, Luan X, Truog WE, Walsh MC, Ballard RA: **Accounting for multiple births in neonatal and perinatal trials: systematic review and case study.** *J Pediatr* 2010, **156**:202-8.
28. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG: **Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support.** *J Biomed Inform* 2009, **42**:377-81.
29. Manning WG, Mullahy J: **Estimating log models: to transform or not to transform?** *J Health Econ* 2001, **20**:461-94.
30. Ramsey SD, Willke RJ, Glick H, Reed SD, Augustovski F, Jonsson B, Briggs A, Sullivan SD: **Cost-effectiveness analysis alongside clinical trials II—An ISPOR Good Research Practices Task Force report.** *Value Health* 2015, **18**:161-72.
31. Efron B, Tibshirani R: **An introduction to the bootstrap.** New York: Chapman & Hall; 1993.
32. Fenwick E, O'Brien BJ, Briggs A: **Cost-effectiveness acceptability curves—facts, fallacies and frequently asked questions.** *Health economics* 2004, **13**:405-15.
33. van Kaam A: **Lung-protective ventilation in neonatology.** *Neonatology* 2011, **99**:338-41.
34. Wright CJ, Kirpalani H: **Targeting inflammation to prevent bronchopulmonary dysplasia: can new insights be translated into therapies?** *Pediatrics* 2011, **128**:111-26.

35. Walsh MC, Morris BH, Wrage LA, Vohr BR, Poole WK, Tyson JE, Wright LL, Ehrenkranz RA, Stoll BJ, Fanaroff AA, *et al.*: **Extremely low birthweight neonates with protracted ventilation: mortality and 18-month neurodevelopmental outcomes.** *J Pediatr* 2005, **146**:798-804.
36. Brown GW: **Errors, Types I and II.** *Am J Dis Child* 1983, **137**:586-91.
37. McAdams RM, Hedstrom AB, DiBlasi RM, Mant JE, Nyonyintono J, Otai CD, Lester DA, Batra M: **Implementation of Bubble CPAP in a Rural Ugandan Neonatal ICU.** *Respir Care* 2015, **60**:437-45.
38. Thukral A, Sankar MJ, Chandrasekaran A, Agarwal R, Paul VK: **Efficacy and safety of CPAP in low- and middle-income countries.** *J Perinatol* 2016, **36 Suppl 1**:S21-8.
39. Buckmaster AG, Arnold G, Wright IM, Foster JP, Henderson-Smart DJ: **Continuous positive airway pressure therapy for infants with respiratory distress in non tertiary care centers: a randomized, controlled trial.** *Pediatrics* 2007, **120**:509-18.
40. Jobe AH, Bancalari E: **Bronchopulmonary dysplasia.** *Am J Respir Crit Care Med* 2001, **163**:1723-9.
41. Robertson NJ, McCarthy LS, Hamilton PA, Moss AL: **Nasal deformities resulting from flow driver continuous positive airway pressure.** *Arch Dis Child Fetal Neonatal Ed* 1996, **75**:F209-12.
42. Bell MJ, Ternberg JL, Feigin RD, Keating JP, Marshall R, Barton L, Brotherton T: **Neonatal necrotizing enterocolitis. Therapeutic decisions based upon clinical staging.** *Ann Surg* 1978, **187**:1-7.
43. Bell RS, Graham CB, Stevenson JK: **Roentgenologic and clinical manifestations of neonatal necrotizing enterocolitis. Experience with 43 cases.** *Am J Roentgenol Radium Ther Nucl Med* 1971, **112**:123-34.
44. Pinto-Martin JA, Riolo S, Cnaan A, Holzman C, Susser MW, Paneth N: **Cranial ultrasound prediction of disabling and nondisabling cerebral palsy at age two in a low birth weight population.** *Pediatrics* 1995, **95**:249-54.
45. International Committee for the Classification of Retinopathy of P: **The International Classification of Retinopathy of Prematurity revisited.** *Arch Ophthalmol* 2005, **123**:991-9.

## Figures



**Figure 1**

A schematic showing the primary components of a bubble nCPAP circuit. The differences in conventional and Seattle-PAP bubbler devices are shown. All other elements of the CPAP circuit are identical between the groups.




Figure 2

Study algorithm

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.docx](#)
- [supplement2.doc](#)
- [supplement3.doc](#)