The Relationship of Coffee Consumption and CVD Risk Factors in Elderly Patients with T2DM

Hossein sayed ghavami

Islamic Azad University, Lahijan Branch

Mehran Khoshtinat

Islamic Azad University Roudehen Branch

Sepehr Sadeghi-Farah

Islamic Azad University, Lahijan Branch

Arman Bayati Kalimani

Islamic Azad University, Lahijan Branch

Suzie Ferrie

University of Sydney

Hossein Faraji (✉ hossinfaraji133@yahoo.com)

Isfahan University of Medical Sciences

Research Article

Keywords: Coffee, Cardiovascular Diseases, Diet, Elderly, Diabetes Mellitus

Posted Date: February 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-190702/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Objective: clinical studies suggest increasing prevalence of cardiovascular disease (CVD) risk factors and diabetes among the elderly. Meanwhile, coffee as a food compound can cause adverse effects on these risk factors. The aim of the present study was to examine the relationship between coffee consumption and CVD risk factors in the elderly with type 2 diabetes mellitus (T2DM).

Methods: this cross-sectional study was performed during 2017 on 300 elderly people above 60 years of age with T2DM in Isfahan, Iran. Dietary assessment was performed using a food frequency questionnaire. Coffee consumption was classified into three groups including <1, 1-3, and >3 cups/day. Partial correlation test was used to investigate the relationship between CVD risk factors and usual coffee consumption.

Results: the mean age and body mass index of participants in this study were 70.04 ± 4.87 years and 24.74 ± 3.34 kg/m² respectively. Coffee consumption had a significant inverse relationship with FBG and DBP in the elderly with T2DM (r : - 0.117, 0.134; p : 0.046, 0.022). TG and HDL-C had a significant positive relationship with coffee consumption levels (r : 0.636, 0.128; p : 0.028, 0.029). These results were obtained after controlling for potential confounders.

Conclusion: increasing coffee consumption was linked to improved status of some CVD risk factors including FBG, HDL-C, and DBP in the elderly with T2DM. Nevertheless, increasing coffee consumption was also associated with higher TG level and had no significant effect on other risk factors. Further studies are required to confirm these results.

Introduction

Evidence suggests increasing prevalence of diseases such as diabetes, metabolic syndrome, and cardiovascular disease (CVD) due to increased inflammation following ageing (1). Meanwhile, the most important cause of mortality worldwide is associated with CVD. Researchers believe that 33.7% of the entire global mortality has been associated with CVD and it is estimated that by 2030, around 23.6 million deaths would occur annually again due to this disease (1, 2). CVD is known as one of the major causes of mortality in Iran (3). One of the most important modifiable risk factors for CVD is lifestyle. Increasing the physical activity, consuming a healthy diet (reducing saturated fatty acids, salt, trans fatty acids, and calorie consumption) and not smoking tobacco can lower the risk of developing CVD (4). Nevertheless, there are some specific food components such as coffee in the food basket which can also modify this equation (5).

Drinking coffee as a health-related common beverage has become an interesting subject worldwide in recent decades (6). Considering coffee as a component of a healthy diet, recent evidence indicates reduction in mortality caused by chronic diseases such as CVD following consumption of this compound (7). On the other hand, the caffeine in coffee can function as a physiological stimulus and thus result in side effects such as insulin resistance and hypertension (8). However, a systematic review found that

moderate coffee consumption is inversely related to CVD risk factors (9). The mechanism of effect of coffee on chronic diseases and their common risk factors is not clearly known. Nevertheless, this drink enjoys antioxidant and anti-inflammatory properties which can play a significant role in preventing CVD, cancers, neurological and psychological disorders, as well as diabetes (5, 8). The minerals and polyphenols in coffee such as chlorogenic acid can also affect the blood glucose and lipid metabolism (10). Epidemiological studies suggest that drinking coffee can reduce some risk factors leading to death in type 2 diabetes mellitus (T2DM) and CVD patients (11, 12). On the other hand, some lifestyle factors can confound this relationship. Smoking tobacco can also result in diminished positive effects of coffee (13).

To the best of our knowledge, population studies suggest that a high level of coffee consumption is common among elderly men and smokers (8, 14). Meanwhile, the results of previous studies examining the relationship between coffee consumption and CVD risk factors also show that most of these studies have been conducted in developed countries as well as on individuals with a wide age range (4, 5, 15). The number of studies in elderly populations with T2DM is limited in developing countries such as Iran. Also, considering the trend of increasing average age of Iran's population over the recent decades and increasing rate of mortality due to CVD, attention should be paid to this disease in the elderly (16). Accordingly, the aim of the present study was to explore the relationship between coffee consumption and CVD risk factors among the elderly with T2DM in Iran.

Material And Methods

Study Population

The present study was conducted in 2017 with a cross-sectional design, targeting elderly patients under care in geriatric healthcare centers in Isfahan city, Iran. The sample selection was performed as cluster sampling; based on a sample size calculation formula (prevalence: 0.5, standard error: 0.07, design effect: 1.5, significance level: 0.05), the target number of participants in this study was set at 300. The inclusion criteria were as follows: ability to perform daily activities, living in geriatric healthcare centers, having T2DM, and aged above 60 years. In other words, those with the following characteristics were not included: inability to do daily activities, not living in geriatric healthcare centers, not having T2DM, and age less than 60 years. Note that T2DM diagnosis was made by a skilled medical team and based on the medical history, drugs taken, and laboratory data registered in the patients' file. Next, the participants responded to the questions related to anthropometric information, diet, lifestyle, plus sociodemographic information. Blood samples were taken by a skilled physician from each subject to perform biochemical analyses.

Dietary assessment

The typical dietary intake of each subject was assessed by an experienced dietitian using a 168-item food frequency questionnaire (FFQ). This questionnaire includes a list of typical food items consumed by Iranians. Based on the data obtained, the frequency of food consumption over the last year was also determined on a daily, weekly, and monthly basis for each subject, with the daily consumption of each food item in the last year calculated for every senior participant. Additionally, the level of coffee consumption was quantified for each subject in the last year (cup per day/week/month). Portion sizes of the food items consumed were converted to gram/day through household measurements. To calculate total calorie, micronutrients, and macronutrients, nutrition IV software (Tinuviel Software, Warrington, UK) was employed.

Note that in this study, no question was asked about method of coffee preparation (espresso, filtered, instant, etc.) from the subjects. However, they responded to questions about the method of coffee consumption such as adding extra materials including sugar, artificial sweeteners, milk, or others. The daily coffee consumption was categorized into three groups based on standard cup size (50 ml): less than 1 cup per day (low), 1-3 cups per day (moderate), and more than 3 cups per day (high). The reliability and validity of FFQ have been demonstrated in previous studies (17). The validation of this questionnaire was also measured following comparison of dietary intake obtained from 24-h dietary recall in 12 consecutive months (one year) with the dietary intake obtained from FFQ. The reliability of the questionnaire was also examined by comparing the food groups obtained from two separate FFQs which had been completed in one year. The results obtained from Pearson correlation statistics revealed acceptable agreement between 24-h recall in 12 consecutive months (one year) and FFQ, as well as between two FFQs (completed in one year), respectively. This questionnaire also showed notable reliability with regards to coffee consumption during a one-year period (18, 19). Generally, the above points suggest that FFQ can also present valid assessment for coffee consumption in the long term.

Basic Characteristics

The lifestyle as well as sociodemographic information of patients included age, gender (male/female), smoking tobacco (yes/no), level of education (<primary school; $\geq$ primary school), alcohol consumption (yes/no), use of drugs (yes/no), patient-reported physical activity (low, moderate, high), and history of underlying disease (yes/no; except for T2DM). Anthropometric measurements captured weight, height, waist circumference. The weight of each patient was measured by a standard digital balance (accuracy: 100 g) with minimal clothing and barefoot). The patients' height was measured by a measuring tape (barefoot) with accuracy of 0.5 cm, after which body mass index (BMI) was determined via wt/ht^2 formula. Finally, waist circumference was measured through an inelastic measuring tape and in the thinnest part below the ribs.

Biochemical analyses

The blood samples of each participant were collected after 12 h of fasting. This was performed based on standard protocols and by an experienced medical team. Using a proper kit and via enzymatic-colorimetric glucose oxidase method, the fasting plasma glucose (FPG) was measured. The enzymatic-colorimetric method was utilized to measure the lipid profile including total cholesterol (TC), low-density lipoprotein (LDL-C), high-density lipoprotein (HDL-C), and triglyceride (TG). Plasma homocysteine was also measured by ELISA method and via Axis/homocysteine EIA kit (IBL Co., Germany). Finally, systolic and diastolic blood pressure (SBP and DBP) of patients was measured by a proper sphygmomanometer (Microlife BP AG1-20, Espenstrasse 139/ 9443/widnau/Switzerland) by a trained nursing technician.

Statistical analysis

The variables examined in this study were described according to coffee consumption classification, and reported as number, percentage (%), and Mean $\pm$ SD in relevant tables. The comparison of categorical variables was also done among coffee consumption subgroups via Chi-square. This comparison for continuous variables was performed through One-way ANOVA test and Kruskal-Wallis test. Note that data normality was tested by Kolmogorov-Smirnov test. Also, homogeneity of variances for the data of study was assessed by Leven's statistics. The relationship between each CVD factor and coffee consumption level (cups/day) was captured using partial correlation test. In this relationship test, the levels of energy intake, caffeine consumption, smoking cigarette, physical activity, BMI, gender, added sugar, history of underlying disease (except for T2DM), history of taking drugs, and level of education were also controlled. All of the analyses performed in this study were done by SPSS software (version 22; Chicago, IL, USA), with the significance level considered p -value <0.05 .

Results

The patients participating in this study had a mean age of 70.04 ± 4.87 years, and BMI of 24.74 ± 3.34 kg/m²; out of this value, 52.3% had BMI above 25 kg/m², while 47.7% had BMI below 25 kg/m². From among all participants, 55% were female and 45% were male. In addition, 93.3% of participants did not smoke, and 71.3% reported low physical activity. The mean coffee consumption among the participants was 2.43 ± 1.41 cups/day, equivalent to 121.5 mL coffee drink (1 cup: 50 mL). Note that none of the participants reported alcohol consumption. Also, all participants reported consumption of caffeinated coffee plus added white sugar. In other words, none of them reported consumption of decaffeinated coffee.

The information related to lifestyle factors, caffeine intake, measurements related to each CVD risk factor, as well as the demographic data based on daily coffee consumption are tabulated in Table 1. Moderate coffee consumption (1-3 cups/day) was higher among those with a high level of education ($\geq$ primary school) compared to less educated patients ($<$ primary school) ($P=0.030$). The fasting plasma glucose and diastolic blood pressure were significantly lower among those consuming coffee ($P=0.015$, 14). On the other hand, with increase in coffee consumption, high-density lipoprotein cholesterol increased

significantly ($P=0.034$). Further, the blood triglyceride was significantly lower in those reporting moderate coffee consumption (1-3 cups/day), as compared to the two other groups ($P=0.036$) (Table 1).

Table 2 also reports the relationship between CVD risk factors and coffee consumption among the elderly with T2DM using partial correlation statistics. Results showed that with increase in coffee consumption, FPG and DBP were significantly decreased ($r: -0.117; -0.134, P=0.046; 0.022$). On the other hand, blood TG increased significantly with increase in coffee consumption ($r: 0.636; P=0.028$). The results obtained after adjusting for the variables of weight, BMI, gender, energy intake, caffeine consumption, history of underlying disease (other than T2DM), cigarette smoking, physical activity, added sugar, history of taking drugs, and level of education were also reported (Table 2).

Table 1
characteristics of elderly subjects with T2DM according to coffee consumption subgroups, Isfahan, Iran,
2017

Characteristics [†]		Coffee intake (cups/day)				p- Value *
		< 1	1–3	> 3	Total	
No. of patients, n (%)		83 (27.7)	141 (47)	76 (25.3)	300 (100)	
Age (year)		69.67 ± 4.87	70.02 ± 5.12	70.48 ± 4.37	70.04 ± 4.87	0.576 _a
Body mass index (kg/m ²)		24.78 ± 3.23	24.93 ± 3.42	24.35 ± 3.33	24.74 ± 3.34	0.471 ^a
Waist circumference (cm)		96.65 ± 4.91	96.30 ± 5.17	97.14 ± 6.24	96.61 ± 5.39	0.425 _c
Sex, n (%)	Men	36 (43.4)	68 (48.2)	31 (40.8)	135 (45)	0.541 _b
	Women	47 (56.6)	73 (51.8)	45 (59.2)	165 (55)	
Physical activity level, n (%)	Low	60 (72.3)	103 (73)	51 (67.1)	214 (71.3)	0.641 _b
	Moderate/High	23 (27.7)	38 (27)	25 (32.9)	86 (28.7)	
Smoking status, n (%)	Never	76 (96.1)	131 (92.9)	73 (96.1)	280 (93.3)	0.478 _b
	Smoker	7 (8.4)	10 (7.1)	3 (3.9)	20 (6.7)	
Educational level, n (%)	≥ Primary school	34 (41)	73 (51.8)	47 (61.8)	154 (51.3)	0.030 _b
	< Primary school	49 (59)	68 (48.2)	29 (38.2)	146 (48.7)	
Drug uses, n (%)	No	62 (74.7)	102 (72.3)	53 (69.7)	217 (72.3)	0.784 _b
	Yes	21 (25.3)	39 (27.7)	23 (30.3)	83 (27.7)	
Diseases history €, n (%)	No	48 (57.8)	87 (61.7)	47 (61.8)	182 (60.7)	0.825 _b
	Yes	35 (42.2)	54 (38.3)	29 (38.2)	118 (39.3)	
Caffeine intake (mg/day)		45.87 ± 19.24	46.29 ± 18.28	47.46 ± 20.38	46.47 ± 19.04	0.644 _c
Total energy intake (kcal/day)		2093.19 ± 549.84	2009.05 ± 555.91	2124.43 ± 535.72	2061.56 ± 549.71	0.280 _a
FPG (mg/dL)		157 ± 12.98	151.52 ± 12.18	151.09 ± 13.70	152.93 ± 13.01	0.015 _c
TG (mg/dL)		147.07 ± 22.67	140.42 ± 22.94	147.57 ± 22.98	144.07 ± 23.06	0.036 _a

TC (mg/dL)	180.51 ± 15.70	181.03 ± 15.83	180.87 ± 15.73	180.85 ± 5.72	0.972 _α
LDL-c (mg/dL)	96.36 ± 15.03	94.55 ± 15.06	96.11 ± 15	95.44 ± 15.01	0.620 _α
HDL-c (mg/dL)	47.73 ± 7.30	47.46 ± 6.93	49.92 ± 6.10	48.16 ± 6.89	0.034 _α
Homocysteine (μmol/L)	12.46 ± 2.89	12.42 ± 2.95	12.57 ± 3.52	12.47 ± 3.08	0.944 _α
SBP (mm Hg)	122.28 ± 9.88	125.05 ± 9.85	122.42 ± 10.32	123.62 ± 10.03	0.068 _α
DBP (mm Hg)	78.28 ± 10.16	77.11 ± 9.99	73.78 ± 10.02	76.59 ± 10.15	0.014 _α
<p>Abbreviations: T2DM: Type 2 diabetes mellitus; FPG: Fasting plasma glucose; TG: Triglyceride; TC: Total cholesterol; LDL-c: Low-density lipoprotein cholesterol; HDL-c: High-density lipoprotein cholesterol; SBP: Systolic blood pressure; DBP: Diastolic blood pressure. ^a One-way ANOVA test; ^b Chi-square test; ^c Kruskal-Wallis test.</p> <p>€ Except for type 2 diabetes mellitus; †All values are expressed as mean ± standard deviation (SD) and number (percent). * P-value <0.05 considered as significant level.</p>					

Table 2
Partial correlation of CVD risk factors with coffee consumption among elderly patients

Variables [®]	coffee consumption (cups/day)		
	Mean ± SD	r	P [*]
FPG (mg/dL)	152.93 ± 13.01	-0.117	0.046
TG (mg/dL)	144.07 ± 23.06	0.636	0.028
TC (mg/dL)	180.85 ± 5.72	- 0.009	0.881
LDL-c (mg/dL)	95.44 ± 15.01	-0.006	0.916
HDL-c (mg/dL)	48.16 ± 6.89	0.128	0.029
Homocysteine (μmol/L)	12.47 ± 3.08	0.020	0.739
SBP (mm Hg)	123.62 ± 10.03	-0.014	0.809
DBP (mm Hg)	76.59 ± 10.15	-0.134	0.022
Abbreviations: CVD: Cardiovascular diseases; FPG: Fasting plasma glucose; TG: Triglyceride; TC: Total cholesterol; LDL-c: Low-density lipoprotein cholesterol; HDL-c: High-density lipoprotein cholesterol; SBP: Systolic blood pressure; DBP: Diastolic blood pressure. [®] Adjusted for BMI, sex, age, physical activity, added sugars, total energy intake, smoking, education level, history of diseases (except for type 2 diabetes mellitus), drug uses and intake of caffeine. * P-value <0.05 considered as significant level.			

Discussion

The current cross-sectional study investigated the relationship between coffee consumption and CVD risk factors amongst elderly patients with T2DM. The results showed that there was an inverse relationship between coffee consumption and FBG as well as DBP levels among the elderly with T2DM. On the other hand, coffee consumption was associated with increased TG and HDL-C. These results were obtained after adjusting for potential confounders. In other words, the patients who consumed more than 3 cups of coffee per day had lower FBG and DBP, while higher HDL-C and TG.

Research on examining the relationship between coffee consumption and CVD risk factors has mostly been performed among healthy individuals and with different age ranges across various countries. There are few studies assessing the relationship between coffee consumption and CVD risk factors among the elderly, who are considered as an age group with high coffee consumption (5). The research results suggest that the effects of coffee on individuals can be different. Some of these studies suggest inverse relationship between coffee consumption and CVD risk factors, while others have shown a direct relationship between coffee consumption and CVD risk factors (4, 20, 21). The important point is that some clinical studies have suggested that there is an inverse relationship between coffee consumption

and CVD risk factors in T2DM patients (22, 23). The results of the Nurses' Health Study indicated that coffee consumption leads to diminished risk of developing diabetes (24). Also, the study by Panagiotakos et al. indicated that moderate coffee consumption would also reduce the risk of developing T2DM among the elderly (25). In order to justify these discrepancies, the following points can be mentioned: type of study design, participants (healthy, patient, teenager, young, or old) as well as various dietary factors such as daily coffee consumption and different forms of coffee consumption (4).

Various risk factors have been identified for CVD, with elevated blood sugar being one of the notable factors in this regard. Studies suggest that daily coffee consumption has antidiabetic effects (26). Coffee consumption can also affect blood glucose levels in different ways including blocking alpha-glucosidase 1 and glucose-6-phosphate translocase 1, elevating glucose-like peptide 1 (GLP-1) level and reducing the amount of glucose released from the liver (27). Nevertheless, the effect of coffee and caffeine consumption on glucose homeostasis in the long run has remained understudied. The results of the present research indicated an inverse relationship between coffee consumption and FBG among the elderly with T2DM. In line with our results, the study by Mirmiran et al. showed that coffee consumption would significantly reduce the risk of developing T2DM (28). On the other hand, the results by Yarmolinsky et al. showed that coffee consumption had no significant relationship with FBG and HbA1c, but coffee consumption can significantly lower the 2-h postload glucose level (29). Also, Keirns et al. found that coffee consumption does not have a significant effect on FBG and glycemic response (30). In order to justify the discrepancies across these results, the following hypotheses can be mentioned. Considering the changes of glucose homeostasis in T2DM patients as well as the altered absorption capacity for food among the elderly, the influence of coffee on blood glucose level may also be dependent on food intake and postprandial modifications (29, 31). Further, the role of genetics in investigating the relationship between coffee and blood glucose level is also notable. Genetic diversity in the cytochrome p-450 hepatic enzyme (P450 1A1 (CYP1A2)) can also affect the relationship between coffee and FBS (32). This enzyme has also different alleles as the main factor of coffee metabolism in the liver. Studies show that individuals with slow allele of CYP1A2 * 1F have impairments in their FBG (33, 34).

The mechanism of effect of coffee on the lipid profile is still not fully understood. Some studies suggest that the antioxidants in coffee (e.g. chlorogenic acid, caffeic acid, trigonelline) may decrease lipid oxidation and then cause elevated TC as well as LDL-C (4, 35, 36). On the other hand, one of the results of our study showed that coffee consumption resulted in higher TG and HDL-C, but it had no significant effect on TC and LDL-C. Confirming our results, the study by Alhamhany et al. showed that consumption of green coffee led to elevated HDL-C and VLDL in 35 obese individuals aged 20-55 years (37). The study by Ihim et al. showed that coffee consumption resulted in elevated HDL-C among 30 men 18-30 years of age, but it had no significant effect on TG, LDL-C, and TC (38). On the other hand, the study by Panagiotakos et al. with the aim of examining the relationship between coffee consumption and prevalence of T2DM showed that drinking coffee had no significant effect on TG, LDL-C, TC, and HDL-C. The participants in this study included 937 elderly with the age range of 65-100 years (25). Also, the results of Miranda et al. suggested that coffee consumption among 557 healthy participants in São

Paulo, Brazil had no significant effect on any lipid factors (TC, TG, HDL-C, and LDL-C) (4). There is also a hypothesis with regards to the method of coffee preparation. Unfiltered coffee is packed with oil granules (cafestol, diterpenes and kahweol), which may lead to elevated cholesterol and thus higher LDL, TC, and TG (35, 39). However, coffee filtration results in removal of the oil granules mentioned above, and may lead to an improved lipid profile (4, 40). Since the participants in the present study may have used various methods of coffee preparation, the above hypothesis could be considered as a possible explanation for the mixed results.

Hypertension is known as one of the significant risk factors of CVD as well as diabetes (41). So far, numerous studies have been conducted on examining the effect of coffee consumption on blood pressure; however, their results are contradictory with each other. The present study showed that coffee consumption had an inverse relationship with DBP, but it had no significant relationship with SBP among the elderly with T2DM. In this regard, the results of the study by Miranda et al. indicated that coffee consumption among 557 healthy individuals with the mean age of 45 years can significantly reduce the chance of developing high systolic or diastolic blood pressure (4). Meanwhile, one of the clinical studies conducted on the elderly to examine the relationship between coffee and prevalence of T2DM also reported that increased coffee consumption may lead to higher SBP, but it had no effect on DBP (25). Further, the study by Noordzij et al. also showed that regular coffee consumption can significantly increase both SBP and DBP (42). One of the important hypotheses to examine the discrepant results discussed above is related to caffeine. Evidence suggests that caffeine consumption causes significant elevation of blood pressure, but coffee consumption does not have such an effect (43). One of the reasons mentioned with regards to this hypothesis is related to the other chemical compounds in coffee, which may neutralize the effect of caffeine on elevating blood pressure. This blood pressure lowering effect may occur through vasodilation by elevating nitric oxide (NO) level and improving vascular endothelial function (44, 45). These chemical compounds include potassium, magnesium, caffeic acid, p-coumaric acids, ferulic acid, and chlorogenic acid (46). Since coffee consumption among the elderly population is relatively high (5), it can be concluded that coffee consumption in this group of people can also reduce blood pressure.

The present study had the following limitations: 1) considering the subject of the present study, assessment of coffee consumption in most previous studies involved evaluating caffeinated and non-caffeinated coffee; however, all participants in our study reported only caffeinated coffee consumption, which may somehow reduce the validity of the final conclusion. 2) the design of the present study was cross-sectional, which can increase recall biases especially in assessing dietary intake and dietary habits in the elderly. 3) considering the customs in Iran, none of the participants reported alcohol consumption. In other words, some participants may deny having alcohol when they do have some alcohol intake. This non-report may have been intentional; hence, alcohol consumption may have affected each of the outcomes examined in our study and subjected the final conclusion to bias. The strong points of this study are as follows: 1) this study has been one of the rarest studies based on the elderly population in Iran which examined the relationship between an important range of CVD risk factors and coffee

consumption simultaneously. 2) the results of the present study have been obtained by controlling the most important potential confounders, which itself can have increased the validity of this study.

Conclusions

The current study indicated that increasing coffee consumption among the elderly with T2DM had a positive relationship with changes in HDL and TG, and a negative relationship with FBG and DBP, but it had no effect on other CVD risk factors in these individuals. Generally, this study showed that increasing coffee consumption may reduce the major and important CVD risk factors among the elderly with T2DM. Nevertheless, further studies seem to be required in order to confirm the results of the present study.

Declarations

Acknowledgements

Not applicable.

Funding

Nil.

Conflicts of interest

There are no conflicts of interest.

Ethics approval and consent to participate

This study was approved by the ethics committee of Isfahan University of Medical Sciences (Ir.mui.rec.1395.1.200), with all subjects signing a written informed consent form.

Consent for publication

Not applicable.

Methodology statement and availability of data

All research methods were carried out in accordance with relevant guidelines and regulations and the datasets used for this study are available from the corresponding author.

Authors' contributions

- 1) HGh and MKh: Study designing, drafted the manuscript and final checking.
- 2) SF and HF: Manuscript editing, contributed in the conceptualization of the study and conducted all statistical analyses.
- 3) SSF and ABK: Contributed in preparation of the study protocol and outcomes assessing.

References

1. Darooghegi Mofrad M, Namazi N, Larijani B, Surkan PJ, Azadbakht L. Association of the Elderly Dietary Index with cardiovascular disease risk factors in elderly men: A cross-sectional study. *Nutrition & Dietetics*. 2019;76(5):580-8.
2. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Executive summary: heart disease and stroke statistics—2015 update: a report from the American Heart Association. *Circulation*. 2015;131(4):434-41.
3. Talaei M, Sarrafzadegan N, Sadeghi M, Oveisgharan S, Marshall T, Thomas GN, et al. Incidence of cardiovascular diseases in an Iranian population: the Isfahan Cohort Study. *Archives of Iranian medicine*. 2013;16(3).
4. Miranda AM, Steluti J, Fisberg RM, Marchioni DM. Association between coffee consumption and its polyphenols with cardiovascular risk factors: A population-based study. *Nutrients*. 2017;9(3):276.
5. Torres-Collado L, Garcia-de-la-Hera M, Navarrete-Muñoz E, Notario-Barandiaran L, Gonzalez-Palacios S, Zurriaga O, et al. Coffee consumption and mortality from all causes of death, cardiovascular disease and cancer in an elderly Spanish population. *European journal of nutrition*. 2019;58(6):2439-48.
6. de Melo Pereira GV, de Carvalho Neto DP, Júnior AIM, do Prado FG, Pagnoncelli MGB, Karp SG, et al. Chemical composition and health properties of coffee and coffee by-products. *Advances in Food and Nutrition Research*. 91: Elsevier; 2020. p. 65-96.
7. Grosso G, Godos J, Galvano F, Giovannucci EL. Coffee, caffeine, and health outcomes: an umbrella review. *Annual review of nutrition*. 2017;37:131-56.
8. Torres-Collado L, García-de la Hera M, Navarrete-Muñoz EM, Compañ-Gabucio LM, Gonzalez-Palacios S, Vioque J. Coffee drinking and associated factors in an elderly population in Spain. *International journal of environmental research and public health*. 2018;15(8):1661.

9. Ding M, Bhupathiraju SN, Satija A, van Dam RM, Hu FB. Long-term coffee consumption and risk of cardiovascular disease: a systematic review and a dose–response meta-analysis of prospective cohort studies. *Circulation*. 2014;129(6):643-59.
10. Pimpley V, Patil S, Srinivasan K, Desai N, Murthy PS. The chemistry of chlorogenic acid from green coffee and its role in attenuation of obesity and diabetes. *Preparative Biochemistry & Biotechnology*. 2020:1-10.
11. Shin S, Lim J, Lee H-W, Kim CE, Kim S-A, Lee J-k, et al. Association between the prevalence of metabolic syndrome and coffee consumption among Korean adults: results from the Health Examinees study. *Applied Physiology, Nutrition, and Metabolism*. 2019;44(12):1371-8.
12. Voskoboinik A, Koh Y, Kistler PM. Cardiovascular effects of caffeinated beverages. *Trends in cardiovascular medicine*. 2019;29(6):345-50.
13. Bjørngaard JH, Nordestgaard AT, Taylor AE, Treur JL, Gabrielsen ME, Munafò MR, et al. Heavier smoking increases coffee consumption: findings from a Mendelian randomization analysis. *International journal of epidemiology*. 2017;46(6):1958-67.
14. Lieberman HR, Stavinoha T, McGraw S, White A, Hadden L, Marriott BP. Caffeine use among active duty US Army soldiers. *Journal of the Academy of Nutrition and Dietetics*. 2012;112(6):902-12. e4.
15. Grosso G, Stepaniak U, Micek A, Stefler D, Bobak M, Pajak A. Coffee consumption and mortality in three Eastern European countries: results from the HAPIEE (Health, Alcohol and Psychosocial factors In Eastern Europe) study. *Public health nutrition*. 2017;20(1):82-91.
16. Mehri N, Messkoub M, Kunkel S. Trends, determinants and the implications of population aging in Iran. *Ageing International*. 2020:1-17.
17. Bijani A, Esmaili H, Ghadimi R, Babazadeh A, Rezaei R, Cumming RG, et al. Development and validation of a Semi-quantitative food frequency questionnaire among older people in north of Iran. *Caspian journal of internal medicine*. 2018;9(1):78.
18. Mirmiran P, Esfahani FH, Mehrabi Y, Hedayati M, Azizi F. Reliability and relative validity of an FFQ for nutrients in the Tehran lipid and glucose study. *Public health nutrition*. 2010;13(5):654-62.
19. Malmir H, Shayanfar M, Mohammad-Shirazi M, Tabibi H, Sharifi G, Esmailzadeh A. Tea and coffee consumption in relation to glioma: a case-control study. *European journal of nutrition*. 2019;58(1):103-11.
20. Grioni S, Agnoli C, Sieri S, Pala V, Ricceri F, Masala G, et al. Espresso coffee consumption and risk of coronary heart disease in a large Italian cohort. *PLoS One*. 2015;10(5):e0126550.

21. Liu J, Sui X, Lavie CJ, Hebert JR, Earnest CP, Zhang J, et al., editors. Association of coffee consumption with all-cause and cardiovascular disease mortality. *Mayo clinic proceedings*; 2013; 88(10):1066-74.
22. Carlström M, Larsson SC. Coffee consumption and reduced risk of developing type 2 diabetes: a systematic review with meta-analysis. *Nutrition reviews*. 2018;76(6):395-417.
23. Elhadad MA, Karavasiloglou N, Wulaningsih W, Tsilidis KK, Tzoulaki I, Patel CJ, et al. Metabolites, Nutrients, and Lifestyle Factors in Relation to Coffee Consumption: An Environment-Wide Association Study. *Nutrients*. 2020;12(5):1470.
24. Salazar-Martinez E, Willett WC, Ascherio A, Manson JE, Leitzmann MF, Stampfer MJ, et al. Coffee consumption and risk for type 2 diabetes mellitus. *Annals of internal medicine*. 2004;140(1):1-8.
25. Panagiotakos DB, Lionis C, Zeimbekis A, Makri K, Bountziouka V, Economou M, et al. Long-term, moderate coffee consumption is associated with lower prevalence of diabetes mellitus among elderly non-tea drinkers from the Mediterranean Islands (MEDIS Study). *The review of diabetic studies: RDS*. 2007;4(2):105.
26. van Dam RM, Hu FB, Willett WC. Coffee, caffeine, and health. *New England Journal of Medicine*. 2020;383(4):369-78.
27. Kondo Y, Goto A, Noma H, Iso H, Hayashi K, Noda M. Effects of coffee and tea consumption on glucose metabolism: a systematic review and network meta-analysis. *Nutrients*. 2019;11(1):48.
28. Mirmiran P, Carlström M, Bahadoran Z, Azizi F. Long-term effects of coffee and caffeine intake on the risk of pre-diabetes and type 2 diabetes: findings from a population with low coffee consumption. *Nutrition, Metabolism and Cardiovascular Diseases*. 2018;28(12):1261-6.
29. Yarmolinsky J, Mueller NT, Duncan BB, Bisi Molina MdC, Goulart AC, Schmidt MI. Coffee consumption, newly diagnosed diabetes, and other alterations in glucose homeostasis: a cross-sectional analysis of the longitudinal study of adult health (ELSA-Brasil). *PLoS One*. 2015;10(5):e0126469.
30. Keirns B, Elliott D, Sciarrillo C, Koemel N, Poindexter K, Emerson S. Effect of Black Coffee on Fasting Metabolic Markers and an Abbreviated Fat Tolerance Test. *Current Developments in Nutrition*. 2020;4(Supplement_2):639.
31. Borrer A, Zieff G, Battaglini C, Stoner L. The effects of postprandial exercise on glucose control in individuals with type 2 diabetes: a systematic review. *Sports Medicine*. 2018;48(6):1479-91.
32. Palatini P, Benetti E, Mos L, Garavelli G, Mazzer A, Cozzio S, et al. Association of coffee consumption and CYP1A2 polymorphism with risk of impaired fasting glucose in hypertensive patients. *European journal of epidemiology*. 2015;30(3):209-17.

33. Cornelis MC, Byrne EM, Esko T, Nalls MA, Ganna A, Paynter N, et al. Genome-wide meta-analysis identifies six novel loci associated with habitual coffee consumption. *Molecular psychiatry*. 2015;20(5):647.
34. van Dam R. Can 'omics' studies provide evidence for causal effects of coffee consumption on risk of type 2 diabetes? *Journal of internal medicine*. 2018;283(6):588-90.
35. Cano-Marquina A, Tarín J, Cano A. The impact of coffee on health. *Maturitas*. 2013;75(1):7-21.
36. Yen W-J, Wang B-S, Chang L-W, Duh P-D. Antioxidant properties of roasted coffee residues. *Journal of Agricultural and Food Chemistry*. 2005;53(7):2658-63.
37. Alhamhany NN, Alassady EH. Does green coffee has a positive effect on body mass index and lipid profile in a sample of obese people. *Journal of Pharmaceutical Sciences and Research*. 2018;10(3):627-30.
38. Ihim A, Nwanua M, Ogbodo E, Meludu S. Effect of Coffee Consumption on Blood Glucose and Lipid Profile Levels in Male Students at Nnamdi Azikiwe University, Nnewi Campus, Anambra State. *J Med Lab Sci*. 2019;29(2):10-20.
39. Cai L, Ma D, Zhang Y, Liu Z, Wang P. The effect of coffee consumption on serum lipids: a meta-analysis of randomized controlled trials. *European journal of clinical nutrition*. 2012;66(8):872-7.
40. Ranheim T, Halvorsen B. Coffee consumption and human health—beneficial or detrimental?—Mechanisms for effects of coffee consumption on different risk factors for cardiovascular disease and type 2 diabetes mellitus. *Molecular nutrition & food research*. 2005;49(3):274-84.
41. Kim YT, Chung HJ, Park BR, Kim YY, Lee JH, Kang DR, et al. Risk of Cardiovascular Disease and Chronic Kidney Disease According to 2017 Blood Pressure Categories in Diabetes Mellitus. *Hypertension*. 2020;76(3):766-75.
42. Noordzij M, Uiterwaal CS, Arends LR, Kok FJ, Grobbee DE, Geleijnse JM. Blood pressure response to chronic intake of coffee and caffeine: a meta-analysis of randomized controlled trials. *Journal of Hypertension*. 2005; 23: 921-928.
43. Mesas AE, Leon-Muñoz LM, Rodriguez-Artalejo F, Lopez-Garcia E. The effect of coffee on blood pressure and cardiovascular disease in hypertensive individuals: a systematic review and meta-analysis. *The American journal of clinical nutrition*. 2011;94(4):1113-26.
44. Zhao Y, Wang J, Ballevre O, Luo H, Zhang W. Antihypertensive effects and mechanisms of chlorogenic acids. *Hypertension Research*. 2012;35(4):370-4.
45. Chrysant SG. The impact of coffee consumption on blood pressure, cardiovascular disease and diabetes mellitus. *Expert review of cardiovascular therapy*. 2017;15(3):151-6.

46. de Abreu Pinheiro F, Elias LF, de Jesus Filho M, Modolo MU, Rocha JdCG, Lemos MF, et al. Arabica and Conilon coffee flowers: Bioactive compounds and antioxidant capacity under different processes. *Food chemistry*. 2020;336:127701.