The Impact of COVID-19 on Patients with Hematological Malignancies: The Mixed-Method Analysis of an Israeli National Survey

Ilana Levy (✉ ilana.e.levy@gmail.com)

Bnai Zion Medical Center <https://orcid.org/0000-0002-8923-1467>

Giora Sharf

Halil Haor non-profit organization

Shlomit Norman

AMEN

Tamar Tadmor

Bnei Zion Hospital

Research Article

Keywords: Hematological malignancies, COVID-19, Survey, Coping, Depression, Mixed-methods.

Posted Date: May 27th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-191174/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The COVID-19 pandemic required reevaluation of the therapeutic approach and added emotional stress for patients with hematological malignancies at high risk of contracting the virus. We aimed to evaluate how it affected such patients during the second lockdown in Israel.

Methods: This national survey included Hebrew-speaking patients with hematological malignancy. This included three tools with 28 items of socio-demographic and medical baseline characteristics, management of hematological disease, and evaluation of emotional coping during COVID-19 pandemic; the Hebrew version of the Patient Health Questionnaire 9; and 3 qualitative open-ended questions. Data was analyzed by mixed methods which combined both quantitative and qualitative thematic analyses.

Results: 408 patients responded to the survey. The management of their hematological disease included a decrease in the number of visits to the hematology clinic (37.0%), delay of some treatment schedules (9.1%), and prescription of replacement therapies permitting less visits to the clinic (2.2%). The frequency and intensity of "feeling afraid" regarding COVID-19 infection was increased (mean $\pm$ SD: 4 $\pm$ 1 to 5 $\pm$ 2 in a 1-7 Likert scale), and a high rate of depression was recorded, which appeared to be more evident in patients with chronic myeloid leukemia (CML) ($p < 0.001$).

Conclusion: The management of hematological malignancies during pandemics should always take into consideration patients' fears, as well as the development of depression related to isolation and loneliness, in addition to the high risk of severe disease. Patients with CML had a high rate of depression which obviously needs to be managed very carefully during and after the COVID-19 pandemic.

Introduction

The Coronavirus 2019 (COVID-19) pandemic first erupted in late 2019 and rapidly spread throughout the world.¹ Patients with comorbidities, including hemato-oncological diseases have an increased risk of developing serious illness and greater mortality.^{2,3} Therefore, in Israel since the beginning of the pandemic, two national population lockdowns have been completed, and patients with chronic illnesses including hemato-oncological malignancies were advised to minimize social activities in order to prevent infection.⁴ Optimal approach in terms of patients' visits to the hematology clinic, and decisions regarding changing treatment to mostly oral drugs was also under investigation worldwide.⁵ In addition, emotional aspects received more attention and assumed greater importance during the pandemic in both the general population⁶ and infected COVID-19 patients.⁷ One should take into consideration that even before the pandemic, depression seemed to be prevalent among patients with hematological malignancy, estimated at 54% in a Greek study, and associated with decreased quality of life (QOL) and cognitive functioning.⁸ Suicide attempts appear to be prevalent among cancer patients in general and patients with hematological malignancies in particular.⁹ Thus, it is not surprising that during the COVID-19 pandemic, a 36% rate of anxiety and 31% rate of depression was recorded among lymphoma outpatients.¹⁰ Coping styles associated with better QOL in cancer patients included emotional support and acceptance,¹¹ while

living alone and insufficient social support seem to be associated with treatment non-adherence in patients with hematological malignancies.¹² However in this respect, a recent study showed that patients with chronic conditions such as inflammatory arthritis seem to have a better capacity to successfully cope with stressful events during COVID-19 pandemic.¹³ Nevertheless, this point was not truly evaluated among patients with hematological malignancies.

Therefore, the aim of the present study was to evaluate the demographic and clinical factors associated with the emotional conditions of patients with hematological malignancies during the COVID-19 second lockdown in Israel.

Methods

Study design: This is a cross-sectional survey conducted among patients with hematological malignancy during the second COVID-19 lockdown in Israel. The study was reviewed and approved by the Bnai Zion Medical Center Institutional Review Board in accordance with the declaration of Helsinki (approval number BNZ-20-0148). Surveys were collected between October 1 and October 21, 2020. They were diffused in Google Forms platform and sent to patients with hematological malignancies via social networks, email, telephone, and non-profit patient organizations' chat groups (Halil Haor, AMEN). Four reminders to fill in the survey in Hebrew were sent in these platforms during the study period.

Participants: Hebrew-speaking patients with hematological malignancy were eligible to participate in the study. No informed consent signing was required since the survey was anonymous, and it was made clear to all involved before filling in the questionnaire that the data will be used for statistical analyses and research purposes.

Questionnaires: The questionnaires were composed of 13 socio-demographic and medical baseline questions, 11 questions on management of hematological disease during COVID-19 pandemic, and 4 questions referring to fears during COVID-19 pandemic in a 1-7 Likert scale. Patients then answered the Patient Health Questionnaire 9 (PHQ-9) which is a validated tool (translation to Hebrew has been validated as well), assessing the degree of depression as well as the impact of their symptoms on their lives.¹⁴ The total 0-27 PHQ-9 score categorizes depression severity as minimal or none (0-4 score), mild (5-9 score), moderate (10-14 score), moderately severe (15-19 score) or severe (20-27 score). Finally, three open-ended questions about difficulties to be handled, supportive resources, and suggestions to improve the current inconveniences/difficulties were filled by the patients. The analysis of these quantitative and qualitative data was based on mixed methods.

Statistical analysis: Descriptive statistics were calculated for all demographic and clinical data, including means with standard deviation (SD) and median with ranges (minimum-maximum) for all continuous data, and proportions for discrete and nominal data. For comparing normally distributed variables between more than two study groups, we used the ANOVA test. For comparing discrete and nominal variables, we used the independent Chi-square test and Fisher exact test. Bonferroni adjustment was used

for multiple comparisons. Multivariate linear regression analysis was performed after collinearity; interaction and confounding analysis of normally distributed Likert scale survey questions, in order to neutralize the effect of the different sociodemographic and medical covariates. Independent variables included age, sex, hematological disease, time from hematological disease diagnosis, familial status, country of birth, religion, education, occupation, and region of follow-up. The adjusted unstandardized regression coefficients (β) were given with confidence interval (CI), while a $p < 0.05$ was considered as statistically significant. Data analysis was performed using IBM SPSS version 22 statistics software.

Thematic analysis: Open-ended questions were analyzed using thematic analysis method, which is a variant of qualitative text analysis and examines the occurrence of themes or concepts in texts.¹⁵⁻¹⁷ The analysis was based on open coding procedure, creating categories and abstraction.¹⁸ The codes were derived by means of inductive analysis, which is used in cases where no previous studies dealt with the same phenomenon or when such studies are fragmentary.¹⁹

Results

Baseline characteristics: Among the total 1206 patients that were approached, 408 answered the survey (response rate 34%). Among them, 113 (27.8%) had chronic myeloid leukemia (CML), 92 (22.7%) multiple myeloma (MM), 75 (18.5%) chronic lymphocytic leukemia (CLL), 67 (16.5%) lymphoma, 41 (10.1%) had a myeloproliferative neoplasm (MPN), 14 (3.4%) had acute leukemia, and 4 (1.0%) hairy cell leukemia. Their baseline characteristics are described in Table 1.

Management of hematological malignancy during COVID-19 pandemic: Fifty-six (13.9%) responders did not visit the hematological clinic at any time during the COVID-19 pandemic, while 37 (9.1%) had their treatment delayed because of the pandemic. When asking patients about their visits to the hematological clinic since the beginning of the COVID-19 pandemic, 65 (15.9%) claimed that their visits had dropped significantly, 86 (21.1%) had their visits slightly decreased, 253 (62.0%) stated that their visits' frequency was unchanged, and 4 (1.0%) had visits slightly increased. Further information on management of hematological malignancies during the pandemic is presented in Table 2.

Fears during COVID-19 pandemic: We asked patients to evaluate their fears during the COVID-19 pandemic in a 1-7 Likert scale (1: almost never, 7: almost always).

Concerning the statement "I am afraid to contract COVID-19", the mean $\pm$ SD answer was 5 $\pm$ 2, and the median [minimum-maximum] was 6 [1-7]; and multivariate linear regression analysis showed that women tended to be more afraid than men to contract COVID-19 infection ($\beta=0.49$ [0.15; 0.83], $p=0.005$).

As for the statement "I am anxious around COVID-19 pandemic in my hematology clinic", the mean answer was 4 $\pm$ 2, the median was 4 [1-7], and multivariate linear regression analysis showed that women ($\beta=0.58$ [0.18; 0.97], $p=0.004$) and younger people ($\beta=-0.02$ [-0.04; -0.003], $p=0.020$) tended to be more anxious around COVID-19 infection in the hematology clinic; every year of age decreased the level of anxiety by 0.02 in the 1-7 Likert scale.

Concerning the statement "The hospital responds to my demands in terms of anxiety around COVID-19 pandemic" (1: not at all, 7: in a significant way), the mean answer was 5 ± 2 , the median was 6 [1-7], and multivariate linear regression analysis showed that divorced and widowed (as compared to married and single) patients ($\beta = -0.45$ [-0.77; -0.13], $p = 0.005$) and patients with longer disease duration ($\beta = 0.03$ [0.003; 0.05], $p = 0.028$) felt that the hospital responded better to their demands in terms of anxiety around the pandemic.

Finally, to the statement "If I contract COVID-19, my chances to have serious disease are (1: much lower, 5: much higher) than the general population", the mean answer was 4 ± 1 , the median was 5 [2-5], and multivariate linear regression analysis showed that women ($\beta = 0.22$ [0.09; 0.36], $p = 0.001$), older patients ($\beta = 0.009$ [0.003; 0.02], $p = 0.003$) and patients with a shorter disease duration ($\beta = -0.011$ [-0.02; 0.00], $p = 0.040$) felt that COVID-19 infection may be more serious if they contract it, compared to the general population.

Depression according to PHQ-9 questionnaire: The mean PHQ-9 score was 9 ± 6 , and its median was 8 [0-24]. Among the 406 responders to this part of the survey, 91 (22.4%) had no or minimal depression, 143 (35.2%) had mild depression, 100 (24.6%) had moderate depression, 49 (12.1%) had moderately severe depression, and 23 (5.7%) had severe depression (Fig. 1A). When comparing depression categorization between hematological malignancies, it should be noted that CML patients had a significantly more severe degree of depression, with 14 (12.4%) patients classified as having severe depression ($\text{Chi-2} = 22.85$, $p < 0.001$) (Fig. 1B).

Open-ended questions: Six categories of burdens/difficulties patients were dealing with during the COVID-19 pandemic were described. When sub-analyzing responses by patients' conditions, those with lymphoma and multiple myeloma mostly had difficulties dealing with isolation and loneliness, while patients with CML, MPN and CLL mostly described the physical manifestations of their disease and its treatment. Furthermore, ten categories of ways to cope with the different difficulties were described by the 357 responders that stated a difficulty in the preceding open-ended question. Family or friends were underlined by 75 (21.0%) of them and "being occupied" by 74 (20.7%). Finally, another ten categories of solutions that may have helped improving the above-mentioned difficulties were defined by patients. While 98 (27.5%) could not think of any potential solution, 59 (16.5%) thought of a life without COVID-19 and 57 (16.0%) believed the restoration of familial and social activities may well improve such difficulties. Quotations and keywords of the thematic analysis are presented in Table 3.

Discussion

The present study relates to important factors in the life of patients with hematological malignancy during COVID-19 second lockdown in Israel. While the patient management during this period was adapted to the circumstances, responders still disclosed frequent fears, accompanied by a relatively high level of depression. The difficulties they experienced during this period differed amongst the various

hematological malignancies, as well as the way they coped with such difficulties and the potential solutions they recommended as help.

How to manage hematological malignancies during COVID-19 pandemic? The current survey showed that a relatively large proportion of patients with hematological malignancies did not visit or delayed their visits to the hematology clinic, and some even had their treatments delayed or received alternative treatments instead. Such practices are in accordance with the guidelines for management of hematological malignancies during the pandemic.⁵ However, a high level of depression in the survey responders was recorded, and qualitative analysis showed that patients were more frequently worried about isolation or loneliness than about being infected with COVID-19, and that their main advice for improving their coping with such difficulties, was restoration of routine social life. In the light of our study results, it seems that management of patients with hematological malignancies should be individualized during pandemics, and that the importance of isolation to prevent contagion should be balanced with the potential psychological and physical consequences of loneliness.

Fears concerning the COVID-19 pandemic: According to our survey results, women were especially afraid to contract COVID-19, more anxious being around COVID-19 in the hematology clinic, and felt that their chances to contract severe disease if infected with COVID-19 were higher. In addition, although older patients justifiably felt that they were more at risk to contract severe disease, they generally felt safe in the hematology clinic around COVID-19 compared to younger patients. Similarly, a recent Italian study showed that outpatients with lymphoma who were at risk of anxiety and post-traumatic stress disorder during the COVID-19 pandemic were mostly women and younger patients.¹⁰ These at-risk populations should be targeted and their needs should be more precisely and individually investigated in order to improve patients' wellbeing in the hematology clinic during such stressful times.

Depression and difficulties during the pandemic: While a high rate of depression was defined for hematological malignancies, even higher than that recorded among lymphoma outpatients in Italy during the pandemic,¹⁰ it seems that patients with CML were most at risk of severe depression during this period. However, it is still unclear whether this predominance of CML is related specifically to the pandemic. Indeed, thematic analysis showed that these patients displayed apparent COVID-19-unrelated difficulties, essentially physical manifestations of the disease or its treatment. These findings support the fact that the high prevalence of severe depression in patients with CML may not be related to the COVID-19 pandemic or lockdown. An earlier study, published before the COVID-19 pandemic, reinforce our data since both patients and their physicians listed fatigue, muscle cramps, swelling, personal worries, and uncertainty about health condition in the future, as the most important QOL issues in CML.²⁰ These findings show that despite the pivotal implementation of treatment with tyrosine kinase inhibitors, CML patients still experience major physical and psychological issues related both to the disease itself and its treatment. Future studies should compare the prevalence of depression among patients with CML during and before/after the pandemic. In this respect it is important that better symptom management and psychological support should be introduced in hematology clinics to help manage these issues in the CML population.

Strengths and limitations: Our study is a national survey which applied to patients with all types of hematological malignancies, permitting a cost-effective method of collecting data from a large sample of patients. Another strength of our survey is its reliability that relates to its standardized diffusion to the participants. Finally, the mixed-methods analysis of both quantitative and qualitative data allowed us to approach the study question from different points of view, which facilitates integration of comprehensive data. However, some limitations should also be stressed. Indeed, the generalizability of our study results are limited because they are based only on Hebrew-speaking patients living in Israel, as well as their recruitment by social networks and non-profit organizations which as mentioned in the results, mainly include Jewish people and patients with basic academic education. Larger international studies should be performed to test the generalizability of our findings. Moreover, the reports of emotional aspects might have been influenced by some data which were not reported in the current study, including number of treatments, relapses/remission, performance status, etc. Future studies should include these clinical data in order to more reliably examine the factors influencing emotional stress among patients with hematological malignancies. Finally, the recruitment of survey responders via social media may have led to selection bias reflected by the low response rate since these media specifically cater to the younger demographic which may not coincide with the age categories of patients affected by hematological malignancies, and to patients with high socioeconomic level since patients with low per capita income as well as patients with low education have little access to social networks and research.

Conclusion: The management of hematological malignancies during pandemics can still be improved and should take into account all the high risks of severe disease among immunosuppressed patients, patients' fear relating to the COVID-19 pandemic and the high level of depression encountered in this population, which seems to be mostly related to individual isolation and loneliness. Unrelated to COVID-19, the care of patients with CML should be improved in terms of symptom management and psychological support.

Declarations

Funding: The research received no funding.

Conflicts of interest/Competing interests: The authors declare there were no conflict of interest.

Ethics approval: The study was approved by the Institutional Review Board of Bnai Zion Medical Center, in accordance with the Declaration of Helsinki.

Consent to participate: No informed consent signing was required since the survey was anonymous, and it was made clear to all involved before filing in the questionnaire that the data will be used for statistical analyses and research purposes.

Consent for publication: N/A.

Availability of data and material: All data are stored in a repository and can be made available as demand.

Authors' contributions: All authors contributed in constructing the study design, patient recruitment, article writing and reviewing.

References

1. Lake MA. What we know so far: COVID-19 current clinical knowledge and research. *Clin Med J R Coll Physicians London*. 2020;20(2):124-127. doi:10.7861/clinmed.2019-coron
2. Yigenoglu TN, Ata N, Altuntas F, et al. The outcome of COVID-19 in patients with hematological malignancy. *J Med Virol*. 2020;(July):1-6. doi:10.1002/jmv.26404
3. Piñana JL, Martino R, García-García I, et al. Risk factors and outcome of COVID-19 in patients with hematological malignancies. *Exp Hematol Oncol*. 2020;9(1):1-16. doi:10.1186/s40164-020-00177-z
4. Staff T. Ministers agree on full lockdown, more severe than Israel's first. The Times of Israel. <https://www.timesofisrael.com/ministers-said-to-agree-on-full-lockdown-more-severe-than-countrys-first/>. Published 2020.
5. Terpos E, Engelhardt M, Cook G, et al. Management of patients with multiple myeloma in the era of COVID-19 pandemic: a consensus paper from the European Myeloma Network (EMN). *Leukemia*. 2020;34(8):2000-2011. doi:10.1038/s41375-020-0876-z
6. Fu W, Wang C, Zou L, et al. Psychological health, sleep quality, and coping styles to stress facing the COVID-19 in Wuhan, China. *Transl Psychiatry*. 2020;10(225):1-9. doi:10.1038/s41398-020-00913-3
7. Samrah SM, Al-Mistarehi AH, Aleshawi AJ, et al. Depression and coping among covid-19-infected individuals after 10 days of mandatory in-hospital quarantine, irbid, jordan. *Psychol Res Behav Manag*. 2020;13:823-830. doi:10.2147/PRBM.S267459
8. Bellali T, Manomenidis G, Meramveliotaki E, Minasidou E, Galanis P. The impact of anxiety and depression in the quality of life and psychological well-being of Greek hematological cancer patients on chemotherapy. *Psychol Heal Med*. 2020;25(2):201-213. doi:10.1080/13548506.2019.1695864
9. Zaorsky NG, Zhang Y, Tuanquin L, Bluethmann SM, Park HS, Chinchilli VM. Suicide among cancer patients. *Nat Commun*. 2019;10(1):1-7. doi:10.1038/s41467-018-08170-1
10. Romito F, Dellino M, Loseto G, et al. Psychological Distress in Outpatients With Lymphoma During the COVID-19 Pandemic. *Front Oncol*. 2020;10(July):1-6. doi:10.3389/fonc.2020.01270
11. Ryan D, Nipp MD, Areej El-Jawahri MD, Joel N. Fishbein BA, et al. The Relationship Between Coping Strategies, Quality of Life, and Mood in Patients with Incurable Cancer. *Cancer*. 2016;122(13):2110-2116. doi:10.1002/cncr.30025.The
12. Bouwman L, Eeltink CM, Visser O, Janssen JJWM, Maaskant JM. Prevalence and associated factors of medication non-adherence in hematological-oncological patients in their home situation. *BMC Cancer*. 2017;17(739):1-8. doi:10.1186/s12885-017-3735-1

13. Ciaffi J, Brusi V, Lisi L, et al. Living with arthritis: a “training camp” for coping with stressful events? A survey on resilience of arthritis patients following the COVID-19 pandemic. *Clin Rheumatol*. 2020;39(11):3163-3170. doi:10.1007/s10067-020-05411-x
14. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: Validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16(9):606-613. doi:10.1046/j.1525-1497.2001.016009606.x
15. Boyatzis R. *Transforming Qualitative Information*.; 1998.
16. Roberts CW, Roberts BCW. *Text Analysis for the Social Sciences: Methods for Drawing Statistical Inferences from Texts and Transcripts*.; 1997.
17. Guest G, MacQueen KM, Namey EE. *Applied Thematic Analysis*.; 2012.
18. Corbin J, Strauss A. *Basics of Qualitative Research*.; 2015.
19. Elo S, Kyngäs H. The qualitative content analysis process. *J Adv Nurs*. 2008;62(1):107-115.
20. Efficace F, Breccia M, Saussele S, et al. Which health-related quality of life aspects are important to patients with chronic myeloid leukemia receiving targeted therapies and to health care professionals? *Ann Hematol*. 2012;91(9):1371-1381. doi:10.1007/s00277-012-1458-6

Tables

Table 1: Baseline characteristics

Baseline characteristics		Descriptive
Age		
Mean±Standard deviation		60±14
Median [minimum-maximum]		63 [18-90]
Years since diagnosis		
Mean±Standard deviation		7±6
Median [minimum-maximum]		5 [0-40]
Sex	Male – N (%)	192 (47.1%)
	Female – N (%)	216 (52.9%)
Familial status	Married / in couple – N (%)	322 (78.9%)
	Divorced – N (%)	36 (8.8%)
	Single – N (%)	32 (7.8%)
	Widow – N (%)	18 (4.4%)
Number of children		
Mean±Standard deviation		3±1
Median [minimum-maximum]		3 [0-7]
I live	With family – N (%)	356 (87.7%)
	Alone – N (%)	48 (11.8%)
	Nursing help – N (%)	2 (0.5%)
Religion	Jewish – N (%)	396 (98.3%)
	Arab (Muslim, Christian, Druze) – N (%)	7 (1.7%)
Country of birth	Israel – N (%)	289 (71.9%)
	Europe – N (%)	42 (10.4%)
	Former USSR – N (%)	27 (6.7%)
	America – N (%)	23 (5.7%)
	Africa – N (%)	12 (3.0%)

	Asia & Middle East – N (%)	9 (2.2%)
Education	Academic – N (%)	279 (68.9%)
	Professional – N (%)	28 (6.9%)
	Finished high-school – N (%)	66 (16.3%)
	Did not finish high-school – N (%)	32 (7.9%)
Occupation	Retiree – N (%)	140 (34.4%)
	Work normally or almost normally – N (%)	121 (29.7%)
	Do not work due to hematological condition – N (%)	69 (17.0%)
	Work up to 50% - N (%)	32 (7.9%)
	Do not work due to COVID-19 pandemic – N (%)	24 (5.9%)
	Do not work for other reasons – N (%)	21 (5.2%)
Hematological disease	Chronic myeloid leukemia – N (%)	113 (27.8%)
	Lymphoma – N (%)	67 (16.5%)
	Multiple Myeloma – N (%)	92 (22.7%)
	Myeloproliferative disease – N (%)	41 (10.1%)
	Chronic lymphocytic leukemia – N (%)	75 (18.5%)
	Acute leukemia – N (%)	14 (3.4%)
	Hairy Cell Leukemia – N (%)	4 (1.0%)
Region of Hematological Follow-up	Center of Israel – N (%)	247 (60.7%)
	Northern Israel – N (%)	106 (26.0%)
	Southern Israel – N (%)	17 (4.2%)
	Jerusalem region – N (%)	36 (8.8%)
	Abroad – N (%)	1 (0.2%)
Treatment for hematological disease	No treatment – N (%)	125 (30.6%)
	Oral – N (%)	218 (53.4%)

USSR: Union of Soviet Socialist Republics.

Table 2: Hematological management during COVID-19 pandemic

Hematological management		N (%)
During COVID-19 pandemic I come to hematological visits	Personal vehicle or walking	249 (61.8%)
	Public transport	13 (3.2%)
	Taxi	11 (2.7%)
	Accompanied by family	74 (18.4%)
	I do not visit hematological clinic	56 (13.9%)
I was infected with COVID-19		6 (1.5%)
If yes, I was hospitalized		3 (50.0%)
A family member was infected with COVID-19		48 (11.8%)
I was supposed to receive treatment but it was delayed because of COVID-19 pandemic		37 (9.1%)
If yes, delayed by	< 1 week	1 (2.9%)
	1 week - 1 month	13 (37.1%)
	> 1 month	21 (60.0%)
Treatment delay was	In my demand	8 (22.2%)
	Physician decision	13 (36.1%)
	Shared decision	13 (36.1%)
	Other	2 (5.6%)
I was supposed to have bone marrow transplantation but it was delayed because of COVID-19 pandemic		2 (0.5%)
I received an alternative treatment for my hematological disease due to COVID-19 pandemic		9 (2.2%)
If yes, it was	In my demand	1 (16.7%)
	Physician decision	2 (33.3%)
	Shared decision	3 (50.0%)

Table 3: Thematic analysis of open-ended questions

Themes	Frequency – N (%)	Quotations	Keywords (mentions)
1st question: Burdens/difficulties patients were dealing with during the COVID-19 pandemic			
Loneliness and isolation	113 (27.8%)	<p>"Impossibility to meet family and friends."</p> <p>"No human interaction."</p> <p>"Loneliness far from children, family and friends."</p> <p>"Impossibility to go out, visit family and children, or travel abroad."</p>	<p>"Friends" (34)</p> <p>"Family" (23)</p> <p>"Lockdown" (20)</p> <p>"Loneliness" (7)</p>
Physical manifestations related to the hematological condition or its treatment	109 (26.8%)	<p>"Significant fatigue related to the disease."</p> <p>"Side effects after each therapy."</p> <p>"Significant muscle pain and weakness related to treatment with a new drug."</p>	<p>"Fatigue" (37)</p> <p>"Pain" (27)</p> <p>"Weakness" (14)</p> <p>"Insomnia" (7)</p>
Fears related to the COVID-19 pandemic	49 (12.1%)	<p>"Uncertainty about what may happen if I am infected with corona."</p> <p>"Fear about the pandemic."</p> <p>"Fear to be infected with corona."</p>	<p>"Infected" (39)</p> <p>"Corona" (23)</p>
Uncertainty about the future	31 (7.6%)	<p>"Fear of the unknown."</p> <p>"Psychological difficulty and fear of what may be in the future."</p> <p>"Uncertainty about when it will end."</p>	<p>"Uncertainty" (14)</p> <p>"Anxiety" (8)</p>
Fears related to the hematological malignancy	30 (7.4%)	<p>"Uncertainty and non-understanding about the future of my hematological disease."</p> <p>"Uncertainty about the efficacy of the treatment I receive."</p> <p>"Fear that the response to the treatment will not be good enough."</p>	<p>"Treatment" (11)</p> <p>"Disease" (8)</p> <p>"Health" (5)</p> <p>"Uncertainty" (5)</p>

		"Uncertainty about my health."	
Related to work and economic issues	19 (4.6%)	"Uncertainty about the future in an economical and professional point of view." "I did not come back to work."	"Work" (9) "Economy" (5) "Busy" (3)
No difficulty or no answer	51 (12.6%)		
2nd question: Main ways to cope with the mentioned difficulties*			
Family or friends	75 (21.0%)	"The necessity to take care of my children." "Conversations with my family." "Family and friends' support."	"Family" (34) "Children" (15) "Friends" (10)
Being occupied	74 (20.7%)	"The fact that I continue working as usual." "Distractions such as work, reading, sport..." "I am cooking a lot and painting almost every day."	"Reading" (19) "Work" (14) "Occupy" (6)
No way to cope to difficulties	37 (10.7%)		
3rd question: Main solutions that may have helped improving the mentioned difficulties**			
Life without COVID-19	59 (16.5%)	"The possibility to go out and meet family normally." "That corona is temporary." "Vaccination for corona."	"Lockdown" (18) "Corona" (13) "Vaccinate" (9)
Restoration of familial and social activities	57 (16.0%)	"Going out and meeting friends and family." "Meeting friends, family, children." "The opening of swimming-pools and sport areas."	"Friends" (17) "Meeting" (8) "Family" (7)
No potential solution	98 (27.5%)		

* Other ways to cope with the mentioned difficulties: Patient's personality (optimism, faith), Substance use (drugs or cannabis), Technology (e.g. internet, telephone), Psychological or complementary medicine treatments, Physical activity, Rest, Protection against COVID-19, Hematologist's support

** Other potential solutions: Conversations and psychological help, Better consideration from the system and the physicians, Better health status and normal test results, Holidays and vacations, Emotional support, Substance use (drugs, cannabis), Work, Physiotherapy

Figures

Figure 1

A. Categorization of depression according to PHQ-9 responding B. Categorization of depression in different hematological malignancies CML: Chronic myeloid leukemia; CLL: chronic lymphocytic leukemia; HCL: Hairy cell leukemia; MM: Multiple myeloma; MPN: Myeloproliferative neoplasm.