Gastric and Urinary Bladder Pressures Correlate with Intra-abdominal Pressure in Patients with Morbid Obesity

Shadi Hamoud

Rambam Health Care Campus

Siham Abdelgani

Rambam Health Care Campus

Michal Mekel

Rambam Health Care Campus

Safa Kinaneh

Technion Israel Institute of Technology

Ahmad Mahajna (✉ a_mahajna@rmc.gov.il)

Rambam Health Care Campus

Research article

Keywords: intra-abdominal pressure, gastric pressure, urinary bladder pressure, laparoscopy

Posted Date: May 8th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-19159/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Intra-abdominal pressure (IAP) affects cardio-respiratory and hemodynamic parameters and can be measured directly or indirectly by measuring gastric or urinary bladder pressure. The aim of this study was to investigate the correlation between IAP, gastric pressure and urinary bladder pressure in patients with morbid obesity, at normal and elevated levels of IAP in two positions. As well, to examine the effects of increasing IAP and patient's position on hemodynamic and respiratory parameters.

Methods

Twelve patients undergoing bariatric surgery were included. IAP, gastric pressure, and urinary bladder pressure were measured while patients were in the supine position and after 45° anti-Trendelenburg tilt. Mean inspiratory pressure, peak inspiratory pressure, and tidal volume were recorded and assessed.

Results

In supine position; directly measured IAP was 9.1 ± 1.8 mmHg, compared to 10 ± 3.6 and 8.9 ± 2.9 mmHg in the stomach and bladder, respectively. Increasing IAP to 15 mmHg resulted in an increased gastric pressure of 17 ± 3.8 mmHg, and urinary bladder pressure of 14.8 ± 3.9 mmHg. Gastric and urinary bladder pressures strongly correlated with IAP ($R = 0.875$ and 0.847 , respectively). With 45° anti-Trendelenburg tilt; directly measured IAP was 9.4 ± 2.2 mmHg, and pressures of 10.8 ± 3.8 mmHg and 9.2 ± 3.8 mmHg were measured in the stomach and the bladder, respectively. Increasing IAP to 15 mmHg resulted in elevating gastric and bladder pressures to 16.6 ± 5.3 and 13.3 ± 4 mmHg, respectively. Gastric and urinary bladder pressures had good correlation with IAP ($R = 0.843$ and 0.819 , respectively). Changing patient position from supine to 45° anti-Trendelenburg position resulted in decreased mean and peak inspiratory pressures, and increased tidal volume.

Conclusions

Basal IAP is high in patients with morbid obesity. IAP shows positive correlation to gastric and urinary bladder pressures at both normal and elevated levels of IAP. Anti-Trendelenburg tilt of mechanically ventilated morbidly obese patients resulted in favorable effects on respiratory parameters.

Background

Increased intra-abdominal pressure is a common clinical condition that can lead to severe morbidity if continued without appropriate treatment. Abdominal compartment syndrome (ACS) is a serious

complication that can cause multi-organ failure (MOF) and death. ACS is defined as a sustained IAP > 20 mmHg (with or without an abdominal perfusion pressure < 60 mmHg) that is associated with new organ dysfunction or failure¹.

There has been a marked improvement in management, diagnosis and treatment of intra-abdominal hypertension (IAH) and ACS based on The consensus definitions of the World Society of the Abdominal Compartment Syndrome (WSACS) that were published in 2006^{1,2} followed by the clinical practice guidelines published in 2007³ and updated in 2013⁴. In addition, many surveys were conducted to determine the current state of awareness and knowledge of medical stuffs and the use of evidence-based medicine regarding IAH and ACS^{5,6}.

ACS can be primary or secondary depending on whether the etiology is related to the abdominal-pelvic cavity and requiring specific intervention of a target organ in case of primary while in secondary ACS there is no abdominal disease that requires specific surgical correction, but the high abdominal pressure is associated with an organic dysfunction that requires immediate surgical decompression. Common causes to ACS include: abdominal or pelvic trauma, intra-abdominal hemorrhage, retroperitoneal hematoma or edema. Other conditions as bowel obstruction, ascites, and necrotizing pancreatitis may lead to ACS as well⁷⁻¹⁰.

Sustained IAP results in elevated intrathoracic pressure which compromises pulmonary function, dynamics, increases afterload, decreased venous return and cardiac output. perfusion to the kidneys and intestinal mucosa is severely reduced¹¹⁻¹⁴.

Early diagnosis and treatment of IAH is essential to avoid MOF and death. There is still controversy regarding the ideal method for measuring IAP. A variety of studies suggest to measure frequently IAP in critically ill patients. Current practice assesses IAP indirectly through the measurement of intra vesicular pressure, however, only few studies have been performed to establish the actual relationship between IAP and urinary bladder pressure (UBP)¹⁵⁻¹⁷. In obese patients no studies have validated this technique.

The aims of this study were to assess the basal IAP and to investigate the correlation between IAP, gastric pressure (GP) and urinary bladder pressure (UBP), in patients with morbid obesity (body mass index- BMI > 40 kg/m²) who underwent bariatric surgery. The measurements were performed at normal and elevated levels of IAP in two positions: supine and 45 degrees anti-trendelenburg tilt. The effect of increasing IAP and change in the patient position on hemodynamic parameters (mean arterial pressure and heart rate) and respiratory parameters (mean inspiratory pressure, peak inspiratory pressure and tidal volume) were evaluated as well.

Methods

Patients

Male and female patients aged ≥ 18 years with morbid obesity were included in the study, while undergoing bariatric surgery in the Department of General Surgery at Rambam Health Care Campus (Rambam). Main exclusion criteria were any contraindications to laparoscopic surgery or urethral catheterization; known intra-abdominal adhesions or ventral hernia due to previous surgery; chronic obstructive pulmonary disease ($\text{PaCO}_2 > 50$ mmHg, $\text{FEV}_1 < 1$ liter); or marked left ventricular dysfunction (left ventricular ejection fraction $< 25\%$).

Experimental Protocol

The study was approved by Rambam's ethics committee (ethics number: 0019 – 12), and consent was obtained from all the participants. After general anesthesia was induced, a nasogastric tube and urinary Foley catheter were inserted. The study was conducted in two stages: (1) while patients were in the supine position, IAP was measured and adjusted for 5 minutes, and then (2) IAP was increased to 15 mmHg by insufflation of CO_2 for 5 minutes. Gastric and urinary bladder pressures were measured together with hemodynamic and respiratory parameters at each stage. Subsequently, the patient was up-tilted to 45° anti-Trendelenburg position and pressure and hemodynamic measurements were repeated.

Measurement of intra-abdominal pressure

The pressure within the abdominal cavity was measured directly using an automatic CO_2 insufflation measurement device (KARL STORZ Endoskope 264320 20, Tuttlingen, Germany).

Gastric pressure

Following insertion of the nasogastric tube, the stomach was drained and filled with 50 ml of normal saline. A rigid pressure tube was connected to the gastric tube using a male to male connector at one end, and to the monitor through a pressure transducer at the other end. The system was flushed with normal saline and pressure transducer zeroed at the mid axillary line.

Urinary bladder pressure

A Foley catheter was placed into the urinary bladder. The bladder was drained and filled with 50 ml of sterile normal saline. The drainage tube was clamped just beyond the aspiration port, and a 16-gauge needle connected to the rigid pressure tube was inserted into the port. The tube was connected to the monitor by a pressure transducer. The system was flushed with normal saline and the pressure transducer zeroed using the symphysis pubis as the zero reference point.

Statistical analysis

Sample size calculation was based on the <http://biomath.info/power/prt.htm> for paired samples. We assumed that the mean difference between the two measurements will be 2 units with 2.5 SD. In this case we needed to recruit less than 15 patients. Variables are presented as mean and standard deviation.

Statistical analysis included the χ^2 test for categorical variables and Student t-test for continuous variables. P-value < 0.05 was considered statistically significant.

Results

Of 13 patients included in the study, one was subsequently excluded due to a huge ventral hernia after previous open cholecystectomy. Table 1 summarizes participants' characteristics. There were eight females and four males with an average age of 35 ± 11 years, and average BMI of 43 ± 5.8 kg/m². The American Society of Anaesthesiologist (ASA) physical status score was I for eight patients and II for four patients.

Table 1
Characteristics of patients.

	n = 12	Range
Males	4	
Females	8	
Average age (years, range)	35 ± 12	18–55
Weight (kilograms)	120 ± 20	102–150
Height (meters)	1.66 ± 0.9	1.54–1.80
BMI (kg/m ²)	43 ± 5.8	37–55

Correlation between directly measured IAP, GP and UBP

When subjects were in the supine position; the directly measured IAP was 9.1 ± 1.8 mmHg, compared to pressures of 10 ± 3.6 mmHg and 8.9 ± 2.9 mmHg in the stomach and urinary bladder, respectively. Increasing IAP to 15 mmHg resulted in an increased GP of 17 ± 3.8 mmHg, and increased UBP to 14.8 ± 3.9 mmHg (Figs. 1 & 2). The correlation coefficients of GP and UBP to IAP were 0.875 and 0.847, respectively (Table 2). After elevating patients to 45 degrees anti-Trendelenburg position, direct IAP increased to 9.4 ± 2.2 , compared to 10.8 ± 3.8 and 9.2 ± 3.8 mmHg in the stomach and the urinary bladder, respectively. Increasing IAP to 15 mmHg resulted in an increased GP of 16.6 ± 5.3 mmHg, and of the UBP to 13.3 ± 4.9 mmHg, while GP and UBP strongly correlated with IAP ($R = 0.843$ and 0.819 , respectively) (Table 2).

Table 2
correlation between IAP, GP and UBP in supine and with 45 degree anti-trendelenburg elevation, in baseline and with wlvvated IAP to 15 mmHg.

	Position	Mean (SD) by group of IAP and significance of position's diff.				p-value (IAP: baseline Vs 15 mmHg)	Correlation with	
	Supine	9.1 (1.8)	p-	15.0 (0.00)	p-	NA	IAP (all groups)	
IAP	45 Degr.	9.4 (2.2)	value ^a	15.0 (0.00)	value ^b	NA	R	p-value
GI	Supine	10.0 (3.6)		17.00 (3.8)		< 0.001	0.875	< 0.001
	45 Degr.	10.8 (3.8)	0.713	16.6 (5.3)	0.531	< 0.001	0.843	< 0.001
Bladder	Supine	8.9 (2.9)		14.8 (3.9)		< 0.001	0.847	< 0.001
	45 Degr.	9.2 (3.8)	0.686	13.3 (4.9)	0.326	< 0.001	0.819	< 0.001
MAP	Supine	87.0 (11)		104.0 (11.0)		< 0.01	0.634	< 0.001
	45 Degr.	99.0 (20.0)	0.760	106.0 (14.0)	0.695	0.33	0.291	0.085
HR	Supine	75 (13)		75 (16)		0.71	0.184	0.284
	45 Degr.	78 (17)	0.718	81 (17)	0.181	0.25	0.152	0.377
MIP	Supine	11 (2)		13 (3)		< 0.001	0.407	0.014
	45 Degr.	10 (2)	0.788	12 (3)	0.021	< 0.001	0.447	0.006
PIP	Supine	27 (1)		31 (2)		< 0.005	0.777	< 0.001
	45 Degr.	24 (3)	0.792	28 (3)	0.004	< 0.001	0.748	< 0.001
Tv	Supine	0.63 (0.1)		0.52 (0.1)		0.01	-0.495	0.002
	45 Degr.	0.65 (0.1)	0.770	0.62 (0.1)	0.016	0.130	-0.366	0.028
a.- by comparing of correlations with IAP.								
b.- by Wilcoxon Signed Ranks Test								

Effect of increased IAP on mean arterial pressure (MAP) and heart rate (HR)

In the supine position, increasing the IAP to 15 mmHg resulted in an elevation of the MAP from 87 ± 11 mmHg to 104 ± 11 mmHg ($p < 0.01$). After 45° anti-Trendelenburg tilt, a similar increase of IAP resulted in an elevated MAP from 99 ± 20 mmHg to 106 ± 14 mmHg ($p = 0.33$, Table 2). In the supine position, increasing IAP to 15 mmHg caused no change in the HR (75 ± 13 bpm and 75 ± 16 bpm, $p = 0.71$). Also after 45° anti-Trendelenburg tilt, a similar increase of IAP resulted in non-significant elevation of the HR from 78 ± 17 bpm to 81 ± 17 bpm ($p = 0.25$, Table 2).

Effect of increased IAP on mean inspiratory pressure (MIP), peak inspiratory pressure (PIP) and Vt

In the supine position, increasing IAP to 15 mmHg resulted in MIP elevation from 11 ± 2 mmHg to 13 ± 3 mmHg ($p < 0.001$). After elevating the patients to the 45° anti-Trendelenburg position, a similar increase of the IAP resulted in MIP elevation from 10 ± 2.4 mmHg to 12 ± 3 mmHg ($p < 0.001$, Table 2). Similarly, in the supine position, increasing the IAP to 15 mmHg resulted in an increase of PIP from 27 ± 1 mmHg to 31 ± 2 mmHg ($p < 0.005$). After tilting patients to 45° anti-Trendelenburg position, a similar IAP increase resulted in PIP elevation from 24 ± 3 mmHg to 28 ± 3 mmHg ($p < 0.001$, Table 2).

In the supine position, increasing IAP to 15 mmHg resulted in a decreased Vt from 0.60 ± 0.1 lit to 0.52 ± 0.1 lit ($p = 0.01$). After a tilt to 45° anti-Trendelenburg position, a similar increase of the IAP decreased the Vt from 0.65 ± 1 lit to 0.62 ± 1 lit ($p = 0.13$, Table 2). While patients were with IAP of 15 mmHg, changing the patient position from supine to the 45° anti-Trendelenburg position resulted in decreased MIP (from 13 ± 3 to 12 ± 3 mmHg, $p = 0.021$) and PIP (from 31.2 ± 2.3 to 28.1 ± 3.3 mmHg, $p = 0.004$), and increased Vt (from 0.52 ± 0.1 to 0.62 ± 1 lit, $p = 0.016$).

Discussion

Morbid obesity has been proclaimed by the WHO Statement as the epidemic of the 21st Century^{18,19}, and is associated with considerable morbidity and mortality. With the widespread success of damage control laparotomy, ACS has become a virtual epidemic in trauma centers throughout the world^{20–22}. Increasing numbers of critically ill patients are obese; therefore the special consideration of IAP and ACS in this patient population has become significantly more relevant.

There is an exponential increase in studies on IAP and ACS in non-obese patients in the literature, but very few studies include the measurement of IAP in obese patients.

The diagnosis of IAH or ACS is heavily dependent on the reproducibility of the IAP measurement technique. Over time, literature has suggested many methods to assess IAP. The ideal tool is still

controversial¹⁵. Malbrain MLNG et al. showed in their study a poor correlation between IAP and abdominal perimeter²³. Other studies have shown that a clinical estimation of IAP examiner's feeling of the tenseness of the abdomen is not accurate, with a low sensitivity²⁴.

Consequently, the IAP needs to be measured with a more accurate and reliable tool. Different direct and indirect measurement methods have been reported^{25,26}. Traditionally, the urinary bladder pressure has been used as the method of choice for measuring the IAP. The technique was originally described by Kron et al.⁸. It is safe, minimally invasive, and has minimal side effects and complications.

Another useful site for measuring IAP is in the stomach through a nasogastric tube, which can be used when the patient has no Foley catheter in place, or when bladder pressure measurement is not possible due to absence of free contractility of the bladder wall²⁷. GP measurement is cheap, does not interfere with urine output, and has no risk of infection. Both methods are ideal for screening and monitoring of critically ill patients.

Several studies have addressed the validation of indirect versus direct measurements of the IAP during laparoscopy in non-obese patients. Yol et al. compared bladder pressure with direct insufflation pressure during laparoscopic cholecystectomy in 40 patients, and obtained a positive correlation between the two measurements ($R = 0.973$, $P < 0.0001$)²⁸. Likewise, Fusco et al. compared direct laparoscopic insufflation pressure with bladder pressure in patients undergoing laparoscopy, and demonstrated a good correlation in pressure values across the IAP range from 0 to 25 mmHg between the direct and indirect measurement methods²⁹. There have been no studies comparing the direct IAP with urinary bladder or gastric pressures in patients with morbid obesity.

In the current study, we found that the baseline IAP measured in the urinary bladder and stomach of patients with morbid obesity was higher than values reported in non-obese subjects, consistent with the literature. Sugerman et al. showed that the urinary bladder pressure was significantly higher in obese compared to non-obese patients (18 ± 0.7 vs. 7 ± 1.5 cmH₂O, respectively), and concluded that increased IAP, as reflected in urinary bladder pressure, contributes to health risks associated with severe obesity³⁰. Similarly, Lambert et al. concluded that IAP is elevated in patients with morbid obesity, and increased IAP is a function of central obesity associated with increased morbidity³¹.

The chronic elevation of IAP can explain why many severely obese patients, especially those with sleep apnea and hypoventilation, have found they must sleep in the sitting position, presumably to lower the effects of increased IAP on their thoracic cavity. In our study, we found that changing the patients' position from supine to 45° anti-Trendelenburg elevated position decreased the inspiratory pressures and increased Vt, findings that can be translated to a decrease in respiratory effort. Furthermore, increasing IAP from baseline to 15 mmHg resulted in a rise in the MAP with no change in the HR, both in the supine position and after a tilt of 45° anti-Trendelenburg. Various mechanisms, such as venous compression caused by elevated IAP (with compression of the abdominal vasculature and organs), and the

pharmacological action of the absorbed CO₂, have been proposed to explain these transient adverse hemodynamic effects^{32,33}. It has also been shown that increased IAP during carbon dioxide pneumoperitoneum is associated with increased mean arterial blood pressure and systemic vascular resistance^{34,35}.

In accordance with Lambert et al. who demonstrated elevated IAP in patients with morbid obesity compared to non-obese patients³¹, we also found that baseline IAP is high in patients with morbid obesity, and obtained a good correlation between the direct IAP measured by laparoscopic insufflation route on the one hand, and urinary bladder and gastric pressures measured in morbidly obese patients on the other hand, both at normal and elevated levels of IAP. In addition, we found that changing the patients' position from the supine position to 45° anti-Trendelenburg elevation position caused decreased mean and peak inspiratory pressures, while increasing Vt. Therefore, patients with increased IAP are expected to benefit from the anti-Trendelenburg position as it improves respiratory parameters without hemodynamic compromise.

Conclusions

Our study demonstrated that in obese patients there is good correlation between directly measured IAP and indirectly measured pressures through the stomach or the urinary bladder. Changing the position of the patient with increased IAP to the anti-Trendelenburg position is helpful in improving respiratory parameters without interference to the patient's hemodynamic status.

Abbreviations

ACS
Abdominal compartment syndrome
ASA
American Society of Anaesthesiologist
GP
gastric pressure
HR
heart rate
IAH
intra-abdominal hypertension
IAP
Intra-abdominal pressure
MAP
mean arterial pressure
MIP
mean inspiratory pressure

PIP
peak inspiratory pressure
Vt
tidal volume
WSACS
World Society of the Abdominal Compartment Syndrome

Declarations

Ethics approval and consent to participate: The study was approved by Rambam's ethics committee (ethics number: 0019-12), and written consent was obtained from all the participants.

Consent of publication- Not applicable.

Availability of data and materials. All data generated or analysed during this study are included in this published article.

Competing Interests. The authors declares that they have no competing interests.

Funding. This study had no financial support.

Authors' contributions. AH and MM designed the study and performed the intra-operative measurements. SH, SA and AH participated in manuscript writing and preparing for publication. SH, SK and AH performed data collection and analysis.

all authors have read and approved the manuscript

Acknowledgements. Authors thank Mrs. Ronit Leiba for her valuable assistance in the statistical analysis.

Trial Registration. The study was retrospectively registered in the NIH registry. Registration number is pending.

References

1. Malbrain ML, Cheatham ML, Kirkpatrick A, Sugrue M, Parr M, De Waele J, et al. Results from the International Conference of Experts on Intra-abdominal Hypertension and Abdominal Compartment Syndrome. I. Definitions. Intensive Care Med. 2006;32(11):1722–1732.
2. Malbrain ML, Cheatham ML. Definitions and pathophysiological implications of intra-abdominal hypertension and abdominal compartment syndrome. Am Surg. 2011;77(Suppl 1):6–11.
3. Cheatham ML, Malbrain ML, Kirkpatrick A, Sugrue M, Parr M, De Waele J, Balogh Z, Leppaniemi A, Olvera C, Ivatury R, D'Amours S, Wendon J, Hillman K, Wilmer A. (2007) Results from the international

- conference of experts on intra-abdominal hypertension and abdominal compartment syndrome. II. Recommendations. *Intensive Care Med* 33:951–962.
4. Kirkpatrick AW, Roberts DJ, De Waele J, Jaeschke R, Malbrain ML, De Keulenaer B, Duchesne J, Bjorck M, Leppaniemi A, Ejike JC, et al. Intra-abdominal hypertension and the abdominal compartment syndrome: updated consensus definitions and clinical practice guidelines from the World Society of the Abdominal Compartment Syndrome. *Intensive Care Med*. 2013;39(7):1190–206. doi:10.1007/s00134-013-2906-z.
 5. 10.5114/ait.2019.87648
Wise R,2, RR,3, BA,5, RD, De Waele J7, Kirkpatrick A8, De Keulenaer B9,10, Malbrain M11,12, The Abdominal Compartment Society FTW. Awareness and knowledge of intra-abdominal hypertension and abdominal compartment syndrome: results of a repeat, international, cross-sectional survey. *Anaesthesiol Intensive Ther*. 2019;51(3):186–199. doi: 10.5114/ait.2019.87648.
 6. De Waele JJ, Cheatham ML, Malbrain ML, Kirkpatrick AW, Sugrue M, Balogh Z, Ivatury R, De Keulenaer B, Kimball EJ. (2009) Recommendations for research from the international conference of experts on intra-abdominal hypertension and abdominal compartment syndrome. *Acta Clin Belg* 64:203–209.
 7. Saggi BH, Sugerman HJ, Ivatury RR, Bloomfield GL. Abdominal compartment syndrome. *J Trauma*. 1998;45(3):597–609.
 8. Kron IL, Harman PK, Nolan SP. The Measurement of Intra-Abdominal Pressure as a Criterion for Abdominal Re-Exploration. *Ann Surg*. 1984;199(1):28–30.
 9. Burch JM, Moore EE, Moore FA, Franciose R. The abdominal compartment syndrome. *Surg Clin North Am*. 1996;76(4):833–42.
 10. Fietsam R Jr, Villalba M, Glover JL, Clark K. Intra-abdominal compartment syndrome as a complication of ruptured abdominal aortic aneurysm repair. *Am Surg*. 1989;55(6):396–402.
 11. Schein M, Wittmann DH, Aprahamian CC, Condon RE. The abdominal compartment syndrome: the physiological and clinical consequences of elevated intra-abdominal pressure. *J Am Coll Surg*. 1995;180(6):745–53.
 12. Rasmussen IB, Berggren U, Arvidsson D, Ljungdahl M, Haglund U. Effects of pneumoperitoneum on splanchnic hemodynamics: an experimental study in pigs. *Eur J Surg*. 1995;161(11):819–26.
 13. Simon RJ, Friedlander MH, Ivatury RR, DiRaimo R, Machiedo GW. Hemorrhage lowers the threshold for intra-abdominal hypertension-induced pulmonary dysfunction. *J Trauma*. 1997;42(3):398–403. discussion 404 – 395.
 14. Sugrue M, Jones F, Deane SA, Bishop G, Bauman A, Hillman K, et al. Intra-abdominal hypertension is an independent cause of postoperative renal impairment. *Arch Surg*. 1999;134(10):1082–5.
 15. MLNG M. Intra-abdominal Pressure in the Intensive Care Unit: Clinical Tool or Toy?. *Yearbook of Intensive Care and Emergency Medicine* 2001. 2001;2001.
 16. MLNG M. Abdominal Perfusion Pressure as a Prognostic Marker in Intra-abdominal Hypertension. *Yearbook of Intensive Care and Emergency Medicine* 2002. 2002;2002:792–814.

17. Al-Abassi AA, Al Saadi AS, Ahmed F. Is intra-bladder pressure measurement a reliable indicator for raised intra-abdominal pressure? A prospective comparative study. *BMC Anesthesiol.* 2018;19(1):69. 18(.
18. Organization WH. Obesity: preventing and managing the global epidemic Report of a WHO Consultation (WHO Technical Report Series 894). 2000:252.
19. Deitel M. The International Obesity Task Force and "Globesity". *Obes Surg.* 2002;12(5):613–4.
20. Morris JA Jr, Eddy VA, Blinman TA, Rutherford EJ, Sharp KW. The staged celiotomy for trauma. Issues in unpacking and reconstruction. *Ann Surg.* 1993;217(5):576–84. discussion 584 – 576.
21. Raeburn CD, Moore EE, Biffl WL, Johnson JL, Meldrum DR, Offner PJ, et al. The abdominal compartment syndrome is a morbid complication of postinjury damage control surgery. *Am J Surg.* 2001;182(6):542–6.
22. Hong JJ, Cohn SM, Perez JM, Dolich MO, Brown M, McKenney MG, et al. Prospective study of the incidence and outcome of intra-abdominal hypertension and the abdominal compartment syndrome. *Brit J Surg.* 2002;89(5):591–6.
23. Malbrain MLNG, De laet I, Van Regenmortel N, Schoonheydt K, Dits H, et al. Can the abdominal perimeter be used as an accurate estimation of intra-abdominal pressure? *Crit Care Med.* 2009;37(1):316–9.
24. Sugrue M, Bauman A, Jones F, Bishop G, Flabouris A, Parr M, et al. Clinical examination is an inaccurate predictor of intraabdominal pressure. *World J Surg.* 2002;26(12):1428–31.
25. Malbrain ML. Different techniques to measure intra-abdominal pressure (IAP): time for a critical re-appraisal. *Intensive Care Med.* 2004;30(3):357–71.
26. Wise RD, Rodseth RN, Correa-Martin L, Sanchez Margallo F, Becker P, Castellanos G, et al. Correlation between different methods of intraabdominal pressure monitoring in varying intraabdominal hypertension models. *S Afr J Crit Care.* 2017;33(1):15–8.
27. Collee GG, Lomax DM, Ferguson C, Hanson GC. Bedside measurement of intra-abdominal pressure (IAP) via an indwelling naso-gastric tube: clinical validation of the technique. *Intensive Care Med.* 1993;19(8):478–80.
28. Yol S, Kartal A, Tavli S, Tatkan Y. Is urinary bladder pressure a sensitive indicator of intra-abdominal pressure? *Endoscopy.* 1998;30(9):778–80.
29. Fusco MA, Martin RS, Chang MC. Estimation of intra-abdominal pressure by bladder pressure measurement: validity and methodology. *J Trauma.* 2001;50(2):297–302.
30. Sugerman H, Windsor A, Bessos M, Wolfe L. Intra-abdominal pressure, sagittal abdominal diameter and obesity comorbidity. *J Intern Med.* 1997;241(1):71–9.
31. Lambert DM, Marceau S, Forse RA. Intra-abdominal pressure in the morbidly obese. *Obes Surg.* 2005;15(9):1225–32.
32. Ho HS, Saunders CJ, Gunther RA, Wolfe BM. Effector of hemodynamics during laparoscopy: CO₂ absorption or intra-abdominal pressure? *J Surg Res.* 1995;59(4):497–503.

33. Horvath KD, Whelan RL, Lier B, Viscomi S, Barry L, Buck K, et al. The effects of elevated intraabdominal pressure, hypercarbia, and positioning on the hemodynamic responses to laparoscopic colectomy in pigs. *Surg Endosc.* 1998;12(2):107–14.
34. Koivusalo AM, Lindgren L. Effects of carbon dioxide pneumoperitoneum for laparoscopic cholecystectomy. *Acta Anaesthesiol Scand.* 2000;44(7):834–41.
35. Lindberg F, Bergqvist D, Björck M, Rasmussen I. Renal hemodynamics during carbon dioxide pneumoperitoneum: an experimental study in pigs. *Surg Endosc.* 2003;17(3):480–4.

Figures

Figure 1

Figure 1

Correlation between directly measured Intra-abdominal Pressure (mmHg) and Gastric Pressure (mmHg) in two positions; Supine and 45° anti-Trendelenburg position. Values are presented as mean ± SD.

Figure 2

Figure 2

Correlation between directly measured Intra-abdominal Pressure (mmHg) and Urinary Bladder Pressure (mmHg) in two positions; Supine and 45° anti-Trendelenburg position. Values are presented as mean $\pm$ SD.