

Statistical Distribution of Novel coronavirus in Iran

Elham Gholami

Tehran University of Medical Sciences

Kamyar Mansori

Zanjan University of Medical Sciences

Mojtaba Soltani-Kermanshahi (✉ msoltani@semums.ac.ir)

Semnan University of Medical Sciences and Health Services <https://orcid.org/0000-0002-6607-6249>

Research article

Keywords: COVID-19, Infection, Distribution, Weibull

Posted Date: April 1st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-19181/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background The Corona Virus Disease-2019 (COVID-19) pandemic by 2019-nCoV is spreading worldwide, and by March 27, 2020, 199 countries, including Iran, have been affected. Prevention and control of infection is the most important priority for the public health. The behavior prediction of COVID-19 is an important problem.

Methods In this research, we compared the different distributions of COVID-19 cases based on the daily reported data of Iran. We used 36 initial data on new cases and deaths with confirmed 2019-nCoV infection of Iran based on official reports from governmental institutes. We used the three types of continuous distribution known as Normal, Lognormal and Weibull.

Results Our study showed that the weibull distribution was the best fit with the data. But the Parameters of distribution were different between new cases and daily deaths data.

Conclusion According to the mean and median of the best fitted distribution, we can expect to pass the peak of the disease. The death rate is decreasing. We can see the similar behaviors of covid-19 in both Iran and China in the long run.

Background

Emerging infectious diseases, such as severe acute respiratory syndrome (SARS) and Zika virus disease, give a significant warning to public health [1-3]. Despite significant medical researches however, how, when and where new diseases appear are yet an origin of considerable uncertainty [4]. On 31 December 2019, the Wuhan Municipal Health Commission in Wuhan City, Hubei province, China reported a cluster of pneumonia cases with unspecified etiology that had a history of exposure to Wuhan's Huanan Seafood Wholesale Market (a wholesale fish and live animal market selling different animal species). On 9 January 2020, China CDC reported that a novel coronavirus (2019-nCoV) had been detected as the causative agent and the genome sequence was made publicly available. Sequence analysis displayed that the newly-identified virus is belonging to the SARS-CoV clade [5]. In an effort to prevent the prevalence, travel limitations were imposed on Wuhan from 23 January, and have since expanded to 12 other cities, and large social gatherings annulled [6, 7]. The Corona Virus Disease-2019 (COVID-19) pandemic by 2019-nCoV is spreading worldwide, and by March 27, 2020, 199 countries, including Iran, have been affected. According to worldwide statistics, the mortality rate is 3.4%. Early symptoms of COVID-19 involves pneumonia, fever, myalgia and fatigue. To date, no successful vaccine or antiviral agents has been clinically approved for COVID-19. Therefore, prevention and control of infection is the most important priority for the public health [8,9].

During the [2019–20 coronavirus pandemic](#), [Iran](#) reported the first authenticated cases of [SARS-CoV-2](#) infections on 19 February 2020 in [Qom](#). [10] As of 27 March 2020, pursuant to Iranian health authorities, there had been 2378 [COVID-19](#) deaths in Iran with more than 32,000 confirmed infections. Also, in the same date, Iran had the fourth highest in term of the number of COVID-19 deaths after [Mainland China](#) ,

Italy and Spain and the first rank in Western Asia. Accordingly, the mean age and sex ratio (man/female) of patients were 59 years and 1.4 respectively; in Iran. Among the deaths related to SARS-CoV-2, 59% were males and 41% females. In early March 2020, non-Iranian-government sources stated assessment of the numbers of SARS-CoV-2 infections that were much higher than official values [11-15].

There are considerable uncertainties in assessing the risk of this disease, due to lack of detailed epidemiological analyses. Extensive researches into the 2019-nCoV are needed to fully elucidate its pathway and pathogenic mechanisms, and to identify potential therapeutic targets, which can be effective in developing the common preventive and therapeutic measures. The behavior prediction of COVID-19 is an important problem. Therefore, in this research, we compared the different distributions of COVID-19 cases based on the daily reported data.

Methods

We used information on new cases and deaths with confirmed 2019-nCoV infection of Iran based on official reports from governmental institutes [13, 9]. We collected the data either directly from governmental websites or from news sites that directly quoted governmental statements. The data were collected in real time, and thus may be updated as more details on cases become publicly available. The arranged data are available as the Online Supplementary Material (file S1). The latest update to the dataset was on 27 March 2020 for cases reported through 26 March.

We performed a bootstrap method, based on case resampling, to compute the 95% confidence intervals (CI). We used the three types of continuous distribution known as Normal, Lognormal and Weibull. Akaike Information Criterion (AIC) was used to identify the best fit model. We also present the median and mean of the best fitted model to know the peak.

Normal Distribution

A normal (or Gaussian or Gauss or Laplace–Gauss) distribution is a type of continuous probability distribution for a real-valued random variable. The general form of its probability density function is

$$f(x) = \frac{1}{\sqrt{2\pi} \sigma} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

The parameter μ is the mean or expectation of the distribution (and also its median and mode); and σ is its standard deviation. The variance of the distribution is σ^2 . A random variable with a Gaussian distribution is said to be normally distributed and is called a normal deviate. [16]

Lognormal Distribution

A positive random variable X is log-normally distributed if the logarithm of X is normally distributed [17],

$$\text{Ln}(X) \sim N(\mu, \sigma^2)$$

Weibull Distribution

The probability density function of a Weibull random variable is:

$$f(x; \lambda, k) = \begin{cases} \left(\frac{k}{\lambda}\right) \left(\frac{x}{\lambda}\right)^{k-1} e^{-\left(\frac{x}{\lambda}\right)^k} & x \geq 0 \\ 0 & x < 0 \end{cases}$$

Where $k > 0$ is the shape parameter and $\lambda > 0$ is the scale parameter of the distribution [18].

All statistical analyses were performed using package fitdistrplus in R (version 3.6.3) [19].

Results

We used 36 daily data from 21 Feb 2020 of Iran. The smallest number of new cases was 11 (22Feb2020) and the highest number was 2926 (27 March 2020). On the other hand, the smallest number of daily deaths was 2 (21, 22 and 23 Feb 2020) and the highest number was 157 (26 March 2020). The Figure1 shows the histogram of daily new cases. The graph also shows three fitted distributions on the data.

Table 1 presents the goodness of fit of three distributions. Accordingly the best fit belongs to Weibull distribution. With assumption of Weibull distribution for daily new cases we can say the mean, median and mode of the data are 897.60, 624.61 and 4.59; respectively.

Table1. Goodness of fit of three distributions (Normal, Lognormal and Weibull) on Daily new Covid-19 infections in Iran

Distribution	Parameters	Estimation	AIC
Normal	Mean	897.97	576.61
	Sd	688.10	
Lognormal	Meanlog	6.13	580.88
	Sdlog	1.58	
Weibull	Shape	1.00	565.61
	Scale	899.48	

AIC: Akaike Information Criterion

The figure2 shows the histogram of daily deaths. The graph also shows three fitted distributions on the data.

Table 2 presents the goodness of fit of three distributions. Accordingly the best fit belongs to Weibull distribution. With assumption of Weibull distribution for daily deaths, we can say the mean, median and mode of data are 66.44, 41.11 and zero; respectively.

Table2. Goodness of fit of three distributions (Normal, Lognormal and Weibull) on Daily new Covid-19 deaths in Iran

Distribution	Parameters	Estimation	AIC
Normal	Mean	66.00	398.38
	Sd	57.89	
Lognormal	Meanlog	3.44	381.86
	Sdlog	1.48	
Weibull	Shape	0.88	376.76
	Scale	62.35	

AIC: Akaike Information Criterion

Discussion

On the 11 March 2020, WHO Director said “In the days and weeks ahead, we expect to see the number of cases, the number of deaths, and the number of affected countries climb even higher. We have therefore made the assessment that COVID-19 can be characterized as a pandemic” [20]. In the present study, we demonstrated the distribution of Covid-19 data in Iran with various types of continuous distributions. One of the distributions we used was the Weibull distribution. The Weibull distribution is related to a number of other probability distributions; in particular, it [interpolates](#) between the [exponential distribution](#) ($k = 1$) and the [Rayleigh distribution](#) ($k = 2$ and). Its [complementary cumulative distribution function](#) is a [stretched exponential function](#) [21]. We used real daily data (positive infections and deaths) reported by Ministry of Health [13, 9]. Our study showed that the weibull distribution was the best fit with the data. But the Parameters of distribution were different between new cases and daily deaths data. The shape parameter on the daily deaths data was less than one; then the rate is decreasing (fortunately). On the other side the shape parameter on the new cases data was approximately one. Thus as we say, the weibull distribution can change to exponential distribution. We could not find any similar works to compare our results. If we use China covid- 19 data (63 daily data), we can see the same distribution. On the other side, If we only use 36 initial China covid-19 data (similar to our data), the Lognormal distribution has the best fit. Then we can see the similar behaviors of covid-19 in both countries in the long run.

According to the mean and median of the best fitted distribution, we can expect to pass the peak of the disease. But, we are on the vacation of Nowruz in Iran, now. Contrary to government warnings, many people are on the travelling. On the other side the World Health Organization (WHO) reported an incubation period for COVID-19 between 2 and 10 days [22]. Then we can expect other peaks if we do not follow the health instructions. There are considerable uncertainties in assessing the risk of this event, due to lack of detailed epidemiological analyses

Conclusion

Our study showed that the weibull distribution was the best fit with the data. The death rate is decreasing. According to the results, we can expect to pass the peak of the disease. We can see the similar behaviors of covid-19 in both Iran and China in the long run.

Abbreviations

AIC: Akaike Information Criterion

SARS: Severe Acute Respiratory Syndrome

COVID-19: Corona Virus Disease-2019

CI: Confidence Intervals

WHO: World Health Organization

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

All authors have approved the manuscript and agree with its submission to the *Journal of BMC Infectious Diseases*.

Availability of data and materials

The data that support the findings of this study are available from:
<https://www.worldometers.info/coronavirus/>

Funding

Not applicable

Competing Interests

Not applicable

Authors' contributions

Study concept and design: Mrs. Golami, Dr. Soltani and Dr. Mansori. Acquisition of data: Mrs. Gholami. Analysis and interpretation of data: Dr. Soltani. Drafting of the manuscript: Dr. Mansori. Critical revision

of the manuscript for important intellectual content: Mrs. Gholami, Dr. Soltani and Dr. Mansori. Statistical analysis: Dr. Soltani.

Acknowledgements

The authors are thankful to Worldometers webmaster for providing data for this project.

References

1. Drosten, C. et al. Identification of a novel coronavirus in patients with severe acute respiratory syndrome. *N. Engl. J. Med.* 348, 1967–1976 (2003).
2. Wolfe, N. D., Dunavan, C. P. & Diamond, J. Origins of major human infectious diseases. *Nature* 447, 279–283 (2007).
3. Ventura, C. V., Maia, M., Bravo-Filho, V., Gis, A. L. & Belfort, R. Jr. Zika virus in Brazil and macular atrophy in a child with microcephaly. *Lancet* 387, 228 (2016).
4. Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, Hu Y, Tao ZW, Tian JH, Pei YY, Yuan ML. A new coronavirus associated with human respiratory disease in China. *Nature*. 2020 Feb 3:1-5.
5. European Centre for Disease Prevention and Control , Risk assessment: Outbreak of acute respiratory syndrome associated with a novel coronavirus, Wuhan, China; first update, [March 27, 2020]. Available from: <https://www.ecdc.europa.eu/en/publications-data/risk-assessment-outbreak-acute-respiratory-syndrome-associated-novel-coronavirus>
6. New York Times (2020) Toll From Outbreak Climbs in China as Infections Reach Europe and Australia. Available at: <https://www.nytimes.com/2020/01/24/world/asia/china-coronavirus.html> (Accessed: 27 March 2020).
7. Read JM, Bridgen JR, Cummings DA, Ho A, Jewell CP. Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. *medRxiv*. 2020 Jan 1.
8. Farnoosh G, Alishiri G, Zijoud SR, Dorostkar R, Farahani AJ. Understanding the 2019-novel Coronavirus (2019-nCoV) and Coronavirus Disease (COVID-19) Based on Available Evidence-A Narrative Review. *Journal of Military Medicine*. 2020 Jan;22(1):1-1.
9. Worldometer, COVID-19 Coronavirus Pandemic (Accessed: 27 March 2020). Available at: <https://www.worldometers.info/coronavirus/>
10. "Iran Reports Its First 2 Cases of the New Coronavirus". *New York Times*. Archived from the original on 19 February 2020. Retrieved 19 February 2020.
11. Wikipedia, 2020 coronavirus pandemic in Iran (Accessed: 27 March 2020). Available at: https://en.wikipedia.org/wiki/2020_coronavirus_pandemic_in_Iran
12. Aftabnews agency, Corona patients in Iran (Accessed: 27 March 2020). Available at: <https://aftabnews.ir/fa/news/641753> [in Persian]
13. Ministry of Health, Some characteristic of Covid19 in Iran (Accessed: 27 March 2020). Available at: <http://behdasht.gov.ir/uploads/factsheet%204.%2026.12%20-%20fa.pdf> [in Persian]

14. Aftabnews agency, We have not yet reached the peak of Covid19 (Accessed: 27 March 2020). Available at: <https://aftabnews.ir/fa/news/642341> [in Persian]
15. Ministry of Health, Daily reports of Covid19 in Iran (Accessed: 27 March 2020). Available at: <http://behdasht.gov.ir/index.jsp?siteid=1&fkeyid=&siteid=1&pageid=54782&newsview=201158> [in Persian]
16. Weisstein, Eric W. "Normal Distribution Function". MathWorld.
17. Johnson, Norman L.; Kotz, Samuel; Balakrishnan, N. (1994), "14: Lognormal Distributions", Continuous univariate distributions. Vol. 1, Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics (2nd ed.), New York: John Wiley & Sons, ISBN978-0-471-58495-7, MR1299979
18. Papoulis, Athanasios Papoulis; Pillai, S. Unnikrishna (2002). Probability, Random Variables, and Stochastic Processes (4th ed.). Boston: McGraw-Hill. ISBN0-07-366011-6.
19. Delignette-Muller ML, Dutang C. fitdistrplus: An R package for fitting distributions. Journal of statistical software. 2015 Mar 20;64(4):1-34.
20. World Health Organization, WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020 (Accessed: 27 March 2020). Available at: <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19--11-march-2020>
21. "Rayleigh Distribution – MATLAB & Simulink – MathWorks Australia". www.mathworks.com.au.
22. Novel Coronavirus (2019-nCoV) Situation Report-7 - World Health Organization (WHO), January 27, 2020.

Figures




Figure 1

Histogram and three fitted distributions (Normal, Lognormal and Weibull) on the daily infections with Covid-19 in Iran.




Figure 2

Histogram and three fitted distributions (Normal, Lognormal and Weibull) on the daily deaths with Covid-19 in Iran

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [corona.iran.dat](#)