

# Knowledge and Attitude on Prevention of Novel COVID-19 Among Community Health Workers in Nepal.

Amul Shrestha (✉ [shresthamul98@gmail.com](mailto:shresthamul98@gmail.com))

IFHR (Innovative foundation for health and research) <https://orcid.org/0000-0001-7892-4279>

**Tek Bahadur Thapa**

IFHR (Innovative Foundation for Health and Research)

**Mahendra Giri**

IFHR, Nepal

**Sanjiv Kumar**

IFHR, Nepal

**Sakil Dhobi**

IFHR, Nepal

**Harikali Thapa**

IFHR, Nepal

**Pawan Prakash Dhami**

IFHR, Nepal

**Arun Shahi**

IFHR, Nepal

**Ajita Ghimire**

IFHR, Nepal

---

## Research article

**Keywords:** Knowledge and attitude, community health care workers (HCW's), enabling factors, palika (Rural/ Urban Municipalities and Metropolitan city)

**Posted Date:** February 1st, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-192110/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at BMC Public Health on July 19th, 2021. See the published version at <https://doi.org/10.1186/s12889-021-11400-9>.

# Abstract

**1. Background:** Emerging and reemerging pathogens are global challenges for public health and the pandemic of Coronavirus disease 2019 is a reemerging case of an infectious disease caused by Severe Acute Respiratory Syndrome-Coronavirus-2. Health care workers worldwide are at higher risk worldwide and the situation is the same in Nepal. The knowledge and attitude of health workers will certainly mark the outcome towards this pandemic. So, this study aims to assess the knowledge and attitude of community health workers towards the prevention of COVID-19 virus.

**2. Methods:** A descriptive cross-sectional study was carried out among the community health workers of various provinces of Nepal. A semi-structured, self-administered questionnaire was prepared in Google Form and circulated to the sampled health workers through various social media platforms like Facebook, Messenger, Instagram, G-mails and messengers. Responses containing anonymized data were collected and analyzed using SPSS-version-20. The results were interpreted and checked with various demographic and enabling factors using chi-square test and logistic regression model. Also, ethical approval was taken from NHRC (ERB protocol registration number: **360/2020P**) prior to the conduction of study.

**3. Results:** Out of the total sample size of 450, we took 399 responses taking into consideration all the inclusion criteria. So, from 399 valid responses, 230 (47.6%) were males and 169 (42.4%) were females. 380 (95.2%) employed participants thought that wearing PPE will reduce the chances of getting COVID-19, majority of the participants 80.5% (321) responded that COVID-19 will successfully be controlled and staffs receiving excellent support from palika had high knowledge level. Knowledge level was significantly associated with the enabling factor "support from palika" ( $P=0.045$ ).

**4. Conclusions:** Proper and adequate knowledge and attitude towards COVID-19 is paramount in the prevention and control of SARS-CoV2 for everyone and especially the HCWs who are the frontline workers. There is a need to orient more HCWs regarding COVID-19, preventive measures. 45.9% of the respondents had no availability of COVID-19 response medical items for prevention of COVID-19 at their respective health facilities which is a serious issue which needs to be addressed very quickly to fight against the pandemic.

## Introduction

Emerging and reemerging pathogens are global challenges for public health [1]. Corona viruses are enveloped RNA viruses that are distributed broadly among humans, other mammals, and birds and causes respiratory, enteric, hepatic, and neurologic diseases [2, 3]. Six coronavirus species are known to cause human disease [4]. Four viruses 229E, OC43, NL63, and HKU1 are prevalent and typically cause common cold symptoms in immunocompetent individuals [4]. The two other strains, severe acute respiratory syndrome coronavirus (SARS-CoV) and Middle East respiratory syndrome (MERS) coronavirus (MERS-CoV) are zoonotic in origin and have been linked to sometimes fatal illness [5]. Corona virus

disease 2019 (COVID-19) is an infectious respiratory disease which is caused by newly recognized corona virus [6]. Given the high prevalence and wide distribution of corona viruses, the large genetic diversity and frequent recombination of their genomes, and increasing human–animal interface activities, novel corona viruses are likely to emerge periodically in humans owing to frequent cross-species infections and occasional spillover events [5, 11].

In late December 2019, several local health facilities reported clusters of patients with pneumonia of unknown cause that were epidemiologically linked to a seafood and wet animal wholesale market in Wuhan, Hubei Province, China [12]. In January 2020, the outbreak of the 2019 novel coronavirus (2019-nCoV) in China spread progressively to other countries [13, 14], with WHO declaring it a Public Health Emergency of International Concern [15]. Among the affected 200 countries globally (where 2,074,529 confirm cases and 139,378 deaths were reported as of April; and 23,560 confirm cases and 586 deaths South East Asia region including Nepal [16]. On 23rd January 2020, the first case was reported for COVID-19 in Nepal and now (21st Sept 2020), the data hiked at 64,122 cases with 46,233 recovered and 411 deaths [17].

Health workers are at the front line of the COVID-19 outbreak response and as such are exposed to hazards that put them at risk of infection [18]. Community health workers are one of the apex part of the health system of Nepal. After the decentralization, local government health system purely depends on community health workers. They are the pioneer for the prevention, promotion, and curative activities at the community level. So, their knowledge and attitude play a pivotal role for preventing COVID-19 at the community level and this study aims to find out the knowledge and attitude on prevention of novel coronavirus among community health workers in Nepal

## Methods

A descriptive cross-sectional study design was carried out to find out the status of knowledge and attitude of community health care workers towards the prevention of novel corona virus-2019. The study was conducted among the community health workers who are working at Health post and Primary Health Care level health institutions during the pandemic of COVID-19 in all the seven provinces of Nepal.

A semi-structured, self-administered questionnaire was prepared in google form and circulated to the sampled health workers of all 7 provinces. The online forms were disseminated via various social media platforms like Facebook, messenger, Instagram, g-mails, messengers, WhatsApp, and Viber. The informed consent was also attached along with the google forms before the beginning of the survey. Informed consent was taken from each respondent before the data collection and they were provided full rights to withdraw the study at any instance of time they wish. Also, ethical approval was taken from NHRC (ERB protocol registration number: 360/2020 P) prior to the conduction of study.

The sample size was calculated using the formula for the descriptive cross-sectional study, i.e.  $N = (Z_{\alpha/2})^2 p * q / d^2$  at 95% CI and 4% allowable error limit. With  $p = 75\%$ , our sample size was calculated to be 450.

The sampling of the respondents was carried out via convenience sampling technique (non-probability sampling method). For maintaining quality control and quality assurance of this study, designed questionnaires was pre-tested for content validity and tools were modified after pre-test and from observation field note. After the distribution of the questionnaire, 420 responses were obtained from the health care workers out of which 399 met the inclusion criteria which was 95% of the total response and 88% of the total sample size.

After the data collection, it was checked for completeness. The information will automatically be transferred to excel sheet and then transferred to SPSS. Responses containing anonymized data was collected and analyzed using SPSS-version 20. The results were interpreted in terms of percentage response, knowledge and attitude score and chi-square test and logistic regression was adopted to find the association between knowledge and attitude with various demographic and enabling factors.

## Results

Out the total sample size of 450, we took 399 responses taking into consideration all the inclusion criteria. Responses from health workers other than HP, PHC of Government of Nepal, not residing in Nepal, not working at the time of COVID-19 pandemic, were excluded. So, from 399 valid response, 230 (47.6%) were males and 169 (42.4%) were females. So, the gender of the participants was almost evenly distributed. Regarding the marital status of the respondents, more than half of them were married (61.7%) followed by unmarried (37.6%). Among the total respondents participated, nearly 1/3rd was HA (Health Assistant) (29.6%) and the least were ANM and staff nurses, both 4.8%. Majority of the respondent (61.2%) said that Television and radio was the media through which they firstly got the information about COVID-19. Nearly 2/5 (40.1%) of the respondents were working at Bagmati province while the least (3%) respondents were from Sudurpaschim province. Regarding the education level, 41.4% of the respondents had completed PCL (Proficiency Certificate level) and only 11.3% had completed their master level study. Nearly half (46.1%) of the respondents responded that their monthly income was between 25 k-35 k which was the average scale of government employee of Nepal of 5/6th level. The information is showed in Table 1.

Table 1  
Socio-demographic characteristics of participants (n = 399)

Variables	Category	Frequency	Percentage
Age	18–27	177	44.4
	28–37	137	34.3
	38–47	63	15.8
	48 and above	22	5.5
		<b>399</b>	<b>100.0</b>
Gender	Male	230	57.6
	Female	169	42.4
		<b>399</b>	<b>100</b>
Marital status	Married	246	61.7
	Unmarried	150	37.6
	Divorced	1	0.3
	Widow	2	0.5
			<b>100</b>
Designation	HA	118	29.6
	AHW	87	21.8
	ANM	97	24.3
	Nursing staff	19	4.8
	PHO	19	4.8
	Others	59	14.8
		<b>399</b>	<b>100</b>
Firstly, known Information about COVID-19	Social medias (FB/Twitter/ YouTube/ Instagram)	137	34.3
	Television/ radio	244	61.2
	Friends and relatives	7	1.8
	scientific journals	11	2.8
		<b>399</b>	<b>100</b>
Working province	Province 2	123	30.8

Variables	Category	Frequency	Percentage
	Bagmati	160	40.1
	Gandaki	6	1.5
	Karnali	98	24.6
	Sudurpaschim	12	3.0
		<b>399</b>	<b>100</b>
Education level	Master	45	11.3
	Bachelor	102	25.6
	PCL	165	41.4
	TSLC	87	21.8
		<b>399</b>	<b>100</b>
Monthly income	25K-35K	184	46.1
	35K-45K	92	23.1
	45K-55K	62	15.5
	Above 55K	61	15.3
		<b>399</b>	<b>100</b>
Orientation on COVID-19	Yes	82	20.6
	No	317	79.4
		<b>399</b>	<b>100</b>
Extra allowance provided	Yes	94	23.6
	No	305	76.4
		<b>399</b>	<b>100</b>
Working hours in a week	<=40	119	29.8
	> 40	280	70.2
		<b>399</b>	<b>100</b>
Support from palika	Excellent	17	4.3
	Good	251	62.9
	Poor	79	19.8
	very Poor	52	13.0

<b>Variables</b>	<b>Category</b>	<b>Frequency</b>	<b>Percentage</b>
		<b>399</b>	<b>100</b>
Palika level COVID-19 Response team	Yes	281	70.4
	No	23	5.8
	Don't know	95	23.8
		<b>399</b>	<b>100</b>

Table 2  
Frequency related to knowledge of participants

<b>Variables</b>	<b>Category</b>	<b>Frequency</b>	<b>Percentage</b>
COVID-19 is caused by	Virus	397	99.5
	Fungus	1	0.3
	Parasite	1	0.3
		<b>399</b>	<b>100</b>
COVID-19 is transmitted by close contact with infected person?	Yes	393	98.5
	No	6	1.5
		<b>399</b>	<b>100</b>
The first case of novel coronavirus was identified in	Beijing	17	4.3
	Shanghai	22	5.5
	Wuhan, Hubei	360	90.2
		<b>399</b>	<b>100</b>
Which is true about COVID-19?	All age group	87	21.8
	Mild in children	9	2.3
	Older and pre-existing medical condition	94	23.6
	All of the above	209	52.4
		<b>399</b>	<b>100</b>
Symptoms of COVID-19 are:	Fever	7	1.8
	Cough	3	0.8
	Shortness of breath	41	10.3
	All the above	348	87.2
		<b>399</b>	<b>100</b>
The COVID-19 virus spreads via respiratory droplets of infected individuals	Yes	395	99
	No	4	1
		<b>399</b>	<b>100</b>

<b>Variables</b>	<b>Category</b>	<b>Frequency</b>	<b>Percentage</b>
The isolation period for COVID-19 is:	1 week	3	0.8
	2 weeks	282	70.7
	3 weeks	74	18.5
	More than 4 weeks	40	10
		<b>399</b>	<b>100</b>
How to prevent from COVID-19?	Clean hands with soap and water/ sanitizer	57	14.3
	Cover mouth and nose when coughing and sneezing	6	1.5
	Avoid close contact with those who show signs of flu	5	1.3
	Properly cook meats, eggs before eating	3	0.8
	All the above	328	82.2
		<b>399</b>	<b>100</b>
COVID-19 vaccines are available in the market now	Yes	54	13.5
	No	345	86.5
		<b>399</b>	<b>100</b>
Health care workers are at high risk of COVID-19	Yes	389	97.5
	No	10	2.5
		<b>399</b>	<b>100</b>
PPE includes:	Gloves	16	4
	Gloves, apron	20	5
	Gloves, apron, face mask	11	2.7
	Gloves, apron, face mask and eye protection	352	88.3
		<b>399</b>	<b>100</b>

Table 1: Frequency regarding the various enabling factors had different responses. It was found from the study that very minimum (20.6%) employee has received orientation on COVID-19 and also very few staffs (23.6%) received extra allowance during the situation of COVID-19 and the rest did not get any allowance which might demotivated them in their performance. Regarding the working hours of health staffs, more than 2/3 (70.2%) said that they had been working for more than 40 hours in a week due to the situation of COVID-19. Support from respective palikas was found good (62.9%) which might have motivate them to work efficiently; maximum respondents (70.4%) said that they had “palika level COVID-19 response team” for the surveillance of situation of COVID-19.

In the knowledge related section shown in Table 2, almost all, 99.5% (397) participants said that COVID-19 is caused by virus and maximum, 98.5% (393) responded that COVID-19 is transmitted by close contact with infected people. Regarding the knowledge about the first case of novel corona virus identified place, it was found that majority of the participants, 90.2% (360) know that it was found firstly in Wuhan, China. Furthermore, regarding the knowledge about symptoms of COVID-19, 87.2% (348) said that fever, cough, shortness of breath was the symptoms of corona virus. Regarding the isolation period of COVID-19, majority of the participants, 70.7% (282) were well known about it and 82.2% (328) had right knowledge about prevention from COVID-19. Remarkably, all 100% of the respondents were well known about the fact that antibiotic was not the first line treatment of COVID-19. A total of 88.3% (352) participants correctly said that PPE includes gloves, apron, face mask and eye protection.

Table 3  
Frequency related to attitude variables of participants

Variables	Categories	Frequency	Percentage
Do you think you will get ill from COVID-19	Yes	224	56.1
	No	175	43.9
		<b>399</b>	<b>100</b>
If you get COVID-19, will you accept isolation in a Health facility?	Yes	367	92
	No	32	8
		<b>399</b>	<b>100</b>
If COVID-19 vaccine were available, would you take it?	Yes	353	88.5
	No	46	11.5
		<b>399</b>	<b>100</b>
Do you think wearing PPE reduces the chance of getting COVID-19?	Yes	380	95.2
	No	19	4.8
		<b>399</b>	<b>100</b>
Do you think lockdown is the best way to prevent COVID-19?	Yes	369	92.5
	No	30	7.5
		<b>399</b>	<b>100</b>
Do you feel bad when people don't use masks while coming for treatment?	Yes	367	92
	No	32	8
		<b>399</b>	<b>100</b>

Variables	Categories	Frequency	Percentage
Do you think COVID-19 will successfully be controlled?	Yes	321	80.5
	No	78	19.5
		<b>399</b>	<b>100</b>

Regarding the attitude of the respondents, 56% (224) had the feeling that they will get infected with COVID-19 at some time of the pandemic period, 92%(367) accepted that they will be in isolation in health facility if they are infected with COVID-19. Only 11.5% (46) participants refuse to take the vaccine if it was available in the market. Furthermore, 95.2% (380) employed participants thought that wearing PPE will reduce the chances of getting COVID-19. Also, regarding the COVID - 19 prevention, 92.5% (369) participants thought that lockdown was the best way. Regarding the attitude and perception towards the control of COVID-19, majority of the participants 80.5%(321) responded that COVID-19 will successfully be controlled followed by 19% (78) who thought that it won't be controlled at the earliest. This is illustrated in Table 3.

In bivariate analysis (chi-square test) Table 4, it was found that gender was significantly associated with knowledge level as male tend to have more knowledge level ( $P < 0.002$ ). Also, marital status was found to be significantly associated with the knowledge scores, for married knowledge level found to be high ( $P = 0.008$ ) compared to the other categories. It was also found that participants working at Bagmati province had high level of knowledge ( $P < 0.001$ ). Participants who were designated as "HA" and education level- PLC complete was significantly associated with the knowledge level. Similarly, people with monthly income 25 k-35 k was found to have more knowledge level ( $P < 0.005$ ) and also the significant association with knowledge level was found with the enabling factor "support from palika" ( $P < 0.045$ ). Staffs receiving excellent support from palika had high knowledge level.

Table 4  
Association of Knowledge with demographic and enabling factors

Variables	Categories	Knowledge		P-value
		High	Low	
Age	18–27	113	64	0.924
	28–37	88	49	
	38–47	28	25	
	48 and above	13	9	
Gender	Male	160	70	0.002
	Female	92	77	
Marital status	Married	142	104	0.008
	Unmarried	108	42	
	Divorced	1	0	
	Widow	1	1	
Designation	HA	81	37	< 0.001
	AHW	56	31	
	ANM	41	56	
	Nursing staff	17	2	
	PHO	16	3	
	Others	41	18	
Firstly, known Information about COVID-19	Social medias (FB/Twitter/YouTube/ Instagram )	101	36	< 0.001
	Television/ radio	134	110	
	Friends and relatives	6	1	
	scientific journals	11	0	
working province	Province 2	86	37	< 0.001
	Bagmati	111	49	
	Gandaki	5	1	
	Karnali	45	53	

Variables	Categories	Knowledge	P-value	
	Sudurpaschim	5	7	
Education level	Master	29	16	< 0.001
	Bachelor	68	34	
	PCL	121	44	
	TSLC	34	53	
Monthly income	25K-35K	118	66	0.005
	35K-45K	45	47	
	45K-55K	44	18	
	Above 55K	45	16	
Orientation on COVID-19	Yes	50	32	0.646
	No	202	115	
Working hours in a week	<=40	74	45	0.793
	> 40	178	102	
Extra allowance provided	Yes	57	37	0.562
	No	195	110	
Support from palika	Excellent	10	7	0.045
	Good	153	98	
	Poor	60	19	
	very Poor	29	23	
Sufficient equipment and medicines available at health facilities	Yes	131	121	0.259
	No	85	62	
Palika level COVID-19 Response team	Yes	179	102	0.879
	No	15	8	
	Don't know	58	37	

On the other hand, gender, marital status, working province and presence/ absence of palika level COVID-19 response team had significant association with the attitude score. Male participants ( $P < 0.001$ ), HA ( $P < 0.001$ ), staffs working at Bagmati province ( $P < 0.01$ ) and participants working in palika having “palika

level COVID-19 response team” ( $P < 0.001$ ) was found to have positive attitude toward the prevention of COVID-19 compared to others Table 5.

Table 5  
Association with Attitude and demographic/ enabling factors

Variables	Categories	Attitude		P-value
		Positive	Negative	
Age	18–27	123	54	0.26
	28–37	84	53	
	38–47	39	24	
	48 and above	13	5	
Gender	Male	167	63	< 0.001
	Female	96	73	
Marital status	Married	154	92	0.12
	Unmarried	108	42	
	Divorced	0	1	
	Widow	1	1	
Designation	HA	90	28	< 0.001
	AHW	56	31	
	ANM	49	48	
	Nursing staff	11	8	
	PHO	16	3	
	Others	41	18	
Firstly, known Information about COVID-19	Social medias (FB/Twitter/ YouTube/ Instagram )	104	33	0.002
	Television/ radio	144	100	
	Friends and relatives	6	1	
	scientific journals	9	2	
working province	Province 2	86	37	0.01
	Bagmati	114	46	
	Gandaki	5	1	
	Karnali	50	48	

Variables	Categories	Attitude		P-value
	Sudurpaschim	8	4	
Education level	Master	31	14	0.09
	Bachelor	67	35	
	PCL	117	48	
	TSLC	48	39	
Monthly income	25K-35K	128	56	0.184
	35K-45K	52	40	
	45K-55K	42	20	
	Above 55K	41	20	
Orientation on COVID-19	Yes	54	28	0.99
	No	209	108	
Working hours in a week	<=40	82	37	0.411
	> 40	181	99	
Extra allowance provided	Yes	60	34	0.626
	No	203	102	
Support from palika	Excellent	11	6	0.309
	Good	164	87	
	Poor	58	21	
	very Poor	30	22	
Sufficient equipment and medicines available at health facilities	Yes	123	60	0.615
	No	140	76	
Palika level COVID-19 Response team	Yes	191	90	0.001
	No	21	2	
	Don't know	51	44	

## Stepwise logistic regression:

We performed forward stepwise logistic regression to identify whether the demographic and enabling factors had significant association with knowledge and attitude status of the participants. In Table 6, the logistic regression analysis revealed that the odds of knowledge level was 2 times higher (AOR = 1.913 at 95% CI: 1.266–2.891) compared to the female participants. Knowledge level was significantly associated with the designation of the respondents as nursing staffs had 2 folds more knowledge (AOR = 2.243, 95%CI = 1.006-5.00) compared to those of the HA (Ref-HA). Similarly, the variables education level and monthly income of the participants were also found to have statistically significant relationship with the knowledge level.

Table 6

Effect of socio-demographic and enabling factors on Knowledge and attitude (logistic regression model)

<b>Variables</b>	<b>Categories</b>	<b>Adjusted Odds ratio (AOR)</b>	<b>95% CI</b>	<b>P-value</b>
Gender	Male	1.913	1.266–2.891	0.002
	Female	Ref		
Designation	HA	Ref		
	AHW	1.213	0.563–2.613	0.622
	ANM	0.991	0.450–2.181	0.982
	Nursing staff	2.243	1.006–5.000	0.048
	PHO	0.279	0.055–1.429	0.126
	Others	0.386	0.092–1.620	0.193
Education level	Master	Ref		
	Bachelor	0.465	0.228–0.950	0.036
	PCL	0.778	0.311–1.949	0.592
	TSLC	0.305	0.161–0.579	< 0.001
Monthly income	25K-35K	Ref		
	35K-45K	1.338	0.647–2.766	0.431
	45K-55K	2.523	1.157–5.500	0.020
	Above 55K	0.979	0.415–2.312	0.962
<b>Variable</b>	<b>Categories</b>	<b>Adjusted Odds ratio (AOR)</b>	<b>95% CI</b>	<b>p-value</b>
Gender	Male	1.884	1.216–2.918	0.005
	Female	Ref		

Variables	Categories	Adjusted Odds ratio (AOR)	95% CI	P-value
Firstly, known Information about COVID-19	Social medias (FB/Twitter/ YouTube/ Instagram)	Ref		
	Television/ radio	0.301	0.035–2.580	0.273
	Friends and relatives	0.367	0.076–1.768	0.211
	scientific journals	0.475	0.293–0.768	0.002
Palika level COVID-19 Response team	Yes	Ref		
	No	1.451	0.882–2.386	0.142
	Do not know	0.195	0.044–0.863	0.031

Furthermore, attitude status of the respondents was also found to have significant association with the variables like gender, firstly known information about COVID-19 and presence/ absence of palika level COVID-19 response team. Here, it was found that male tend to have positive attitude which is 2 times more (AOR:1.884, 95% CI: 1.216–2.918) compared to those of the females (ref-female). The other two variable are shown in the Table 6.

## Discussion

The study investigated the knowledge and attitude of community health care providers who were working during the time of COVID-19 pandemic at health facilities of Nepal. The result showed that gender, marital status, designation, information about COVID-19, working province, education level, monthly income, support from palika were found statistically significant with knowledge level and attitude status was associated with gender, designation, information about COVID-19, working province and palika level COVID-19 response team. HCWs are the frontline workers in the management of suspected and potential cases of the COVID-19. Their knowledge and attitude will likely have an important bearing on the course and containment of the pandemic.

Our findings revealed that most of the participants (63.2%) were knowledgeable about the prevention of COVID-19. The finding is consistent with other studies done at china, Indian and Saudi Arabia [21–24]. The high level of knowledge might be due to the information network of the current modern world and higher education level of the people. On the other hand, the attitude was also found positive among more than 2/3rd (65.9%) of the participants which was also correlated with various studies conducted at China India, Vietnam and Saudi Arabia [21–24]. The gender of the respondents was much evenly distributed as

47.6% male and 42.4% females which we considered a good gender response rate. Nearly half (46.1%) of the respondents responded that their monthly income was between 25 k-35 k which was the average scale of government employee of Nepal of 5/6th level and our almost respondents fall on this category. Regarding the enabling factors, it was found that very less (20.6%) participants had received orientation on COVID-19 but maximum (70.4%) said that they had palika level COVID-19 response team for the surveillance of the situation of COVID-19.

Regarding the knowledge level, nearly all the participants (98.5%) know that COVID-19 is transmitted by close contact with the infected people and 90% were correct regarding the place of outbreak of Corona virus i.e. Wuhan, China. Furthermore, regarding the knowledge about symptoms of COVID-19, 87.2% (348) said that fever, cough, shortness of breath was the symptoms of corona virus which was a very good response rate. The high rate might be because of the field and area they were working i.e. medical field. Regarding the isolation period of COVID-19, majority of the participants, 70.7% (282) were well known about it and 82.2% (328) had correct knowledge about prevention from COVID-19. This finding was consistent with a recent study done in Nepal [25]. Remarkably, all 100% of the respondents were well known about the fact that antibiotic was not the first line treatment of COVID-19.

Concerning the attitude of the respondents, more than half of the respondents (56%) had the feeling that they will be infected with Corona virus at some point of time of the pandemic. The positive point is that majority of the respondents (92%) accepted that they will stay at isolation if they get infected with corona virus. This is a very positive attitude of the respondents and especially the health workers. On the other hand, still 8% did not accept it, which is likely to be related to a lack of knowledge within the HCW's about current and important prevention and isolation strategies. A study from Taiwan also exhibit a similar result [26]. Majority of the respondents (80.5%) were positive that COVID-19 will successfully be controlled which is a positive attitude toward any health problems. Positive attitudes and high confidence in the control of COVID-19 can be explained by the government's unprecedented actions and prompt response in taking stringent control and precautionary measures against COVID-19, to safeguard citizens and ensure their well-being. These measures include the lockdown, and the suspension of all domestic and international flights, schools and universities, and the stepwise shutdown and prohibitory orders imposed.

In the study it was found that gender was associated with both knowledge level and attitude of the respondents. Male were found to have more knowledge and positive attitude toward the prevention of COVID-19 ( $p < 0.002$ ). This was contrary to the finding of the study conducted at India and China [27, 28]. The reason for male to be more in this category might be that in Nepal, still male get higher chance to get exposed in various field and orientation programs.

Television and radio were significantly associated with the knowledge and attitude of the respondents as higher level of knowledge was associated with those who get information via TV and radio. This might be due to the reason that TV and radio are the reliable medias [30] and only broadcast the information based on evidences despite some of the social medias like Facebook, You-tube, Instagram etc. Also, education

level ( $p < 0.001$ ), working province ( $p < 0.001$ ) and monthly income ( $p < 0.005$ ) are some of the demographic characteristics which have association with the knowledge level and regarding the enabling factors only “support from palika” is found to be associated with the knowledge level ( $p < 0.045$ ). Since this, study assessed only limited demographic variables, it is recommended that we include more demographic as well as socio-cultural variables in further studies.

This study also highlighted the area of some enabling factors which might be responsible for the knowledge and attitude of the HCW's. Most of the respondents (79.4%) had not received any orientation regarding COVID-19. This might have serious impact on the knowledge level of the HCW's [20, 21]. Another finding of this study revealed that 70% of the respondents work for more than 40 hours in a week and nearly 3/5th (76.4%) is not paid extra allowance during their extra hours of services. So, this will really demotivate the HCW's toward their dedicated services. This issue should be addressed by the concerned authority as soon as possible. Many of the palikas, HCW's work have a palika level COVID-19 response team (70.4%) followed by 5.8% not having and 23.8% were not aware of it. HCW's not being aware of the palika level COVID-19 response team clarifies that some of them are still not serious and concerned about their roles and responsibilities.

Nevertheless, the study findings revealed that the knowledge and attitude of the respondents/ HCW's working at various health facilities at different provinces of Nepal was found to be good despite some had low level of knowledge and negative attitude toward the prevention of COVID-19. The logistic regression (forward stepwise) also revealed several factors being associated with the knowledge and attitude of the HCW's. among then some were found matching with various studies conducted at national/ international level [19, 20, 21], whereas some were found to contradict with the various study findings [25–28] .

Our study also has some notable limitations. First, as those who had no internet access could not take part in the survey as the questionnaire were circulated through google forms. Also, the study could not take many variables of prime focus into consideration due to shorten the length of the questionnaire as it was self-administered and online based. Furthermore, since the attitudes are based on the healthcare workers' knowledge and availability of specialized logistics like PPE for maintaining the appropriate biosafety along with their perception of the healthcare system, their understanding of institutional preparedness are solely based on their own observation and perception.

Finally, the study aims to explore the ground reality of the knowledge and attitude of HCW's toward the prevention from COVID-19 in Nepal. This is an utmost need in the context of this pandemic to diminish the situation from worsening further. This study will eventually bridge the gap between the current situation of COVID-19 of the HCW's and the policy makers at the local, province and federal level of GoN. There is a need to educate and orient the HCW's regarding COVID-19 via different method and medias appropriate at the current situation of pandemic to upgrade the knowledge level of the HCW's and also building the HCW's confidence on existing health system regarding appropriate and timely containment

of the pandemic so improving the trust and reliability between government policy makers and the HCW's is of paramount importance.

## Conclusion

Proper and adequate knowledge and attitude towards COVID-19 is a paramount in the prevention and control of SARS-COV2. And it is utmost important to the frontline health workers who are daily exposed to the virus. Knowledge of the disease is considered the first steppingstone to any health education activity that is implemented. Knowing the causes and transmission sources of a disease, increases the likelihood that people will become more aware of the spread of communicable diseases, and of the preventive measures to slow the transmission. Although most of the study participants possess a good knowledge and positive attitude toward the prevention of COVID-19. Still there is a need to orient more HCW's regarding COVID-19, preventive measures. Also, very less (45.9%) of the respondents had no availability of COVID-19 response medical items for prevention. So, it is a prime concern that the authoritative body take the responsibility of their HCW's and supply the necessary medical items for them to fight against the current pandemic of COVID-19.

## Abbreviations

COVID-19

Corona Virus Disease-19

SARS

Severe Acute Respiratory Syndrome

HCW's

Health Care Workers

PPE

Personal Protective equipment

PCL

Proficiency Certificate Level

TSLC

Technical School Leaving Certificate

HP

Health Post

PHCC

Primary Health Care Center

GoN

Government of Nepal

## Declarations

**Availability of data and materials.**

The datasets generated and /or analyzed during the current study are not publicly available but are available from the corresponding author on reasonable request.

### **Acknowledgement:**

The author would firstly like to thank the ERC (Ethical Review Committee) of NHRC (Nepal Health Research Council) for their support, cooperation, and permission to conduct the research. Also we owe our deepest gratitude to all the Health Care Worker of Nepal who had continually been devoted toward the service and even in this time of COVID-19 pandemic and especially to those who provided their valuable time to participate in the study. Also, our sincere thanks go to all the team of IFHR (Innovative foundation for Health and Research) who had been committed to make this research work a success. And last but not the least to all those who directly/ indirectly supported us to distribute the form to the respective participants and feedback while designing the questionnaire. We really appreciate for all your love and support.

### **Funding:**

The study was not funded by any organization or institution.

### **Authors information:**

#### **Affiliations:**

Amul Shrestha (AS), Tek Bahadur Thapa (TBT), Mahendra Giri (MG), Sanjiv Kumar (SK), Sakil Ali (SA), Harikala Thapa (HT), Pawan Prakash Dhimi (PPD), Arun Sahi (AS) and Ajita Ghimire (AG), all are involve in the organization IFHR (Innovative Foundation for Health and Research).

#### **Contribution:**

AS, TBT, MG, SK, SA, HT, PPD conceptualize the study. AS designed the methodology of the study and the background; the literature search of the study was done by TBT, MG and SA. PPD, HT, AS and AG designed the questionnaire and prepare the google form to circulate to the participants. Also, AS and SK worked out on the workplan for the conduction of the study. The ethical letter for the research was collected by MG and TBT. After the data collection, PPD, MG worked on the data cleaning and processing. Following this the data analysis, the chi-square testing/ logistic regression model for assessing the variables and the first draft of the manuscript was written by AS with inputs and reviews from TBT, MG, SK, SA, PPD, HT, AS and AG. Finally, all author read and approved the final manuscript.

#### **\*Corresponding author:**

Correspondence: Amul Shrestha

ORCID- Amul Shrestha: <https://orcid.org/0000-0001-7892-4279>

## Ethics declarations:

### Ethics approval and consent to participate

Ethical approval was taken from NHRC (ERB protocol registration number: **360/2020P**) prior to the conduction of study and all the participants were provided informed consent form before the start of the survey via google forms.

### Consent for publication

Not applicable.

### Competing interest:

The authors declare that they have no competing interests.

### Author Details:

<sup>1</sup> Innovative Foundation for Health and Research (IFHR), Chabahil, Kathmandu-Bagmati Province, Nepal.

## References

1. Bi Y, Shi W, Chen J, Chen Q, Ma Z, Wong G, Tian W, Yin R, Fu G, Yang Y, et al **CASCIRE surveillance network and work on avian influenza viruses**. *Sci. China Life Sci.*, 60 (2017), pp. 1386 – 139.
2. Almazan F, Gonzalez JM, Penzes Z, Izeta A, Calvo E, Plana-Duran J, Enjuanes L. Engineering the largest RNA virus genome as an infectious bacterial artificial chromosome. *Proc. Natl. Acad. Sci. USA*. 2000;97:5516–5521.
3. Masters PS, Perlman S. Coronaviridae. In: Knipe DM, Howley PM, eds. *Fields virology*. 6th ed. Lippincott Williams & Wilkins, 2013:825 – 58.
4. Su S, Wong G, Shi W, et al. Epidemiology, genetic recombination, and pathogenesis of coronaviruses. *Trends Microbiol.* 2016;24:490–502.
5. Zhong NS, et al. Epidemiology and cause of severe acute respiratory syndrome (SARS) in Guangdong, People's Republic of China, in February, 2003. *Lancet*. 2003;362:1353–8.
6. Zhong NS, Zheng BJ, Li YM, et al. Epidemiology and cause of severe acute respiratory syndrome (SARS) in Guangdong, People's Republic of China, in February, 2003. *Lancet*. 2003;362:1353–8.
7. Ksiazek TG, Erdman D, Goldsmith CS, et al. A novel coronavirus associated with severe acute respiratory syndrome. *N Engl J Med*. 2003;348:1953–66.
8. Drosten C, Günther S, Preiser W, et al. Identification of a novel coronavirus in patients with severe acute respiratory syndrome. *N Engl J Med*. 2003;348:1967–76.
9. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med*. 2012;367:1814–20.

10. Coronavirus [Internet]. World Health Organization. World Health Organization; [cited 2020Apr15]. Available from: [https://www.who.int/health-topics/coronavirus#tab=tab\\_1](https://www.who.int/health-topics/coronavirus#tab=tab_1).
11. Wong G, Liu W, Liu Y, Zhou B, Bi Y, Gao GF. MERS, SARS, and Ebola: the role of super-spreaders in infectious disease. *Cell Host Microbe*. 2015;18:398–401.
12. Report of clustering pneumonia of unknown etiology in Wuhan City. Wuhan Municipal Health Commission 2019, <http://wjw.wuhan.gov.cn/front/web/shoeDetail/201912310898>.
13. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; published online Jan 24. [http://dx.doi.org/10.1016/S0140-6736\(20\)30183-5](http://dx.doi.org/10.1016/S0140-6736(20)30183-5).
14. Chan JF-W, Yuan S, Kok K-H, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020; published online Jan 24. [http://dx.doi.org/10.1016/S0140-6736\(20\)30154-9](http://dx.doi.org/10.1016/S0140-6736(20)30154-9).
15. WHO. Statement on the second meeting of the International Health Regulations. (2005) Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV). 2020. [https://www.who.int/news-room/detail/30-01-2020-statement-on-the-secondmeeting-of-the-international-healthregulations-\(2005\)-emergency-committeeregarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-secondmeeting-of-the-international-healthregulations-(2005)-emergency-committeeregarding-the-outbreak-of-novel-coronavirus-(2019-ncov)) (accessed Jan 2, 2020).
16. World Health Organization. Infection prevention and control during health care when COVID-19 is suspected [https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-\(ncov\)-infection-is-suspected-20200125](https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-(ncov)-infection-is-suspected-20200125).
17. Coronavirus disease. (COVID-19) outbreak updates & resource materials, MOHP/HEDMU/HEOC-Nepal; <https://heoc.mohp.gov.np/update-on-novel-corona-virus-covid-19/>; <https://covid19.mohp.gov.np/>.
18. Country. & Technical Guidance - Coronavirus disease (COVID-19), WHO/Global/Region- <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance-publications>.
19. Coronavirus [Internet]. World Health Organization. World Health Organization; [cited 2020Apr15]. Available from: [https://www.who.int/health-topics/coronavirus#tab=tab\\_1](https://www.who.int/health-topics/coronavirus#tab=tab_1).
20. COVID-19 Strategy. Update March 23rd. [cited 2020Apr15]. Available from: <https://www.options-it.com/blog/2020/03/23/covid-19-strategy-update-march-23rd/>.
21. Giao H, Ngoc Han NT, Khanh TV, Ngan VK. Knowledge and attitude toward COVID-19 among healthcare workers at District 2 Hospital, Ho Chi Minh City. *Asian Pacific Journal of Tropical Medicine*. 2020;Mar29;:1–6.
22. Malik YS, Sircar S, Bhat S, Vinodhkumar OR, Tiwari R, Sah R, et al. Emerging coronavirus disease (COVID-19), a pandemic public health emergency with animal linkages: Current status update. Preprints doi:10.20944/preprints202003.0343.v1. [Online] Available from [https://www.preprints.org/manuscript/202003.0343/download/final\\_file](https://www.preprints.org/manuscript/202003.0343/download/final_file).

23. Al-Mohrej OA, Al-Shirian SD, Al-Otaibi SK, Tamim HM, Masuadi EM, Fakhoury HM. Is the Saudi public aware of Middle East respiratory syndrome? *J Infect Public Health*. 2016;9:259–66. doi:10.1016/j.jiph.2015.10.003.
24. Aldowyan N, Abdallah AS, El-Gharabawy R, Knowledge. Attitude and Practice (KAP) Study about Middle East Respiratory Syndrome Coronavirus (MERS-CoV) among Population in Saudi Arabia. *International Archives of Medicine*. 2017. doi:10.3823/2524.
25. Prabina Ghimire. Samridh Dhungel Anil Pokhrel. Knowledge, Attitude and Practice of healthcare workers Towards Coronavirus Disease 2019 (COVID-19) Pandemic. DOI: <https://doi.org/10.33314/jnhrc.v18i2.2658>. Available from: <https://www.jnhrc.com.np/index.php/jnhrc/article/view/2658/950>.
26. Deng JF, Olowokure B, Kaydos-Daniels SC, Chang HJ, Barwick RS, Lee ML, et al. Severe acute respiratory syndrome (SARS): Knowledge, attitudes, practices and sources of information among physicians answering a SARS fever hotline service. *Public Health*. 2006;120(1):15–9.
27. Han NThiN, Van Khanh T, Ngan VK, Van Tam V, An PL. Knowledge and attitude toward COVID-19 among healthcare workers at District 2 Hospital, Ho Chi Minh City Huynh. doi: 10.4103/1995-7645.280396. Available at: [https://www.apjtm.org/temp/AsianPacJTropMed136260-2683971\\_072719.pdf](https://www.apjtm.org/temp/AsianPacJTropMed136260-2683971_072719.pdf).
28. Sonam Maheshwari PK, Gupta RS., Pravesh Rawat. Knowledge, attitude, and practice towards coronavirus disease 2019 (COVID-19) among medical students: A cross-sectional study. Doi: 10.4103/2221-6189.283886.
29. Mohammed K, Al-Hanawi K, Angawi N. Alshareef et.al. Knowledge, Attitude and Practice Toward COVID-19 Among the Public in the Kingdom of Saudi Arabia: A Cross-Sectional Study. <https://doi.org/10.3389/fpubh.2020.00217>. Available at: <https://www.frontiersin.org/articles/10.3389/fpubh.2020.00217/full>.
30. BBC news services- “Radio is the most trusted source of news and information, says Ofcom” Available at: <https://www.bbc.com/news/10121759>.