Soft subalgebras and ideals of BCK|BCI-algebras based on $\mathbb{N}$ -structures

Hashem Bordbar (✉ Hashem.bordbar@ung.si)

University of Nova Gorica <https://orcid.org/0000-0003-3871-217X>

Rajab Ali Borzooei

Shahid Beheshti University

Arsham Borumand Saeid

Shahid Bahonar University of Kerman

Young Bae Bae Jun

Gyeongsang National University

Research Article

Keywords: $\mathbb{N}$ -ideal of types $(\mathbb{N}, \mathbb{N})$ and $(\mathbb{N}, \mathbb{N} \vee \mathbb{Q})$, soft $\mathbb{N}$ -set, soft $\mathbb{N}$ q-set, soft $\mathbb{N} \vee \mathbb{Q}$ -set, soft $\mathbb{N}$ -subalgebra, soft $\mathbb{N}$ -ideal

Posted Date: May 3rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-193896/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Soft Computing on August 20th, 2021. See the published version at <https://doi.org/10.1007/s00500-021-06078-5>.

Soft subalgebras and ideals of BCK/BCI -algebras based on $\mathcal{N}$ -structures

HASHEM BORDBAR^{1,*}, R.A. BORZOOEI², A. BORUMAND SAEID³ AND YOUNG BAE JUN⁴

¹*Center for Information Technology and Applied Mathematics,
University of Nova Gorica, Slovenia*
e-mail: hashem.bordbar@ung.si

²*Department of Mathematics
Shahid Beheshti University, Tehran, Iran,*
e-mail: borzooei@sbu.ac.ir

³*Department of Pure Mathematics, Faculty of Mathematics and Computer,
Shahid Bahonar University of Kerman, Kerman, Iran,*
e-mail: arsham@uk.ac.ir

⁴*Department of Mathematics Education,
Gyeongsang National University, Jinju 52828, Korea*
e-mail: skywine@gmail.com

Abstract The notions of $\mathcal{N}$ -ideal of types $(\in, \in)$ and $(\in, \in \vee q)$, soft $\mathcal{N}_{\in}$ -set, soft $\mathcal{N}_q$ -set, soft $\mathcal{N}_{\in \vee q}$ -set, soft $\mathcal{N}$ -subalgebra and soft $\mathcal{N}$ -ideal in BCK/BCI -algebra are introduced, and several properties are investigated. Characterizations of $\mathcal{N}$ -subalgebra of types $(\in, \in)$ and $(\in, \in \vee q)$, $\mathcal{N}$ -ideal of types $(\in, \in)$ and $(\in, \in \vee q)$, soft $\mathcal{N}$ -subalgebra and soft $\mathcal{N}$ -ideal are discussed.

Mathematics Subject Classification (2010): 06F35, 03G25.

Keywords: $\mathcal{N}$ -ideal of types $(\in, \in)$ and $(\in, \in \vee q)$, soft $\mathcal{N}_{\in}$ -set, soft $\mathcal{N}_q$ -set, soft $\mathcal{N}_{\in \vee q}$ -set, soft $\mathcal{N}$ -subalgebra, soft $\mathcal{N}$ -ideal.

*Corresponding author.

1 Introduction

The generalization of the crisp set to fuzzy sets relied on spreading positive information that fit the crisp point $\{1\}$ into the interval $[0, 1]$. Because no negative meaning of information is suggested, we now feel a need to deal with negative information. To do so, we also feel a need to supply mathematical tools. To attain such an object, Jun et al. [6] introduced a new function which is called a negative-valued function and constructed $\mathcal{N}$ -structures. They applied $\mathcal{N}$ -structures to BCK/BCI -algebras, and discussed $\mathcal{N}$ -subalgebras and $\mathcal{N}$ -ideals in BCK/BCI -algebras. Jun et al. [7] considered closed ideals in BCH -algebras based on $\mathcal{N}$ -structures. Bordbar et al. [3], [8] applied these notions. To obtain a more general form of an $\mathcal{N}$ -subalgebra in BCK/BCI -algebras, Jun et al. [5] defined the notions of $\mathcal{N}$ -subalgebras of types $(\in, \in)$, $(\in, q)$, $(\in, \in \vee q)$, $(q, \in)$, (q, q) and $(q, \in \vee q)$, and investigated related properties. They provided a characterization of an $\mathcal{N}$ -subalgebra of type $(\in, \in \vee q)$, and considered conditions for an $\mathcal{N}$ -structure to be an $\mathcal{N}$ -subalgebra of type $(q, \in \vee q)$. Also for more information about soft algebraic structures, please refer to [1], [2] and [9].

In this paper, we introduce $\mathcal{N}$ -ideal of types $(\in, \in)$ and $(\in, \in \vee q)$, soft $\mathcal{N}$ -set, soft $\mathcal{N}_{\in}$ -set, soft $\mathcal{N}_q$ -set, soft $\mathcal{N}_{\in \vee q}$ -set, soft $\mathcal{N}$ -subalgebra and soft $\mathcal{N}$ -ideal in BCK/BCI -algebra, and investigate several properties. We consider characterizations of $\mathcal{N}$ -subalgebra of types $(\in, \in)$ and $(\in, \in \vee q)$, $\mathcal{N}$ -ideal of types $(\in, \in)$ and $(\in, \in \vee q)$, soft $\mathcal{N}$ -subalgebra and soft $\mathcal{N}$ -ideal.

2 Preliminaries

Let $K(\tau)$ be the class of all algebras with type $\tau = (2, 0)$. By a BCI -algebra we mean a system $X := (X, *, 0) \in K(\tau)$ in which the following axioms hold:

- (i) $((x * y) * (x * z)) * (z * y) = 0$,
- (ii) $(x * (x * y)) * y = 0$,
- (iii) $x * x = 0$,
- (iv) $x * y = y * x = 0 \implies x = y$

for all $x, y, z \in X$. If a BCI -algebra X satisfies $0 * x = 0$ for all $x \in X$, then we say that X is a BCK -algebra. We can define a partial ordering $\leq$ by

$$(\forall x, y \in X) (x \leq y \iff x * y = 0).$$

In a BCK/BCI -algebra X , the following hold:

$$(a1) \quad (\forall x \in X) (x * 0 = x),$$

$$(a2) \quad (\forall x, y, z \in X) ((x * y) * z = (x * z) * y).$$

for all $x, y, z \in X$.

A non-empty subset S of a BCK/BCI -algebra X is called a *subalgebra* of X if $x * y \in S$ for all $x, y \in S$. A subset I of a BCK/BCI -algebra X is called an *ideal* of X if

$$0 \in I, \tag{2.1}$$

$$(\forall x, y \in X) (x * y \in I, y \in I \Rightarrow x \in I). \tag{2.2}$$

We refer the reader to the books [4] and [10] for further information regarding BCK/BCI -algebras.

For any family $\{a_i \mid i \in \Lambda\}$ of real numbers, we define

$$\bigvee \{a_i \mid i \in \Lambda\} := \begin{cases} \max\{a_i \mid i \in \Lambda\} & \text{if } \Lambda \text{ is finite,} \\ \sup\{a_i \mid i \in \Lambda\} & \text{otherwise.} \end{cases}$$

$$\bigwedge \{a_i \mid i \in \Lambda\} := \begin{cases} \min\{a_i \mid i \in \Lambda\} & \text{if } \Lambda \text{ is finite,} \\ \inf\{a_i \mid i \in \Lambda\} & \text{otherwise.} \end{cases}$$

For two real numbers a_1 and a_2 , $\bigvee \{a_1, a_2\}$ and $\bigwedge \{a_1, a_2\}$ are also denoted by $a_1 \vee a_2$ and $a_1 \wedge a_2$, respectively.

Denote by $\mathcal{F}(X, [-1, 0])$ the collection of functions from a set X to $[-1, 0]$. We say that an element of $\mathcal{F}(X, [-1, 0])$ is a *negative-valued function* from X to $[-1, 0]$ (briefly, $\mathcal{N}$ -function on X). By an $\mathcal{N}$ -structure we mean an ordered pair (X, f) of X and an $\mathcal{N}$ -function f on X .

Let (X, f) be an $\mathcal{N}$ -structure in which f is given by

$$f(y) = \begin{cases} 0 & \text{if } y \neq x, \\ \alpha & \text{if } y = x \end{cases}$$

where $\alpha \in [-1, 0)$. In this case, f is denoted by x_α and we call (X, x_α) a *point $\mathcal{N}$ -structure*. For any $\mathcal{N}$ -structure (X, g) , we say that a point $\mathcal{N}$ -structure (X, x_α) is an $\mathcal{N}_\infty$ -subset (resp., $\mathcal{N}_q$ -subset) of (X, g) if $g(x) \leq \alpha$ (resp., $g(x) + \alpha + 1 < 0$). If a point $\mathcal{N}$ -structure (X, x_α) is an $\mathcal{N}_\infty$ -subset of (X, g) or an $\mathcal{N}_q$ -subset of (X, g) , we say (X, x_α) is an $\mathcal{N}_{\infty q}$ -subset of (X, g) .

An $\mathcal{N}$ -structure (X, f) is called an $\mathcal{N}$ -subalgebra of type $(\in, \in)$ (resp., type $(\in, \in \vee q)$) (see [5]) if whenever two point $\mathcal{N}$ -structures (X, x_α) and (X, y_β) are $\mathcal{N}_\in$ -subsets of (X, f) then the point $\mathcal{N}$ -structure $(X, (x * y)_{\vee\{\alpha, \beta\}})$ is an $\mathcal{N}_\in$ -subset (resp., $\mathcal{N}_{\in \vee q}$ -subset) of (X, f) .

3 Soft $\mathcal{N}$ -subalgebras and soft $\mathcal{N}$ -ideals

In what follows, let X denote a BCK/BCI -algebra unless otherwise specified. For a subset Δ of $[-1, 0]$, a pair $(\mathcal{A}, \Delta)$ is called a *soft $\mathcal{N}$ -set* over X , where $\mathcal{A} : \Delta \rightarrow \mathcal{P}(X)$ is a mapping.

Given an $\mathcal{N}$ -structure (X, f) and $\Delta \subseteq [-1, 0]$, we define two mappings:

$$\mathcal{A}_\in : \Delta \rightarrow \mathcal{P}(X), \alpha \mapsto \{x \in X \mid (X, x_\alpha) \text{ is an } \mathcal{N}_\in\text{-subset of } (X, f)\} \quad (3.1)$$

$$\mathcal{A}_q : \Delta \rightarrow \mathcal{P}(X), \alpha \mapsto \{x \in X \mid (X, x_\alpha) \text{ is an } \mathcal{N}_q\text{-subset of } (X, f)\} \quad (3.2)$$

Then $(\mathcal{A}_\in, \Delta)$ and $(\mathcal{A}_q, \Delta)$ are soft $\mathcal{N}$ -sets over X . If $\mathcal{A}_\in(\alpha) \neq \emptyset$ (resp., $\mathcal{A}_q(\alpha) \neq \emptyset$) for $\alpha \in \Delta$, then we say $(\mathcal{A}_\in, \Delta)$ (resp., $(\mathcal{A}_q, \Delta)$) is a *soft $\mathcal{N}_\in$ -set* (resp., *soft $\mathcal{N}_q$ -set*) over X . A *soft $\mathcal{N}_{\in \vee q}$ -set* over X is defined to be the union of a soft $\mathcal{N}_\in$ -set and a soft $\mathcal{N}_q$ -set over X , and is denoted by $(\mathcal{A}_{\in \vee q}, \Delta)$ where $\mathcal{A}_{\in \vee q}(\alpha) = \mathcal{A}_\in(\alpha) \cup \mathcal{A}_q(\alpha)$ for all $\alpha \in \Delta$.

Definition 3.1. A soft $\mathcal{N}$ -set $(\mathcal{A}, \Delta)$ over X is called a *soft $\mathcal{N}$ -subalgebra* over X if it satisfies:

$$(\forall \alpha \in \Delta) (\mathcal{A}(\alpha) \neq \emptyset \Rightarrow \mathcal{A}(\alpha) \text{ is a subalgebra of } X). \quad (3.3)$$

Theorem 3.2. Given an $\mathcal{N}$ -structure (X, f) and $\Delta = [-1, 0]$, the soft $\mathcal{N}_\in$ -set $(\mathcal{A}_\in, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X if and only if (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in)$.

Proof. Assume that $(\mathcal{A}_\in, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X . If (X, f) is not an $\mathcal{N}$ -subalgebra of type $(\in, \in)$, then there exist $a, b \in X$ and $t \in \Delta$ such that (X, a_t) and (X, b_t) are $\mathcal{N}_\in$ -subsets of (X, f) , but $(X, (a * b)_t)$ is not an $\mathcal{N}_\in$ -subset of (X, f) . Hence $a, b \in \mathcal{A}_\in(t)$ and $a * b \notin \mathcal{A}_\in(t)$, which shows that $\mathcal{A}_\in(t)$ is not a subalgebra of X . This is a contradiction, and therefore (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in)$.

Conversely, suppose that (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in)$ and let $\alpha \in \Delta$ be such that $\mathcal{A}_\in(\alpha) \neq \emptyset$. If $x, y \in \mathcal{A}_\in(\alpha)$, then (X, x_α) and (X, y_α) are $\mathcal{N}_\in$ -subsets of (X, f) . Thus $(X, (x * y)_\alpha) = (X, (x * y)_{\vee\{\alpha, \alpha\}})$ is an $\mathcal{N}_\in$ -subset of (X, f) , that is, $x * y \in \mathcal{A}_\in(\alpha)$. Hence $\mathcal{A}_\in(\alpha)$ is a subalgebra of X for all $\alpha \in \Delta$, and thus $(\mathcal{A}_\in, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X . $\square$

Lemma 3.3 ([5]). *An $\mathcal{N}$ -structure (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in)$ if and only if the following inequality is valid.*

$$(\forall x, y \in X) \left(f(x * y) \leq \bigvee \{f(x), f(y)\} \right). \quad (3.4)$$

Theorem 3.4. *Given an $\mathcal{N}$ -structure (X, f) and $\Delta = [-1, 0)$, the soft $\mathcal{N}_q$ -set $(\mathcal{A}_q, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X if and only if (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in)$.*

Proof. Assume that (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in)$ and let $\alpha \in \Delta$ be such that $\mathcal{A}_q(\alpha) \neq \emptyset$. If $x, y \in \mathcal{A}_q(\alpha)$, then (X, x_α) and (X, y_α) are $\mathcal{N}_q$ -subsets of (X, f) , and so $f(x) + \alpha + 1 < 0$ and $f(y) + \alpha + 1 < 0$. It follows from (3.4) that

$$f(x * y) + \alpha + 1 \leq \bigvee \{f(x), f(y)\} + \alpha + 1 < 0$$

and so that $(X, (x * y)_\alpha) = (X, (x * y)_{\bigvee \{\alpha, \alpha\}})$ is an $\mathcal{N}_q$ -subset of (X, f) . Hence $x * y \in \mathcal{A}_q(\alpha)$, and thus $\mathcal{A}_q(\alpha)$ is a subalgebra of X for all $\alpha \in \Delta$ with $\mathcal{A}_q(\alpha) \neq \emptyset$. Therefore $(\mathcal{A}_q, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X .

Conversely, suppose that the soft $\mathcal{N}_q$ -set $(\mathcal{A}_q, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X , and assume that $f(a * b) > \bigvee \{f(a), f(b)\}$ for some $a, b \in X$. Then there exists $t \in \Delta$ such that

$$f(a * b) + t + 1 \geq 0 \text{ and } \bigvee \{f(a), f(b)\} + t + 1 < 0.$$

It follows that (X, a_t) and (X, b_t) are $\mathcal{N}_q$ -subsets of (X, f) but $(X, (a * b)_t)$ is not an $\mathcal{N}_q$ -subset of (X, f) . This is a contradiction, and hence $f(x * y) \leq \bigvee \{f(x), f(y)\}$ for all $x, y \in X$. Therefore (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in)$ by Lemma 3.3. $\square$

Theorem 3.5. *Given an $\mathcal{N}$ -structure (X, f) and the soft $\mathcal{N}_\in$ -set $(\mathcal{A}_\in, \Delta)$ with $\Delta = [-1, -0.5)$, the following are equivalent:*

- (1) $(\mathcal{A}_\in, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X .
- (2) $(\forall x, y \in X) (\bigwedge \{f(x * y), -0.5\} \leq \bigvee \{f(x), f(y)\})$.

Proof. Assume that the soft $\mathcal{N}_\in$ -set $(\mathcal{A}_\in, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X . Then $\mathcal{A}_\in(\alpha)$ is a subalgebra of X for all $\alpha \in \Delta$ with $\mathcal{A}_\in(\alpha) \neq \emptyset$. If there exist $a, b \in X$ such that

$$\bigwedge \{f(a * b), -0.5\} > t := \bigvee \{f(a), f(b)\},$$

then $t \in \Delta$, and (X, a_t) and (X, b_t) are $\mathcal{N}_\in$ -subsets of (X, f) , that is, $a, b \in \mathcal{A}_\in(t)$, but $(X, (a * b)_t)$ is not an $\mathcal{N}_\in$ -subset of (X, f) , i.e., $a * b \notin \mathcal{A}_\in(t)$. This is a contradiction, and so $\bigwedge \{f(x * y), -0.5\} \leq \bigvee \{f(x), f(y)\}$ for all $x, y \in X$.

Conversely, suppose that (2) is valid. Let $x, y \in \mathcal{A}_\epsilon(\alpha)$ for every $\alpha \in \Delta$. Then (X, x_α) and (X, y_α) are $\mathcal{N}_\epsilon$ -subsets of (X, f) , and so

$$\bigwedge \{f(x * y), -0.5\} \leq \bigvee \{f(x), f(y)\} \leq \alpha < -0.5.$$

It follows that $(X, (x * y)_\alpha)$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) , that is, $x * y \in \mathcal{A}_\epsilon(\alpha)$. Thus $\mathcal{A}_\epsilon(\alpha)$ is a subalgebra of X , and therefore $(\mathcal{A}_\epsilon, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X . $\square$

Lemma 3.6 ([5]). *An $\mathcal{N}$ -structure (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in \vee q)$ if and only if it satisfies:*

$$(\forall x, y \in X) \left(f(x * y) \leq \bigvee \{f(x), f(y), -0.5\} \right). \quad (3.5)$$

Theorem 3.7. *Given an $\mathcal{N}$ -structure (X, f) and a soft $\mathcal{N}_\epsilon$ -set $(\mathcal{A}_\epsilon, \Delta)$, the following assertions are equivalent:*

- (1) (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in \vee q)$.
- (2) $(\mathcal{A}_\epsilon, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X for $\Delta = [-0.5, 0)$.

Proof. Assume that (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in \vee q)$. Let $x, y \in X$ and $\alpha \in \Delta$ be such that $x, y \in \mathcal{A}_\epsilon(\alpha)$. Then (X, x_α) and (X, y_α) are $\mathcal{N}_\epsilon$ -subsets of (X, f) . It follows from (3.5) that

$$(\forall x, y \in X) \left(f(x * y) \leq \bigvee \{f(x), f(y), -0.5\} \leq \bigvee \{\alpha, -0.5\} = \alpha \right).$$

This shows that $(X, (x * y)_\alpha)$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) . Thus $x * y \in \mathcal{A}_\epsilon(\alpha)$, and so $(\mathcal{A}_\epsilon, \Delta)$ is a soft $\mathcal{N}$ -subalgebra over X .

Conversely, suppose that the soft $\mathcal{N}_\epsilon$ -set $(\mathcal{A}_\epsilon, \Delta)$ with $\Delta = [-0.5, 0)$ is a soft $\mathcal{N}$ -subalgebra over X . Assume that (3.5) is not valid. Then

$$f(a * b) > t \geq \bigvee \{f(a), f(b), -0.5\}$$

for some $t \in \Delta$ and $a, b \in X$. It follows that (X, a_t) and (X, b_t) are $\mathcal{N}_\epsilon$ -subsets of (X, f) , and so that $a, b \in \mathcal{A}_\epsilon(t)$. But $f(a * b) > t$ induces that $(X, (a * b)_t)$ is not an $\mathcal{N}_\epsilon$ -subset of (X, f) . This is a contradiction, and thus $f(x * y) \leq \bigvee \{f(x), f(y), -0.5\}$ for all $x, y \in X$. Using Lemma 3.6, we know that (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in \vee q)$. $\square$

Theorem 3.8. *Let $(\mathcal{A}_\epsilon, \Delta)$ be a soft $\mathcal{N}_\epsilon$ -set over X . If $\Delta = [-0.5, 0)$, then for any subalgebra S of X there exists an $\mathcal{N}$ -subalgebra (X, f) of type $(\in, \in \vee q)$ such that $\mathcal{A}_\epsilon(\alpha) = S$ for all $\alpha \in \Delta$.*

Proof. Take an $\mathcal{N}$ -subalgebra (X, f) in which f is given as follows:

$$f : X \rightarrow [-1, 0], \quad x \mapsto \begin{cases} \alpha \in \Delta & \text{if } x \in S, \\ 0 & \text{if otherwise.} \end{cases}$$

Obviously, $\mathcal{A}_\epsilon(\alpha) = S$ for all $\alpha \in \Delta$. Assume that

$$f(a * b) > \bigvee \{f(a), f(b), -0.5\}$$

for some $a, b \in X$. Then $f(a * b) = 0$ and $\bigvee \{f(a), f(b), -0.5\} = \alpha$ since $|\text{Im}(f)| = 2$. It follows that $f(a) = \alpha = f(b)$ so that $a, b \in S$. But $a * b \notin S$ since $f(a * b) = 0$. This is impossible, and so

$$f(x * y) \leq \bigvee \{f(x), f(y), -0.5\}$$

for all $x, y \in X$. Therefore (X, f) is an $\mathcal{N}$ -subalgebra of type $(\in, \in \vee q)$ by Lemma 3.6. $\square$

Definition 3.9. An $\mathcal{N}$ -structure (X, f) is called an $\mathcal{N}$ -ideal of type $(\in, \in)$ (resp., type $(\in, \in \vee q)$) if the following assertions are valid.

- (1) If a point $\mathcal{N}$ -structure (X, x_α) is an $\mathcal{N}_\epsilon$ -subset of (X, f) , then $(X, 0_\alpha)$ is an $\mathcal{N}_\epsilon$ -subset (resp., $\mathcal{N}_{\in \vee q}$ -subset) of (X, f) .
- (2) If two point $\mathcal{N}$ -structures $(X, (x * y)_\alpha)$ and (X, y_β) are $\mathcal{N}_\epsilon$ -subsets of (X, f) then the point $\mathcal{N}$ -structure $(X, x_{\alpha \vee \beta})$ is an $\mathcal{N}_\epsilon$ -subset (resp., $\mathcal{N}_{\in \vee q}$ -subset) of (X, f) .

Lemma 3.10. Let (X, f) be an $\mathcal{N}$ -structure. Then $f(0) \leq f(x)$ for all $x \in X$ if and only if $(X, 0_\alpha)$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) whenever (X, x_α) is an $\mathcal{N}_\epsilon$ -subset of (X, f) for all $x \in X$ and $\alpha \in \Delta = [-1, 0)$.

Proof. Assume that $f(0) \leq f(x)$ for all $x \in X$ and let $\alpha \in \Delta$ be such that (X, x_α) is an $\mathcal{N}_\epsilon$ -subset of (X, f) . Then $f(0) \leq f(x) \leq \alpha$, and so $(X, 0_\alpha)$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) .

Conversely, suppose that $(X, 0_\alpha)$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) when (X, x_α) is an $\mathcal{N}_\epsilon$ -subset of (X, f) for all $x \in X$ and $\alpha \in \Delta = [-1, 0)$. If we take $\beta = f(x)$ for any $x \in X$, then (X, x_β) is an $\mathcal{N}_\epsilon$ -subset of (X, f) , and thus $(X, 0_\beta)$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) . Hence $f(0) \leq \beta = f(x)$ for all $x \in X$. $\square$

Lemma 3.11. Given an $\mathcal{N}$ -structure (X, f) , the following are equivalent.

- (1) $f(x) \leq \bigvee \{f(x * y), f(y)\}$ for all $x, y \in X$.
- (2) For any $x, y \in X$ and $\alpha, \beta \in \Delta = [-1, 0)$, if $(X, (x * y)_\alpha)$ and (X, y_β) are $\mathcal{N}_\epsilon$ -subsets of (X, f) , then $(X, x_{\alpha \vee \beta})$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) .

Proof. Assume that $f(x) \leq \bigvee \{f(x * y), f(y)\}$ for all $x, y \in X$. Let $x, y \in X$ and $\alpha, \beta \in \Delta$ be such that $(X, (x * y)_\alpha)$ and (X, y_β) are $\mathcal{N}_\in$ -subsets of (X, f) . Then $f(x * y) \leq \alpha$ and $f(y) \leq \beta$, which imply that

$$f(x) \leq \bigvee \{f(x * y), f(y)\} \leq \alpha \vee \beta.$$

Hence $(X, x_{\alpha \vee \beta})$ is an $\mathcal{N}_\in$ -subset of (X, f) .

Conversely, suppose (2) is valid. If we take $\alpha = f(x * y)$ and $\beta = f(y)$, then $(X, (x * y)_\alpha)$ and (X, y_β) are $\mathcal{N}_\in$ -subsets of (X, f) . It follows that $(X, x_{\alpha \vee \beta})$ is an $\mathcal{N}_\in$ -subset of (X, f) and so that

$$f(x) \leq \alpha \vee \beta = \bigvee \{f(x * y), f(y)\}.$$

This completes the proof. $\square$

Combining Lemmas 3.10 and 3.11, we have the following theorem.

Theorem 3.12. *An $\mathcal{N}$ -structure (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in)$ if and only if the following inequality is valid.*

$$(\forall x, y \in X) \left(f(0) \leq f(x) \leq \bigvee \{f(x * y), f(y)\} \right). \quad (3.6)$$

Definition 3.13. A soft $\mathcal{N}$ -set $(\mathcal{A}, \Delta)$ over X is called a *soft $\mathcal{N}$ -ideal* over X if it satisfies:

$$(\forall \alpha \in \Delta) (\mathcal{A}(\alpha) \neq \emptyset \Rightarrow \mathcal{A}(\alpha) \text{ is an ideal of } X). \quad (3.7)$$

Example 3.14. Let $X = \{0, a, b, c, d\}$ be a *BCK*-algebra with the binary operation $*$ in Table 1.

Given $\Delta = [-1, 0]$, let $(\mathcal{A}, \Delta)$ be a soft $\mathcal{N}$ -set over X in which $\mathcal{A}$ is given as follows:

$$\mathcal{A} : \Delta \rightarrow \mathcal{P}(X), \alpha \mapsto \begin{cases} \{0\} & \text{if } \alpha = -1, \\ \{0, 3\} & \text{if } \alpha \in (-1, -0.8), \\ \{0, 4\} & \text{if } \alpha \in (-1, -0.8], \\ \{0, 2, 3\} & \text{if } \alpha \in (-0.8, -0.6], \\ \{0, 3, 4\} & \text{if } \alpha \in (-0.6, -0.4], \\ \{0, 1, 2, 3\} & \text{if } \alpha \in (-0.4, -0.2], \\ \{0, 1, 3, 4\} & \text{if } \alpha \in (-0.2, 0] \end{cases}$$

It is routine to verify that $(\mathcal{A}, \Delta)$ is a soft $\mathcal{N}$ -ideal over X .

Table 1: Tabular representation of the binary operation $*$

$*$	0	a	b	c	d
0	0	0	0	0	0
a	a	0	a	a	a
b	b	b	0	b	b
c	c	c	c	0	c
d	d	d	d	d	0

Theorem 3.15. *Given an $\mathcal{N}$ -structure (X, f) and the soft $\mathcal{N}_\infty$ -set $(\mathcal{A}_\infty, \Delta)$, the following are equivalent:*

- (1) $(\mathcal{A}_\infty, \Delta)$ is a soft $\mathcal{N}$ -ideal over X for $\Delta = [-1, 0)$.
- (2) (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in)$.

Proof. Assume that $(\mathcal{A}_\infty, \Delta)$ is a soft $\mathcal{N}$ -ideal over X for $\Delta = [-1, 0)$. If there exists $a \in X$ such that $f(0) > f(a)$, then we can take $\alpha \in \Delta$ such that $f(0) > \alpha \geq f(a)$. Thus $(X, 0_\alpha)$ is not an $\mathcal{N}_\infty$ -subset of (X, f) , and so $0 \notin \mathcal{A}_\infty(\alpha)$. This is a contradiction, and so $f(0) \leq f(x)$ for all $x \in X$. Suppose that there exist $a, b \in X$ such that $f(a) > \bigvee \{f(a*b), f(b)\}$. Taking $\beta = \bigvee \{f(a*b), f(b)\}$ implies that $(X, (a*b)_\beta)$ and (X, b_β) are $\mathcal{N}_\infty$ -subsets of (X, f) , that is, $a*b \in \mathcal{A}_\infty(\beta)$ and $b \in \mathcal{A}_\infty(\beta)$. Since $\mathcal{A}_\infty(\beta)$ is an ideal of X , it follows that $a \in \mathcal{A}_\infty(\beta)$. Hence (X, a_β) is an $\mathcal{N}_\infty$ -subset of (X, f) , and so $f(a) \leq \beta$. This is a contradiction, and therefore $f(x) \leq \bigvee \{f(x*y), f(y)\}$ for all $x, y \in X$. Hence (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in)$ by Theorem 3.12.

Conversely, assume that (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in)$ and let $\alpha \in \Delta$ be such that $\mathcal{A}_\infty(\alpha) \neq \emptyset$. Then there exists $x \in \mathcal{A}_\infty(\alpha)$, and so (X, x_α) is an $\mathcal{N}_\infty$ -subset of (X, f) . It follows that $f(0) \leq f(x) \leq \alpha$ and so that $(X, 0_\alpha)$ is an $\mathcal{N}_\infty$ -subset of (X, f) , that is, $0 \in \mathcal{A}_\infty(\alpha)$. Let $x*y \in \mathcal{A}_\infty(\alpha)$ and $y \in \mathcal{A}_\infty(\alpha)$. Then $(X, (x*y)_\alpha)$ and (X, y_α) are $\mathcal{N}_\infty$ -subsets of (X, f) . Thus $f(x*y) \leq \alpha$ and $f(y) \leq \alpha$. It follows from (3.6) that $f(x) \leq \bigvee \{f(x*y), f(y)\} \leq \alpha$. Hence $(X, x_\alpha) = (X, x_{\alpha \vee \alpha})$ is an $\mathcal{N}_\infty$ -subset of (X, f) , and so $x \in \mathcal{A}_\infty(\alpha)$. Thus $\mathcal{A}_\infty(\alpha)$ is an ideal of X for all $\alpha \in \Delta$, and therefore $(\mathcal{A}_\infty, \Delta)$ is a soft $\mathcal{N}$ -ideal over X for $\Delta = [-1, 0)$. $\square$

Theorem 3.16. *The soft $\mathcal{N}_q$ -set $(\mathcal{A}_q, \Delta)$ is a soft $\mathcal{N}$ -ideal over X for $\Delta = [-1, 0)$ if and only if the $\mathcal{N}$ -structure (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in)$.*

Proof. Assume that (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in)$ and let $\alpha \in \Delta$ be such that $\mathcal{A}_q(\alpha) \neq \emptyset$. Then there exists $x \in \mathcal{A}_q(\alpha)$, and so (X, x_α) is an $\mathcal{N}_q$ -subset of (X, f) . If $0 \notin \mathcal{A}_q(\alpha)$, then $(X, 0_q)$ is not an $\mathcal{N}_q$ -subset of (X, f) and so $f(0) + \alpha + 1 \geq 0$. It follows from (3.6) that

$$f(x) + \alpha + 1 \geq f(0) + \alpha + 1 \geq 0,$$

and so that (X, x_α) is not an $\mathcal{N}_q$ -subset of (X, f) . This is a contradiction, and thus $0 \in \mathcal{A}_q(\alpha)$. Let $x * y \in \mathcal{A}_q(\alpha)$ and $y \in \mathcal{A}_q(\alpha)$. Then $(X, (x * y)_\alpha)$ and (X, y_α) are $\mathcal{N}_q$ -subsets of (X, f) . If (X, x_α) is not an $\mathcal{N}_q$ -subset of (X, f) , then $f(x) + \alpha + 1 \geq 0$. It follows from (3.6) that

$$\bigvee \{f(x * y), f(y)\} + \alpha + 1 \geq f(x) + \alpha + 1 \geq 0.$$

Hence $f(x * y) + \alpha + 1 \geq 0$ or $f(y) + \alpha + 1 \geq 0$, that is, $(X, (x * y)_\alpha)$ is not an $\mathcal{N}_q$ -subset of (X, f) or (X, y_α) is not an $\mathcal{N}_q$ -subset of (X, f) . This is a contradiction, and thus $x \in \mathcal{A}_q(\alpha)$. Therefore $(\mathcal{A}_q, \Delta)$ is a soft $\mathcal{N}$ -ideal over X for $\Delta = [-1, 0)$.

Conversely, suppose that the soft $\mathcal{N}_q$ -set $(\mathcal{A}_q, \Delta)$ is a soft $\mathcal{N}$ -ideal over X for $\Delta = [-1, 0)$. If $f(0) > f(a)$ for some $a \in X$, then there exists $\beta \in \Delta$ such that $f(0) + \beta + 1 \geq 0$ and $f(a) + \beta + 1 < 0$. Thus (X, a_β) is an $\mathcal{N}_q$ -subset of (X, f) , and so $(X, 0_\beta)$ is an $\mathcal{N}_q$ -subset of (X, f) . This is a contradiction, and therefore $f(0) \leq f(x)$ for all $x \in X$. Suppose that there exist $a, b \in X$ such that $f(a) > \bigvee \{f(a * b), f(b)\}$. Then $f(a) + \beta + 1 \geq 0$ and $\bigvee \{f(a * b), f(b)\} + \beta + 1 < 0$ for some $\beta \in \Delta$. Thus $f(a * b) + \beta + 1 < 0$ and $f(b) + \beta + 1 < 0$, that is, $(X, (a * b)_\beta)$ and (X, b_β) are $\mathcal{N}_q$ -subsets of (X, f) . Hence $a * b \in \mathcal{A}_q(\beta)$ and $b \in \mathcal{A}_q(\beta)$. Since $\mathcal{A}_q(\beta)$ is an ideal of X , we have $a \in \mathcal{A}_q(\beta)$, that is, (X, a_β) is an $\mathcal{N}_q$ -subset of (X, f) . This is a contradiction, and hence $f(x) \leq \bigvee \{f(x * y), f(y)\}$ for all $x, y \in X$. Using Theorem 3.12, (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in)$. $\square$

Theorem 3.17. *Given an $\mathcal{N}$ -structure (X, f) and the soft $\mathcal{N}_\in$ -set $(\mathcal{A}_\in, \Delta)$, the following are equivalent:*

- (1) (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in \vee q)$.
- (2) $(\mathcal{A}_\in, \Delta)$ is a soft $\mathcal{N}$ -ideal over X for $\Delta = [-0.5, 0)$.

Proof. Assume that (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in \vee q)$. We first show that

$$(\forall x \in X) \left(f(0) \leq \bigvee \{f(x), -0.5\} \right). \quad (3.8)$$

Suppose that $f(0) > f(x) > -0.5$. Then $f(0) > \alpha \geq f(x)$ for some $\alpha \in (-0.5, 0)$, which implies that (X, x_α) is an $\mathcal{N}_\epsilon$ -subset of (X, f) , but $(X, 0_\alpha)$ is not an $\mathcal{N}_\epsilon$ -subset of (X, f) . Also, $(X, 0_\alpha)$ is not an $\mathcal{N}_q$ -subset of (X, f) since $f(0) + \alpha + 1 \geq 0$. Thus $(X, 0_\alpha)$ is not an $\mathcal{N}_{\epsilon \vee q}$ -subset of (X, f) , a contradiction. Hence $f(0) \leq f(x)$ for all $x \in X$. Now if $f(x) \leq -0.5$, then $(X, x_{-0.5})$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) and so $(X, 0_{-0.5})$ is an $\mathcal{N}_{\epsilon \vee q}$ -subset of (X, f) . It follows that $f(0) \leq -0.5$ or $f(0) - 0.5 + 1 < 0$ and so that $f(0) \leq -0.5$ because if not, then $f(0) - 0.5 + 1 > 0$, a contradiction. Consequently, the condition (3.8) is valid. Let $\alpha \in \Delta = [-0.5, 0)$. The condition (3.8) implies that $f(0) \leq \bigvee \{f(x), -0.5\}$ for all $x \in \mathcal{A}_\epsilon(\alpha)$, and so

$$f(0) \leq \bigvee \{f(x), -0.5\} \leq \bigvee \{\alpha, -0.5\} = \alpha,$$

that is, $(X, 0_\alpha)$ is an $\mathcal{N}_\epsilon$ -subset of (X, f) . Hence $0 \in \mathcal{A}_\epsilon(\alpha)$. Now we show that

$$(\forall x, y \in X) \left(f(x) \leq \bigvee \{f(x * y), f(y), -0.5\} \right). \quad (3.9)$$

If $\bigvee \{f(x * y), f(y)\} > -0.5$, then $f(x) \leq \bigvee \{f(x * y), f(y)\}$. Otherwise, there exists $\beta \in (-0.5, 0)$ such that $f(x) > \beta \geq \bigvee \{f(x * y), f(y)\}$. It follows that $(X, (x * y)_\beta)$ and (X, y_β) are $\mathcal{N}_\epsilon$ -subsets of (X, f) , but (X, x_β) is not an $\mathcal{N}_\epsilon$ -subset of (X, f) . Also (X, x_β) is not an $\mathcal{N}_q$ -subset of (X, f) because $f(x) + \beta + 1 > 2\beta + 1 > 0$. Hence (X, x_β) is not an $\mathcal{N}_{\epsilon \vee q}$ -subset of (X, f) , a contradiction. If $\bigvee \{f(x * y), f(y)\} \leq -0.5$, then $(X, (x * y)_{-0.5})$ and $(X, y_{-0.5})$ are $\mathcal{N}_\epsilon$ -subsets of (X, f) . Thus $(X, x_{-0.5})$ is an $\mathcal{N}_{\epsilon \vee q}$ -subset of (X, f) , and so $f(x) \leq -0.5$ or $f(x) - 0.5 + 1 < 0$. It follows that $f(x) \leq -0.5$ because if $f(x) > -0.5$, then $f(x) - 0.5 + 1 > 0$ which is a contradiction. Therefore $f(x) \leq \bigvee \{f(x * y), f(y), -0.5\}$ for all $x, y \in X$. Let $x, y \in X$ be such that $x * y \in \mathcal{A}_\epsilon(\alpha)$ and $y \in \mathcal{A}_\epsilon(\alpha)$. Then $(X, (x * y)_\alpha)$ and (X, y_α) are $\mathcal{N}_\epsilon$ -subsets of (X, f) , and so $f(x * y) \leq \alpha$ and $f(y) \leq \alpha$. Using (3.9), we have

$$f(x) \leq \bigvee \{f(x * y), f(y), -0.5\} \leq \bigvee \{\alpha, -0.5\} = \alpha.$$

Thus (X, x_α) is an $\mathcal{N}_\epsilon$ -subset of (X, f) , and hence $x \in \mathcal{A}_\epsilon(\alpha)$. Therefore $(\mathcal{A}_\epsilon, \Delta)$ is a soft $\mathcal{N}$ -ideal over X for $\Delta = [-0.5, 0)$.

Conversely, suppose that (2) is valid. If $f(0) > \bigvee \{f(a), -0.5\}$ for some $a \in X$, then there exists $\alpha \in \Delta$ such that $f(0) > \alpha \geq \bigvee \{f(a), -0.5\}$. Then $\alpha \in \Delta$ and (X, a_α) is an $\mathcal{N}_\epsilon$ -subset of (X, f) . But $(X, 0_\alpha)$ is not an $\mathcal{N}_\epsilon$ -subset of (X, f) , i.e., $0 \notin \mathcal{A}_\epsilon(\alpha)$. This is a contradiction, and so $f(0) \leq \bigvee \{f(x), -0.5\}$ for all $x \in X$. Let $x \in X$ and $\alpha \in \Delta$ be such that (X, x_α) is an $\mathcal{N}_\epsilon$ -subset of (X, f) . Then $f(x) \leq \alpha$. Suppose that $(X, 0_\alpha)$ is not an $\mathcal{N}_\epsilon$ -subset of (X, f) . Then $f(0) > \alpha$. If $f(x) > -0.5$, then $f(0) \leq \bigvee \{f(x), -0.5\} = f(x) \leq \alpha$

which is impossible. Thus $f(x) \leq -0.5$ and so

$$f(0) + \alpha + 1 < 2f(0) + 1 \leq 2 \bigvee \{f(x), -0.5\} + 1 = 0,$$

that is, $(X, 0_\alpha)$ is an $\mathcal{N}_q$ -subset of (X, f) . Hence $(X, 0_\alpha)$ is an $\mathcal{N}_{\in \vee q}$ -subset of (X, f) . Assume that there exist $a, b \in X$ such that

$$f(a) > \bigvee \{f(a * b), f(b), -0.5\}.$$

Taking $\alpha = \bigvee \{f(a * b), f(b), -0.5\}$ implies that $\alpha \in \Delta$, and $(X, (a * b)_\alpha)$ and (X, b_α) are $\mathcal{N}_\in$ -subsets of (X, f) . But (X, a_α) is not an $\mathcal{N}_\in$ -subset of (X, f) . This is a contradiction, and thus $f(x) \leq \bigvee \{f(x * y), f(y), -0.5\}$ for all $x, y \in X$. Let $x, y \in X$ and $\alpha, \beta \in \Delta$ be such that $(X, (x * y)_\alpha)$ and (X, y_β) are $\mathcal{N}_\in$ -subsets of (X, f) , and suppose that $(X, x_{\alpha \vee \beta})$ is not an $\mathcal{N}_\in$ -subset of (X, f) . Then $f(x * y) \leq \alpha$, $f(y) \leq \beta$ and $f(x) > \alpha \vee \beta$. If $\bigvee \{f(x * y), f(y)\} > -0.5$, then

$$f(x) \leq \bigvee \{f(x * y), f(y), -0.5\} = \bigvee \{f(x * y), f(y)\} \leq \alpha \vee \beta.$$

which is a contradiction. Thus $\bigvee \{f(x * y), f(y)\} \leq -0.5$, and so

$$f(x) + (\alpha \vee \beta) + 1 < 2f(x) + 1 \leq 2 \bigvee \{f(x * y), f(y), -0.5\} + 1 = 0,$$

which shows that $(X, x_{\alpha \vee \beta})$ is an $\mathcal{N}_q$ -subset of (X, f) . Thus $(X, x_{\alpha \vee \beta})$ is an $\mathcal{N}_{\in \vee q}$ -subset of (X, f) . Consequently, (X, f) is an $\mathcal{N}$ -ideal of type $(\in, \in \vee q)$. $\square$

Conclusion

We have introduced $\mathcal{N}$ -ideal of types $(\in, \in)$ and $(\in, \in \vee q)$, soft $\mathcal{N}_\in$ -set, soft $\mathcal{N}_q$ -set, soft $\mathcal{N}_{\in \vee q}$ -set, soft $\mathcal{N}$ -subalgebra and soft $\mathcal{N}$ -ideal in BCK/BCI -algebra.

We have considered characterizations of $\mathcal{N}$ -subalgebra of types $(\in, \in)$ and $(\in, \in \vee q)$, $\mathcal{N}$ -ideal of types $(\in, \in)$ and $(\in, \in \vee q)$, soft $\mathcal{N}$ -subalgebra and soft $\mathcal{N}$ -ideal.

4 Compliance with Ethical Standards

Conflict of Interest: Author declares that he has no conflict of interest.

Ethical approval: This article does not contain any studies with human participants or animals performed by the author.

Informed consent: Informed consent was obtained from all individual participants included in the study.

Author contributions: Conceptualization: Young Bae Jun, Hashem Bordbar; Methodology: Young Bae Jun, Hashem Bordbar; Formal analysis and investigation: Hashem Bordbar, Rajab Ali Borzooei, Borumand Saeid; Writing - original draft preparation: Hashem Bordbar; Writing - review and editing: Young Bae Jun; Resources: Hashem Bordbar; Supervision: Young Bae Jun.

References

- [1] H. Bordbar, R. A. Borzooei, Y. B. Jun *Uni-Soft Commutative Ideals and Closed Uni-Soft Ideals in BCI-Algebras* New Mathematics and Natural Computation, Volume 14(2) (2018), Pages 235-247.
- [2] H. Bordbar, H. Harizavi, Y. B. Jun, *Uni-Soft Ideals in Coresiduated Lattices* Sigma J Eng and Nat Sci 9 (1), 2018, 69-75
- [3] H. Bordbar, M.M. Zahedi, Y. B. Jun, *Ideals of IS-algebras based on $\mathcal{N}$ -Structures* Kragujevac Journal of Mathematics, Volume 42(4) (2018), Pages 631–641.
- [4] Y. S. Huang, *BCI-algebra*, Science Press, Beijing, 2006.
- [5] Y. B. Jun, M. S. Kang and C. H. Park, *$\mathcal{N}$ -subalgebras in BCK/BCI-algebras based on point $\mathcal{N}$ -structures*, Int. J. Math. Math. Sci. Volume 2010, Article ID 303412, 9 pages.
- [6] Y. B. Jun, K. J. Lee and S. Z. Song, *$\mathcal{N}$ -ideals of BCK/BCI-algebras*, J. Chungcheong Math. Soc. **22** (2009), 417–437.
- [7] Y. B. Jun, M. A. Öztürk and E. H. Roh, *$\mathcal{N}$ -structures applied to closed ideals in BCH-algebras*, Int. J. Math. Math. Sci. Volume 2010, Article ID 943565, 9 pages.
- [8] Y. B. Jun, F. Smarandache, H. Bordbar *Neutrosophic $\mathcal{N}$ -Structures Applied to BCK/BCI-algebras*, Information 2017, 8(4), 128
- [9] Y. B. Jun, S. Z. Song, H. Bordbar *Int-Soft Ideals of Pseudo MV-algebra*, Bulletin of the Section of Logic Volume 47/1 (2018), pp. 1–14
- [10] J. Meng and Y. B. Jun, *BCK-algebras*, Kyungmoon Sa Co., Seoul, 1994.